Where Have You Gone, Joe DiMaggio? Probability and Hitting Streaks

Phil Yates

Department of Mathematical Sciences DePaul University

February 16, 2018



Player	BA	HR	RBI	AB	Н	OBP	SLG	WAR
Player A	.357	30	125	541	193	.440	.643	9.1
Player B	.406	37	120	456	185	.553	.735	10.6

Player			l I					
Player A	.357	30	125	541	193	.440	.643	9.1
Player B	.406	37	120	456	185	.553	.735	10.6

Who are these players?

Player			l I					
Player A	.357	30	125	541	193	.440	.643	9.1
Player B	.406	37	120	456	185	.553	.735	10.6

Who are these players?



Player	BA	HR	RBI	AB	Н	OBP	SLG	WAR
Player A								
Player B	.406	37	120	456	185	.553	.735	10.6

Who are these players?



Joe DiMaggio (Player A – won the AL MVP) and Ted Williams (Player B)

Longest Hitting Streaks in MLB History

Year	Name	Team	Games
1941	Joe DiMaggio	New York Yankees	56
1896-97	Willie Keeler	Baltimore Orioles	45
1978	Pete Rose	Cincinnati Reds	44
1894	Bill Dahlen	Chicago Colts (Cubs)	42
1922	George Sisler	St. Louis Browns	41
1911	Ty Cobb	Detroit Tigers	40
1987	Paul Molitor	Milwaukee Brewers	39
2005-06	Jimmy Rollins	Philadelphia Phillies	38
1945	Tommy Holmes	Boston Braves	37
1896-97	Gene DeMontreville	Washington Senators	36

Source: MLB.com

What is a hitting streak?

• the number of consecutive official games in which a player appears and gets at least one base hit

- the number of consecutive official games in which a player appears and gets at least one base hit
- the streak is ended when a player has at least 1 plate appearance and no hits

- the number of consecutive official games in which a player appears and gets at least one base hit
- the streak is ended when a player has at least 1 plate appearance and no hits
- the streak is not terminated if all official plate appearances result in a base on balls (a walk), hit by pitch, defensive interference or a sacrifice bunt

- the number of consecutive official games in which a player appears and gets at least one base hit
- the streak is ended when a player has at least 1 plate appearance and no hits
- the streak is not terminated if all official plate appearances result in a base on balls (a walk), hit by pitch, defensive interference or a sacrifice bunt
- the streak shall terminate if the player has a sacrifice fly and no hit Source: Official Rules, Major League Baseball, 10.23 Guidelines For Cumulative Performance Records

Basic concepts of probability needed to analyze hitting streaks:

Sample Space

All possible outcomes in an experiment. Denoted by S

Basic concepts of probability needed to analyze hitting streaks:

Sample Space

All possible outcomes in an experiment. Denoted by S

Experiment: A batter has four at bats in a single game. Each at bat can be a hit (H) or an out (O).

Basic concepts of probability needed to analyze hitting streaks:

Sample Space

All possible outcomes in an experiment. Denoted by S

Experiment: A batter has four at bats in a single game. Each at bat can be a hit (H) or an out (O).

Sample Space 5: 16 total outcomes

Basic concepts of probability needed to analyze hitting streaks:

Sample Space

All possible outcomes in an experiment. Denoted by S

Experiment: A batter has four at bats in a single game. Each at bat can be a hit (H) or an out (O).

Sample Space 5: 16 total outcomes

"No hits in four at bats" – 0000

Basic concepts of probability needed to analyze hitting streaks:

Sample Space

All possible outcomes in an experiment. Denoted by S

Experiment: A batter has four at bats in a single game. Each at bat can be a hit (H) or an out (O).

Sample Space 5: 16 total outcomes

- "No hits in four at bats" 0000
- "One hit in four at bats" HOOO, OHOO, OOHO, OOOH

Basic concepts of probability needed to analyze hitting streaks:

Sample Space

All possible outcomes in an experiment. Denoted by S

Experiment: A batter has four at bats in a single game. Each at bat can be a hit (H) or an out (O).

Sample Space 5: 16 total outcomes

- "No hits in four at bats" OOOO
- "One hit in four at bats" HOOO, OHOO, OOHO, OOOH
- "Two hits in four at bats" HHOO, HOHO, HOOH, OHHO, OHOH, OOHH

Basic concepts of probability needed to analyze hitting streaks:

Sample Space

All possible outcomes in an experiment. Denoted by S

Experiment: A batter has four at bats in a single game. Each at bat can be a hit (H) or an out (O).

Sample Space 5: 16 total outcomes

- "No hits in four at bats" OOOO
- "One hit in four at bats" HOOO, OHOO, OOHO, OOOH
- "Two hits in four at bats" HHOO, HOHO, HOOH, OHHO, OHOH, OOHH
- "Three hits in four at bats" HHHO, HHOH, HOHH, OHHH

Yates (DePaul) Hitting Streaks 16 February 2018 5 / 21

Basic concepts of probability needed to analyze hitting streaks:

Sample Space

All possible outcomes in an experiment. Denoted by S

Experiment: A batter has four at bats in a single game. Each at bat can be a hit (H) or an out (O).

Sample Space 5: 16 total outcomes

- "No hits in four at bats" OOOO
- "One hit in four at bats" HOOO, OHOO, OOHO, OOOH
- "Two hits in four at bats" HHOO, HOHO, HOOH, OHHO, OHOH, OOHH
- "Three hits in four at bats" HHHO, HHOH, HOHH, OHHH
- "Four hits in four at bats" HHHH

Basic concepts of probability needed to analyze hitting streaks:

Complement of an Event A

The set of all outcomes that are in the sample space S but are not in the event A. Typically denoted as \overline{A} or A^c .

Basic concepts of probability needed to analyze hitting streaks:

Complement of an Event A

The set of all outcomes that are in the sample space S but are not in the event A. Typically denoted as \overline{A} or A^c .

Basic concepts of probability needed to analyze hitting streaks:

Complement of an Event A

The set of all outcomes that are in the sample space S but are not in the event A. Typically denoted as \overline{A} or A^c .

Event A: A player gets at least one hit in four at bats in a single game.

• "One hit in four at bats" - HOOO, OHOO, OOHO, OOOH

Yates (DePaul) Hitting Streaks 16 February 2018 6 / 21

Basic concepts of probability needed to analyze hitting streaks:

Complement of an Event A

The set of all outcomes that are in the sample space S but are not in the event A. Typically denoted as \overline{A} or A^c .

- "One hit in four at bats" HOOO, OHOO, OOHO, OOOH
- "Two hits in four at bats" HHOO, HOHO, HOOH, OHHO, OHOH, OOHH

Basic concepts of probability needed to analyze hitting streaks:

Complement of an Event A

The set of all outcomes that are in the sample space S but are not in the event A. Typically denoted as \overline{A} or A^c .

- "One hit in four at bats" HOOO, OHOO, OOHO, OOOH
- "Two hits in four at bats" HHOO, HOHO, HOOH, OHHO, OHOH, OOHH
- "Three hits in four at bats" HHHO, HHOH, HOHH, OHHH

Basic concepts of probability needed to analyze hitting streaks:

Complement of an Event A

The set of all outcomes that are in the sample space S but are not in the event A. Typically denoted as \overline{A} or A^c .

- "One hit in four at bats" HOOO, OHOO, OOHO, OOOH
- "Two hits in four at bats" HHOO, HOHO, HOOH, OHHO, OHOH, OOHH
- "Three hits in four at bats" HHHO, HHOH, HOHH, OHHH
- "Four hits in four at bats" HHHH

Basic concepts of probability needed to analyze hitting streaks:

Complement of an Event A

The set of all outcomes that are in the sample space S but are not in the event A. Typically denoted as \overline{A} or A^c .

- "One hit in four at bats" HOOO, OHOO, OOHO, OOOH
- "Two hits in four at bats" HHOO, HOHO, HOOH, OHHO, OHOH, OOHH
- "Three hits in four at bats" HHHO, HHOH, HOHH, OHHH
- "Four hits in four at bats" HHHH

Basic concepts of probability needed to analyze hitting streaks:

Complement of an Event A

The set of all outcomes that are in the sample space S but are not in the event A. Typically denoted as \overline{A} or A^c .

Event A: A player gets at least one hit in four at bats in a single game.

- "One hit in four at bats" HOOO, OHOO, OOHO, OOOH
- "Two hits in four at bats" HHOO, HOHO, HOOH, OHHO, OHOH, OOHH
- "Three hits in four at bats" HHHO, HHOH, HOHH, OHHH
- "Four hits in four at bats" HHHH

Complement Ac: A player gets zero hits in four at bats in a single game

"No hits in four at bats" – OOOO

Basic concepts of probability needed to analyze hitting streaks:

Rules of Probability

Complement Rule:

$$P(A^c) = 1 - P(A)$$

Multiplication Rule for Independent Events:

Let A and B be two independent events. Then $P(A \text{ and } B) = P(A) \times P(B)$.

Basic concepts of probability needed to analyze hitting streaks:

Rules of Probability

Complement Rule:

$$P(A^c) = 1 - P(A)$$

Multiplication Rule for Independent Events:

Let A and B be two independent events. Then $P(A \text{ and } B) = P(A) \times P(B)$.

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in a single game?

Basic concepts of probability needed to analyze hitting streaks:

Rules of Probability

Complement Rule:

$$P(A^c) = 1 - P(A)$$

Multiplication Rule for Independent Events:

Let A and B be two independent events. Then $P(A \text{ and } B) = P(A) \times P(B)$.

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in a single game?

P(at least one H in 4 at bats) = 1 - P(zero H's in 4 at bats)

Basic concepts of probability needed to analyze hitting streaks:

Rules of Probability

Complement Rule:

$$P(A^c) = 1 - P(A)$$

Multiplication Rule for Independent Events:

Let A and B be two independent events. Then $P(A \text{ and } B) = P(A) \times P(B)$.

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in a single game?

P(at least one H in 4 at bats) = 1 - P(zero H's in 4 at bats)= 1 - P(OOOO)

Basic concepts of probability needed to analyze hitting streaks:

Rules of Probability

Complement Rule:

$$P(A^c) = 1 - P(A)$$

Multiplication Rule for Independent Events:

Let A and B be two independent events. Then $P(A \text{ and } B) = P(A) \times P(B)$.

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in a single game?

```
P(\text{at least one H in 4 at bats}) = 1 - P(\text{zero H's in 4 at bats})
= 1 - P(\text{OOOO})
= 1 - (0.700 \times 0.700 \times 0.700 \times 0.700)
```

Basic concepts of probability needed to analyze hitting streaks:

Rules of Probability

Complement Rule:

$$P(A^c) = 1 - P(A)$$

Multiplication Rule for Independent Events:

Let A and B be two independent events. Then $P(A \text{ and } B) = P(A) \times P(B)$.

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in a single game?

```
P(\text{at least one H in 4 at bats}) = 1 - P(\text{zero H's in 4 at bats})
= 1 - P(\text{OOOO})
= 1 - (0.700 \times 0.700 \times 0.700 \times 0.700)
= 1 - 0.700^4
```

Basic concepts of probability needed to analyze hitting streaks:

Rules of Probability

Complement Rule:

$$P(A^c) = 1 - P(A)$$

Multiplication Rule for Independent Events:

Let A and B be two independent events. Then $P(A \text{ and } B) = P(A) \times P(B)$.

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in a single game?

P(at least one H in 4 at bats) = 1 - P(zero H's in 4 at bats) = 1 - P(OOOO) $= 1 - (0.700 \times 0.700 \times 0.700 \times 0.700)$ $= 1 - 0.700^4$ = 0.7599

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

The player needs to get at least one hit in four at bats in the first game AND at least one hit in four at bats in the second game.

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

The player needs to get at least one hit in four at bats in the first game AND at least one hit in four at bats in the second game.

$$0.7599 \times 0.7599 = 0.7599^2 = \mathbf{0.5774}$$

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

The player needs to get at least one hit in four at bats in the first game AND at least one hit in four at bats in the second game.

$$0.7599 \times 0.7599 = 0.7599^2 = \mathbf{0.5774}$$

Five consecutive games?

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

The player needs to get at least one hit in four at bats in the first game AND at least one hit in four at bats in the second game.

$$0.7599 \times 0.7599 = 0.7599^2 = \mathbf{0.5774}$$

Five consecutive games? $0.7599^5 = 0.2534$

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

The player needs to get at least one hit in four at bats in the first game AND at least one hit in four at bats in the second game.

$$0.7599 \times 0.7599 = 0.7599^2 = \mathbf{0.5774}$$

Five consecutive games? $0.7599^5 = 0.2534$

Ten consecutive games?

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

The player needs to get at least one hit in four at bats in the first game AND at least one hit in four at bats in the second game.

$$0.7599 \times 0.7599 = 0.7599^2 = \mathbf{0.5774}$$

Five consecutive games? $0.7599^5 = 0.2534$

Ten consecutive games? $0.7599^{10} = 0.0642$

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

The player needs to get at least one hit in four at bats in the first game AND at least one hit in four at bats in the second game.

$$0.7599 \times 0.7599 = 0.7599^2 = \mathbf{0.5774}$$

Five consecutive games? $0.7599^5 = 0.2534$

Ten consecutive games? $0.7599^{10} = 0.0642$

Twenty consecutive games?

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

The player needs to get at least one hit in four at bats in the first game AND at least one hit in four at bats in the second game.

$$0.7599 \times 0.7599 = 0.7599^2 = \mathbf{0.5774}$$

Five consecutive games? $0.7599^5 = 0.2534$

Ten consecutive games? $0.7599^{10} = 0.0642$

Twenty consecutive games? $0.7599^{10} = 0.004122$

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

The player needs to get at least one hit in four at bats in the first game AND at least one hit in four at bats in the second game.

$$0.7599 \times 0.7599 = 0.7599^2 = \mathbf{0.5774}$$

Five consecutive games? $0.7599^5 = 0.2534$

Ten consecutive games? $0.7599^{10} = 0.0642$

Twenty consecutive games? $0.7599^{10} = 0.004122$

Fifty-six consecutive games?

What is the probability that a player with a 0.300 batting average gets at least one hit in four at bats in two consecutive games?

The player needs to get at least one hit in four at bats in the first game AND at least one hit in four at bats in the second game.

$$0.7599 \times 0.7599 = 0.7599^2 = \mathbf{0.5774}$$

Five consecutive games? $0.7599^5 = 0.2534$

Ten consecutive games? $0.7599^{10} = 0.0642$

Twenty consecutive games? $0.7599^{10} = 0.004122$

Fifty-six consecutive games? $0.7599^{56} = 0.0000002101$

In 1941, Joe DiMaggio had 541 at bats in 139 games.

3.892 at bats per game — let's round to 4 at bats (can't have a fraction of an at bat)

What is the probability that Joe DiMaggio, a 0.357 hitter in 1941, gets at least one hit in four at bats in fifty-six consecutive games?

In 1941, Joe DiMaggio had 541 at bats in 139 games.

3.892 at bats per game — let's round to 4 at bats (can't have a fraction of an at bat)

What is the probability that Joe DiMaggio, a 0.357 hitter in 1941, gets at least one hit in four at bats in fifty-six consecutive games?

$$P(H) = 0.357 \Rightarrow P(O) = 0.643$$

In 1941, Joe DiMaggio had 541 at bats in 139 games.

3.892 at bats per game — let's round to 4 at bats (can't have a fraction of an at bat)

What is the probability that Joe DiMaggio, a 0.357 hitter in 1941, gets at least one hit in four at bats in fifty-six consecutive games?

$$P(H) = 0.357 \Rightarrow P(O) = 0.643$$

Probability of at least one H in 4 at bats in one game?

In 1941, Joe DiMaggio had 541 at bats in 139 games.

3.892 at bats per game — let's round to 4 at bats (can't have a fraction of an at bat)

What is the probability that Joe DiMaggio, a 0.357 hitter in 1941, gets at least one hit in four at bats in fifty-six consecutive games?

$$P(H) = 0.357 \Rightarrow P(O) = 0.643$$

Probability of at least one H in 4 at bats in one game? $1 - 0.643^4 = 0.8290599$

Fifty-six consecutive games?

In 1941, Joe DiMaggio had 541 at bats in 139 games.

3.892 at bats per game — let's round to 4 at bats (can't have a fraction of an at bat)

What is the probability that Joe DiMaggio, a 0.357 hitter in 1941, gets at least one hit in four at bats in fifty-six consecutive games?

$$P(H) = 0.357 \Rightarrow P(O) = 0.643$$

Probability of at least one H in 4 at bats in one game? $1 - 0.643^4 = 0.8290599$

Fifty-six consecutive games? $0.8290599^{56} = 0.00002759$

One big assumption that we made in the previous calculation?

One big assumption that we made in the previous calculation?

Joe DiMaggio had 4 at bats in every single game during the streak! One big assumption that we made in the previous calculation?

Joe DiMaggio had 4 at bats in every single game during the streak!

Why is this a problem? Let us assume he had 8 at bats in 2 games.

Game 1 AB's	Game 2 AB's	Prob. of H's in both games
1	7	$(1-0.643^1) \times (1-0.643^7) = 0.3408$
2	6	$(1 - 0.643^2) \times (1 - 0.643^6) = 0.5451$
3	5	$(1-0.643^3) \times (1-0.643^5) = 0.6535$
4	4	$(1-0.643^4) \times (1-0.643^4) = 0.6873$
5	3	$(1-0.643^5) \times (1-0.643^3) = 0.6535$
6	2	$(1 - 0.643^6) \times (1 - 0.643^2) = $ 0.5451
1	7	$(1 - 0.643^7) \times (1 - 0.643^1) = 0.3408$

By assuming the same number of at bats in each game, it **inflates** or potentially **overestimates** the likelihood of a hitting streak

By assuming the same number of at bats in each game, it **inflates** or potentially **overestimates** the likelihood of a hitting streak

Solution?

By assuming the same number of at bats in each game, it **inflates** or potentially **overestimates** the likelihood of a hitting streak

Solution? Vary the at bats for each game.

During the 56 game hitting streak, DiMaggio had:

- 3 games with 2 at bats
- 11 games with 3 at bats
- 26 games with 4 at bats
- 16 games with 5 at bats

Source: Cliff Blau, a member of SABR (Society for American Baseball Research)

$$(1-0.643^2)^3$$

$$\left(1-0.643^2\right)^3 \times \left(1-0.643^3\right)^{11}$$

$$\left(1-0.643^2\right)^3 \, \times \left(1-0.643^3\right)^{11} \, \times \left(1-0.643^4\right)^{26}$$

$$\left(1-0.643^2\right)^3 \, \times \left(1-0.643^3\right)^{11} \, \times \left(1-0.643^4\right)^{26} \, \times \left(1-0.643^5\right)^{16}$$

When varying the at bats based on his **actual** at bats during the hitting streak, we can find the probability of a 56 game hitting streak:

$$\left(1-0.643^2\right)^3 \, \times \left(1-0.643^3\right)^{11} \, \times \left(1-0.643^4\right)^{26} \, \times \left(1-0.643^5\right)^{16}$$

Probability?

When varying the at bats based on his actual at bats during the hitting streak, we can find the probability of a 56 game hitting streak:

$$\left(1-0.643^2\right)^3 \, \times \left(1-0.643^3\right)^{11} \, \times \left(1-0.643^4\right)^{26} \, \times \left(1-0.643^5\right)^{16}$$

Probability? **0.00007993**

Notice this is much smaller than 0.00002759 – the probability of the 56 game hitting streak when assuming 4 at bats per game

Hitting Streaks 16 February 2018 12 / 21 When varying the at bats based on his **actual** at bats during the hitting streak, we can find the probability of a 56 game hitting streak:

$$\left(1-0.643^2\right)^3 \, \times \left(1-0.643^3\right)^{11} \, \times \left(1-0.643^4\right)^{26} \, \times \left(1-0.643^5\right)^{16}$$

Probability? 0.000007993

Notice this is much smaller than 0.00002759 – the probability of the 56 game hitting streak when assuming 4 at bats per game

Problem?

When varying the at bats based on his **actual** at bats during the hitting streak, we can find the probability of a 56 game hitting streak:

$$\left(1-0.643^2\right)^3 \, \times \left(1-0.643^3\right)^{11} \, \times \left(1-0.643^4\right)^{26} \, \times \left(1-0.643^5\right)^{16}$$

Probability? 0.000007993

Notice this is much smaller than 0.00002759 – the probability of the 56 game hitting streak when assuming 4 at bats per game

Problem? This probability is specific to these 56 games and not necessarily any 56 consecutive games over the course of an **entire** baseball season.

Let's say we have a player that has a 0.333 batting average and has 4 at bats each game. The baseball season has 162 games in a season (154 when DiMaggio played). How can we estimate the probability of this player having a hitting streak as long as Joe DiMaggio's 56 game hitting streak?

Let's say we have a player that has a 0.333 batting average and has 4 at bats each game. The baseball season has 162 games in a season (154 when DiMaggio played). How can we estimate the probability of this player having a hitting streak as long as Joe DiMaggio's 56 game hitting streak?

We can use dice (or some other chance mechanism) to "simulate" the season!

Let's say we have a player that has a 0.333 batting average and has 4 at bats each game. The baseball season has 162 games in a season (154 when DiMaggio played). How can we estimate the probability of this player having a hitting streak as long as Joe DiMaggio's 56 game hitting streak?

We can use dice (or some other chance mechanism) to "simulate" the season!

Sample space for a single 6 sided die?

Let's say we have a player that has a 0.333 batting average and has 4 at bats each game. The baseball season has 162 games in a season (154 when DiMaggio played). How can we estimate the probability of this player having a hitting streak as long as Joe DiMaggio's 56 game hitting streak?

We can use dice (or some other chance mechanism) to "simulate" the season!

Sample space for a single 6 sided die? $S = \{1, 2, 3, 4, 5, 6\}$.

Let's say we have a player that has a 0.333 batting average and has 4 at bats each game. The baseball season has 162 games in a season (154 when DiMaggio played). How can we estimate the probability of this player having a hitting streak as long as Joe DiMaggio's 56 game hitting streak?

We can use dice (or some other chance mechanism) to "simulate" the season!

Sample space for a single 6 sided die? $S = \{1, 2, 3, 4, 5, 6\}$.

The purple numbers are "hits" and the gold numbers are "outs."

Let's say we have a player that has a 0.333 batting average and has 4 at bats each game. The baseball season has 162 games in a season (154 when DiMaggio played). How can we estimate the probability of this player having a hitting streak as long as Joe DiMaggio's 56 game hitting streak?

We can use dice (or some other chance mechanism) to "simulate" the season!

Sample space for a single 6 sided die? $S = \{1, 2, 3, 4, 5, 6\}$.

The purple numbers are "hits" and the gold numbers are "outs."

Each person should roll the die four times. That would be a "game." If a 1 or 2 occurs in **any** of the four rolls of a die, the player had a hit in that game.

Let's say we have a player that has a 0.333 batting average and has 4 at bats each game. The baseball season has 162 games in a season (154 when DiMaggio played). How can we estimate the probability of this player having a hitting streak as long as Joe DiMaggio's 56 game hitting streak?

We can use dice (or some other chance mechanism) to "simulate" the season!

Sample space for a single 6 sided die? $S = \{1, 2, 3, 4, 5, 6\}$.

The purple numbers are "hits" and the gold numbers are "outs."

Each person should roll the die four times. That would be a "game." If a 1 or 2 occurs in **any** of the four rolls of a die, the player had a hit in that game.

Problem?

Let's say we have a player that has a 0.333 batting average and has 4 at bats each game. The baseball season has 162 games in a season (154 when DiMaggio played). How can we estimate the probability of this player having a hitting streak as long as Joe DiMaggio's 56 game hitting streak?

We can use dice (or some other chance mechanism) to "simulate" the season!

Sample space for a single 6 sided die? $S = \{1, 2, 3, 4, 5, 6\}$.

The purple numbers are "hits" and the gold numbers are "outs."

Each person should roll the die four times. That would be a "game." If a 1 or 2 occurs in **any** of the four rolls of a die, the player had a hit in that game.

Problem? This is just one simulated "season." To estimate the probability we would need to repeat this process thousands of times!

Assuming consecutive at bats are independent of one another and the batting average ("probability of getting a hit in an at bat") is the same for every at bat, the number of hits in a game follows a **Binomial Probability Distribution** with:

Assuming consecutive at bats are independent of one another and the batting average ("probability of getting a hit in an at bat") is the same for every at bat, the number of hits in a game follows a **Binomial Probability Distribution** with:

 \bullet *n* – number of trials – the number of at bats in a given game

Assuming consecutive at bats are independent of one another and the batting average ("probability of getting a hit in an at bat") is the same for every at bat, the number of hits in a game follows a **Binomial Probability Distribution** with:

- n − number of trials − the number of at bats in a given game

Assuming consecutive at bats are independent of one another and the batting average ("probability of getting a hit in an at bat") is the same for every at bat, the number of hits in a game follows a **Binomial Probability Distribution** with:

- \bullet *n* number of trials the number of at bats in a given game
- \bullet p probability of success the probability of getting a hit in an at bat

For each player in a given season, n will vary from game to game while p will be treated as constant.

Assuming consecutive at bats are independent of one another and the batting average ("probability of getting a hit in an at bat") is the same for every at bat, the number of hits in a game follows a **Binomial Probability Distribution** with:

- n number of trials the number of at bats in a given game
- \bullet p probability of success the probability of getting a hit in an at bat

For each player in a given season, n will vary from game to game while p will be treated as constant.

How do we do this?

Assuming consecutive at bats are independent of one another and the batting average ("probability of getting a hit in an at bat") is the same for every at bat, the number of hits in a game follows a **Binomial Probability Distribution** with:

- n number of trials the number of at bats in a given game
- \bullet p probability of success the probability of getting a hit in an at bat

For each player in a given season, n will vary from game to game while p will be treated as constant.

How do we do this? Using a computer program (statisticians love R!)

First read in the hits and at bats for DiMaggio's 1941 season into R.

First read in the hits and at bats for DiMaggio's 1941 season into R. Then run the streaks program. What's the program do?

• Runs *k* simulated "baseball seasons." Here the season is DiMaggio's 1941 season. We will set the number of simulates seasons to be 10,000.

- Runs *k* simulated "baseball seasons." Here the season is DiMaggio's 1941 season. We will set the number of simulates seasons to be 10,000.
- Let $n_1, n_2, \ldots, n_{139}$ be the **actual** at bats during DiMaggio's 1941 season. A simulated baseball season will sample **with replacement** these at bats 139 times.

- Runs *k* simulated "baseball seasons." Here the season is DiMaggio's 1941 season. We will set the number of simulates seasons to be 10,000.
- Let $n_1, n_2, \ldots, n_{139}$ be the **actual** at bats during DiMaggio's 1941 season. A simulated baseball season will sample **with replacement** these at bats 139 times.
- For each "game" in the simulated baseball season, we will randomly generate the number of hits in each game by using a Binomial Distribution, where the probability of success is p=0.357.

- Runs *k* simulated "baseball seasons." Here the season is DiMaggio's 1941 season. We will set the number of simulates seasons to be 10,000.
- Let $n_1, n_2, \ldots, n_{139}$ be the **actual** at bats during DiMaggio's 1941 season. A simulated baseball season will sample **with replacement** these at bats 139 times.
- For each "game" in the simulated baseball season, we will randomly generate the number of hits in each game by using a Binomial Distribution, where the probability of success is p=0.357.
- We will look at the longest streak in each simulated baseball season. This streak is the number of consecutive "games" with a hit. We will have 10,000 of these longest streaks!

- Runs *k* simulated "baseball seasons." Here the season is DiMaggio's 1941 season. We will set the number of simulates seasons to be 10,000.
- Let $n_1, n_2, \ldots, n_{139}$ be the **actual** at bats during DiMaggio's 1941 season. A simulated baseball season will sample **with replacement** these at bats 139 times.
- For each "game" in the simulated baseball season, we will randomly generate the number of hits in each game by using a Binomial Distribution, where the probability of success is p=0.357.
- We will look at the longest streak in each simulated baseball season. This streak is the number of consecutive "games" with a hit. We will have 10,000 of these longest streaks!
- Estimate the probability of having a hitting streak of 56 games by counting the number of simulated seasons with a streak of 56 consecutive games or longer and divide by the number of simulated seasons here 10,000.

10,000 Simulations of DiMaggio's 1941 Season:

Method	Max	40+	<i>50+</i>	56+
Constant AB's	75	57	8	2
Varying AB's	57	41	2	1

Estimated probability of DiMaggio hitting safely in at least 56 straight games:

10,000 Simulations of DiMaggio's 1941 Season:

Method	Max	40+	50+	56+
Constant AB's	75	57	8	2
Varying AB's	57	41	2	1

Estimated probability of DiMaggio hitting safely in at least 56 straight games: $\frac{1}{10.000} = 0.0001$

But really we are interested in the probability of there ever being a 56-game hitting streak by **any** player achieving it over a given 56-game stretch. How do we do this?

Rockoff and Yates (2009, 2011) analyzed play-by-play data from Retrosheet.org for:

- National League only: 1911, 1921, 1922, 1953
- American and National League: 1920-1929, 1954-2007

Since then they have added 1930-1952 (both leagues), 1953 (American League), and 2008-2017 (both leagues). These seasons are not included in the analysis about to be discussed.

Simulation & Retrosheet Data

The process for hitter i in season j who plays in k games that season:

- $\bullet \ \mathbf{AB}_{ij} = (AB_{ij1}, AB_{ij2}, \dots, AB_{ijk})$
- The number of hits a player *i* in season *j* gets in game *k* (assuming that at-bats over the course of a single game are independent of each other):

$$H_{ijk} \sim \text{Binomial}(AB_{ijk}, p_{ij})$$

 A simulated season's worth of at-bats are the at-bats in each season sampled with replacement.

$$\boldsymbol{\mathsf{AB}}_{ij}^* = \left(AB_{ij1}^*, AB_{ij2}^*, \dots, AB_{ijk}^* \right)$$

• If m seasons are simulated, then for player i and season j:

$$\mathbf{AB}_{ij}^1, \mathbf{AB}_{ij}^2, \dots, \mathbf{AB}_{ij}^m$$

ullet The number of hits a player gets in each game in the $\emph{m}^{ ext{th}}$ simulated season is

$$\mathbf{H}_{ij}^{*m} \sim \mathsf{Binomial}\left(\mathbf{AB}_{ij}^{*m}, p_{ij}\right)$$

• Any run of hits in \mathbf{H}_{ij}^{*m} that are greater than zero denotes a hitting streak. The simulations will keep track of each player's maximum hitting streak in any given simulated season.

Top 10 Maximum Hitting Streaks:

		Simulated Season							
Player	Year	40+	50+	56+	Min	Q1	Q2	Q3	Max
Harry Heilmann	1921	34	9	5	10	18	22	27	91
Rogers Hornsby	1922	54	10	3	11	19	23	29	89
Felipe Alou	1966	4	3	2	9	14	16	21	75
Julio Franco	1991	3	1	1	8	14	16	20	74
Alex Rodriguez	1996	10	3	2	9	15	18	22	72
Rogers Hornsby	1921	21	3	1	6	14	17	22	71
Ichiro Suzuki	2004	34	8	5	11	18	22	27	69
Jimmy Rollins	2007	2	1	1	7	13	15	18	64
Rogers Hornsby	1922	23	4	2	7	15	18	23	63
Ralph Garr	1974	13	3	2	9	15	18	22	63

Hitting Streaks in 18,607,000 Simulated Player-Seasons:

Max	40+	50+	56+		
91	2237	284	85		

Results 1000 Simulated Baseball Histories

Hitting Streaks in 1000 Simulated Baseball Histories:

Max	40+	50+	56+
91	252	165	70

The estimated probability of a single player having a hitting streak of at least 56 games:

$$\frac{85}{18,607,000} = 0.000004568 = 0.0004568\%$$

The estimated probability of a hitting streak of at least 56 games occurring at **some point** in baseball history (technically 64 seasons of AL-NL and 4 seasons of just NL):

$$\frac{70}{1000} = 0.07 = 7\%$$

One of the big assumptions to the previous simulations and probability calculations?

One of the big assumptions to the previous simulations and probability calculations?

 The batting average ("probability of getting a hit") is constant for every single at bat over the course of the season

Solution?

One of the big assumptions to the previous simulations and probability calculations?

 The batting average ("probability of getting a hit") is constant for every single at bat over the course of the season

Solution?

• Vary the batting average over the course of a simulated season!

One of the big assumptions to the previous simulations and probability calculations?

 The batting average ("probability of getting a hit") is constant for every single at bat over the course of the season

Solution?

- Vary the batting average over the course of a simulated season!
- There are a variety of ways to do this!

One of the big assumptions to the previous simulations and probability calculations?

 The batting average ("probability of getting a hit") is constant for every single at bat over the course of the season

Solution?

- Vary the batting average over the course of a simulated season!
- There are a variety of ways to do this!

Another huge assumption in these calculations?

One of the big assumptions to the previous simulations and probability calculations?

 The batting average ("probability of getting a hit") is constant for every single at bat over the course of the season

Solution?

- Vary the batting average over the course of a simulated season!
- There are a variety of ways to do this!

Another huge assumption in these calculations?

• At bats are independent of one another. Why might this not be a good assumption to make? Is it a reasonable assumption to make?