

Analyzing the Feature Importance of Different Variables on the Price of Ikea Products

Philip Krück, Johannes Pein

Hamburg School of Business Administration

B.Sc. Business Informatics (18A-BI)

Digital Toolbox: Data Business

Lecturer: Ulf Köther

Group Number: 7

Matriculation Numbers: 3938 (P.Krück), 4001 (J.Pein)

04.12.2020

Contents

List of Figures	iv
List of Abbreviations	v
1 Introduction	1
2 Theoretical Background & Research Question	2
2.1 Data Set	2
2.2 Theoretical Background	2
2.2.1 Random Forest Basics	2
2.2.2 Overfitting	3
2.2.3 Feature Importance	3
2.3 Research Question	4
3 Methods	5
3.1 Data Cleaning and Transformation	5
3.2 Exploratory Data Analysis	7
3.2.1 Step 1: Outliers in Price and Independent Variables	7
3.2.2 Step 2: Homogeneity of Price	8
3.2.3 Step 3: Normality	9
3.2.4 Step 4: Missing Values	9
3.2.5 Step 5: Collinearity between Independent Variables	10
3.2.6 Step 6: Relationship between Independent Variables and Price	11
3.2.7 Step 7: Interactions	11
3.2.8 Step 8: Independence Observations of Response Variable Price	14
3.3 Random Forest Regression Model	15
4 Results	17

5 Discussion	19
5.1 Feature Importance	19
5.2 Conclusion	21

Appendices

A Plots	24
B Code Chunks	29
C Declaration of Honor	30
D Bibliography	32

List of Figures

3.1	Boxplots for Price in Euro Based on Independent Variables	8
3.2	Homogeneity of Category for Selected Combinations of Name and Category	9
3.3	Histogram of Numerical Variables with a Bin Width of 100 Euro . .	10
3.4	Percentage of Missing Values	11
3.5	Relationship of Price and Old Price	12
3.6	Relationship of Price and Size in Cubic Meters	12
4.1	Mean Feature Importance of the Predictor Variables on the Response Variable	18
A.1	Price Distribution per Designer	25
A.2	Price Distribution per Category	25
A.3	Price Distribution Other Colors	26
A.4	Price Distribution Sellable Online	26
A.5	Relationship between Independent Variables and Price	27
A.6	Coplot of Size and Price Split by Designer and Name	28

List of Abbreviations

R	Statistical Programming Language
VIF	Variance Inflation Factor
MSE	Mean Squared Error
IQR	Interquartile Range

Introduction

This project report is an examination at the Hamburg School of Business Administration in the module ‘Data Business’ as a part of a Bachelor of Science degree program. The students were given a data set and the task was to first explore the data and then choose a research question which was to be answered scientifically with the help of the statistical programming language R. The authors of this report were given a data set which contains Ikea products with different features, such as price, dimensional measures, name, category and designers of the product.

Theoretical Background & Research Question

2.1 Data Set

by P. Krück

The data set was obtained by a kaggle.com user (Reem Abdulrahman) by the means of webscraping techniques from the Saudi Arabian Ikea website in the furniture category on the 20th of April 2020. Noteworthy features include the name, category, price in Saudi Riyals, the designer and dimensions (width, height and depth). The data set has 13 variables and 2962 distinct observations after the removal of duplicates.

2.2 Theoretical Background

2.2.1 Random Forest Basics

by J. Pein

In order to analyze the feature importance in relation to the price variable, a

random forest regression model was chosen. A random forest consists of many decision trees, which predicts the response variable based on a majority decision process. In standard decision trees, each node is split to achieve the best performing model. In random forests however, the nodes are randomly split. Compared to linear regression, random forests not only take the mean and covariance structure into account, but also include deeper aspects of the data¹ resulting in a more advanced and robust model. To learn more about random forests, please see Breiman.²

2.2.2 Overfitting

by P. Krück

In statistical modelling, overfitting refers to the phenomenon where an analysis model corresponds too closely to a given data set and thus fails to generalize to new data or future observations.

2.2.3 Feature Importance

by J. Pein

There are different ways to measure feature importance. In this analysis permuting feature importance by a random forest algorithm is used. This algorithm leaves each feature out once while leaving all others unchanged, at each step calculating the mean squared error (MSE) of the predictions. This is done for every tree, calculating the overall MSE of each feature for the whole model.³

¹Grömping, “Variable Importance Assessment in Regression: Linear Regression versus Random Forest,” 317.

²“Random Forests.”

³Ibid., 23.

2.3 Research Question

by P. Krück

This paper explores the following research question:

How important are the different features of Ikea products in regard to their price?

The motivating forces for this research question are the possible implications for price determination of new items.

3.1 Data Cleaning and Transformation

by P. Krück

To examine the given data set properly, the authors first had to restructure and reformat it. This initial data cleaning step included type conversion, value mutation, addition of newly calculated fields and the removal of irrelevant columns.

Concretely, `name`, `category` and `designer` were converted to categorical variables. In the `designer` column, blank strings and values prefixed by “IKEA of Sweden” were converted to missing values (`NA`). Furthermore, both the price and old price were converted to double values and the currency was changed from Saudi Arabian Riyals to Euros based on the exchange rate from the time the data set was obtained by the author (See section 2.2).

Interestingly, the data set had a peculiarity where some rows were exact duplicates except for values in the `category` vector. The authors considered

multiple approaches to handle these data duplications without losing information about the category of an item.

One considered option was to merge the two category values into one column value via comma separation (e.g. "a" and "b" converts to "a, b"). However, this approach leads to the creation of many combinatorial categories with a low count of items per category. Additionally, it reduces the item count per category where the category is not comma separated. Overall this would lead to having many small categories which increases the difficulty in applying a regression model due to overfitting (See section 2.2.2).

The second option was to create separate columns for the different values of `category`. The data set would then have observations with category one, two and three. While no information is lost utilizing this approach, most observations in the second and third category column would contain missing values, thus increasing the difficulty of analysis using a predefined model (See section 3.3).

The authors chose the option of selecting the observations where the category count occurred most frequently when considering duplicates. The most frequent categories could be retained without including more column vectors into the data set as in option two.

To better facilitate the comparison of the different sizes of furniture items, the size in cubic meters was computed based on the depth, width and height values, and added as a column vector for further analysis.

Finally, the authors only selected columns that could have a potential impact on outcome of the analysis (See section 2.3) for further investigation. A detailed com-

Table 3.1: Initial Data Set formatting.

name	category	price	old_price	sellable_online	other_colors	designer	...
PAX...	Wardrobes	20450	No old price	TRUE	No	IKEA
ELV...	Wardrobes	7500	SR 820	TRUE	No	Ehlén...	...
ELV...	Wardrobes	15720	SR 1,755	TRUE	No	Ehlén...	...
ELV...	Wardrobes	9240	SR 1,050	TRUE	No	Ehlén...	...
ELV...	Wardrobes	27450	SR 3,130	TRUE	No	Ehlén...	...
ELV...	Wardrobes	12310	SR 1,535	TRUE	No	Ehlén...	...

Table 3.2: Transformed Data Set formatting.

name	category	price_eur	old_price_eur	sellable_online	other_colors	designer	size_m3
PAX...	Wardrobes	501.78	NA	TRUE	FALSE	Ehlén...	3.12
ELV...	Wardrobes	184.03	201.2	TRUE	FALSE	Ehlén...	NA
ELV...	Wardrobes	385.72	430.62	TRUE	FALSE	Ehlén...	NA
ELV...	Wardrobes	226.72	257.64	TRUE	FALSE	Ehlén...	NA
ELV...	Wardrobes	673.54	768.01	TRUE	FALSE	Ehlén...	NA
ELV...	Wardrobes	302.05	376.64	TRUE	FALSE	Ehlén...	NA

parison of the initial vs. transformed data structure can be seen in tables 3.1 and 3.2.

3.2 Exploratory Data Analysis

by P. Krück

The following sections explore our data based on the eight step data exploration protocol proposed by Zuur et al.¹

3.2.1 Step 1: Outliers in Price and Independent Variables

by P. Krück

Outliers of the chosen variables can be observed for each variable (see figure 3.1). Web scraping code is written in a generic form which makes it generalizable to all applied pages. This removes the occurrence of observation errors in the Ikea data set. Additionally, it is also unlikely that the outliers are due to falsely scraped items. To verify this hypothesis, the authors randomly tested outlier observations and manually checked them against the ikea website information. With a high

¹Zuur, “A protocol for data exploration to avoid common statistical problems.”




Figure 3.1: Boxplots for Price in Euro Based on Independent Variables

level of statistical confidence, the authors proceeded with further analysis without removing outliers as observation errors.

3.2.2 Step 2: Homogeneity of Price

by P. Krück

The homogeneity (homoscedasticity) of variance for price is explored by the means of conditional boxplotting. Within each name, and within each category the variance is heterogenous (see fig. 3.2). However, looking at both name and category in conjunction, it is possible to explore homoscedasticity of variance for price.

Due to the limited scope and length of this paper, the authors were not able to inspect all variable combinations for the three categorical plus two logical variables of the data set ($2^5 = 32$).




Figure 3.2: Homogeneity of Category for Selected Combinations of Name and Category

3.2.3 Step 3: Normality

by P. Krück

No numerical variables (`price`, `old_price` and `size_m3`) are arranged along a normal distribution (see fig. 3.3), but rather follow an exponential decay (e^{-x}).

3.2.4 Step 4: Missing Values

by P. Krück

All variables were examined for missing values. Only `designer`, `size_m3` and `old_price_eur` have missing values with percentages of 3.44%, 45.9% and 81% respectively (see fig. 3.4). The missing values for `designer` were deliberately set to `NA` by the authors in the case where the values contained digits, which is likely a scraping error. The `NA` values for the size can be explained due to the computation




Figure 3.3: Histogram of Numerical Variables with a Bin Width of 100 Euro

Table 3.3: Variance Inflation Factors for Numerical Variables

price_eur	old_price_eur	size_m3
77.97	78.54	2.36

of this column vector. `size_m3` is the product of `depth`, `width` and `height`. If one of those three values is missing, the end result is also a missing value. In contrast, the absence of the old price variables is due to the fact that most items are not on sale and thus do not have a missing value.

3.2.5 Step 5: Collinearity between Independent Variables

by P. Krück

The old price has a rather high variance inflation factor (VIF) which corresponds to high multicollinearity (see table 3.3). Contrarily, size has a low VIF which translates to low multicollinearity among the other independent variables (see table 3.3).




Figure 3.4: Percentage of Missing Values

3.2.6 Step 6: Relationship between Independent Variables and Price

by P. Krück

Inspecting the relationship between the independent variables and price, a strong correlation between `old_price_eur` and `size_m3` can be observed, while none can be detected for the other variables (see A.5). `old_price_eur` has a linear relationship (see fig. 3.5) whereas `size_m3` begins with a linear relationship as well, which is only disturbed by an outlier. (see fig. 3.6).

3.2.7 Step 7: Interactions

by P. Krück

The interactions between different variables are explored by the use of con-




Figure 3.5: Relationship of Price and Old Price




Figure 3.6: Relationship of Price and Size in Cubic Meters

ditioning plots (coplots). Utilizing this form of plotting the relationship of two numerical variables is explored by creating a matrix of plots subdivided by two categorical variables. In the given data set there are three numerical and three categorical variables which can be explored in this form of interaction. For the numerical variables, `old_price_eur` has such a strong relationship with `price` (see fig 3.5). A more detailed breakdown by the categorical variables would not reveal new information. This leaves the exploration of `size_m3` and `price_eur` subdivided by designer, name and category resulting in $\binom{3}{2} = 3$ combinations of coplots.

Interaction of size and price coplotted by designer and name

by P. Krück

It is unlikely that there is an interaction between size and price split by name and designer as is indicated by the non-parallelism of the fitted lines in the coplot (see figure A.6).

Problems with Coplot Development

by P. Krück

This section describes a programming error the authors ran into regarding coplotting.

Unfortunately, the authors of this papers weren't able to fully explore all combinations. Plotting designer and name works (see above) while the other two options would not plot properly. The authors could not fully debug the problem with these plots. The linear model predicted infinite values for some of the coplotted combinations for both amalgamations that would not render correctly. Dropping

all NA values left 354 observations and the coplot would correctly render for the combination of `name` and `category` while for `name` and `designer` it would not. The number of observations for `name` and `category` is rather low considering the additional categorical subdivision which lead the authors to discard it as an insignificant research finding. Still there seemed to be infinite values outputted by the linear method. The authors hypothesized that those values were caused by a division by 0 of the internal algorithm mechanics. This however proved to be wrong after applying the respective filters. The code for the two problematic plots can be viewed in Appendix chapter B.

The inclined reader is encouraged to dabble with the code. Sending any hints or even a solution to fix the code and fully render the plots would be highly appreciated by the authors.²

3.2.8 Step 8: Independence Observations of Response Variable Price

by P. Krück

The independence of observations of the response variables assumes that “[...] information from any one observation should not provide information on another after the effects of other variables have been accounted for.”³ The data cleaning step left the observed data set in a tidy format which implies that the observations are independent of each other.⁴

²Any questions, hints or solutions may kindly be sent to philip.krueck@myhsba.de.

³Ibid., 11, ll.23–26.

⁴“There are three interrelated rules which make a dataset tidy: Each variable must have its own column. Each observation must have its own row. Each value must have its own cell.” (, Wickham, *R for Data Science*, 149, ll.4–7.)

3.3 Random Forest Regression Model

by J. Pein

To analyze the feature importance of different variables on the price variable, a random forest model was chosen. In the given data set, categorical variables with lots of different values can be observed. Including all of those values in, for example, a linear regression model, would most likely lead to overfitting (see section 2.2.2). This is because the linear model would learn the immanent data structure of the sample, but would not be able to generalize from this. Thus, the authors chose the random forest model, because according to Breiman⁵ it is robust against overfitting.

In this analysis, the R *randomForest* package was used, which is based on the original Breiman and Cutler's Fortran code for random forest regression. To reproduce the results, the further data preparation steps are described here. These steps are based on the cleaned ikea data set which is described in section 3.1. This data set is then transformed further allowing it to be used with the *randomForest* package. To learn more about how random forests work see chapter 2, to learn more about the *randomForest* package see Liaw and Wiener.⁶

First, the variable `old_price_eur` is removed from the cleaned ikea data set, due to a very high correlation and relationship to the response variable `price_eur` (price) analyzed in section 3.2.5 and section 3.2.6. Then, the `designers` and `names`, which are not part of the 50 `designers` and 49 `names` with the highest number of occurrences, are grouped in the `other` value. This is because the `randomForest` method does not allow categorical variables with more than 53 predictors. The

⁵“Random Forests,” 29.

⁶“Classification and Regression by randomForest.”

last step deals with the missing values in the data. As described in section 3.2.4, there are many missing values in the `size_m3` and `designer` variables. To apply the `randomForest` method of the *randomForest* package on the data, those missing values are treated using three different approaches. In the first approach the rows with missing values are deleted, reducing the total number of rows by approximately 50%. In the second approach the missing values are dummy coded with a value of -1000. The third approach uses the `na.roughfix = na.omit` argument, which is the built-in way of the *randomForest* package to deal with missing values. After preparing the data, the `randomForest` method of the *randomForest* package is applied to the data with the number of trees set to 2000 and importance set to `TRUE`, training the random forest model with the prepared ikea data set.

```
randomForest(price_eur ~ ., rf_ikea, ntree=2000, importance=TRUE)
```

Then the `importance` method of the *randomForest* package is used to calculate the feature importances, which are computed by permuting feature importance, which was introduced in section 2.2.3. The three different approaches of dealing with the missing values in the data set lead to different results, so the authors chose to calculate the mean result of the three approaches. The result of this analysis is presented in the following chapter.

4

Results

by J. Pein

In this chapter, the result of the analysis of the feature importance of different features on the response variable price of Ikea products are presented.

As described in section 3.3, the feature importance was calculated using permuting feature importance of the *randomForest* R package. In this analysis, feature importance is derived from the percentage increase of the mean squared error (MSE) of the overall random forest regression model in regard to the response variable `price_eur`. A larger percentage increase of the MSE implies greater feature importance. Conversely, a lower percentage increase of the MSE translates to less feature importance.

Thus, as can be seen in figure 4.1, the most important feature is `size_m3` with an increase of the MSE of 182%. The second, third and fourth most important features are `designer` with an MSE increase of 120%, `name` with an increase of 114% and `category` with an increase of 105%. The fifth most important feature




Figure 4.1: Mean Feature Importance of the Predictor Variables on the Response Variable

is `other_colors` with a MSE increase of 78% and the least important feature is `sellable_online` with a 9% increase.

These results are further discussed in the following chapter.

5

Discussion

by J. Pein

In this chapter, the results are discussed in connection with the research question. The question that the results were supposed to answer is the following:

How important are the different features of Ikea products in regard to their price?

5.1 Feature Importance

The `size_m3` variable is the most important feature. Probably the main reason for this is the size of a product being closely linked to its material cost. Big items are generally more costly to produce, thus leading to a higher selling price and vice versa. Due to the high correlation to the price variable described in section 3.2.5 and section 3.2.6, it is worth discussing in future research whether or not to include this variable in a possible predictive analysis model .

The `designer` variable is the second most important feature. When looking at the price distribution per designer, it can be clearly seen that the interquartile

range (IQR) of price varies for each designer. In addition, the IQR often is smaller than 300€, thus showing a tendency towards a certain price range, which might be the reason for the relatively high feature importance of the designer variable. For the plot, see figure A.1.

The **name** variable includes around 50 different product names with partly small numbers of occurrences. But, as can be seen in figure 3.2, certain product lines (names) tend to be more expensive than others. *LIDTHULT*, for example, is the most expensive product line in each of the categories *beds*, *chairs* and *sofas & armchairs* while *HEMNES* is on the lower end of the price scale. This behavior explains a relatively high feature importance of the **name** variable.

Category is another feature with a relatively high importance. One reason for this is the different category's price distributions showing a clear tendency towards certain price segments (see figure A.2), i.e. wardrobes and beds are generally more expensive than chairs.

The feature importance of **other_colors** is the second lowest, but still considerable. This still relatively high feature importance might be due to the difference of the mean price and the relatively small IQR (see: figure A.3). On the other hand, there is a large overlapping area within the IQR in the two expressions of **other_colors** possibly reducing the feature importance.

The very low feature importance of the **sellable_online** variable is probably because the low number of occurrences of a product being sellable online. Only around 0.6% of the products are sellable online.

5.2 Conclusion

The data used in this paper was scraped from the Saudi Arabian Ikea website, thus this analysis mainly focusses on Ikea products in the Saudi Arabian market. To analyze the geographically independent feature importances, more data should be scraped from other international Ikea websites. The research question of this paper, *How important are the different features of Ikea products in regard to their price?*, could thus not be answered for the global Ikea product market, but for the Saudi Arabian market only.

Also, in further research, the results presented on the feature importance of the predictor variables on the response variable price should be validated by other techniques than the random forest model used in this analysis to achieve unbiased results.

Furthermore, based on this analysis a predictive model could be developed which predicts the price of Ikea products in the Saudi Arabian Market based on the features analyzed. This could be used by Ikea internally to analyze if the price of their new product aligns with the prices of the currently available product or by market researchers.

Lastly, the authors were surprised by the high feature importances of the variables **designer** and **name** computed by the random forest model. Both variables have lots different values, suggesting an overfitting of the random forest model. This, however, is objected by scientific research.¹ Exploring what the reason behind the high feature importances of the variables **designer** and **name** and if they are due

¹Breiman, "Random Forests," 29.

to overfitting of a random forest model should be analyzed in future research. If the high feature importance of the variables truly is not due to overfitting, this example backs up the thesis of Grömping² that random forests are able to discover deeper patterns in the data. These patterns are beyond the discussion applied here, showing the power of random forests models.

²“Variable Importance Assessment in Regression: Linear Regression versus Random Forest,” 317.

Appendices

A Plots



Figure A.1: Price Distribution per Designer






Figure A.2: Price Distribution per Category

**Figure A.3:** Price Distribution Other Colors

Price Distribution Sellable Online

**Figure A.4:** Price Distribution Sellable Online




Figure A.5: Relationship between Independent Variables and Price




Figure A.6: Coplot of Size and Price Split by Designer and Name

B

Code Chunks

```
coplot(size_m3 ~ price_eur | fct_lump_n(category, 5) +  
      fct_lump_n(name, 5), data = drop_na(tidy_ikea),  
ylab = "Size in m^3",  
xlab = "Price in Euro", panel = function(x, y, ...) {  
  tmp <- lm(y ~ x, na.action = na.omit)  
  abline(tmp)  
  points(x, y )})
```

```
coplot(size_m3 ~ price_eur | fct_lump_n(designer, 5) +  
      fct_lump_n(category, 5), data = drop_na(tidy_ikea),  
ylab = "Size in m^3",  
xlab = "Price in Euro", panel = function(x, y, ...) {  
  tmp <- lm(y ~ x, na.action = na.omit)  
  abline(tmp)  
  points(x, y )})
```

C

Declaration of Honor

Declaration: We hereby declare that we

1. wrote this project report without the assistance of others;
2. have marked direct quotes used from the literature and the use of ideas of other authors at the corresponding locations in the thesis;
3. have not presented this thesis for any other exam. We acknowledge that a false declaration will have legal consequences.

Hamburg, 01.12.20

(Place, date) (Signatures)

JPlein

We accept that HSBA may check the originality of my work using a range of manual and computer-based techniques, including transferring and storing my submission in a database for the purpose of data-matching to help detect plagiarism.

C. Declaration of Honor

31

Hamburg, 01.12.20

(Place, date) (Signatures)

Hein

D Bibliography

Breiman, Leo. “Random Forests.” *Machine Learning*, no. 45 (2001). <https://link.springer.com/content/pdf/10.1023/A:1010933404324.pdf>.

Grömping, Ulrike. “Variable Importance Assessment in Regression: Linear Regression versus Random Forest.” *The American Statistician*, nos. Vol. 63, No. 4 (2009): 308–19. https://prof.beuth-hochschule.de/fileadmin/prof/groemp/downloads/tast_2E2009_2E08199.pdf.

Liaw, Andy, and Matthew Wiener. “Classification and Regression by randomForest.” *R News*, nos. Vol. 2/3, December 2002 (2001). https://www.researchgate.net/publication/228451484_Classification_and_Regression_by_RandomForest.

Wickham. *R for Data Science*. O’Reilly Media, 2017.

Zuur. “A protocol for data exploration to avoid common statistical problems.” *British Ecological Society*, 2010. doi:10.1145/1738826.1738829.