

Homework Nr.1

Philipp Stassen

September 26, 2018

Remark. In this solution proposals a ring is always meant to be a commutative ring with 1.

7.1

Exercise 26

Let K be a field and $\nu : K^\times \rightarrow \mathbb{Z}$ be a discrete valuation on K . Let $R = \{x \in K^\times \mid \nu(x) \geq 0\} \cup \{0\}$.

a) Claim: R is a subring of K which contains the identity.

Proof. It suffices to show that R is closed under *addition*, *multiplikation* and *additive inverses* as well as that it contains 0 and 1. Properties as commutativity of $+$ and \cdot , associativity or distributivity are all valid as $R \subset K$ and K is a field.

1. R is closed under *addition*: Let $x, y \in R$.

Case 1: $x + y \neq 0$. Then $\nu(x) \geq 0$ and $\nu(y) \geq 0$. Hence, $\nu(x + y) \geq \min\{\nu(x), \nu(y)\} \geq 0$. Therefore, $x + y \in R$.

Case 2: $x + y = 0$. This is trivial as $0 \in R$.

2. R is closed under *multiplikation*:

Let $x, y \in R$ then $\nu(x) \geq 0$ and $\nu(y) \geq 0$. We have that $\nu(xy) = \nu(x) + \nu(y) \geq 0$. Therefore, $xy \in R$.

3. R contains 0.

4. R contains 1: $\nu(a) = \nu(a1) = \nu(a) + \nu(1)$. Therefore, $\nu(1) = 0$ and $1 \in R$.

5. R is closed under building *additive inverses*:

Observe that we have for any $x \in R$

$$0 = x \cdot 0 = x \cdot (1 + (-1)) = x \cdot 1 + x \cdot (-1) \quad (1)$$

$$\iff -x = (-1) \cdot x \quad (2)$$

and by in particular $1 = -(-1) = (-1) \cdot (-1)$. Furthermore, it is

$$0 = \nu(1) = \nu((-1) \cdot (-1)) = 2\nu(-1) \quad (3)$$

Therefore, $\nu(-1) = 0$. Now we can conclude that for any $x \in K$ holds $\nu(-x) = \nu(x \cdot (-1)) = \nu(x) + \nu(-1) = \nu(x)$. Hence, if $x \in R$ then also $-x \in R$.

□

b) Claim: For every nonzero element $x \in K$ either x or x^{-1} is in R .

Proof. Let $x \in K^\times$, as K is a field every nonzero element is a unit. As ν is a discrete valuation we have

$$0 = \nu(1) = \nu(x \cdot x^{-1}) = \nu(x) + \nu(x^{-1}) \quad (4)$$

Hence, $\nu(x) = -\nu(x^{-1})$ and $\nu(x) \geq 0$ or $\nu(x^{-1}) \geq 0$. This implies that either $x \in R$ or $x^{-1} \in R$. □

c) $x \in R$ is a unit iff $\nu(x) = 0$.

Proof. " \implies " Let $x \in R$ be a unit. Then there is $b \in R$ such that $xb = 1$. We have that $\nu(x) \geq 0$ and $\nu(b) \geq 0$ as $x, b \in R$.

Furthermore,

$$0 = \nu(1) = \nu(xb) = \nu(x) + \nu(b). \quad (5)$$

This implies that $\nu(x) = \nu(b) = 0$.

" \impliedby " Let $\nu(x) = 0$. We have that

$$0 = \nu(1) = \nu(x \cdot x^{-1}) = \nu(x) + \nu(x^{-1}) = \nu(x^{-1}). \quad (6)$$

Hence, $\nu(x^{-1}) \geq 0$ and $x^{-1} \in R$. This makes x a unit of R . □

Chapter 7.4

Exercise 37

Let R be a local ring. Let \mathfrak{m} be the maximal ideal of R .

Claim: Every element of $R - \mathfrak{m} = \{r \in R \mid r \notin \mathfrak{m}\}$ is a unit.

Proof. Assume $r \in R$ is a nonunit and $r \notin \mathfrak{m}$. Then the ideal that is generated from \mathfrak{m} and r - let's call it M - is strict larger than \mathfrak{m} (as it contains r) but also $M \subsetneq R$ as $1 \notin M$. This contradicts the maximality of \mathfrak{m} . □

Let $M = \{r \in R \mid r \text{ is not a unit}\}$ be the ideal of nonunit forms. Claim: M is the unique maximal ideal in R .

Proof. Assume there is an ideal $\mathfrak{m} \supsetneq M$ then there exists $r \in \mathfrak{m}$ such that $r \notin M$. Hence, r is a unit and $1 \in \mathfrak{m}$. Therefore, $\mathfrak{m} = R$ and \mathfrak{m} is not a proper ideal. This implies that M is a maximal ideal.

M is also unique, as any proper ideal $I \subsetneq R$ is also contained in M . If there was an Ideal $I \not\subseteq M$ then by the same argument as before we already have $I = R$. \square

Exercise 40

Lemma. Let $\mathfrak{p} \subsetneq R$ be a prime ideal then $\mathfrak{N}(R) \subseteq \mathfrak{p}$.

Proof. Let $r \in \mathfrak{N}(R)$ and $n \in \mathbb{N}$ such that $0 = r^n = r^{n-1}r$, we want to show that $r \in \mathfrak{p}$. As $r^n = 0 \in \mathfrak{p}$ we have that either $r \in \mathfrak{p}$ or $r^{n-1} \in \mathfrak{p}$. Wlog $r^{n-1} \in \mathfrak{p}$ (otherwise we are done). Now we can induct on $n \in \mathbb{N}$ eventually showing that $r \in \mathfrak{p}$. \square

Claim: Let R be a commutative Ring and let $\mathfrak{N}(R)$ be the nilradical of R . The following are equivalent:

1. R has exactly one prime ideal
2. every element of R is either nilpotent or a unit.
3. $R/\mathfrak{N}(R)$ is a field

Proof. "(i) \implies (ii)" Let $\mathfrak{p} \subsetneq R$ be the only prime ideal.

We know that $\mathfrak{N}(R) = \bigcap_{\mathfrak{q} \text{ prime}} \mathfrak{q}$ ¹. Hence, $\mathfrak{N}(R) = \mathfrak{p}$. Furthermore, every proper ideal is contained in a maximal ideal². Hence, we have $\mathfrak{p} \subseteq \mathfrak{m}$. As every maximal ideal in a commutative ring is prime³ we must have $\mathfrak{p} = \mathfrak{m}$. Now we can conclude that either $r \in \mathfrak{N}(R)$, i.e. r is nilpotent, or $r \in R - \mathfrak{N}(R)$, i.e. r is a unit, (by exercise 37).

"(ii) \implies (iii)" Let every element of R be either nilpotent or a unit. $\mathfrak{N}(R)$ is an ideal. It is also maximal as every proper superset of $\mathfrak{N}(R)$ contains a unit. Hence, $R/\mathfrak{N}(R)$ is a field⁴.

"(iii) \implies (i)" Let $R/\mathfrak{N}(R)$ be a field. Then $\mathfrak{N}(R)$ is a maximal ideal⁵. Let \mathfrak{p} be a prime ideal then $\mathfrak{N}(R) \subseteq \mathfrak{p}$ and as $\mathfrak{N}(R)$ is maximal we have $\mathfrak{p} = \mathfrak{N}(R)$. Therefore $\mathfrak{N}(R)$ is the only prime ideal of R . \square

¹DF §15.2 Proposition 12

²DF §7.4 Proposition 11

³DF §7.4 Corollary 14

⁴DF §7.4 Proposition 12

⁵DF §7.4 Proposition 12

Chapter 15.1

Exercise 2

a) Let R denote the Ring of continuous, real-valued functions on $[0, 1]$. Claim: R is not noetherian.

Proof. We define the sequence of ideals

$$I_n := \{f \in R \mid f(x) = 0, \forall x \leq \frac{1}{n}\} \quad (7)$$

Claim(1): Every I_n is an ideal.

We need to show that

1. I_n is nonempty and $\forall x, y \in I : x - y \in I$

proof. Take $f, g \in I_n$ then $\forall x \leq \frac{1}{n} : (f - g)(x) = 0$. Hence, $f - g \in I_n$.

2. $\forall x \in I, \forall r \in R : r \cdot x \in I$

proof. Take $f \in I_n$ and $r \in R$. Then $(f \cdot r)(x) = f(x)r(x)$. As $f(x) = 0$ for all $x \leq \frac{1}{n}$ we also have that $f(x)r(x) = 0$ for all $x \leq \frac{1}{n}$. Hence, $(f \cdot r) \in I_n$. $\square(1)$

Furthermore, for all $m, n \in \mathbb{N}$ with $m < n$ we have that $I_m \subsetneq I_n$. Hence, $I_0 \subsetneq I_1 \subsetneq \dots \subsetneq$ is an infinite ascending chain of ideals that is not eventually constant. \square

b) Let X be any infinite set and let $R = \{f : X \rightarrow \mathbb{Z}/2\mathbb{Z}\}$ denote the Ring of all functions from X to $\mathbb{Z}/2\mathbb{Z}$. Claim: R is not noetherian.

Proof. As X is infinite we have an injection $\iota : \mathbb{N} \hookrightarrow X$. Now we can define the ideals

$$I_n := \{f \in R \mid f(\iota(m)) = 0, \forall m \geq n\} \quad (8)$$

Claim(1): Every I_n is an ideal.

We need to show that

1. I_n is nonempty and $\forall x, y \in I : x - y \in I$

proof. Take $f, g \in I_n$ then $\forall m \geq n : (f - g)(\iota(m)) = 0$. Hence, $f - g \in I_n$.

2. $\forall x \in I, \forall r \in R : r \cdot x \in I$

proof. Take $f \in I_n$ and $r \in R$. Then $(f \cdot r)(\iota(m)) = f(\iota(m))r(\iota(m))$. As $f(\iota(m)) = 0$ for all $m \geq n$ we also have that $f(\iota(m))r(\iota(m)) = 0$ for all $m \geq n$. Hence, $(f \cdot r) \in I_n$. $\text{qed}(1)$

we have that each I_n is an ideal and that $I_0 \subsetneq \dots \subsetneq \dots$ is an ascending chain of ideals that is not eventually constant. \square

Exercise 3

Claim: Let K be a field. $K(x)$ is not a finitely generated k -algebra.

Proof. Let $g_1, \dots, g_n \in K(x)$ with $g_i = \frac{P_i}{Q_i}$ and $P_i, Q_i \in K[X]$ be a finite set of generators. Define $d := \prod_{i \leq n} Q_i$. Then we have for each g_i that $g_i \in K[X, d^{-1}]$ as

$$g_i = \frac{P_i \cdot \prod_{j \neq i} Q_j}{d^{-1}} \quad (9)$$

with $P_i \cdot \prod_{j \neq i} Q_j \in K[X]$. Hence, $K[g_1, \dots, g_n] \subset K[X, d^{-1}]$.

Notice, that $K[X]$ contains infinitely many irreducible polynomials. This can be easily concluded from the fact that $K[X]$ contains infinitely many primes⁶

Hence, we can take an irreducible polynomial $R \in K[X]$ such that $R \nmid d$.

Claim(1): $\frac{1}{R} \notin K[X, d]$.

Note that - by taking the greatest common divisor - each $p \in K[X, d^{-1}]$ is of the form

$$p = P + p_n d^{-n} + \dots + p_0 = Q d^{-n} \quad (10)$$

with $P, Q \in K[X]$. Assume we had

$$\frac{1}{R} = \frac{Q}{d^n}. \quad (11)$$

Hence, $1 = Q \frac{R}{d^n}$. However, since R is irreducible and $R \nmid d$ this is not possible. \square

Exercise 4

Claim: If R is noetherian then so is $R[[X]]$. Similar to the "degree" of a polynomial one can define the "degree" of a power series:

$$\deg : R[[X]] \rightarrow \mathbb{N} \quad (12)$$

$$\deg \left(\sum_{i=0}^{\infty} r_i X^i \right) := \begin{cases} \min\{i \in \mathbb{N} | r_i \neq 0\}, & \text{if } f \neq 0 \\ 0, & \text{else} \end{cases} \quad (13)$$

Definition. For any power series we define $\text{lead}(a_k + a_{k+1}X + \dots) := a_k$ to be the minimal nonzero coefficient. Furthermore for any ideal $I \subseteq R[[X]]$ let

$$L := \{\text{lead}(P) | P \in I\} \cup \{0\} \quad (14)$$

$$L_d := \{\text{lead}(P) | P \in I \wedge \deg(P) = d\} \cup \{0\} \quad \forall d \geq 1 \quad (15)$$

⁶similar to euclids proof for the conjecture that there are infinitely many primes in \mathbb{Z}

Claim(1): For every $d \geq 1$ L_d is an ideal, i.e. $\forall a, b \in L_d, \forall r \in R : ar - b \in L_d$.

proof. Let $a, b \in L_d$ and $r \in R$. Wlog $ar - b \neq 0$, otherwise there is nothing to show. Take $P, Q \in I$ with $\deg(P) = \deg(Q) = d$ such that $\text{lead}(P) = a$ resp. $\text{lead}(Q) = b$. Then we have $\text{lead}(rP - Q) = ar - b$. qed(1)

Claim(2): L is an ideal.

proof. Let $a, b \in L$ and $r \in R$. Again we may assume that wlog $ar - b \neq 0$. As $a, b \in L$ we can take $P, Q \in I$ such that $\text{lead}(P) = a$ respectively $\text{lead}(Q) = b$. Furthermore, we define $d := \deg(P)$ and $e := \deg(Q)$. Therefore, it follows that $\text{lead}(rPx^e - Qx^d) = ar - b$ qed(2)

proof of claim. As R is noetherian we have that L and all the L_d are finitely generated; let $\{a_1, \dots, a_n\}$ and $\{a_{d,1}, \dots, a_{d,n_d}\}$ be generators for L and L_d . Choose $P_i, Q_{d,i} \in I$ such that $\text{lead}(P_i) = a_i$, respectively $\deg(Q_{d,i}) = d$ and $\text{lead}(Q_{d,i}) = a_{d,i}$. We define $e_i := \deg(P_i)$, $N := \max_{i=1}^n e_i$ and finally

$$I' := \langle \{P_i, | 1 \leq i \leq n\} \cup \{Q_{d,i} | 0 \leq d \leq N, 1 \leq i \leq n_d\} \rangle_{R[[X]]}. \quad (16)$$

Clearly I' is finitely generated and $I' \subseteq I$. It remains to show that $I' = I$.

Claim(3): $I \subseteq I'$.

proof. Let $V \in I$ such that $\deg(V) = d$ and $\text{lead}(V) = a$.

Case 1: $d \leq N$. Let $a = \sum_{i=1}^{n_d} r_i a_{d,i}$. Furthermore, let

$$Q_d = \sum_{i=1}^{n_d} r_i Q_{d,i}, \quad (17)$$

then clearly $Q_d \in I'$, $\deg(Q_d) = d$ and $\text{lead}(Q_d) = \sum_{i=1}^{n_d} r_i a_{d,i} = a$. Hence, we can define $V_{d+1} := V - Q_d$ and thus have $\deg(V_{d+1}) \geq d+1$ as well as $V_{d+1} \in I$. We iterate the procedure for $d \leq N$ and get

$$V_{N+1} = V - \sum_{i=d}^N Q_i \quad (18)$$

with $V_{N+1} \in I$ and $\deg(V_{N+1}) \geq N+1$. As we have that $\sum_{i=d}^N Q_i \in I'$ it follows that $V \in I'$ iff $V_{N+1} \in I'$. In conclusion, we reduced the problem to Case 2 as we may assume without loss of generality that $\deg(P) > N$.

Case 2: $d > N$. Let $a_d = \sum_{i=1}^n r_{i,d} a_i$. Then we can define

$$P'_d = \sum_{i=1}^n r_{i,d} P_i X^{d-e_i} \quad (19)$$

and hence get $P'_d \in I'$, $\deg(P'_d) = d$ and $\text{lead}(P'_d) = a_d$. We can proceed as before to get an infinite series

$$\sum_{j=d}^{\infty} P'_j = \sum_{j=d}^{\infty} \sum_{i=1}^n r_{i,d} P_i X^{j-e_i} \quad (20)$$

$$= \sum_{i=1}^n P_i \sum_{j=d}^{\infty} r_{i,d} X^{j-e_i} \quad (21)$$

$$= \sum_{i=1}^n P_i h_i \quad \text{with} \quad (22)$$

$$h_i := \sum_{j=d}^{\infty} r_{i,d} X^{j-e_i}. \quad (23)$$

Now as $h_i \in R$ we have that $\sum_{i=1}^n P_i h_i \in I'$. Finally, we can conclude that $V \in I'$ by the fact that the series of coefficients of V and $\sum_{j=d}^{\infty} P'_j$ is by construction the same and hence $V = \sum_{j=d}^{\infty} P'_j$. \square

Exercise 5

Let M be a noetherian R -module and $\varphi : M \rightarrow M$ an R -module endomorphism of M .

Lemma. Let M, N be R -modules and $\varphi : M \rightarrow N$ a R -module-morphism. Then $\ker(\varphi)$ is a submodule of M .

Proof. We need to verify that $\ker(\varphi)$ is nonempty and $\forall x, y \in \ker(\varphi), \forall r \in R : x + ry \in \ker(\varphi)$. Clearly $\ker(\varphi) \neq \emptyset$ as $0 \in \ker(\varphi)$. Now let $x, y \in \ker(\varphi)$ and $r \in R$. We have that $\varphi(x) = \varphi(y) = 0$. Hence, $\varphi(x + ry) = \varphi(x) + r\varphi(y) = 0 + r0 = 0$. Therefore $x + ry \in \ker(\varphi)$. \square

Claim: There is an $n \in \mathbb{N}$ such that $\ker(\varphi^n) \cap \text{im}(\varphi^n) = 0$.

Proof. As $\varphi(0) = 0$ and we have that $\ker(\varphi) \subset \ker(\varphi^2) \subset \dots$ is an increasing chain of submodules. As M is noetherian it is eventually constant and there exists $N \in \mathbb{N}$ such that $\ker(\varphi^N) = \ker(\varphi^m)$ for all $m \geq N$.

In particular $\ker(\varphi^N) = \ker(\varphi \circ \varphi^N)$. Hence, $\ker(\varphi|_{\text{im}(\varphi^N)}) = 0$ and $\varphi|_{\text{im}(\varphi^N)}$ is injective. Hence, if $x \in \text{im}(\varphi^{N+1})$ and $\varphi(x) = 0$ then $x = 0$. This implies that $\text{im}(\varphi^{N+1}) \cap \ker(\varphi^{N+1}) = 0$. \square

Claim: If φ is surjective then φ is an isomorphism.

Proof. If $\varphi : M \rightarrow M$ is surjective then so is φ^n for any $n \in \mathbb{N}$. As before we have $N \in \mathbb{N}$ such that $\varphi|_{\text{im}(\varphi^N)}$ is injective. Considering that φ^N is surjective we can conclude that $\text{im}(\varphi^N) = M$ and therefore $\varphi|_{\text{im}(\varphi^N)} = \varphi$ is both, surjective and injective, and thus an isomorphism. \square

Exercise 6

Consider the R -modules M, M', M'' and the following exact sequence; I will refer to it as \mathcal{S} from now on.

$$0 \longrightarrow M' \xrightarrow{f} M \xrightarrow{g} M'' \longrightarrow 0$$

Claim: M is a noetherian R -module iff M' and M'' are noetherian R -modules.

Proof. " \implies " Let M be a noetherian R -module.

Let $M'_1 \subseteq M'_2 \subseteq \dots$ be an increasing sequence of submodules of M' . Then $f(M'_1) \subseteq f(M'_2) \subseteq \dots$ is an increasing sequence of submodules of M . As M is noetherian there exists $N \in \mathbb{N}$ such that $f(M'_N) = f(M'_m)$ for all $m \geq N$. As f is injective we also have $M'_N = M'_m$ for all $m \geq N$. Hence, the (arbitrary) sequence of R -submodules in M' is eventually constant and thus M' noetherian.

Let $M''_1 \subseteq M''_2 \subseteq \dots$ be an increasing sequence of R -submodules of M'' . It follows that $g^{-1}(M''_1) \subseteq g^{-1}(M''_2) \subseteq \dots$ is an increasing sequence R -submodules of M . As M is noetherian the latter one is eventually constant, let's say from index N onwards. As g is surjective we have that $g(g^{-1}(M''_i)) = M''_i$. Hence, we can conclude that $M''_1 \subseteq M''_2 \subseteq \dots$ is constant from N onwards as well.

qed(\implies)

" \impliedby " Let M' and M'' be noetherian R -modules.

Let $M_1 \subseteq M_2 \subseteq \dots$ be an increasing sequence of submodules of M . This implies that $f^{-1}(M_1) \subseteq \dots$ is an increasing sequence in M' and $g(M_1) \subseteq \dots$ is an increasing sequence in M'' . The latter ones are eventually constant as M' and M'' are noetherian; let us assume from index N' resp. N'' onwards. We define $N := \max(N', N'')$. If we prove that $M_n \subseteq M_N$ for any $n \geq N$ we can conclude that $M_1 \subseteq \dots$ is eventually constant and thus M noetherian (as the sequence was arbitrary).

$$\begin{array}{ccccc}
 & & y & & \\
 & & \cap & & \\
 f^{-1}(M_N) & \xleftarrow{f} & M_N & \xrightarrow{g} & g(M_N) \\
 \parallel & & \cap & & \parallel \\
 f^{-1}(M_n) & \xleftarrow{f} & M_n & \xrightarrow{g} & g(M_n) \\
 \psi & & \psi & & \\
 f^{-1}(x-y) & & x & &
 \end{array}$$

Figure 1:

Let $x \in M_n$. As $g \upharpoonright_{M_n} \rightarrow g(M_n)$ there exists $y \in M_N$ such that $g(y) = g(x)$ (see also fig. 1). It follows that $x - y \in \ker(g) \cap M_n$. As \mathcal{S} is exact we have that $x - y \in \text{im}(f) \cap M_n$ and thus may consider $f^{-1}(x - y) \in M'_n = M'_N$. As f is injective we have that $f(f^{-1}(x - y)) = x - y \in M_N$. But this means that $x = y + (x - y) \in M_N$ and as $x \in M_n$ was arbitrary we can conclude $M_n \subset M_N$. \square

Exercise 11

Let R be a commutative ring in which all prime ideals are finitely generated.

a) Claim: If the collection of ideals that are not finitely generated is nonempty, then it contains a maximal element I . As a consequence we have that R/I is a noetherian ring.

Proof. Let $L := \{J \subset R \mid J \text{ is not f.g. ideal}\}$. We want to use the *Lemma of Zorn* to deduce that L possesses a maximal element. Clearly L is partially ordered by \subset . We need to find an upper bound in L for every chain in L . Let $J_1 \subset J_2 \subset \dots$ be a chain in L . Clearly $\bigcup_{i>0} J_i$ is an upper bound; it remains to show that $\bigcup_{i>0} J_i \in L$.

Claim(1): $\bigcup_{i>0} J_i$ is an ideal.

proof. Take $x, y \in \bigcup_{i>0} J_i$ then there is an $n \in \mathbb{N}$ such that $x, y \in J_n$. Hence, $x - y \in J_n \subset \bigcup_{i>0} J_i$.

Furthermore, let $x \in \bigcup_{i>0} J_i$ and $r \in R$. Again there exists $n \in \mathbb{N}$ such that $x \in J_n$. Hence, $r \cdot x \in J_n \subset \bigcup_{i>0} J_i$. qed(1)

Claim(2): $\bigcup_{i>0} J_i$ is not finitely generated, i.e. $\bigcup_{i>0} J_i \in L$.

proof. Assume $\bigcup_{i>0} J_i$ was finitely generated by the generators g_1, \dots, g_m . Take $n \in \mathbb{N}$ such that $g_1, \dots, g_m \in J_n$. But this implies that $J_n \subseteq \bigcup_{i>0} J_i \subseteq J_n$ and J_n would be finitely generated. Hence, $J_n \notin L$. \nmid qed(2)

By *Zorn's Lemma* we can conclude that L possesses a maximal element; let us call it I .

As I is an ideal and R a ring we have a canonical Ringhomomorphism $\phi : R \rightarrow R/I$ that provides a Ring structure on R/I . Now let $J \subseteq R/I$ be a non-trivial ideal. Then $\phi^{-1}(J) \supset I$. As I is maximal in L we have that $\phi^{-1}(J)$ is either finitely generated or equal to I . However, $\phi^{-1}(J) = I$ is not possible as then $J = (0)$. Hence, $\phi^{-1}(J)$ must be f.g. and therefore also J is. As every ideal of R/I is finitely generated we have that R/I is noetherian. \square

b) Claim: There are f.g. ideals J_1 and J_2 such that $J_1 J_2 \supseteq I$ and $J_1 J_2 \subseteq I$.

Proof. We know that I is not a prime ideal as any prime ideal in R is finitely generated. Hence, we can find $x, y \in R$ such that $x, y \notin I$ but $xy \in I$. Let $\mathfrak{J}_x := (\phi(x))_{R/I}$ denote the R/I -ideal generated by $\phi(x)$. We define $J_1 := \phi^{-1}(\mathfrak{J}_x)$ and $J_2 := \phi^{-1}(\mathfrak{J}_y)$. As $x, y \notin I$ we have that $J_1, J_2 \supseteq I$.

Claim(1): $\mathfrak{J}_x \mathfrak{J}_y = (\phi(xy))_{R/I}$

proof.

$$\mathfrak{J}_x \mathfrak{J}_y = \left\{ \sum_{i \leq n} a_i b_i \mid a_i \in \mathfrak{J}_x \wedge b_i \in \mathfrak{J}_y \right\} \quad (24)$$

$$= \left\{ \sum_{i \leq n} (r_i \phi(x))(s_i \phi(y)) \mid r_i, s_i \in R/I \right\} \quad (25)$$

$$= \left\{ \phi(x) \phi(y) \sum_{i \leq n} r_i s_i \mid r_i, s_i \in R/I \right\} \quad (26)$$

$$= \{ \phi(xy) r \mid r \in R/I \} \quad (27)$$

$$= (\phi(xy))_{R/I} \quad (28)$$

qed(1)

Claim(2): Let $\phi : R_1 \rightarrow R_2$ be a ringhomomorphism and I_1, I_2 ideals of R_2 . Then $\phi^{-1}(I_1 I_2) \supseteq \phi^{-1}(I_2) \phi^{-1}(I_1)$.

proof. Let $x \in \phi^{-1}(I_1) \phi^{-1}(I_2)$ such that $x = ab$ with $a \in \phi^{-1}(I_1)$ and $b \in \phi^{-1}(I_2)$. Hence, $\phi(x) = \phi(ab) = \phi(a) \phi(b) \in I_1 I_2$. Therefore, $x \in \phi^{-1}(I_1 I_2)$.

qed(2)

As we have $xy \in I$ we can now conclude

$$I = \phi^{-1}((0)_{R/I}) \supseteq \phi^{-1}((\phi(xy))_{R/I}) = \phi^{-1}(\mathfrak{J}_x \mathfrak{J}_y) \quad (29)$$

$$\supseteq \phi^{-1}(\mathfrak{J}_x) \phi^{-1}(\mathfrak{J}_y) \quad (30)$$

$$= J_1 J_2 \quad (31)$$

Clearly $J_1 J_2 = (\{\sum_{i \leq n} a_i b_i \mid a_i \in J_1 \wedge b_i \in J_2\})_R$ is finitely generated as J_1 and J_2 are. \square

c)

Remark. Let R be a ring and I an ideal. Then I can be naturally viewed as a R -module by simply interpreting the ring multiplication as module product. Indeed, as ideals are abelian groups in $+$ and closed under multiplication they satisfy the R module axioms.

Recall that for ideals I, J of some ring R with $J \supset I$ we have that J/I is an ideal of R/I by the *third isomorphism theorem for rings*⁷. Hence, J/I always forms an abelian group with the induced binary operation $+_{R/I}$.

Claim: $I/J_1 J_2$ is a finitely generated R/I -submodule of $J_1/J_1 J_2$.

Proof. It suffices to prove that $J_1/J_1 J_2$ is a f.g. R/I module. Indeed, as R/I is noetherian we can deduce by a previous exercise⁸ that $I/J_1 J_2 \subset J_1/J_1 J_2$ is f.g..

We define an action on $J_1/J_1 J_2$

$$\star : R/I \times J_1/J_1 J_2 \rightarrow J_1/J_1 J_2 \quad (32)$$

$$[r]_I \star [j]_{J_1 J_2} := [r \cdot j]_{J_1 J_2} \quad (33)$$

⁷DF § 7.3 Theorem 8 (2)

⁸DF §15.1 Exercise 8

where "·" denotes the ring-multiplication in R .

Claim(1): \star is well defined.

proof. We have for $i \in I$ and $l \in J_1 J_2$

$$[r + i]_I \star [j + l]_{J_1 J_2} = [(r + i) \cdot (j + l)]_{J_1 J_2} \quad (34)$$

$$= [r \cdot j + r \cdot l + i \cdot j + i \cdot l]_{J_1 J_2} \quad (35)$$

$$= [r \cdot j]_{J_1 J_2}. \quad (36)$$

The last equality follows from the fact that $r \cdot l, i \cdot j, i \cdot l \in J_1 J_2$ qed(1)

Claim(2): $J_1/J_1 J_2$ is a R/I -module

proof.

1. As J_1 and $J_1 J_2$ are ideals in R we have that $(J_1/J_1 J_2, +_{R/I})$ is abelian.
2. for $[r]_I, [s]_I \in R/I$ and $j \in J_1/J_1 J_2$ we have

$$([r]_I + [s]_I) \star [j]_{J_1 J_2} = [r + s]_I \star [j]_{J_1 J_2} \quad (37)$$

$$= [(r + s) \cdot j]_{J_1 J_2} \quad (38)$$

$$= [r \cdot j]_{J_1 J_2} + [s \cdot j]_{J_1 J_2} \quad (39)$$

$$= [r]_I \star [j]_{J_1 J_2} + [s]_I \star [j]_{J_1 J_2} \quad (40)$$

3. for $[r]_I, [s]_I \in R/I$ and $j \in J_1/J_1 J_2$ we have

$$([r]_I \cdot_{R/I} [s]_I) \star [j]_{J_1 J_2} = [r \cdot s]_I \star [j]_{J_1 J_2} \quad (41)$$

$$= [(r \cdot s) \cdot j]_{J_1 J_2} \quad (42)$$

$$= [r \cdot (s \cdot j)]_{J_1 J_2} \quad (43)$$

$$= [r]_I \star [s \cdot j]_{J_1 J_2} \quad (44)$$

$$= [r]_I \star ([s]_I \star [j]_{J_1 J_2}) \quad (45)$$

4. for $[r]_I \in R/I$ and $j, l \in J_1/J_1 J_2$ we have

$$[r]_I \star ([j]_{J_1 J_2} + [l]_{J_1 J_2}) = [r]_I \star [j + l]_{J_1 J_2} \quad (46)$$

$$= [(r) \cdot (j + l)]_{J_1 J_2} \quad (47)$$

$$= [r \cdot j]_{J_1 J_2} + [r \cdot l]_{J_1 J_2} \quad (48)$$

$$= [r]_I \star [j]_{J_1 J_2} + [r]_I \star [l]_{J_1 J_2} \quad (49)$$

5. for $[j]_{J_1 J_2} \in J_1/J_1 J_2$

$$[1]_I \star [j]_{J_1 J_2} = [1 \cdot j]_{J_1 J_2} = [j]_{J_1 J_2}. \quad (50)$$

qed(2)

Claim(3): $J_1/J_1 J_2$ is finitely generated as R/I module.

proof. Let $\{g_1, \dots, g_n\}$ be a set of generators for J_1 and $[j]_{J_1 J_2} \in J_1/J_1 J_2$ be arbitrary. It follows that

$$[j]_{J_1 J_2} = \left[\sum_{i=1}^n r_i \cdot g_i \right]_{J_1 J_2} \quad (51)$$

$$= \sum_{i=1}^n [r_i \cdot g_i]_{J_1 J_2} \quad (52)$$

$$= \sum_{i=1}^n [r_i]_I \star [g_i]_{J_1 J_2}. \quad (53)$$

Hence, $\{[g_1]_{J_1 J_2}, \dots, [g_n]_{J_1 J_2}\}$ is a generating set for $J_1/J_1 J_2$. Therefore, $J_1/J_1 J_2$ is finitely generated. qed(3)

By the preliminary remarks we can conclude the claim □

d) Claim: (c) implies that I is finitely generated over R .

Remark. Recall that any R/I -module M with action $\cdot : R/I \times M \rightarrow M$ can be transformed into a R -module in a canonical way; use the projection $\phi : R \rightarrow R/I$ to define the action

$$\cdot_R : R \times M \rightarrow M \quad (54)$$

$$r \cdot_R m := \phi(r) \cdot m \quad (55)$$

Clearly if M is f.g. as R/I module it is also f.g. as R module. In fact the same elements that generate M as R/I -module will also do it as R -module.

Lemma 0.1. *Let M be a R -module and $N \subset M$ be an R -submodule of M . If N and M/N are finitely generated then so is M .*

Proof. Observe that we have the exact sequence in fig. 2.

$$0 \longrightarrow N \xrightarrow{\iota} M \xrightarrow{\phi} M/N \longrightarrow 0$$

Figure 2: Exact sequence

Let $\{g_1, \dots, g_n\}$ be the generators for N and $\{[h_1]_N, \dots, [h_m]_N\}$ for M/N . We can use the projection map $\phi : M \rightarrow M/N$ to get a set $\{\tilde{h}_1, \dots, \tilde{h}_m\} \subset M$ such that $\phi(\tilde{h}_1) = [h_1]_N$.

Now take $x \in M$ arbitrary, then $\phi(x) = [x]_N = \sum_{i \leq m} r_i [h_i]_N$. Define $x' := \sum_{i \leq m} r_i \tilde{h}_i \in M$, then clearly $x - x' \in \ker(\phi) = \text{im}(\iota)$. Hence, we can write $x - x' = \sum_{j \leq n} r_j \iota(g_j)$. All together we have $x = x' + x - x' = \sum_{i \leq m} r_i \tilde{h}_i + \sum_{j \leq n} r_j \iota(g_j)$ and as x was arbitrary we can conclude that M is generated by $\{\tilde{h}_1, \dots, \tilde{h}_m, \iota(g_1), \dots, \iota(g_n)\}$. □

We can now prove that I is finitely generated over R .

proof of claim. We have that I is a R -module and J_1J_2 is a R -submodule of I . As both J_1J_2 and I/J_1J_2 are finitely generated as R -modules we can conclude by the previous lemma that I is finitely generated. \square

Assuming that R is a ring in which all prime ideals are finitely generated we have shown (in **(a)**) that if there was an ideal which is not finitely generated then we would also have an inclusion maximal ideal which is not finitely generated, which we named I . Now in **(d)** we deduced that such an I would in fact be finitely generated(∇). Hence, we must reject the premise of **(a)** and end up with

The collections of ideals of R that are not finitely generated is empty. (56)

For a ring this is equivalent to being noetherian. \square