Book 1 — Lecture 1 — Interlude 1 — Exercise 1

Plotting Four Introductory Functions

Prompt. Using a graphing calculator or a program like Mathematica, plot each of the following functions:

$$\begin{aligned} f(t) &= t^4 + 3t^3 - 12t^2 + t - 6, \\ g(x) &= \sin(x) - \cos(x), \\ \theta(\alpha) &= e^\alpha + \alpha \ln(\alpha), \\ x(t) &= \sin^2(t) - \cos(t). \end{aligned}$$

Notes Before Plotting

- $\theta(\alpha)$ is only defined for $\alpha > 0$, because $\ln(\alpha)$ needs a positive argument.
- $f(t)$ is a quartic polynomial, so it grows to $+\infty$ for large positive and negative t.
- $g(x)$ and $x(t)$ are bounded oscillatory functions built from sine/cosine terms.

Example Plotting Windows

- $t \in [-4, 4]$ for $f(t)$
- $x \in [-2\pi, 2\pi]$ for $g(x)$
- $\alpha \in [0.05, 4]$ for $\theta(\alpha)$ (avoid $\alpha = 0$)
- $t \in [-2\pi, 2\pi]$ for $x(t)$

These are starting points; zooming in/out helps identify turning points and global behavior.

Result. A useful first pass is to use windows $t \in [-4, 4]$, $x \in [-2\pi, 2\pi]$, and $\alpha \in [0.05, 4]$.

Example. For $\theta(\alpha) = e^\alpha + \alpha \ln(\alpha)$, the logarithm term enforces $\alpha > 0$, so avoid $\alpha = 0$ in plotted ranges.

Static Plot Figure