

Internal absoluteness

Philipp Schlicht, University of Bristol

23 September 2019



Partially funded by the EU's Horizon 2020 programme, grant No 794020

Overview

- ▶ We study *internal absoluteness* (IA) between $M[g]$ and V , where $M \prec H_\theta$ and $g \in V$ is generic over M .
- ▶ Here we only consider projective absoluteness.
- ▶ The story is that these or similar principles were used in proofs of Steel and Woodin (see Steel 2008) for the tree production lemma
- ▶ **Our motivation:**
We re-discovered such principles for an application (2018) to selectors for ideals. and a connection with universally Baire sets.
- ▶ **Our aims:**
Fix a class of forcings.
 - ▶ (How) can IA be characterized via descriptive set theory?
 - ▶ How is IA related to other notions of absoluteness?
 - ▶ What is the consistency strength of IA?
- ▶ This is recent work in progress with Sandra Müller.

Forcing \leftrightarrow descriptive set theory

Zapletal (2008): TFAE for forcings of the form $\mathbb{P} = \text{Borel}/I$, where I is a σ -ideal

<i>Forcing</i>	<i>Descriptive set theory</i>
\mathbb{P} is proper	For every countable transitive $M \prec H_\theta$ and every $A \in \mathbb{P} \cap M$, the set of \mathbb{P} -generic reals over M in A is I -positive
\mathbb{P} is ω^ω - bounding	Every Borel I -positive set contains a compact I -positive subset, and continuous reading of names

The second equivalence assumes that \mathbb{P} is proper. Zapletal proved a large number of such characterisations.

Forcing \leftrightarrow descriptive set theory

Ikegami (2010): TFAE for proper tree forcings \mathbb{P}

<i>Forcing</i>	<i>Descriptive set theory</i>
1-step Σ_3^1 - \mathbb{P} -absoluteness	\mathbb{P} -measurability of Δ_2^1 sets
1-step Σ_4^1 - \mathbb{P} -absoluteness	\mathbb{P} -measurability of Δ_3^1 sets

Restrictions:

1. The second equivalence assumes that **every real has a sharp**, but there's **no inner model with a Woodin cardinal**.
2. There's **no full equivalence** in ZFC: projective 1-step Cohen absoluteness doesn't imply that all projective sets have the Baire property.

Forcing classes

We consider the classes:

1. All forcings
2. **Proper** forcings
3. Simply definable proper forcings on the reals: forcings of the form *Borel*/ I for σ -ideals I ; tree forcings
4. Simply definable **ccc** forcings on the reals

1-step and 2-step absoluteness

Suppose that \mathcal{F} is a class of forcings and Λ a class of formulas $\varphi(., x)$ (where x ranges over certain parameters).

Definition (1-step absoluteness)

1-step \mathcal{F} - Λ -absoluteness means: if $\mathbb{P} \in \mathcal{F}$ and G is \mathbb{P} -generic over V , then $V \prec_{\Lambda} V[G]$ holds.

Definition (2-step absoluteness)

2-step \mathcal{F} - Λ -absoluteness means: If $\mathbb{P} \in \mathcal{F}$ and G, H are \mathbb{P} -generic filters over V with $V[G] \subseteq V[H]$, then $V[G] \prec_{\Lambda} V[H]$ holds.

If \mathcal{F} contains the trivial forcing, then 2-step implies 1-step absoluteness.

Why 2-step absoluteness?

Consider the class of all forcings.

Theorem (Martin, Solovay, Woodin)

2-step Σ_3^1 -absoluteness $\iff \forall X \text{ } X^\# \text{ exists}$, where X denotes sets of ordinals.

Theorem (Feng, Magidor, Woodin 1992)

1-step Σ_3^1 -absoluteness is equiconsistent with the existence of a regular cardinal κ with $H_\kappa \prec_{\Sigma_2} V$ (weaker than a Mahlo cardinal).

Here 2-step-absoluteness is strictly stronger.

This is not always the case:

Theorem (Steel, Woodin)

The first-order theory $\text{Th}(L(\mathbb{R}))$ cannot be changed by forcing $\iff \text{AD}^{L(\mathbb{R})}$ holds in all generic extensions, assuming Ord is measurable in an outer model.

The proof shows that 1-step and 2-step absoluteness are equivalent for formulas of the form $\varphi^{L(\mathbb{R})}$ with real parameters.

Internal absoluteness

Definition (Internal absoluteness)

Let \mathcal{F} be a class of forcings, Λ a class of formulas and θ an uncountable regular cardinal.

1. For any countable $M \prec H_\theta$, $\mathbf{IA}_{\mathcal{F},\Lambda}^M$ denotes the statement:


$$M[g] \prec_\Lambda H_\theta$$

holds for all $\mathbb{P} \in \mathcal{F} \cap \text{Def}(M)$ and all \mathbb{P} -generic filters $g \in V$ over M .

2. $\mathbf{IA}_{\mathcal{F},\Lambda}^\theta : \Longleftrightarrow \mathbf{IA}_{\mathcal{F},\Lambda}^M$ holds for club many $M \in [H_\theta]^\omega$.

We write \mathbf{IA} if Λ is the set of projective formulas with real parameters, \mathcal{F} is the class of all forcings, and $\mathbf{IA}_{\mathcal{F},\Lambda}^\theta$ holds for **all regular** θ with H_θ sufficiently elementary in V .

\mathbf{IA} implies 1-step and 2-step absoluteness:



Separating internal from 1-step and 2-step absoluteness

Fact

$\text{IA}_{\mathbb{C}}$ implies the *Baire property* for all projective sets.

Proof.

Take a countable $M \prec H_{\theta}$. Note that the set of Cohen reals x over M is comeager.

Let $\varphi(x)$ define a projective set.

The set of Cohen conditions p with $p \Vdash_{\mathbb{C}}^M \varphi(\dot{x})$, where \dot{x} is a name for the Cohen real, defines an open set witnessing the Baire property. \square

Projective *1-step and 2-step* absoluteness hold in $L^{Add(\omega, \omega_1)}$.

But not all projective sets have the Baire property in this model, so internal Cohen absoluteness $\text{IA}_{\mathbb{C}}$ fails there.

Uniformization up to meager

We now consider Cohen forcing.

Proposition (Müller, S.)

TFAE:

1. $\text{IA}_{\mathbb{C}}$
2. *Every projective relation can be uniformized on a **comeager set**.*

Shelah (1984) proved the consistency of 2. relative to ZFC.

Proof sketch.

1 \Rightarrow 2: Assume $\text{IA}_{\mathbb{C}}$. Let $R = \{(x, y) \mid \varphi(x, y, z)\}$.

Let $M \prec H_\theta$ with $z \in M$ and $\sigma \in M$ be a name for the Cohen real (all names are nice).

Let $\tau \in M$ with $1 \Vdash \exists y \varphi(\sigma, y, z) \Rightarrow \varphi(\sigma, \tau, z)$.

$\text{IA}_{\mathbb{C}}$ implies $M[x] \models \varphi(x, \tau^x, z) \iff V \models \varphi(x, \tau^x, z)$ for Cohen reals $x \in V$ over M .

The continuous function $x \mapsto \tau^x$ uniformizes R on the set of Cohen reals over M . \square

Uniformization up to meager

Proposition (Müller, S.)

TFAE:

1. IA_C
2. Every projective relation can be uniformized on a *comeager set*.

Proof sketch.

2 \Rightarrow 1: Take a Σ_{n+1}^1 -formula $\exists y \varphi(x, y)$, where $n \geq 1$.

For a nice name σ and any $i \in \omega$, let $A_\sigma = \{(x, y) \mid \varphi(\sigma^x, y)\}$.

Let $M \prec H_\theta$.

Claim: For any Cohen real $x \in V$ over M , $\exists y \varphi(\sigma^x, y)$ is downwards absolute from V to $M[x]$.

To see this, let f_σ be a uniformization of A_σ up to meager. Since the existence of f_σ is projective, we can take f_σ to be projectively defined in M .

Suppose that $\exists y \varphi(u, y)$ holds in V .

Recall that x is Cohen generic over M iff $x \in A$ for every comeager Borel set with a code in M . So $x \in \text{dom}(f_\sigma)$ and $f_\sigma(x) \in M[x]$. □

Forcings of the form $Borel/I$

Let I be a σ -ideal on a Polish space X . A binary relation R on X is called *total* if $\text{proj}(R) = X$.

Definition (Uniformization and regularity)

1. $U_I : \iff$ If R is a projective total relation on X , then for any Borel set $A \notin I$, $R \upharpoonright A$ has a **projective subfunction** with domain $B \notin I$.
2. $R_I : \iff$ If A is a projective set and $B \notin I$ is a Borel set, then there is some Borel set $C \notin I$ with $C \subseteq B$ and either $C \cap A = \emptyset$ or $C \subseteq A$.

Clearly $U_I \implies R_I$.

Why don't we uniformize everywhere except for a set in I ? This would be much stronger.

Forcings of the form $Borel/I$

Theorem (Müller, S.)

The following statements are equivalent for proper forcings of the form $\mathbb{P} = Borel/I$:

1. $\mathbf{IA}_{\mathbb{P}}$
2. 1-step \mathbb{P} -absoluteness holds and \mathbf{R}_I holds for all projective sets.
3. \mathbf{U}_I

<i>Forcing</i>	<i>Descriptive set theory</i>
$\mathbf{IA}_{\mathbb{P}}$	\mathbf{U}_I for projective sets
1-step \mathbb{P} -absoluteness	\mathbf{R}_I for projective sets

$\not\Rightarrow$

$\Leftarrow ?$

We have seen $1. \Leftrightarrow 3.$ for Cohen forcing. $1. \Rightarrow 2.$ is clear.

Consequences of internal Cohen absoluteness

Fact (Müller, S.)

$\text{IA}_{\mathbb{C}}$ implies:

1. 1-step and 2-step Cohen *absoluteness*
2. The *Baire property* for projective sets
3. For every real x , the set of Cohen reals over $L[x]$ is comeager
4. For every real x , there is a real dominating $L[x]$
5. Projective *uniformization up to meager*

Obtaining internal absoluteness

$\mathsf{IA}_{\mathbb{C}}$ is consistent relative to ZFC, by the consistency of projective uniformization up to meager (Shelah 1984).

In many cases, $\mathsf{IA}_{\mathbb{P}}$ follows directly from large cardinals:

Proposition (Müller, S.)

PD implies that any sufficiently absolute Axiom A forcing of the form $\mathbb{P} = \text{Borel}/I$ satisfies $\mathsf{IA}_{\mathbb{P}}$.

This is based on work of Fabiana Castiblanco.

The connection with universally Baire sets

Let $A \subseteq \omega^\omega$.

Definition (Feng, Magidor, Woodin)

A is *universally Baire* if for every continuous $f: X \rightarrow \omega^\omega$, $f^{-1}(A)$ has the Baire property.

A is *absolutely complemented* if there are trees S, T with $A = p[S]$ such that

$$\omega^\omega = p[S] \cup p[T]$$

holds in all generic extensions.

Feng, Magidor and Woodin proved the equivalence of these notions:

<i>Forcing</i>	<i>Descriptive set theory</i>
A is absolutely complemented	A is universally Baire

The connection with universally Baire sets

Definition

We call a tree T on $\omega \times \lambda$ *absolute* if for all regular θ such that H_θ is sufficiently elementary in V , there is a club of $M \in [H_\theta]^\omega$ such that for all generic extensions $M[g] \subseteq V$:

$$p[T]^{M[g]} = p[T] \cap M[g].$$

Definition

We call a formula $\varphi(x)$ *\mathbb{P} -treeable* if there is an absolute tree T on $\omega \times \lambda$, for some λ , such that

$$p[T] = \{x \in {}^\omega\omega \mid \varphi(x)\}$$

holds in every \mathbb{P} -generic extension.

Moreover, *treeable* means \mathbb{P} -treeable for all forcings \mathbb{P} .

Fact

Every treeable formula is internally absolute.

From internally absolute to treeable

Proposition (essentially Steel or Woodin)

If $\varphi(x)$ is internally absolute, then $\varphi(x)$ is treeable.

Proof sketch.

The tree for $\varphi(x)$ searches for

1. A countable transitive model M of ZFC^-
2. An elementary $j: M \rightarrow H_\theta$
3. A generic filter g over M with $x \in M[g]$ and $M[g] \models \varphi(x)$


□

Internal projective absoluteness \iff Projective treeability

Internally absolute and universally Baire

The following are equivalent for a formula $\varphi(x)$:

1. φ is internally absolute
2. $\{x \mid \varphi(x)\}$ is universally Baire, with a tree projecting to $\{x \mid \varphi(x)\}$ in all generic extensions.



Absoluteness from absorption

How does one prove internal absoluteness?

Next is a well-known property that is used in several proofs of absoluteness.

Definition

Let M be an inner model and κ a cardinal. (M, V) has the *$<\kappa$ -absorption* property for a forcing \mathbb{P} if:

For any \mathbb{P} -generic extension $V[H]$ of V and any real $x \in V[H]$, there exist:

1. a forcing $\mathbb{Q} \in M$ with $|\mathbb{Q}|^M < \kappa$ and
2. a \mathbb{Q} -generic filter $I \in V[H]$ over M

with $x \in M[I]$.

For instance, Schindler (2001) used this property to obtain $L(\mathbb{R})$ -absoluteness for proper forcings from a remarkable cardinal.

Internal absoluteness from absorption

Lemma (Müller, S.)

Suppose that κ is inaccessible and G is $\text{Col}(\omega, <\kappa)$ -generic for V . Suppose that $(V, V[G])$ has the $<\kappa$ -absorption property for \mathbb{P} .

Then every $L(\mathbb{R})$ -formula with real and ordinal parameters is \mathbb{P} -treeable.

The proof combines previous arguments for absoluteness with the construction of a tree projecting to $\{x \mid \varphi(x)\}$.

Thus one can, in many cases, prove internal absoluteness in a similar way as 1-step absoluteness.

Some open questions

For Cohen forcing, our next goals are:

Question

- ▶ Does IA_C fail in Shelah's "first" model of the Baire property for all projective sets?
- ▶ For Cohen forcing, does 2-step imply 1-step projective absoluteness?

Similar questions appear for forcings of the form Borel/I .

For the class of all forcings, the following are long-standing open questions:

Question (Feng, Magidor, Woodin, Wilson)

- ▶ If every projective set is universally Baire, does *internal* projective absoluteness hold?
- ▶ If every projective set is universally Baire, does *1-step* projective absoluteness hold? Does the converse implication hold?

Literature



Qi Feng, Menachem Magidor, and Hugh Woodin. "Universally Baire sets of reals." Set theory of the continuum. Springer, New York, NY, 1992. 203-242.



Saharon Shelah. "Can you take Solovay's inaccessible away?." Israel Journal of mathematics 48.1 (1984): 1-47.



John Steel. The derived model theorem, 2008



Jindrich Zapletal. Forcing idealized. Vol. 174. Cambridge: Cambridge University Press, 2008.



Daisuke Ikegami. "Forcing absoluteness and regularity properties." Annals of Pure and Applied Logic 161.7 (2010): 879-894.



Sandra Müller, Philipp Schlicht. Internal absoluteness, in preparation

Thank you for listening!