

## OPEN PROBLEMS ON GENERALISED BAIRE SPACES 2020

The *Fifth Workshop on Generalised Baire Spaces* took place at the School of Mathematics of the University of Bristol on 3-4 February, 2020, see <https://philippschlicht.github.io/meetings/generalizedbairespaces2020>. This is an incomplete list of some open problems that were discussed at the workshop.<sup>1</sup>

### 1. GENERALIZED BAIRE SPACES

**1.1. Claudio Agostini: Winning tactics.** A *tactic* in a game of length  $\omega$  is a strategy that depends only on the previous move of the opponent. For separable regular Hausdorff spaces, the following conditions are equivalent:

- (1) There is a compatible complete metric.
- (2) There is a winning tactic for II in the strong Choquet game.
- (3) There is a winning strategy for II in the strong Choquet game.

Notice that tactics do not make sense for longer games, since moves at limit times don't have a predecessor. But one can define the following natural variant for the strong Choquet game  $G_\kappa(X)$  of length  $\kappa$  for  $X$ . A tactic for II in  $G_\kappa(X)$  is defined as a strategy with the following properties:

- (1) For successor times, the reply of II depends only on the preceding move.
- (2) For a play of limit length consisting of  $\vec{U} = \langle U_i \mid i < \alpha \rangle$  and  $\vec{x} = \langle x_i \mid i < \alpha \rangle$ , the reply of II depends only on  $\bigcap_{i < \alpha} U_i$  and (possibly)  $\alpha$ .

**Question 1.1.** Suppose that  $X$  is a regular Hausdorff space of weight  $\leq \kappa$ . Does the existence of a winning strategy for player II in  $G_\kappa(X)$  imply the existence of a winning tactic?

**1.2. Dorottya Sziraki: The open colouring axiom.** The *open colouring axiom* for  $X$ ,  $\text{OCA}_\kappa(X)$ , states that any open graph  $G$  on  $X$  either has a  $\kappa$ -coloring or else contains a complete subgraph of size  $\kappa^+$ .  $\text{OCA}_\kappa$  states that  $\text{OCA}_\kappa(X)$  holds for all  $X \subseteq {}^\kappa \kappa$ .

**Question 1.2.** Is  $\text{OCA}_\kappa$  consistent with ZFC?

The *open graph dichotomy*  $\text{OGD}_\kappa(X)$  for  $X$  states that any open graph on  $X$  either has a  $\kappa$ -colouring or else contains a  $\kappa$ -perfect complete subgraph. The *perfect set property*  $\text{PSP}_\kappa(X)$  for  $X \subseteq {}^\kappa \kappa$  states that  $|X| \leq \kappa$  or  $X$  has a  $\kappa$ -perfect subsets.

**Question 1.3.** Does  $\text{PSP}_\kappa(\text{closed})$  imply  $\text{OGD}_\kappa(\text{closed})$ ?

**Fact.**  $\text{OGD}_\kappa(\text{closed}) \Rightarrow \text{OGD}_\kappa(\Sigma_1^1(\kappa)) \Rightarrow \text{PSP}_\kappa(\Sigma_1^1(\kappa))$ .

**Fact.** After forcing with  $\text{Col}(\kappa, < \lambda)$ , where  $\lambda > \kappa$  is inaccessible,  $\text{OGD}_\kappa$  holds for all sets definable from  $\kappa$ -sequences of ordinals.

**Question 1.4.** Can we force  $\text{PSP}_\kappa$  with some forcing other than  $\text{Col}(\kappa, < \lambda)$ ?

---

<sup>1</sup>If you have an answer, comment or further question, please email Philipp Schlicht at philipp.schlicht@bristol.ac.uk.

## 2. CONNECTIONS WITH MODEL THEORY

## 2.1. Jan Dobrowolski: Polish groups.

**Fact.** Suppose that  $X$  is Polish,  $E$  is an equivalence relation on  $X$  such that

- (1) the equivalence class  $[x]_E$  of any  $x \in X$  is closed and
- (2) the saturation  $[U]_E$  of any open subset  $U$  of  $X$  is Borel.

Then  $X/E$ , with the Effros Borel structure, is standard Borel, and the natural projection  $\pi: X \rightarrow X/E$  has a Borel measurable right inverse  $f$  with  $\pi \circ f = \text{id}$ .

In particular, this holds for any Polish group  $H$  and the equivalence relation on  $H$  induced by a closed subgroup  $G \leq H$ .

**Question 2.1.** Does the analogue of the previous statement about Polish groups hold for  $\kappa^\kappa$ ?

The product of two closed subgroups  $G, H$  of  $\text{Sym}(\omega)$  has the property of Baire.

**Question 2.2.** Does the analogue of the previous statement hold for  $\text{Sym}(\kappa)$  with the bounded topology?

## 2.2. Rosario Mennuni: Omitting types.

**Disclaimer:** The answer to this question might be easy or already known.

The most basic version of the Omitting Type Theorem states:

**Theorem.** Let  $T$  be a consistent first-order theory in a countable language,  $x = (x_0, \dots, x_{n-1})$  an  $n$ -tuple of variables, and  $\pi(x)$  a partial  $n$ -type over  $\emptyset$  such that for no formula  $\phi(x)$  consistent with  $T$  we have  $T \cup \{\phi(x)\} \vdash \pi(x)$ . Then there is a countable  $M \models T$  such that no element of  $M^n$  realises  $\pi(x)$ .

A better version states (see [2, Theorem 10.3]):

**Theorem.** Let  $T$  be a consistent first-order theory in a countable language, and for every  $n \in \omega$  let  $A_n$  be a meagre subset of the space  $S_n(T)$  of  $n$ -types over  $\emptyset$ . Then there is a countable  $M \models T$  such that, for every  $n \in \omega$  and  $p(x) \in A_n$ , no element of  $M^n$  realises  $p(x)$ .

The basic version can be generalised to what is called the  $\kappa$ -omitting types theorem (see [1, Theorem 2.2.19]).

**Theorem.** Let  $T$  be a consistent first-order theory in a language of size  $\kappa$ ,  $x = (x_0, \dots, x_{n-1})$  an  $n$ -tuple of variables, and  $\pi(x)$  a partial  $n$ -type over  $\emptyset$  such that for no set of formulas  $\Phi(x)$  of size  $< \kappa$  consistent with  $T$  we have  $T \cup \Phi(x) \vdash \pi(x)$ . Then there is  $M \models T$  with  $|M| \leq \kappa$  and such that no element of  $M^n$  realises  $\pi(x)$ .

**Question 2.3.** Let  $L$  be a language of size  $\kappa$ , and let  $T$  be a consistent  $L$ -theory. Equip each  $S_n(T)$  with the topology generated by declaring partial types of size  $< \kappa$  to induce open sets, and replace “meagre” with “ $\kappa$ -meagre”. For which cardinals  $\kappa$  does Theorem 2.2 generalise in the fashion of Theorem 2.2?

## REFERENCES

- [1] C.C. CHANG, H.J. KEISLER. *Model Theory*, Third edition, Studies in Logic and the Foundations of Mathematics 73. North-Holland Publishing Co., Amsterdam, (1990).
- [2] B. POIZAT. *A Course in Model Theory*. Universitext. Springer (2000).

3. COMBINATORICS OF  $\kappa^\kappa$ 3.1. **Adrian Mathias, Vera Fischer.**

**Question 3.1.** Do analogues to some classical results for mad families on  $\omega$  hold for mad families on  $\kappa$ ?

3.2. **Johannes Schürz.**

**Question 3.2.** Is there a forcing that adds a  $\kappa$ -dominating real, but no  $\kappa$ -Cohen real?

3.3. **Sarka Stejskalova.**

**Question 3.3.** Is  $\mathfrak{u}_\kappa < 2^\kappa$  consistent for the successor  $\kappa$  of a regular cardinal?