

Uniformization results and Feldman-Moore Theorem in generalized descriptive set theory

Luca Motto Ros

Department of Mathematics “G. Peano”
University of Turin, Italy
`luca.mottoros@unito.it`
<https://sites.google.com/site/lucamottoros/>


Workshop “Generalised Baire Space and Large Cardinals”
Bristol, February 8–10, 2024

Uniformization theorems

Given $P \subseteq X \times Y$, a **uniformization** of P is a subset $P^* \subseteq P$ such that for all $x \in X$

$$\exists y P(x, y) \iff \exists! y P^*(x, y).$$

Equivalently, P^* is the graph of a function f (called **uniformizing function**) with domain $\text{proj}_X(P)$ such that $f(x) \in P_x$ for every $x \in A$.



Fact

Let X, Y be standard Borel spaces. A set $P \subseteq X \times Y$ has a Borel uniformization if and only if $\text{proj}_X(P)$ is Borel and there is a Borel uniformizing function $f: \text{proj}_X(P) \rightarrow Y$ for P .

Fact

Let X, Y be standard Borel spaces. A set $P \subseteq X \times Y$ has a Borel uniformization if and only if $\text{proj}_X(P)$ is Borel and there is a Borel uniformizing function $f: \text{proj}_X(P) \rightarrow Y$ for P .

It is easy to show that there are Borel sets without Borel uniformizations.

General theme

Suppose that X, Y are Polish or standard Borel spaces, and that $P \subseteq X \times Y$ is Borel. Under which conditions there is a Borel uniformization of P ?

Fact

Let X, Y be standard Borel spaces. A set $P \subseteq X \times Y$ has a Borel uniformization if and only if $\text{proj}_X(P)$ is Borel and there is a Borel uniformizing function $f: \text{proj}_X(P) \rightarrow Y$ for P .

It is easy to show that there are Borel sets without Borel uniformizations.

General theme

Suppose that X, Y are Polish or standard Borel spaces, and that $P \subseteq X \times Y$ is Borel. Under which conditions there is a Borel uniformization of P ?

Today we are interested in “small section” uniformization results: if all the vertical sections of P are sufficiently small, then there is a Borel uniformization of P .

“Small section” uniformization results

Theorem (Lusin-Novikov)

Let X, Y be standard Borel spaces, and $P \subseteq X \times Y$ a Borel set with **countable** vertical sections. Then P has a Borel uniformization.

“Small section” uniformization results

Theorem (Lusin-Novikov)

Let X, Y be standard Borel spaces, and $P \subseteq X \times Y$ a Borel set with **countable** vertical sections. Then P has a Borel uniformization. Moreover, $P = \bigcup_{n \in \omega} P_n$ with each P_n a Borel set with vertical sections of size 1.

“Small section” uniformization results

Theorem (Lusin-Novikov)

Let X, Y be standard Borel spaces, and $P \subseteq X \times Y$ a Borel set with **countable** vertical sections. Then P has a Borel uniformization. Moreover, $P = \bigcup_{n \in \omega} P_n$ with each P_n a Borel set with vertical sections of size 1.

Theorem (???)

Let X be a standard Borel space, Y a Polish space, and $P \subseteq X \times Y$ a Borel set with **compact** vertical sections P_x . Then the map $x \mapsto P_x$ from X to $K(Y)$ (endowed with the Vietoris topology) is Borel.

“Small section” uniformization results

Theorem (Lusin-Novikov)

Let X, Y be standard Borel spaces, and $P \subseteq X \times Y$ a Borel set with **countable** vertical sections. Then P has a Borel uniformization. Moreover, $P = \bigcup_{n \in \omega} P_n$ with each P_n a Borel set with vertical sections of size 1.

Theorem (???)

Let X be a standard Borel space, Y a Polish space, and $P \subseteq X \times Y$ a Borel set with **compact** vertical sections P_x . Then the map $x \mapsto P_x$ from X to $K(Y)$ (endowed with the Vietoris topology) is Borel. Therefore P has a Borel uniformization.

“Small section” uniformization results

Theorem (Lusin-Novikov)

Let X, Y be standard Borel spaces, and $P \subseteq X \times Y$ a Borel set with **countable** vertical sections. Then P has a Borel uniformization. Moreover, $P = \bigcup_{n \in \omega} P_n$ with each P_n a Borel set with vertical sections of size 1.

Theorem (???)

Let X be a standard Borel space, Y a Polish space, and $P \subseteq X \times Y$ a Borel set with **compact** vertical sections P_x . Then the map $x \mapsto P_x$ from X to $K(Y)$ (endowed with the Vietoris topology) is Borel. Therefore P has a Borel uniformization.

Theorem (Arsenin-Kunugui)

Let X be a standard Borel space, Y a Polish space, and $P \subseteq X \times Y$ a Borel set whose vertical sections P_x are **σ -compact** (= countable unions of compact sets). Then P has a Borel uniformization.

Feldman-Moore Theorem

An equivalence relation E on a standard Borel space X is a **countable Borel equivalence relation** (CBER) if it is Borel as a subset of X^2 , and all its equivalence classes are countable.

Feldman-Moore Theorem

An equivalence relation E on a standard Borel space X is a **countable Borel equivalence relation** (CBER) if it is Borel as a subset of X^2 , and all its equivalence classes are countable.

Feldman-Moore Theorem

Let E be an equivalence relation on a standard Borel space X . Then E is a CBER if and only if it is the orbit equivalence relation induced by a Borel action of a countable (discrete) group G on X .

Equivalently: E is CBER if and only if there is a countable group G acting on X by Borel isomorphisms (in fact, involutions) which generates E .

Feldman-Moore Theorem

An equivalence relation E on a standard Borel space X is a **countable Borel equivalence relation** (CBER) if it is Borel as a subset of X^2 , and all its equivalence classes are countable.

Feldman-Moore Theorem

Let E be an equivalence relation on a standard Borel space X . Then E is a CBER if and only if it is the orbit equivalence relation induced by a Borel action of a countable (discrete) group G on X .

Equivalently: E is CBER if and only if there is a countable group G acting on X by Borel isomorphisms (in fact, involutions) which generates E .

The importance of this theorem in descriptive set theory
cannot be underestimated!

[Study of CBERs, Borel combinatorics, definable paradoxical decompositions, ...]

Hyperfinite equivalence relations

A CBER E is **hyperfinite** if it can be written as a countable increasing union of finite (= all classes are finite) Borel equivalence relations.

Some known facts:

Hyperfinite equivalence relations

A CBER E is **hyperfinite** if it can be written as a countable increasing union of finite (= all classes are finite) Borel equivalence relations.

Some known facts:

- Any CBER that admits a Borel transversal is hyperfinite.
[Easy application of the Feldman-Moore Theorem.]

Hyperfinite equivalence relations

A CBER E is **hyperfinite** if it can be written as a countable increasing union of finite (= all classes are finite) Borel equivalence relations.

Some known facts:

- Any CBER that admits a Borel transversal is hyperfinite.
[Easy application of the Feldman-Moore Theorem.]
- Every CBER on a Polish space is hyperfinite on a comeager set.

Hyperfinite equivalence relations

A CBER E is **hyperfinite** if it can be written as a countable increasing union of finite (= all classes are finite) Borel equivalence relations.

Some known facts:

- Any CBER that admits a Borel transversal is hyperfinite.
[Easy application of the Feldman-Moore Theorem.]
- Every CBER on a Polish space is hyperfinite on a comeager set.
- There are CBERs which are not hyperfinite.
[Consider e.g. the shift-action of \mathbb{F}_2 on $2^{\mathbb{F}_2}$. More generally, every non-amenable countable group admits a Borel action on a standard Borel space which induces a non-hyperfinite CBER.]

Hyperfinite equivalence relations

A CBER E is **hyperfinite** if it can be written as a countable increasing union of finite (= all classes are finite) Borel equivalence relations.

Some known facts:

- Any CBER that admits a Borel transversal is hyperfinite.
[Easy application of the Feldman-Moore Theorem.]
- Every CBER on a Polish space is hyperfinite on a comeager set.
- There are CBERs which are not hyperfinite.
[Consider e.g. the shift-action of \mathbb{F}_2 on $2^{\mathbb{F}_2}$. More generally, every non-amenable countable group admits a Borel action on a standard Borel space which induces a non-hyperfinite CBER.]

Weiss' conjecture

Any Borel action of a countable amenable group on a standard Borel space induces a hyperfinite CBER.

Moving to generalized descriptive set theory (GDST)...

Let κ be an uncountable cardinal such that $2^{<\kappa} = \kappa$.

Moving to generalized descriptive set theory (GDST)...

Let κ be an uncountable cardinal such that $2^{<\kappa} = \kappa$.

In all definitions, ω must be replaced by κ or $\text{cof}(\kappa)$:

Moving to generalized descriptive set theory (GDST)...

Let κ be an uncountable cardinal such that $2^{<\kappa} = \kappa$.

In all definitions, ω must be replaced by κ of $\text{cof}(\kappa)$:

Cantor space ${}^\omega 2$	\rightsquigarrow	generalized Cantor space ${}^\kappa 2$
Baire space ${}^\omega \omega$	\rightsquigarrow	generalized Baire space ${}^{\text{cof}(\kappa)} \kappa$
Borel	\rightsquigarrow	κ^+ -Borel

Moving to generalized descriptive set theory (GDST)...

Let κ be an uncountable cardinal such that $2^{<\kappa} = \kappa$.

In all definitions, ω must be replaced by κ of $\text{cof}(\kappa)$:

Cantor space ${}^\omega 2$	\rightsquigarrow	generalized Cantor space ${}^\kappa 2$
Baire space ${}^\omega \omega$	\rightsquigarrow	generalized Baire space ${}^{\text{cof}(\kappa)} \kappa$
Borel	\rightsquigarrow	κ^+ -Borel
countable	\rightsquigarrow	of size $\leq \kappa?$ or $\leq \text{cof}(\kappa)?$
finite	\rightsquigarrow	of size $< \kappa$ (i.e. " κ -small")? or $< \text{cof}(\kappa)?$
compact	\rightsquigarrow	κ -Lindelöf? or $\text{cof}(\kappa)$ -Lindelöf?

...and so on.

Moving to generalized descriptive set theory (GDST)...

Let κ be an uncountable cardinal such that $2^{<\kappa} = \kappa$.

In all definitions, ω must be replaced by κ of $\text{cof}(\kappa)$:

Cantor space ${}^\omega 2$	\rightsquigarrow	generalized Cantor space ${}^\kappa 2$
Baire space ${}^\omega \omega$	\rightsquigarrow	generalized Baire space ${}^{\text{cof}(\kappa)} \kappa$
Borel	\rightsquigarrow	κ^+ -Borel
countable	\rightsquigarrow	of size $\leq \kappa?$ or $\leq \text{cof}(\kappa)?$
finite	\rightsquigarrow	of size $< \kappa$ (i.e. “ κ -small”)? or $< \text{cof}(\kappa)?$
compact	\rightsquigarrow	κ -Lindelöf? or $\text{cof}(\kappa)$ -Lindelöf?

...and so on.

Can we have “small section” uniformization results and/or an analogue of the Feldman-Moore theorem in GDST?

Interludio: the regular case

Assume that $\kappa > \omega$ is regular.

Interludio: the regular case

Assume that $\kappa > \omega$ is regular.

Theorem (S.-D. Friedman-Hyttinen-Kulikov)

If E is an orbit equivalence relation on ${}^\kappa 2$ induced by a κ^+ -Borel action of a (discrete) group of size at most κ , then $E \leq_B^\kappa E_0^\kappa$, where E_0^κ is defined on 2^κ by $x E_0^\kappa y \iff \exists \alpha < \kappa \forall \beta \geq \alpha (x(\beta) = y(\beta))$.

Interludio: the regular case

Assume that $\kappa > \omega$ is regular.

Theorem (S.-D. Friedman-Hyttinen-Kulikov)

If E is an orbit equivalence relation on ${}^\kappa 2$ induced by a κ^+ -Borel action of a (discrete) group of size at most κ , then $E \leq_B^\kappa E_0^\kappa$, where E_0^κ is defined on 2^κ by $x E_0^\kappa y \iff \exists \alpha < \kappa \forall \beta \geq \alpha (x(\beta) = y(\beta))$.

In particular, if κ is inaccessible, then every orbit equivalence relation induced by a $\leq \kappa$ -sized discrete group is **hyper- κ -small**, i.e. it can be written as an increasing union of size κ of κ^+ -Borel equivalence relations which are κ -small (= all their classes have size $< \kappa$).

Interludio: the regular case

Assume that $\kappa > \omega$ is regular.

Theorem (S.-D. Friedman-Hyttinen-Kulikov)

If E is an orbit equivalence relation on ${}^\kappa 2$ induced by a κ^+ -Borel action of a (discrete) group of size at most κ , then $E \leq_B^\kappa E_0^\kappa$, where E_0^κ is defined on 2^κ by $x E_0^\kappa y \iff \exists \alpha < \kappa \forall \beta \geq \alpha (x(\beta) = y(\beta))$.

In particular, if κ is inaccessible, then every orbit equivalence relation induced by a $\leq \kappa$ -sized discrete group is **hyper- κ -small**, i.e. it can be written as an increasing union of size κ of κ^+ -Borel equivalence relation which are κ -small (= all their classes have size $< \kappa$). However:

Theorem (S.-D. Friedman-Hyttinen-Kulikov)

Assume $V = L$. Then there is a κ^+ -Borel equivalence relation E whose classes have size 2 which is not induced by a κ^+ -Borel action of a (discrete) group of size $\leq \kappa$.

Interludio: the regular case

One can also easily observe that no “small section” uniformization result can hold if we formulate it in terms of *uniformizing functions*.

Interludio: the regular case

One can also easily observe that no “small section” uniformization result can hold if we formulate it in terms of *uniformizing functions*.

Fact

There is a function $f: {}^\kappa 2 \rightarrow {}^\kappa 2$ whose graph $P \subseteq {}^\kappa 2 \times {}^\kappa 2$ is κ^+ -Borel, yet f is not κ^+ -Borel itself.

Interludio: the regular case

One can also easily observe that no “small section” uniformization result can hold if we formulate it in terms of *uniformizing functions*.

Fact

There is a function $f: {}^\kappa 2 \rightarrow {}^\kappa 2$ whose graph $P \subseteq {}^\kappa 2 \times {}^\kappa 2$ is κ^+ -Borel, yet f is not κ^+ -Borel itself.

Open problems (?)

Let $\kappa > \omega$ be regular and such that $2^{<\kappa} = \kappa$.

Interludio: the regular case

One can also easily observe that no “small section” uniformization result can hold if we formulate it in terms of *uniformizing functions*.

Fact

There is a function $f: {}^\kappa 2 \rightarrow {}^\kappa 2$ whose graph $P \subseteq {}^\kappa 2 \times {}^\kappa 2$ is κ^+ -Borel, yet f is not κ^+ -Borel itself.

Open problems (?)

Let $\kappa > \omega$ be regular and such that $2^{<\kappa} = \kappa$.

- 1 Is it consistent that the generalized Feldman-Moore Theorem holds for equivalence relations on ${}^\kappa 2$?

Interludio: the regular case

One can also easily observe that no “small section” uniformization result can hold if we formulate it in terms of *uniformizing functions*.

Fact

There is a function $f: {}^\kappa 2 \rightarrow {}^\kappa 2$ whose graph $P \subseteq {}^\kappa 2 \times {}^\kappa 2$ is κ^+ -Borel, yet f is not κ^+ -Borel itself.

Open problems (?)

Let $\kappa > \omega$ be regular and such that $2^{<\kappa} = \kappa$.

- 1 Is it consistent that the generalized Feldman-Moore Theorem holds for equivalence relations on ${}^\kappa 2$?
- 2 Is there a κ^+ -Borel equivalence relation on ${}^\kappa 2$ with classes of size at most κ which is not κ^+ -Borel reducible to E_0^κ ?

Interludio: the regular case

One can also easily observe that no “small section” uniformization result can hold if we formulate it in terms of *uniformizing functions*.

Fact

There is a function $f: {}^\kappa 2 \rightarrow {}^\kappa 2$ whose graph $P \subseteq {}^\kappa 2 \times {}^\kappa 2$ is κ^+ -Borel, yet f is not κ^+ -Borel itself.

Open problems (?)

Let $\kappa > \omega$ be regular and such that $2^{<\kappa} = \kappa$.

- 1 Is it consistent that the generalized Feldman-Moore Theorem holds for equivalence relations on ${}^\kappa 2$?
- 2 Is there a κ^+ -Borel equivalence relation on ${}^\kappa 2$ with classes of size at most κ which is not κ^+ -Borel reducible to E_0^κ ?
- 3 Can we have (at least consistently) “small section” uniformization results for κ^+ -Borel subsets of ${}^\kappa 2$?

The countable cofinality case

From now on λ is an uncountable cardinal with $\text{cof}(\lambda) = \omega$ satisfying $2^{<\lambda} = \lambda$ (equivalently, λ is strong limit).

The countable cofinality case

From now on λ is an uncountable cardinal with $\text{cof}(\lambda) = \omega$ satisfying $2^{<\lambda} = \lambda$ (equivalently, λ is strong limit).

For such a λ , GDST becomes the theory of λ -**Polish spaces** (= completely metrizable spaces of weight $\leq \lambda$).

The countable cofinality case

From now on λ is an uncountable cardinal with $\text{cof}(\lambda) = \omega$ satisfying $2^{<\lambda} = \lambda$ (equivalently, λ is strong limit).

For such a λ , GDST becomes the theory of λ -**Polish spaces** (= completely metrizable spaces of weight $\leq \lambda$). The right analogue of Borel sets is given by λ^+ -Borel sets or, equivalently, λ -Borel sets. One can also develop a solid theory of standard λ -Borel spaces.

The countable cofinality case

From now on λ is an uncountable cardinal with $\text{cof}(\lambda) = \omega$ satisfying $2^{<\lambda} = \lambda$ (equivalently, λ is strong limit).

For such a λ , GDST becomes the theory of λ -**Polish spaces** (= completely metrizable spaces of weight $\leq \lambda$). The right analogue of Borel sets is given by λ^+ -Borel sets or, equivalently, λ -Borel sets. One can also develop a solid theory of standard λ -Borel spaces.

Recall that in GDST we must replace ω with either λ or $\text{cof}(\lambda)$, so “countable” should be translated to “of size $\leq \lambda$ ” or remain “of size $\leq \omega$ ”. Similarly, “compact” could be replaced by “ λ -Lindelöf” or stay the same.

The countable cofinality case

From now on λ is an uncountable cardinal with $\text{cof}(\lambda) = \omega$ satisfying $2^{<\lambda} = \lambda$ (equivalently, λ is strong limit).

For such a λ , GDST becomes the theory of λ -**Polish spaces** (= completely metrizable spaces of weight $\leq \lambda$). The right analogue of Borel sets is given by λ^+ -Borel sets or, equivalently, λ -Borel sets. One can also develop a solid theory of standard λ -Borel spaces.

Recall that in GDST we must replace ω with either λ or $\text{cof}(\lambda)$, so “countable” should be translated to “of size $\leq \lambda$ ” or remain “of size $\leq \omega$ ”. Similarly, “compact” could be replaced by “ λ -Lindelöf” or stay the same.

We first consider the second option and look at λ -Borel sets with **countable** vertical sections, or with **compact** vertical sections.

Theorem

Let X be standard λ -Borel, Y be λ -Polish, and $P \subseteq X \times Y$ a λ -Borel set with **countable** vertical sections. Then:

Theorem

Let X be standard λ -Borel, Y be λ -Polish, and $P \subseteq X \times Y$ a λ -Borel set with **countable** vertical sections. Then:

- 1 There is a λ -Borel uniformization of P .

[Equivalently: $\text{proj}_X(P)$ is λ -Borel and there is a λ -Borel uniformizing function $f: \text{proj}_X(P) \rightarrow Y$ for P .]

Theorem

Let X be standard λ -Borel, Y be λ -Polish, and $P \subseteq X \times Y$ a λ -Borel set with **countable** vertical sections. Then:

- 1 There is a λ -Borel uniformization of P .
[Equivalently: $\text{proj}_X(P)$ is λ -Borel and there is a λ -Borel uniformizing function $f: \text{proj}_X(P) \rightarrow Y$ for P .]
- 2 The map from $\text{proj}_X(P)$ to the standard λ -Borel space $F(Y)$ sending $x \in \text{proj}_X(P)$ to $\text{cl}(P_x)$ is λ -Borel.

Theorem

Let X be standard λ -Borel, Y be λ -Polish, and $P \subseteq X \times Y$ a λ -Borel set with **countable** vertical sections. Then:

- 1 There is a λ -Borel uniformization of P .
[Equivalently: $\text{proj}_X(P)$ is λ -Borel and there is a λ -Borel uniformizing function $f: \text{proj}_X(P) \rightarrow Y$ for P .]
- 2 The map from $\text{proj}_X(P)$ to the standard λ -Borel space $F(Y)$ sending $x \in \text{proj}_X(P)$ to $\text{cl}(P_x)$ is λ -Borel.
- 3 There is a sequence $(\zeta_n^P)_{n \in \omega}$ of λ -Borel functions $\zeta_n^P: \text{proj}_X(P) \rightarrow Y$ such that for all $x \in \text{proj}_X(P)$ the set $\{\zeta_n^P(x) \mid n \in \omega\}$ is dense in P_x .

Theorem

Let X be standard λ -Borel, Y be λ -Polish, and $P \subseteq X \times Y$ a λ -Borel set with **countable** vertical sections. Then:

- 1 There is a λ -Borel uniformization of P .
[Equivalently: $\text{proj}_X(P)$ is λ -Borel and there is a λ -Borel uniformizing function $f: \text{proj}_X(P) \rightarrow Y$ for P .]
- 2 The map from $\text{proj}_X(P)$ to the standard λ -Borel space $F(Y)$ sending $x \in \text{proj}_X(P)$ to $\text{cl}(P_x)$ is λ -Borel.
- 3 There is a sequence $(\zeta_n^P)_{n \in \omega}$ of λ -Borel functions $\zeta_n^P: \text{proj}_X(P) \rightarrow Y$ such that for all $x \in \text{proj}_X(P)$ the set $\{\zeta_n^P(x) \mid n \in \omega\}$ is dense in P_x .
- 4 There is a sequence $(\varrho_n^P)_{n \in \omega}$ of λ -Borel functions $\varrho_n^P: \text{proj}_X(P) \rightarrow Y$ such that $P_x = \{\varrho_n^P \mid n \in \omega\}$ for all $x \in \text{proj}_X(P)$.

Theorem

Let X be standard λ -Borel, Y be λ -Polish, and $P \subseteq X \times Y$ a λ -Borel set with **countable** vertical sections. Then:

- 1 There is a λ -Borel uniformization of P .
[Equivalently: $\text{proj}_X(P)$ is λ -Borel and there is a λ -Borel uniformizing function $f: \text{proj}_X(P) \rightarrow Y$ for P .]
- 2 The map from $\text{proj}_X(P)$ to the standard λ -Borel space $F(Y)$ sending $x \in \text{proj}_X(P)$ to $\text{cl}(P_x)$ is λ -Borel.
- 3 There is a sequence $(\zeta_n^P)_{n \in \omega}$ of λ -Borel functions $\zeta_n^P: \text{proj}_X(P) \rightarrow Y$ such that for all $x \in \text{proj}_X(P)$ the set $\{\zeta_n^P(x) \mid n \in \omega\}$ is dense in P_x .
- 4 There is a sequence $(\varrho_n^P)_{n \in \omega}$ of λ -Borel functions $\varrho_n^P: \text{proj}_X(P) \rightarrow Y$ such that $P_x = \{\varrho_n^P \mid n \in \omega\}$ for all $x \in \text{proj}_X(P)$.
- 5 The set P can be written as $P = \bigcup_{n \in \omega} P_n$ where the sets P_n are pairwise disjoint λ -Borel sets with vertical sections of size 1.

Countable vertical sections

The proof is radically different from the classical one because we lack a Baire category notion matching well with that of λ -Borel sets. Instead, we use an easy uniformization theorem and a “change of topology” argument.

Countable vertical sections

The proof is radically different from the classical one because we lack a Baire category notion matching well with that of λ -Borel sets. Instead, we use an easy uniformization theorem and a “change of topology” argument.

Proposition

Let X be a standard λ -Borel space, Z a λ -Polish space, and $F \subseteq X \times Z$ a λ -Borel set such that each of its nonempty vertical sections $F_x \subseteq Z$ has an isolated point. Then F has a λ -Borel uniformization.

Countable vertical sections

The proof is radically different from the classical one because we lack a Baire category notion matching well with that of λ -Borel sets. Instead, we use an easy uniformization theorem and a “change of topology” argument.

Proposition

Let X be a standard λ -Borel space, Z a λ -Polish space, and $F \subseteq X \times Z$ a λ -Borel set such that each of its nonempty vertical sections $F_x \subseteq Z$ has an isolated point. Then F has a λ -Borel uniformization.

Let $P \subseteq X \times Y$ be λ -Borel and with countable vertical sections. Pick a closed set $F \subseteq X \times {}^\omega\lambda$ and a λ -Borel isomorphism $f: F \rightarrow P$ such that $\text{proj}_X(w) = \text{proj}_X(f(w))$ for all $w \in F$.

Countable vertical sections

The proof is radically different from the classical one because we lack a Baire category notion matching well with that of λ -Borel sets. Instead, we use an easy uniformization theorem and a “change of topology” argument.

Proposition

Let X be a standard λ -Borel space, Z a λ -Polish space, and $F \subseteq X \times Z$ a λ -Borel set such that each of its nonempty vertical sections $F_x \subseteq Z$ has an isolated point. Then F has a λ -Borel uniformization.

Let $P \subseteq X \times Y$ be λ -Borel and with countable vertical sections. Pick a closed set $F \subseteq X \times {}^\omega\lambda$ and a λ -Borel isomorphism $f: F \rightarrow P$ such that $\text{proj}_X(w) = \text{proj}_X(f(w))$ for all $w \in F$. All nonempty vertical sections F_x of F are closed and countable, hence Polish (in the classical sense!).

Countable vertical sections

The proof is radically different from the classical one because we lack a Baire category notion matching well with that of λ -Borel sets. Instead, we use an easy uniformization theorem and a “change of topology” argument.

Proposition

Let X be a standard λ -Borel space, Z a λ -Polish space, and $F \subseteq X \times Z$ a λ -Borel set such that each of its nonempty vertical sections $F_x \subseteq Z$ has an isolated point. Then F has a λ -Borel uniformization.

Let $P \subseteq X \times Y$ be λ -Borel and with countable vertical sections. Pick a closed set $F \subseteq X \times {}^\omega\lambda$ and a λ -Borel isomorphism $f: F \rightarrow P$ such that $\text{proj}_X(w) = \text{proj}_X(f(w))$ for all $w \in F$. All nonempty vertical sections F_x of F are closed and countable, hence Polish (in the classical sense!). By countability, each F_x has an isolated point, hence we can apply the proposition.

Countable vertical sections

The proof is radically different from the classical one because we lack a Baire category notion matching well with that of λ -Borel sets. Instead, we use an easy uniformization theorem and a “change of topology” argument.

Proposition

Let X be a standard λ -Borel space, Z a λ -Polish space, and $F \subseteq X \times Z$ a λ -Borel set such that each of its nonempty vertical sections $F_x \subseteq Z$ has an isolated point. Then F has a λ -Borel uniformization.

Let $P \subseteq X \times Y$ be λ -Borel and with countable vertical sections. Pick a closed set $F \subseteq X \times {}^\omega\lambda$ and a λ -Borel isomorphism $f: F \rightarrow P$ such that $\text{proj}_X(w) = \text{proj}_X(f(w))$ for all $w \in F$. All nonempty vertical sections F_x of F are closed and countable, hence Polish (in the classical sense!). By countability, each F_x has an isolated point, hence we can apply the proposition. Let F^* be a λ -Borel uniformization of F : then $P^* = f(F^*)$ is a λ -Borel uniformization of P . □ (part 1)

Proposition

Let E be an equivalence relation on a standard λ -Borel space X that can be written as $E = \bigcup_{\alpha < \mu} P_\alpha$ with $\omega \leq \mu \leq \lambda$ and each P_α a λ -Borel set with vertical sections of size 1. Then there is a (discrete) group G of size $\leq \mu$ acting on X by λ -Borel isomorphisms (in fact, involutions) which generates E . If moreover $\mu > \omega$, then E is hyper- μ -small.

Proposition

Let E be an equivalence relation on a standard λ -Borel space X that can be written as $E = \bigcup_{\alpha < \mu} P_\alpha$ with $\omega \leq \mu \leq \lambda$ and each P_α a λ -Borel set with vertical sections of size 1. Then there is a (discrete) group G of size $\leq \mu$ acting on X by λ -Borel isomorphisms (in fact, involutions) which generates E . If moreover $\mu > \omega$, then E is hyper- μ -small.

Generalized Feldman-Moore Theorem

Let X be a standard λ -Borel space. Then E is a countable λ -Borel equivalence relation on X if and only if it is the orbit equivalence relation induced by a λ -Borel action of a countable (discrete) group G on X .

Theorem

Let X be a standard λ -Borel space, Y a λ -Polish space, and $P \subseteq X \times Y$ a λ -Borel set with **compact** vertical sections. Then:

Theorem

Let X be a standard λ -Borel space, Y a λ -Polish space, and $P \subseteq X \times Y$ a λ -Borel set with **compact** vertical sections. Then:

- 1 There is a λ -Borel uniformization of P .

Theorem

Let X be a standard λ -Borel space, Y a λ -Polish space, and $P \subseteq X \times Y$ a λ -Borel set with **compact** vertical sections. Then:

- 1 There is a λ -Borel uniformization of P .
- 2 The map from $\text{proj}_X(P)$ to $K(Y) \subseteq F(Y)$ sending $x \in \text{proj}_X(P)$ to P_x is λ -Borel.

Theorem

Let X be a standard λ -Borel space, Y a λ -Polish space, and $P \subseteq X \times Y$ a λ -Borel set with **compact** vertical sections. Then:

- 1 There is a λ -Borel uniformization of P .
- 2 The map from $\text{proj}_X(P)$ to $K(Y) \subseteq F(Y)$ sending $x \in \text{proj}_X(P)$ to P_x is λ -Borel.
- 3 There is a sequence $(\zeta_n^P)_{n \in \omega}$ of λ -Borel functions $\zeta_n^P: \text{proj}_X(P) \rightarrow Y$ such that for all $x \in \text{proj}_X(P)$ the set $\{\zeta_n^P(x) \mid n \in \omega\}$ is dense in P_x .

Theorem

Let X be a standard λ -Borel space, Y a λ -Polish space, and $P \subseteq X \times Y$ a λ -Borel set with **compact** vertical sections. Then:

- 1 There is a λ -Borel uniformization of P .
- 2 The map from $\text{proj}_X(P)$ to $K(Y) \subseteq F(Y)$ sending $x \in \text{proj}_X(P)$ to P_x is λ -Borel.
- 3 There is a sequence $(\zeta_n^P)_{n \in \omega}$ of λ -Borel functions $\zeta_n^P: \text{proj}_X(P) \rightarrow Y$ such that for all $x \in \text{proj}_X(P)$ the set $\{\zeta_n^P(x) \mid n \in \omega\}$ is dense in P_x .
- 4 There is a sequence $(\varrho_\alpha^P)_{\alpha < 2^{\aleph_0}}$ of λ -Borel maps $\varrho_\alpha^P: \text{proj}_X(P) \rightarrow Y$ such that $P_x = \{\varrho_\alpha^P \mid \alpha < 2^{\aleph_0}\}$ for all $x \in \text{proj}_X(P)$.

[Recall that $\omega < 2^{\aleph_0} < \lambda$ by choice of λ .]

Theorem

Let X be a standard λ -Borel space, Y a λ -Polish space, and $P \subseteq X \times Y$ a λ -Borel set with **compact** vertical sections. Then:

- 1 There is a λ -Borel uniformization of P .
- 2 The map from $\text{proj}_X(P)$ to $K(Y) \subseteq F(Y)$ sending $x \in \text{proj}_X(P)$ to P_x is λ -Borel.
- 3 There is a sequence $(\zeta_n^P)_{n \in \omega}$ of λ -Borel functions $\zeta_n^P: \text{proj}_X(P) \rightarrow Y$ such that for all $x \in \text{proj}_X(P)$ the set $\{\zeta_n^P(x) \mid n \in \omega\}$ is dense in P_x .
- 4 There is a sequence $(\varrho_\alpha^P)_{\alpha < 2^{\aleph_0}}$ of λ -Borel maps $\varrho_\alpha^P: \text{proj}_X(P) \rightarrow Y$ such that $P_x = \{\varrho_\alpha^P \mid \alpha < 2^{\aleph_0}\}$ for all $x \in \text{proj}_X(P)$.

[Recall that $\omega < 2^{\aleph_0} < \lambda$ by choice of λ .]

- 5 The set P can be written as $P = \bigcup_{\alpha < 2^{\aleph_0}} P_\alpha$ where the sets P_α are pairwise disjoint λ -Borel sets with vertical sections of size 1.

Compact vertical sections

The classical proof uses compactification, which is out of reach here because our spaces are not separable. So we had to develop a quite different machinery.

Compact vertical sections

The classical proof uses compactification, which is out of reach here because our spaces are not separable. So we had to develop a quite different machinery.

Corollary

If E is a λ -Borel equivalence relation on a λ -Polish space X and all its classes are compact, then E is λ -smooth (= λ -Borel reducible to identity on a λ -Polish space).

Compact vertical sections

The classical proof uses compactification, which is out of reach here because our spaces are not separable. So we had to develop a quite different machinery.

Corollary

If E is a λ -Borel equivalence relation on a λ -Polish space X and all its classes are compact, then E is λ -smooth (= λ -Borel reducible to identity on a λ -Polish space).

Recall that $\omega < 2^{\aleph_0} < \lambda$.

Corollary (“half” Feldman-Moore Theorem)

Let E be a λ -Borel equivalence relation on a λ -Polish space X such that all its classes are compact. Then E is the orbit equivalence relation induced by a λ -Borel action on X of a (discrete) group G of size 2^{\aleph_0} .

Compact vertical sections

The classical proof uses compactification, which is out of reach here because our spaces are not separable. So we had to develop a quite different machinery.

Corollary

If E is a λ -Borel equivalence relation on a λ -Polish space X and all its classes are compact, then E is λ -smooth (= λ -Borel reducible to identity on a λ -Polish space).

Recall that $\omega < 2^{\aleph_0} < \lambda$.

Corollary (“half” Feldman-Moore Theorem)

Let E be a λ -Borel equivalence relation on a λ -Polish space X such that all its classes are compact. Then E is the orbit equivalence relation induced by a λ -Borel action on X of a (discrete) group G of size 2^{\aleph_0} .

Moreover, E is hyper- 2^{\aleph_0} -small.

What's next?

There are other “smallness conditions” that could be considered here:

What's next?

There are other “smallness conditions” that could be considered here:

- 1 σ -compact

What's next?

There are other “smallness conditions” that could be considered here:

- ① σ -compact
- ② $\text{size} \leq \lambda$

What's next?

There are other “smallness conditions” that could be considered here:

- ① σ -compact
- ② $\text{size} \leq \lambda$
- ③ closed λ -Lindelöf

What's next?

There are other “smallness conditions” that could be considered here:

- ① σ -compact
- ② $\text{size} \leq \lambda$
- ③ closed λ -Lindelöf
- ④ λ -sized union of compact sets

What's next?

There are other “smallness conditions” that could be considered here:

- ① σ -compact
- ② $\text{size} \leq \lambda$
- ③ closed λ -Lindelöf
- ④ λ -sized union of compact sets
- ⑤ λ -sized union of closed λ -Lindelöf sets

What's next?

There are other “smallness conditions” that could be considered here:

- ① σ -compact
- ② $\text{size} \leq \lambda$
- ③ closed λ -Lindelöf
- ④ λ -sized union of compact sets
- ⑤ λ -sized union of closed λ -Lindelöf sets

In the classical setting, the most general result is provided by the σ -compact case. Here instead we have:

What's next?

There are other “smallness conditions” that could be considered here:

- ① σ -compact
- ② $\text{size} \leq \lambda$
- ③ closed λ -Lindelöf
- ④ λ -sized union of compact sets
- ⑤ λ -sized union of closed λ -Lindelöf sets

In the classical setting, the most general result is provided by the σ -compact case. Here instead we have:

Proposition

If a λ -Polish space X is λ -Lindelöf, then X is well-orderable and $|X| \leq \lambda$.

What's next?

There are other “smallness conditions” that could be considered here:

- ① σ -compact
- ② $\text{size} \leq \lambda$
- ③ closed λ -Lindelöf
- ④ λ -sized union of compact sets
- ⑤ λ -sized union of closed λ -Lindelöf sets

In the classical setting, the most general result is provided by the σ -compact case. Here instead we have:

Proposition

If a λ -Polish space X is λ -Lindelöf, then X is well-orderable and $|X| \leq \lambda$.

Hence the more general case is given by ② (which is equivalent to ④ and ⑤).

What's next?

There are other “smallness conditions” that could be considered here:

- ① σ -compact
- ② $\text{size} \leq \lambda$
- ③ closed λ -Lindelöf
- ④ λ -sized union of compact sets
- ⑤ λ -sized union of closed λ -Lindelöf sets

In the classical setting, the most general result is provided by the σ -compact case. Here instead we have:

Proposition

If a λ -Polish space X is λ -Lindelöf, then X is well-orderable and $|X| \leq \lambda$.

Hence the more general case is given by ② (which is equivalent to ④ and ⑤). This is still work in progress...

...more on the blackboard!!

Open problems

Let $\lambda > \omega$ be singular and such that $2^{<\lambda} = \lambda$.

Open problems

Let $\lambda > \omega$ be singular and such that $2^{<\lambda} = \lambda$.

- ① Can we have “large section” uniformization results, too?

[Meta-problem: Are there interesting λ -ideals to play with?]

Open problems

Let $\lambda > \omega$ be singular and such that $2^{<\lambda} = \lambda$.

- 1 Can we have “large section” uniformization results, too?

[Meta-problem: Are there interesting λ -ideals to play with?]

- 2 What about cardinals λ with uncountable cofinality?

[This case seems to be much closer to the regular one, so we mostly expect negative results.]

Open problems

Let $\lambda > \omega$ be singular and such that $2^{<\lambda} = \lambda$.

- 1 Can we have “large section” uniformization results, too?

[Meta-problem: Are there interesting λ -ideals to play with?]

- 2 What about cardinals λ with uncountable cofinality?

[This case seems to be much closer to the regular one, so we mostly expect negative results.]

Thank you for your attention!