

# Baire property and the Ellentuck-Prikry topology

Vincenzo Dimonte

February 4, 2020

Joint work with Xianghui Shi

## Inspiration (Woodin)

I0 is a large cardinal similar to AD

## Inspiration (Woodin)

I0 is a large cardinal similar to AD.

## Motivation

- Proving theorems that reinforce such statement

## Inspiration (Woodin)

I0 is a large cardinal similar to AD.

## Motivation

- Proving theorems that reinforce such statement
- Understanding the deep reasons behind such similarity

## Inspiration (Woodin)

I0 is a large cardinal similar to AD.

## Motivation

- Proving theorems that reinforce such statement
- Understanding the deep reasons behind such similarity

## Definition (Woodin, 1980)

We say that  $I0(\lambda)$  holds iff there is an elementary embedding  $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$  such that  $j \upharpoonright V_{\lambda+1}$  is not the identity

## Inspiration (Woodin)

I0 is a large cardinal similar to AD.

## Motivation

- Proving theorems that reinforce such statement
- Understanding the deep reasons behind such similarity

## Definition (Woodin, 1980)

We say that  $I_0(\lambda)$  holds iff there is an elementary embedding  $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$  such that  $j \restriction V_{\lambda+1}$  is not the identity.

It is a large cardinal

## Inspiration (Woodin)

I0 is a large cardinal similar to AD.

## Motivation

- Proving theorems that reinforce such statement
- Understanding the deep reasons behind such similarity

## Definition (Woodin, 1980)

We say that  $I_0(\lambda)$  holds iff there is an elementary embedding  $j : L(V_{\lambda+1}) \prec L(V_{\lambda+1})$  such that  $j \upharpoonright V_{\lambda+1}$  is not the identity.

It is a large cardinal: if  $I_0(\lambda)$  holds, then  $\lambda$  is a strong limit cardinal of cofinality  $\omega$ , limit of cardinals that are  $n$ -huge for every  $n \in \omega$ .

These are some similarities with AD



These are some similarities with AD:

$L(\mathbb{R})$ under AD	$L(V_{\lambda+1})$ under I0( $\lambda$ )
DC	$DC_\lambda$

These are some similarities with AD:

$L(\mathbb{R})$ under AD	$L(V_{\lambda+1})$ under I0( $\lambda$ )
DC	$DC_\lambda$
$\Theta$ is regular	$\Theta^{L(V_{\lambda+1})}$ is regular

These are some similarities with AD:

$L(\mathbb{R})$ under AD	$L(V_{\lambda+1})$ under I0( $\lambda$ )
DC	$DC_\lambda$
$\Theta$ is regular	$\Theta^{L(V_{\lambda+1})}$ is regular
$\omega_1$ is measurable	$\lambda^+$ is measurable

These are some similarities with AD:

$L(\mathbb{R})$ under AD	$L(V_{\lambda+1})$ under $I0(\lambda)$
DC	$DC_\lambda$
$\Theta$ is regular	$\Theta^{L(V_{\lambda+1})}$ is regular
$\omega_1$ is measurable	$\lambda^+$ is measurable
the Coding Lemma holds	the Coding Lemma holds

These are some similarities with AD:

$L(\mathbb{R})$ under AD	$L(V_{\lambda+1})$ under I0( $\lambda$ )
DC	$DC_\lambda$
$\Theta$ is regular	$\Theta^{L(V_{\lambda+1})}$ is regular
$\omega_1$ is measurable	$\lambda^+$ is measurable
the Coding Lemma holds	the Coding Lemma holds

### Theorem (Laver)

Let  $\langle \kappa_n : n \in \omega \rangle$  be a cofinal sequence in  $\lambda$ . For every  $A \subseteq V_\lambda$ :

- $A$  is  $\Sigma_1^1$ -definable in  $(V_\lambda, V_{\lambda+1})$  iff there is a tree  $T \subseteq \prod_{n \in \omega} V_{\kappa_n} \times \prod_{n \in \omega} V_{\kappa_n}$  whose projection is  $A$ ;
- $A$  is  $\Sigma_2^1$ -definable in  $(V_\lambda, V_{\lambda+1})$  iff there is a tree  $T \subseteq \prod_{n \in \omega} V_{\kappa_n} \times \lambda^+$  whose projection is  $A$

These are some similarities with AD:

$L(\mathbb{R})$ under AD	$L(V_{\lambda+1})$ under I0( $\lambda$ )
DC	$DC_\lambda$
$\Theta$ is regular	$\Theta^{L(V_{\lambda+1})}$ is regular
$\omega_1$ is measurable	$\lambda^+$ is measurable
the Coding Lemma holds	the Coding Lemma holds

### Theorem (Laver)

Let  $\langle \kappa_n : n \in \omega \rangle$  be a cofinal sequence in  $\lambda$ . For every  $A \subseteq V_\lambda$ :

- $A$  is  $\Sigma_1^1$ -definable in  $(V_\lambda, V_{\lambda+1})$  iff there is a tree  $T \subseteq \prod_{n \in \omega} V_{\kappa_n} \times \prod_{n \in \omega} V_{\kappa_n}$  whose projection is  $A$ ;
- $A$  is  $\Sigma_2^1$ -definable in  $(V_\lambda, V_{\lambda+1})$  iff there is a tree  $T \subseteq \prod_{n \in \omega} V_{\kappa_n} \times \lambda^+$  whose projection is  $A$ .

Let us go a bit deeper

These are some similarities with AD:

$L(\mathbb{R})$ under AD	$L(V_{\lambda+1})$ under $I0(\lambda)$
DC	$DC_\lambda$
$\Theta$ is regular	$\Theta^{L(V_{\lambda+1})}$ is regular
$\omega_1$ is measurable	$\lambda^+$ is measurable
the Coding Lemma holds	the Coding Lemma holds

### Theorem (Laver)

Let  $\langle \kappa_n : n \in \omega \rangle$  be a cofinal sequence in  $\lambda$ . For every  $A \subseteq V_\lambda$ :

- $A$  is  $\Sigma_1^1$ -definable in  $(V_\lambda, V_{\lambda+1})$  iff there is a tree  $T \subseteq \prod_{n \in \omega} V_{\kappa_n} \times \prod_{n \in \omega} V_{\kappa_n}$  whose projection is  $A$ ;
- $A$  is  $\Sigma_2^1$ -definable in  $(V_\lambda, V_{\lambda+1})$  iff there is a tree  $T \subseteq \prod_{n \in \omega} V_{\kappa_n} \times \lambda^+$  whose projection is  $A$ .

Let us go a bit deeper.

We define a topology on  $V_{\lambda+1}$



We define a topology on  $V_{\lambda+1}$ : Since  $V_{\lambda+1} = \mathcal{P}(V_\lambda)$ , the basic open sets of the topology are, for any  $\alpha < \lambda$  and  $a \subseteq V_\alpha$ ,

$$O_{(a,\alpha)} = \{b \in V_{\lambda+1} : b \cap V_\alpha = a\}.$$

Theorem (Cramer, 2015)

Suppose I0( $\lambda$ )

We define a topology on  $V_{\lambda+1}$ : Since  $V_{\lambda+1} = \mathcal{P}(V_\lambda)$ , the basic open sets of the topology are, for any  $\alpha < \lambda$  and  $a \subseteq V_\alpha$ ,

$$O_{(a,\alpha)} = \{b \in V_{\lambda+1} : b \cap V_\alpha = a\}.$$

### Theorem (Cramer, 2015)

Suppose  $\text{I0}(\lambda)$ . Then for every  $X \subseteq V_{\lambda+1}$ ,  $X \in L(V_{\lambda+1})$ , either  $|X| \leq \lambda$  or  ${}^\omega\lambda$  can be continuously embedded inside  $X$  ( ${}^\omega\lambda$  with the bounded topology)

We define a topology on  $V_{\lambda+1}$ : Since  $V_{\lambda+1} = \mathcal{P}(V_\lambda)$ , the basic open sets of the topology are, for any  $\alpha < \lambda$  and  $a \subseteq V_\alpha$ ,

$$O_{(a,\alpha)} = \{b \in V_{\lambda+1} : b \cap V_\alpha = a\}.$$

### Theorem (Cramer, 2015)

Suppose  $I_0(\lambda)$ . Then for every  $X \subseteq V_{\lambda+1}$ ,  $X \in L(V_{\lambda+1})$ , either  $|X| \leq \lambda$  or  ${}^\omega\lambda$  can be continuously embedded inside  $X$  ( ${}^\omega\lambda$  with the bounded topology).

This is similar to AD: in fact, under AD every subset of the reals has the Perfect Set Property.

But the proof is completely different: Cramer uses heavily elementary embeddings (inverse limit reflection), while in the classical case involves games.

In recent work, with Motto Ros we clarified the similarity

In recent work, with Motto Ros we clarified the similarity.  
The classical case is

In recent work, with Motto Ros we clarified the similarity.

The classical case is:

- Large cardinals  $\Rightarrow$  every set of reals in  $L(\mathbb{R})$  is homogeneously Suslin

In recent work, with Motto Ros we clarified the similarity.

The classical case is:

- Large cardinals  $\Rightarrow$  every set of reals in  $L(\mathbb{R})$  is homogeneously Suslin
- Every homogeneously Suslin set is determined (so  $L(\mathbb{R}) \models \text{AD}$ )

In recent work, with Motto Ros we clarified the similarity.

The classical case is:

- Large cardinals  $\Rightarrow$  every set of reals in  $L(\mathbb{R})$  is homogeneously Suslin
- Every homogeneously Suslin set is determined (so  $L(\mathbb{R}) \models \text{AD}$ )
- Every determined set has the Perfect Set Property



But there is a shortcut for regularity properties

But there is a shortcut for regularity properties:

- Infinite Woodin cardinals  $\Rightarrow$  every set of reals in  $L(\mathbb{R})$  is weakly homogeneously Suslin

But there is a shortcut for regularity properties:

- Infinite Woodin cardinals  $\Rightarrow$  every set of reals in  $L(\mathbb{R})$  is weakly homogeneously Suslin
- Every weakly homogeneously Suslin set has the Perfect Set Property

## D.-Motto Ros

Let  $\lambda$  be a strong limit cardinal of cofinality  $\omega$ , and let  $\langle \kappa_n : n \in \omega \rangle$  be an increasing cofinal sequence in  $\lambda$

## D.-Motto Ros

Let  $\lambda$  be a strong limit cardinal of cofinality  $\omega$ , and let  $\langle \kappa_n : n \in \omega \rangle$  be a increasing cofinal sequence in  $\lambda$ . Then the following spaces are isomorphic:

- ${}^\lambda 2$ , with the bounded topology

## D.-Motto Ros

Let  $\lambda$  be a strong limit cardinal of cofinality  $\omega$ , and let  $\langle \kappa_n : n \in \omega \rangle$  be a increasing cofinal sequence in  $\lambda$ . Then the following spaces are isomorphic:

- ${}^\lambda 2$ , with the bounded topology;
- ${}^\omega \lambda$ , with the bounded topology, and the discrete topology in every copy of  $\lambda$

## D.-Motto Ros

Let  $\lambda$  be a strong limit cardinal of cofinality  $\omega$ , and let  $\langle \kappa_n : n \in \omega \rangle$  be a increasing cofinal sequence in  $\lambda$ . Then the following spaces are isomorphic:

- ${}^\lambda 2$ , with the bounded topology;
- ${}^\omega \lambda$ , with the bounded topology, and the discrete topology in every copy of  $\lambda$ ;
- $\prod_{n \in \omega} \kappa_n$ , with the bounded topology and the discrete topology in every  $\kappa_n$

## D.-Motto Ros

Let  $\lambda$  be a strong limit cardinal of cofinality  $\omega$ , and let  $\langle \kappa_n : n \in \omega \rangle$  be a increasing cofinal sequence in  $\lambda$ . Then the following spaces are isomorphic:

- ${}^\lambda 2$ , with the bounded topology;
- ${}^\omega \lambda$ , with the bounded topology, and the discrete topology in every copy of  $\lambda$ ;
- $\prod_{n \in \omega} \kappa_n$ , with the bounded topology and the discrete topology in every  $\kappa_n$ ;
- if  $|V_\lambda| = \lambda$ ,  $V_{\lambda+1}$ , with the previously defined topology



## D.-Motto Ros

Let  $\lambda$  be a strong limit cardinal of cofinality  $\omega$ , and let  $\langle \kappa_n : n \in \omega \rangle$  be a increasing cofinal sequence in  $\lambda$ . Then the following spaces are isomorphic:

- ${}^\lambda 2$ , with the bounded topology;
- ${}^\omega \lambda$ , with the bounded topology, and the discrete topology in every copy of  $\lambda$ ;
- $\prod_{n \in \omega} \kappa_n$ , with the bounded topology and the discrete topology in every  $\kappa_n$ ;
- if  $|V_\lambda| = \lambda$ ,  $V_{\lambda+1}$ , with the previously defined topology.

Moreover, they are  $\lambda$ -Polish, i.e., completely metrizable and with a dense subset of cardinality  $\lambda$ .

So, for example, we can rewrite Cramer's result as

So, for example, we can rewrite Cramer's result as:

### Theorem (Cramer, 2015)

Suppose  $I_0(\lambda)$ . Then  $L(\lambda^2) \models \forall X X \subseteq \lambda^2$  has the  $\lambda$ -PSP

So, for example, we can rewrite Cramer's result as:

### Theorem (Cramer, 2015)

Suppose  $I_0(\lambda)$ . Then  $L(\lambda^2) \models \forall X X \subseteq \lambda^2$  has the  $\lambda$ -PSP.

For any  $\lambda$  strong limit of cofinality  $\omega$ , we defined *representable* subsets of  ${}^\omega\lambda$ , a generalization of weakly homogeneously Suslin sets

So, for example, we can rewrite Cramer's result as:

### Theorem (Cramer, 2015)

Suppose  $I_0(\lambda)$ . Then  $L(\lambda^2) \models \forall X X \subseteq \lambda^2$  has the  $\lambda$ -PSP.

For any  $\lambda$  strong limit of cofinality  $\omega$ , we defined *representable* subsets of  ${}^\omega\lambda$ , a generalization of weakly homogeneously Suslin sets.

### D.-Motto Ros

Let  $\lambda$  strong limit of cofinality  $\omega$ . Then every representable subset of  ${}^\omega\lambda$  has the  $\lambda$ -PSP

So, for example, we can rewrite Cramer's result as:

### Theorem (Cramer, 2015)

Suppose  $I_0(\lambda)$ . Then  $L(\lambda^2) \models \forall X X \subseteq \lambda^2$  has the  $\lambda$ -PSP.

For any  $\lambda$  strong limit of cofinality  $\omega$ , we defined *representable* subsets of  ${}^\omega\lambda$ , a generalization of weakly homogeneously Suslin sets.

### D.-Motto Ros

Let  $\lambda$  strong limit of cofinality  $\omega$ . Then every representable subset of  ${}^\omega\lambda$  has the  $\lambda$ -PSP.

Cramer's analysis of  $I_0$  finalizes the similarity with AD

So, for example, we can rewrite Cramer's result as:

### Theorem (Cramer, 2015)

Suppose  $I_0(\lambda)$ . Then  $L(\lambda^2) \models \forall X X \subseteq \lambda^2$  has the  $\lambda$ -PSP.

For any  $\lambda$  strong limit of cofinality  $\omega$ , we defined *representable* subsets of  ${}^\omega\lambda$ , a generalization of weakly homogeneously Suslin sets.

### D.-Motto Ros

Let  $\lambda$  strong limit of cofinality  $\omega$ . Then every representable subset of  ${}^\omega\lambda$  has the  $\lambda$ -PSP.

Cramer's analysis of  $I_0$  finalizes the similarity with AD:

### Theorem (Cramer, to appear)

Suppose  $I_0(\lambda)$ . Then every  $X \subseteq V_{\lambda+1}$ ,  $X \in L(V_{\lambda+1})$  is representable.

- Infinite Woodin cardinals  $\Rightarrow$  every set of reals in  $L(\mathbb{R})$  is weakly homogeneously Suslin
- Every weakly homogeneously Suslin set has the Perfect Set Property



- $I_0 \Rightarrow$  every subset of  $V_{\lambda+1}$  in  $L(V_{\lambda+1})$  is representable
- Every representable set has the  $\lambda$ -Perfect Set Property

This approach is more scalable, as it works even in  $\lambda$ -Polish spaces such that  $\lambda$  does not satisfy I0

This approach is more scalable, as it works even in  $\lambda$ -Polish spaces such that  $\lambda$  does not satisfy I0:

### D.-Motto Ros

Suppose  $I0(\lambda)$ . Then it is consistent that there is  $\kappa$  strong limit of cofinality  $\omega$  such that all the subsets of  ${}^\omega\kappa$  in  $L(V_{\kappa+1})$  have the  $\kappa$ -PSP, and  $\neg I0(\kappa)$ .

The next step would be to analyze the Baire Property.

The most natural thing is to define nowhere dense sets as usual,  $\lambda$ -meager sets as  $\lambda$ -union of nowhere dense sets and  $\lambda$ -comeager sets as complement of  $\lambda$ -meager sets.

The most natural thing is to define nowhere dense sets as usual,  $\lambda$ -meagre sets as  $\lambda$ -union of nowhere dense sets and  $\lambda$ -comeagre sets as complement of  $\lambda$ -meagre sets.

Let  $f : \omega \rightarrow \omega$

Let  $f : \omega \rightarrow \omega$ .

Then  $D_f = \prod_{n \in \omega} \kappa_{f(n)}$  is nowhere dense in  ${}^\omega \lambda$



Let  $f : \omega \rightarrow \omega$ .

Then  $D_f = \prod_{n \in \omega} \kappa_{f(n)}$  is nowhere dense in  ${}^\omega\lambda$ .

But  ${}^\omega\lambda = \bigcup_{f \in {}^\omega\omega} D_f$ , therefore the whole space is  $\lambda$ -meagre (in fact, it is  $\mathfrak{c}$ -meagre), and the Baire property in this setting is just nonsense.

Or is it?

From now on, we work with  $\lambda$  strong limit of cofinality  $\omega$ ,  
 $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable*  
cardinals in  $\lambda$

From now on, we work with  $\lambda$  strong limit of cofinality  $\omega$ ,  
 $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable*  
cardinals in  $\lambda$ .

The space we work in is  $\prod_{n \in \omega} \kappa_n$

From now on, we work with  $\lambda$  strong limit of cofinality  $\omega$ ,  $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable* cardinals in  $\lambda$ .

The space we work in is  $\prod_{n \in \omega} \kappa_n$ .

### Idea

- Baire category is closely connected to Cohen forcing

From now on, we work with  $\lambda$  strong limit of cofinality  $\omega$ ,  $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable* cardinals in  $\lambda$ .

The space we work in is  $\prod_{n \in \omega} \kappa_n$ .

### Idea

- Baire category is closely connected to Cohen forcing
- The space  ${}^\kappa 2$ , with  $\kappa$  regular, is  $\kappa$ -Baire (i.e., every nonempty open set is not  $\kappa$ -meagre) because Cohen forcing on  $\kappa$  is  $< \kappa$ -distributive

From now on, we work with  $\lambda$  strong limit of cofinality  $\omega$ ,  $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable* cardinals in  $\lambda$ .

The space we work in is  $\prod_{n \in \omega} \kappa_n$ .

### Idea

- Baire category is closely connected to Cohen forcing
- The space  ${}^\kappa 2$ , with  $\kappa$  regular, is  $\kappa$ -Baire (i.e., every nonempty open set is not  $\kappa$ -meagre) because Cohen forcing on  $\kappa$  is  $< \kappa$ -distributive
- “Cohen” forcing on  $\lambda$  singular is not  $< \lambda$ -distributive, and this is why  ${}^\lambda 2$  is not  $\lambda$ -Baire

From now on, we work with  $\lambda$  strong limit of cofinality  $\omega$ ,  $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable* cardinals in  $\lambda$ .

The space we work in is  $\prod_{n \in \omega} \kappa_n$ .

### Idea

- Baire category is closely connected to Cohen forcing
- The space  ${}^\kappa 2$ , with  $\kappa$  regular, is  $\kappa$ -Baire (i.e., every nonempty open set is not  $\kappa$ -meagre) because Cohen forcing on  $\kappa$  is  $< \kappa$ -distributive
- “Cohen” forcing on  $\lambda$  singular is not  $< \lambda$ -distributive, and this is why  ${}^\lambda 2$  is not  $\lambda$ -Baire
- But there are other forcings on  $\lambda$  that are  $< \lambda$ -distributive, like Prikry forcing



From now on, we work with  $\lambda$  strong limit of cofinality  $\omega$ ,  $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable* cardinals in  $\lambda$ .

The space we work in is  $\prod_{n \in \omega} \kappa_n$ .

### Idea

- Baire category is closely connected to Cohen forcing
- The space  ${}^\kappa 2$ , with  $\kappa$  regular, is  $\kappa$ -Baire (i.e., every nonempty open set is not  $\kappa$ -meagre) because Cohen forcing on  $\kappa$  is  $< \kappa$ -distributive
- “Cohen” forcing on  $\lambda$  singular is not  $< \lambda$ -distributive, and this is why  ${}^\lambda 2$  is not  $\lambda$ -Baire
- But there are other forcings on  $\lambda$  that are  $< \lambda$ -distributive, like Prikry forcing
- We can try to define Baire category via Prikry forcing instead of Cohen forcing.

## Definition

Let  $\lambda$  be strong limit of cofinality  $\omega$ ,  $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable* cardinals in  $\lambda$ , and fix  $\mu_n$  a measure for each  $\kappa_n$

## Definition

Let  $\lambda$  be strong limit of cofinality  $\omega$ ,  $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable* cardinals in  $\lambda$ , and fix  $\mu_n$  a measure for each  $\kappa_n$ . The *Präkry forcing*  $\mathbb{P}_{\vec{\mu}}$  on  $\lambda$  respect to  $\vec{\mu}$  has conditions of the form  $\langle \alpha_1, \dots, \alpha_n, A_{n+1}, A_{n+2} \dots \rangle$ , where  $\alpha_i \in \kappa_i$  and  $A_i \in \mu_i$

## Definition

Let  $\lambda$  be strong limit of cofinality  $\omega$ ,  $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable* cardinals in  $\lambda$ , and fix  $\mu_n$  a measure for each  $\kappa_n$ . The *Prikry forcing*  $\mathbb{P}_{\vec{\mu}}$  on  $\lambda$  respect to  $\vec{\mu}$  has conditions of the form  $\langle \alpha_1, \dots, \alpha_n, A_{n+1}, A_{n+2} \dots \rangle$ , where  $\alpha_i \in \kappa_i$  and  $A_i \in \mu_i$ .

$\langle \alpha_1, \dots, \alpha_n \rangle$  is the *stem* of the condition

## Definition

Let  $\lambda$  be strong limit of cofinality  $\omega$ ,  $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable* cardinals in  $\lambda$ , and fix  $\mu_n$  a measure for each  $\kappa_n$ . The *Prikry forcing*  $\mathbb{P}_{\vec{\mu}}$  on  $\lambda$  respect to  $\vec{\mu}$  has conditions of the form  $\langle \alpha_1, \dots, \alpha_n, A_{n+1}, A_{n+2} \dots \rangle$ , where  $\alpha_i \in \kappa_i$  and  $A_i \in \mu_i$ .

$\langle \alpha_1, \dots, \alpha_n \rangle$  is the *stem* of the condition.

$\langle \beta_1, \dots, \beta_m, B_{m+1}, B_{m+2} \dots \rangle \leq \langle \alpha_1, \dots, \alpha_n, A_{n+1}, A_{n+2} \dots \rangle$  iff  $m \geq n$  and

- for  $i \leq n$   $\beta_i = \alpha_i$
- for  $n < i \leq m$   $\beta_i \in A_i$
- for  $i > m$   $B_i \subseteq A_i$

## Definition

Let  $\lambda$  be strong limit of cofinality  $\omega$ ,  $\langle \kappa_n : n \in \omega \rangle$  a strictly increasing cofinal sequence of *measurable* cardinals in  $\lambda$ , and fix  $\mu_n$  a measure for each  $\kappa_n$ . The *Prikry forcing*  $\mathbb{P}_{\vec{\mu}}$  on  $\lambda$  respect to  $\vec{\mu}$  has conditions of the form  $\langle \alpha_1, \dots, \alpha_n, A_{n+1}, A_{n+2} \dots \rangle$ , where  $\alpha_i \in \kappa_i$  and  $A_i \in \mu_i$ .

$\langle \alpha_1, \dots, \alpha_n \rangle$  is the *stem* of the condition.

$\langle \beta_1, \dots, \beta_m, B_{m+1}, B_{m+2} \dots \rangle \leq \langle \alpha_1, \dots, \alpha_n, A_{n+1}, A_{n+2} \dots \rangle$  iff  $m \geq n$  and

- for  $i \leq n$   $\beta_i = \alpha_i$
- for  $n < i \leq m$   $\beta_i \in A_i$
- for  $i > m$   $B_i \subseteq A_i$ .

$p \leq^* q$  if  $p \leq q$  and they have the same stem.

## Definition

The *Ellentuck-Prikry  $\vec{\mu}$ -topology* (in short EP-topology) on  $\prod_{n \in \omega} \kappa_n$  is the topology generated by the family  $\{O_p : p \in \mathbb{P}_{\vec{\mu}}\}$

## Definition

The *Ellentuck-Prikry  $\vec{\mu}$ -topology* (in short EP-topology) on  $\prod_{n \in \omega} \kappa_n$  is the topology generated by the family  $\{O_p : p \in \mathbb{P}_{\vec{\mu}}\}$ , where if  $p = \langle \alpha_1, \dots, \alpha_n, A_{n+1}, A_{n+2} \dots \rangle$ , then

$$O_p = \{x \in \prod_{n \in \omega} \kappa_n : \forall i \leq n \ x(i) = \alpha_i, \ \forall i > n \ x(i) \in A_i\}.$$

The EP-topology is a refinement of the bounded topology



## Definition

The *Ellentuck-Prikry  $\vec{\mu}$ -topology* (in short EP-topology) on  $\prod_{n \in \omega} \kappa_n$  is the topology generated by the family  $\{O_p : p \in \mathbb{P}_{\vec{\mu}}\}$ , where if  $p = \langle \alpha_1, \dots, \alpha_n, A_{n+1}, A_{n+2} \dots \rangle$ , then

$$O_p = \{x \in \prod_{n \in \omega} \kappa_n : \forall i \leq n \ x(i) = \alpha_i, \ \forall i > n \ x(i) \in A_i\}.$$

The EP-topology is a refinement of the bounded topology: if a set is open in the bounded topology, it is open also in the EP-topology, but not viceversa

## Definition

The *Ellentuck-Prikry  $\vec{\mu}$ -topology* (in short EP-topology) on  $\prod_{n \in \omega} \kappa_n$  is the topology generated by the family  $\{O_p : p \in \mathbb{P}_{\vec{\mu}}\}$ , where if  $p = \langle \alpha_1, \dots, \alpha_n, A_{n+1}, A_{n+2} \dots \rangle$ , then

$$O_p = \{x \in \prod_{n \in \omega} \kappa_n : \forall i \leq n \ x(i) = \alpha_i, \ \forall i > n \ x(i) \in A_i\}.$$

The EP-topology is a refinement of the bounded topology: if a set is open in the bounded topology, it is open also in the EP-topology, but not viceversa (in fact, many open sets in the EP-topology are nowhere dense in the bounded topology).

There is a connection between the concepts of “open” and “dense” relative to the forcing and relative to the topology

There is a connection between the concepts of “open” and “dense” relative to the forcing and relative to the topology:

$\mathbb{P}_{\vec{\mu}}$  (forcing)

$\prod_{n \in \omega} \kappa_n$  (EP-topology)

There is a connection between the concepts of “open” and “dense” relative to the forcing and relative to the topology:

$$\frac{\mathbb{P}_{\vec{\mu}} \text{ (forcing)}}{O \text{ open}} \rightarrow \frac{\prod_{n \in \omega} \kappa_n \text{ (EP-topology)}}{O = \{x \in \prod_{n \in \omega} \kappa_n : \exists p \in \mathbb{P}_{\vec{\mu}} x \in O_p\} \text{ open}}$$

There is a connection between the concepts of “open” and “dense” relative to the forcing and relative to the topology:

$$\begin{array}{ccc}
 \mathbb{P}_{\vec{\mu}} \text{ (forcing)} & & \prod_{n \in \omega} \kappa_n \text{ (EP-topology)} \\
 \hline
 O \text{ open} & \rightarrow & {}^I O = \{x \in \prod_{n \in \omega} \kappa_n : \\
 & & \exists p \in \mathbb{P}_{\vec{\mu}} x \in O_p\} \text{ open} \\
 \\ 
 {}^I U = & \leftarrow & U \text{ open} \\
 \{p \in \mathbb{P}_{\vec{\mu}} : O_p \subseteq U\} \text{ open} & & 
 \end{array}$$

There is a connection between the concepts of “open” and “dense” relative to the forcing and relative to the topology:

$$\begin{array}{ccc}
 \mathbb{P}_{\vec{\mu}} \text{ (forcing)} & & \prod_{n \in \omega} \kappa_n \text{ (EP-topology)} \\
 \hline
 O \text{ open} & \rightarrow & {}^I O = \{x \in \prod_{n \in \omega} \kappa_n : \\
 & & \exists p \in \mathbb{P}_{\vec{\mu}} x \in O_p\} \text{ open} \\
 \\ 
 {}^I U = & \leftarrow & U \text{ open} \\
 \{p \in \mathbb{P}_{\vec{\mu}} : O_p \subseteq U\} \text{ open} & & \\
 \\ 
 O \text{ open dense} & \rightarrow & {}^I O \text{ open dense}
 \end{array}$$

There is a connection between the concepts of “open” and “dense” relative to the forcing and relative to the topology:

$$\begin{array}{ccc}
 \mathbb{P}_{\vec{\mu}} \text{ (forcing)} & & \prod_{n \in \omega} \kappa_n \text{ (EP-topology)} \\
 \hline
 O \text{ open} & \rightarrow & {}^I O = \{x \in \prod_{n \in \omega} \kappa_n : \\
 & & \exists p \in \mathbb{P}_{\vec{\mu}} x \in O_p\} \text{ open} \\
 \\ 
 {}^I U = & \leftarrow & U \text{ open} \\
 \{p \in \mathbb{P}_{\vec{\mu}} : O_p \subseteq U\} \text{ open} & & \\
 \\ 
 O \text{ open dense} & \rightarrow & {}^I O \text{ open dense} \\
 \\ 
 {}^I U \text{ open dense} & \leftarrow & U \text{ open dense}
 \end{array}$$



There is a connection between the concepts of “open” and “dense” relative to the forcing and relative to the topology:

$$\frac{\mathbb{P}_{\vec{\mu}} \text{ (forcing)}}{O \text{ open}} \rightarrow \frac{\prod_{n \in \omega} \kappa_n \text{ (EP-topology)}}{{}^I O = \{x \in \prod_{n \in \omega} \kappa_n : \exists p \in \mathbb{P}_{\vec{\mu}} x \in O_p\} \text{ open}}$$

$$\begin{array}{ccc} {}_I U = & \leftarrow & U \text{ open} \\ \{p \in \mathbb{P}_{\vec{\mu}} : O_p \subseteq U\} \text{ open} & & \end{array}$$

$$O \text{ open dense} \rightarrow {}^I O \text{ open dense}$$

$${}_I U \text{ open dense} \leftarrow U \text{ open dense}$$

${}_I({}_I U) = U$ , but not viceversa.

## Definition

Let  $X$  be a topological space

## Definition

Let  $X$  be a topological space.

- a set  $A \subseteq X$  is  $\lambda$ -meagre iff it is the  $\lambda$ -union of nowhere dense sets

## Definition

Let  $X$  be a topological space.

- a set  $A \subseteq X$  is  $\lambda$ -meagre iff it is the  $\lambda$ -union of nowhere dense sets
- a set  $A \subseteq X$  is  $\lambda$ -comeagre iff it is the complement of a  $\lambda$ -meagre set

## Definition

Let  $X$  be a topological space.

- a set  $A \subseteq X$  is  $\lambda$ -meagre iff it is the  $\lambda$ -union of nowhere dense sets
- a set  $A \subseteq X$  is  $\lambda$ -comeagre iff it is the complement of a  $\lambda$ -meagre set
- a set  $A \subseteq X$  has the  $\lambda$ -Baire property iff there is an open set  $U$  such that  $A \triangle U$  is  $\lambda$ -meagre

## Definition

Let  $X$  be a topological space.

- a set  $A \subseteq X$  is  $\lambda$ -meagre iff it is the  $\lambda$ -union of nowhere dense sets
- a set  $A \subseteq X$  is  $\lambda$ -comeagre iff it is the complement of a  $\lambda$ -meagre set
- a set  $A \subseteq X$  has the  $\lambda$ -Baire property iff there is an open set  $U$  such that  $A \triangle U$  is  $\lambda$ -meagre
- $X$  is a  $\lambda$ -Baire space iff every nonempty open set in  $X$  is not  $\lambda$ -meagre, i.e., the intersection of  $\lambda$ -many open dense sets is dense.

The key to prove that the space  $\prod_{n \in \omega} \kappa_n$  is  $\lambda$ -Baire resides in this combinatorial property of Prikry forcing

The key to prove that the space  $\prod_{n \in \omega} \kappa_n$  is  $\lambda$ -Baire resides in this combinatorial property of Prikry forcing:

### Strong Prikry condition

Let  $D \subseteq \mathbb{P}_{\vec{\mu}}$  be open dense



The key to prove that the space  $\prod_{n \in \omega} \kappa_n$  is  $\lambda$ -Baire resides in this combinatorial property of Prikry forcing:

### Strong Prikry condition

Let  $D \subseteq \mathbb{P}_{\vec{\mu}}$  be open dense. Then for every  $p \in \mathbb{P}_{\vec{\mu}}$  there are  $p' \leq^* p$  and  $n \in \omega$  such that for every  $q \leq p'$  with stem of length at least  $n$ ,  $q \in D$

The key to prove that the space  $\prod_{n \in \omega} \kappa_n$  is  $\lambda$ -Baire resides in this combinatorial property of Prikry forcing:

### Strong Prikry condition

Let  $D \subseteq \mathbb{P}_{\vec{\mu}}$  be open dense. Then for every  $p \in \mathbb{P}_{\vec{\mu}}$  there are  $p' \leq^* p$  and  $n \in \omega$  such that for every  $q \leq p'$  with stem of length at least  $n$ ,  $q \in D$ .

Topologically: Let  $D \subseteq \prod_{n \in \omega} \kappa_n$  be open dense. Then for every  $p \in \mathbb{P}_{\vec{\mu}}$  there is a  $p' \leq^* p$  such that  $O_{p'} \subseteq D$

The key to prove that the space  $\prod_{n \in \omega} \kappa_n$  is  $\lambda$ -Baire resides in this combinatorial property of Prikry forcing:

### Strong Prikry condition

Let  $D \subseteq \mathbb{P}_{\vec{\mu}}$  be open dense. Then for every  $p \in \mathbb{P}_{\vec{\mu}}$  there are  $p' \leq^* p$  and  $n \in \omega$  such that for every  $q \leq p'$  with stem of length at least  $n$ ,  $q \in D$ .

Topologically: Let  $D \subseteq \prod_{n \in \omega} \kappa_n$  be open dense. Then for every  $p \in \mathbb{P}_{\vec{\mu}}$  there is a  $p' \leq^* p$  such that  $O_{p'} \subseteq D$ .

Coupled with the fact that if  $p \in \mathbb{P}_{\vec{\mu}}$  has stem of length  $n$ , then the intersection of  $< \kappa_n$ -many  $\leq^*$ -extensions of  $p$  is still in  $\mathbb{P}_{\vec{\mu}}$ , we have:

### Proposition (D.-Shi)

The space  $\prod_{n \in \omega} \kappa_n$  with the EP-topology is  $\lambda$ -Baire.

### (Generalized) Mycielski Theorem (D.-Shi)

In  $\prod_{n \in \omega} \kappa_n$  with the EP-topology every  $\lambda$ -comeagre set contains a  $\lambda$ -perfect set

### (Generalized) Mycielski Theorem (D.-Shi)

In  $\prod_{n \in \omega} \kappa_n$  with the EP-topology every  $\lambda$ -comeagre set contains a  $\lambda$ -perfect set.

### Conjecture

All the results in classical descriptive set theory that depend only on Baire category can be generalized to this setting

### (Generalized) Mycielski Theorem (D.-Shi)

In  $\prod_{n \in \omega} \kappa_n$  with the EP-topology every  $\lambda$ -comeagre set contains a  $\lambda$ -perfect set.

### Conjecture

All the results in classical descriptive set theory that depend only on Baire category can be generalized to this setting.

Test case

## (Generalized) Mycielski Theorem (D.-Shi)

In  $\prod_{n \in \omega} \kappa_n$  with the EP-topology every  $\lambda$ -comeagre set contains a  $\lambda$ -perfect set.

## Conjecture

All the results in classical descriptive set theory that depend only on Baire category can be generalized to this setting.

Test case:

## Kuratowski-Ulam Theorem

Let  $X, Y$  be second-countable spaces, and  $A \subseteq X \times Y$  with the Baire property. Then  $A$  is meagre iff  $\{x \in X : \{y \in Y : (x, y) \in A\} \text{ is meagre in } Y\}$  is  $\lambda$ -comeagre in  $X$ .

The key lemma to prove the Kuratowski-Ulam Theorem is the following



The key lemma to prove the Kuratowski-Ulam Theorem is the following:

### Lemma

Let  $X, Y$  be second-countable spaces. Then if  $A \subseteq X \times Y$  is open dense,  $\{x \in X : \{y \in Y : (x, y) \in A\} \text{ is open dense in } Y\}$  is comeagre in  $X$

The key lemma to prove the Kuratowski-Ulam Theorem is the following:

### Lemma

Let  $X, Y$  be second-countable spaces. Then if  $A \subseteq X \times Y$  is open dense,  $\{x \in X : \{y \in Y : (x, y) \in A\} \text{ is open dense in } Y\}$  is comeagre in  $X$ .

### Sketch of proof.

For any  $x \in X$ , let  $A_x = \{y \in Y : (x, y) \in A\}$ , and let  $\langle V_n : n \in \omega \rangle$  be a countable base

The key lemma to prove the Kuratowski-Ulam Theorem is the following:

### Lemma

Let  $X, Y$  be second-countable spaces. Then if  $A \subseteq X \times Y$  is open dense,  $\{x \in X : \{y \in Y : (x, y) \in A\} \text{ is open dense in } Y\}$  is comeagre in  $X$ .

### Sketch of proof.

For any  $x \in X$ , let  $A_x = \{y \in Y : (x, y) \in A\}$ , and let  $\langle V_n : n \in \omega \rangle$  be a countable base. Then  $A_x$  is open, and  $A_x$  is dense iff  $\forall n \in \omega \ A_x \cap V_n \neq \emptyset$

The key lemma to prove the Kuratowski-Ulam Theorem is the following:

### Lemma

Let  $X, Y$  be second-countable spaces. Then if  $A \subseteq X \times Y$  is open dense,  $\{x \in X : \{y \in Y : (x, y) \in A\} \text{ is open dense in } Y\}$  is comeagre in  $X$ .

### Sketch of proof.

For any  $x \in X$ , let  $A_x = \{y \in Y : (x, y) \in A\}$ , and let  $\langle V_n : n \in \omega \rangle$  be a countable base. Then  $A_x$  is open, and  $A_x$  is dense iff  $\forall n \in \omega \ A_x \cap V_n \neq \emptyset$ . But then  $\{x \in X : A_x \text{ is open dense}\} = \bigcap_{n \in \omega} \{x \in X : A_x \cap V_n \neq \emptyset\}$ , a countable intersection of open dense sets, so comeagre.  $\square$

We can see why this proof cannot be generalized

We can see why this proof cannot be generalized:

$\prod_{n \in \omega} \kappa_n$ , with the EP-topology, has a base of cardinality  $2^\lambda$

We can see why this proof cannot be generalized:

$\prod_{n \in \omega} \kappa_n$ , with the EP-topology, has a base of cardinality  $2^\lambda$ , so the set we want to be  $\lambda$ -comeagre is actually the intersection of  $2^\lambda$ -many open dense sets, not  $\lambda$ -many

We can see why this proof cannot be generalized:

$\prod_{n \in \omega} \kappa_n$ , with the EP-topology, has a base of cardinality  $2^\lambda$ , so the set we want to be  $\lambda$ -comeagre is actually the intersection of  $2^\lambda$ -many open dense sets, not  $\lambda$ -many.

The key is still the Strong Prikry condition, in this more general definition



We can see why this proof cannot be generalized:

$\prod_{n \in \omega} \kappa_n$ , with the EP-topology, has a base of cardinality  $2^\lambda$ , so the set we want to be  $\lambda$ -comeagre is actually the intersection of  $2^\lambda$ -many open dense sets, not  $\lambda$ -many.

The key is still the Strong Prikry condition, in this more general definition:

### Strong Prikry condition

Let  $A \subseteq \mathbb{P}_{\vec{\mu}}$  be an *open* set

We can see why this proof cannot be generalized:

$\prod_{n \in \omega} \kappa_n$ , with the EP-topology, has a base of cardinality  $2^\lambda$ , so the set we want to be  $\lambda$ -comeagre is actually the intersection of  $2^\lambda$ -many open dense sets, not  $\lambda$ -many.

The key is still the Strong Prikry condition, in this more general definition:

### Strong Prikry condition

Let  $A \subseteq \mathbb{P}_{\vec{\mu}}$  be an *open* set. Then for any  $p \in \mathbb{P}_{\vec{\mu}}$ , there is a  $p^A \leq^* p$  such that if there is a  $q \leq p^A$  with  $q \in A$  with stem of length  $n$ , then for every  $q \leq p^A$  with stem at least  $n$ ,  $q \in A$ .

We can see why this proof cannot be generalized:

$\prod_{n \in \omega} \kappa_n$ , with the EP-topology, has a base of cardinality  $2^\lambda$ , so the set we want to be  $\lambda$ -comeagre is actually the intersection of  $2^\lambda$ -many open dense sets, not  $\lambda$ -many.

The key is still the Strong Prikry condition, in this more general definition:

### Strong Prikry condition

Let  $A \subseteq \mathbb{P}_{\vec{\mu}}$  be an *open* set. Then for any  $p \in \mathbb{P}_{\vec{\mu}}$ , there is a  $p^A \leq^* p$  such that if there is a  $q \leq p^A$  with  $q \in A$  with stem of length  $n$ , then for every  $q \leq p^A$  with stem at least  $n$ ,  $q \in A$ .

Topologically: Let  $U \subseteq \prod_{n \in \omega} \kappa_n$  be an open set. Then for any  $p \in \mathbb{P}_{\vec{\mu}}$ , there is a  $p^A \leq^* p$  such that either  $O_{p^A} \subseteq A$ , or  $O_{p^A} \cap A = \emptyset$ .

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , let  $1_s = s^\frown \langle \kappa_{m+1}, \kappa_{m+2}, \dots \rangle$

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , let  $1_s = s^\frown \langle \kappa_{m+1}, \kappa_{m+2}, \dots \rangle$ .

Let  $A \subseteq \prod_{n \in \omega} \kappa_n$  be open

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , let  $1_s = s^\frown \langle \kappa_{m+1}, \kappa_{m+2}, \dots \rangle$ .

Let  $A \subseteq \prod_{n \in \omega} \kappa_n$  be open.

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , fix  $1_s^A$  as in the Strong Prikry condition

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , let  $1_s = s^\frown \langle \kappa_{m+1}, \kappa_{m+2}, \dots \rangle$ .

Let  $A \subseteq \prod_{n \in \omega} \kappa_n$  be open.

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , fix  $1_s^A$  as in the Strong Prikry condition.

Then  $A$  is dense iff for all  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$  there is a  $q \leq 1_s^A$  such that  $q \in A$

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , let  $1_s = s^\frown \langle \kappa_{m+1}, \kappa_{m+2}, \dots \rangle$ .

Let  $A \subseteq \prod_{n \in \omega} \kappa_n$  be open.

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , fix  $1_s^A$  as in the Strong Prikry condition.

Then  $A$  is dense iff for all  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$  there is a  $q \leq 1_s^A$  such that  $q \in A$ .

So to test that  $A$  open is dense, we do not need to test it for all the basic open sets, just for a subfamily of them of size  $\lambda$ !



For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , let  $1_s = s^\frown \langle \kappa_{m+1}, \kappa_{m+2}, \dots \rangle$ .

Let  $A \subseteq \prod_{n \in \omega} \kappa_n$  be open.

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , fix  $1_s^A$  as in the Strong Prikry condition.

Then  $A$  is dense iff for all  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$  there is a  $q \leq 1_s^A$  such that  $q \in A$ .

So to test that  $A$  open is dense, we do not need to test it for all the basic open sets, just for a subfamily of them of size  $\lambda$ !

### (Generalized) Kuratowski-Ulam Theorem (D.)

Let  $A \subseteq \prod_{n \in \omega} \kappa_n \times \prod_{n \in \omega} \kappa_n$  be with the  $\lambda$ -Baire property

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , let  $1_s = s^\frown \langle \kappa_{m+1}, \kappa_{m+2}, \dots \rangle$ .

Let  $A \subseteq \prod_{n \in \omega} \kappa_n$  be open.

For any  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$ , fix  $1_s^A$  as in the Strong Prikry condition.

Then  $A$  is dense iff for all  $s \in \bigcup_{m \in \omega} \prod_{n \leq m} \kappa_n$  there is a  $q \leq 1_s^A$  such that  $q \in A$ .

So to test that  $A$  open is dense, we do not need to test it for all the basic open sets, just for a subfamily of them of size  $\lambda$ !

### (Generalized) Kuratowski-Ulam Theorem (D.)

Let  $A \subseteq \prod_{n \in \omega} \kappa_n \times \prod_{n \in \omega} \kappa_n$  be with the  $\lambda$ -Baire property. Then  $A$  is  $\lambda$ -meagre iff  $\{x \in X : A_x \text{ is } \lambda\text{-meagre}\}$  is  $\lambda$ -comeagre.

## Question

Are measurable cardinals necessary?

### Question

Are measurable cardinals necessary?

### Question

What other forcings there are on  $\lambda$  that can generate interesting concepts?

### Question

Are measurable cardinals necessary?

### Question

What other forcings there are on  $\lambda$  that can generate interesting concepts?

### Question

What about  $\lambda$ -universally Baire sets?

Thanks for watching