

# A descriptive main gap theorem

Luca Motto Ros

Department of Mathematics “G. Peano”  
University of Turin, Italy  
`luca.mottoros@unito.it`  
<https://sites.google.com/site/lucamottoros/>

Fifth Workshop on Generalised Baire Spaces  
University of Bristol, 3–4.2.2020

Joint work with Francesco Mangraviti

# The spectrum problem

Given a complete countable first-order theory  $T$ , the *spectrum problem* for  $T$  consists in determining the number

$$I(\kappa, T)$$

of isomorphism types of models of  $T$  of size  $\kappa \geq \aleph_1$ .

# The spectrum problem

Given a complete countable first-order theory  $T$ , the *spectrum problem* for  $T$  consists in determining the number

$$I(\kappa, T)$$

of isomorphism types of models of  $T$  of size  $\kappa \geq \aleph_1$ . Assuming that  $T$  has an infinite model, the obvious bounds are

$$1 \leq I(\kappa, T) \leq 2^\kappa.$$

# The spectrum problem

Given a complete countable first-order theory  $T$ , the *spectrum problem* for  $T$  consists in determining the number

$$I(\kappa, T)$$

of isomorphism types of models of  $T$  of size  $\kappa \geq \aleph_1$ . Assuming that  $T$  has an infinite model, the obvious bounds are

$$1 \leq I(\kappa, T) \leq 2^\kappa.$$

**Remark:** The restriction to uncountable models is because we cannot (yet) deal with the countable case: the Vaught's conjecture, asserting that either  $I(\aleph_0, T) \leq \omega$  or  $I(\aleph_0, T) = 2^{\aleph_0}$ , is still widely open.

# The spectrum problem

Given a complete countable first-order theory  $T$ , the *spectrum problem* for  $T$  consists in determining the number

$$I(\kappa, T)$$

of isomorphism types of models of  $T$  of size  $\kappa \geq \aleph_1$ . Assuming that  $T$  has an infinite model, the obvious bounds are

$$1 \leq I(\kappa, T) \leq 2^\kappa.$$

**Remark:** The restriction to uncountable models is because we cannot (yet) deal with the countable case: the Vaught's conjecture, asserting that either  $I(\aleph_0, T) \leq \omega$  or  $I(\aleph_0, T) = 2^{\aleph_0}$ , is still widely open.

## Main question

Given  $T$ , can one provide non-trivial lower/upper bounds for the spectrum function  $I(\kappa, T)$ ?

# Morley's theorem and classification theory

Recall that a theory  $T$  is  $\kappa$ -categorical if  $I(\kappa, T) = 1$ .

# Morley's theorem and classification theory

Recall that a theory  $T$  is  $\kappa$ -categorical if  $I(\kappa, T) = 1$ .

## Theorem (Morley)

$T$  is  $\kappa$ -categorical for *some* uncountable  $\kappa$  if and only if  $T$  is  $\kappa$ -categorical for *all* uncountable  $\kappa$ .

# Morley's theorem and classification theory

Recall that a theory  $T$  is  $\kappa$ -categorical if  $I(\kappa, T) = 1$ .

## Theorem (Morley)

$T$  is  $\kappa$ -categorical for *some* uncountable  $\kappa$  if and only if  $T$  is  $\kappa$ -categorical for *all* uncountable  $\kappa$ .

What for other theories  $T$ ? To answer that question, Shelah isolated a number of dividing lines for complete countable first-order theories:



# Morley's theorem and classification theory

Recall that a theory  $T$  is  $\kappa$ -categorical if  $I(\kappa, T) = 1$ .

## Theorem (Morley)

$T$  is  $\kappa$ -categorical for *some* uncountable  $\kappa$  if and only if  $T$  is  $\kappa$ -categorical for *all* uncountable  $\kappa$ .

What for other theories  $T$ ? To answer that question, Shelah isolated a number of dividing lines for complete countable first-order theories:

- stable vs unstable

# Morley's theorem and classification theory

Recall that a theory  $T$  is  $\kappa$ -categorical if  $I(\kappa, T) = 1$ .

## Theorem (Morley)

$T$  is  $\kappa$ -categorical for *some* uncountable  $\kappa$  if and only if  $T$  is  $\kappa$ -categorical for *all* uncountable  $\kappa$ .

What for other theories  $T$ ? To answer that question, Shelah isolated a number of dividing lines for complete countable first-order theories:

- stable vs unstable
- superstable vs unsuperstable

# Morley's theorem and classification theory

Recall that a theory  $T$  is  $\kappa$ -categorical if  $I(\kappa, T) = 1$ .

## Theorem (Morley)

$T$  is  $\kappa$ -categorical for *some* uncountable  $\kappa$  if and only if  $T$  is  $\kappa$ -categorical for *all* uncountable  $\kappa$ .

What for other theories  $T$ ? To answer that question, Shelah isolated a number of dividing lines for complete countable first-order theories:

- stable vs unstable
- superstable vs unsuperstable
- NDOP vs DOP (dimensional order property)

# Morley's theorem and classification theory

Recall that a theory  $T$  is  $\kappa$ -categorical if  $I(\kappa, T) = 1$ .

## Theorem (Morley)

$T$  is  $\kappa$ -categorical for *some* uncountable  $\kappa$  if and only if  $T$  is  $\kappa$ -categorical for *all* uncountable  $\kappa$ .

What for other theories  $T$ ? To answer that question, Shelah isolated a number of dividing lines for complete countable first-order theories:

- stable vs unstable
- superstable vs unsuperstable
- NDOP vs DOP (dimensional order property)
- NOTOP vs OTOP (omitting types order property)

# Morley's theorem and classification theory

Recall that a theory  $T$  is  $\kappa$ -categorical if  $I(\kappa, T) = 1$ .

## Theorem (Morley)

$T$  is  $\kappa$ -categorical for *some* uncountable  $\kappa$  if and only if  $T$  is  $\kappa$ -categorical for *all* uncountable  $\kappa$ .

What for other theories  $T$ ? To answer that question, Shelah isolated a number of dividing lines for complete countable first-order theories:

- stable vs unstable
- superstable vs unsuperstable
- NDOP vs DOP (dimensional order property)
- NOTOP vs OTOP (omitting types order property)
- shallow vs deep

# Morley's theorem and classification theory

Recall that a theory  $T$  is  $\kappa$ -categorical if  $I(\kappa, T) = 1$ .

## Theorem (Morley)

$T$  is  $\kappa$ -categorical for *some* uncountable  $\kappa$  if and only if  $T$  is  $\kappa$ -categorical for *all* uncountable  $\kappa$ .

What for other theories  $T$ ? To answer that question, Shelah isolated a number of dividing lines for complete countable first-order theories:

- stable vs unstable
- superstable vs unsuperstable
- NDOP vs DOP (dimensional order property)
- NOTOP vs OTOP (omitting types order property)
- shallow vs deep
- ...

# Morley's theorem and classification theory

Recall that a theory  $T$  is  $\kappa$ -categorical if  $I(\kappa, T) = 1$ .

## Theorem (Morley)

$T$  is  $\kappa$ -categorical for *some* uncountable  $\kappa$  if and only if  $T$  is  $\kappa$ -categorical for *all* uncountable  $\kappa$ .

What for other theories  $T$ ? To answer that question, Shelah isolated a number of dividing lines for complete countable first-order theories:

- stable vs unstable
- superstable vs unsuperstable
- NDOP vs DOP (dimensional order property)
- NOTOP vs OTOP (omitting types order property)
- shallow vs deep
- ...

This gave birth to a beautiful branch of model theory, later called  
**stability theory.**

Shelah's proved that if  $T$  has one of the following properties

- unsuperstable
- DOP (dimensional order property, defined only for superstable theories)
- OTOP (omitting types order property)



Shelah's proved that if  $T$  has one of the following properties

- unsuperstable
- DOP (dimensional order property, defined only for superstable theories)
- OTOP (omitting types order property)

then  $I(\kappa, T) = 2^\kappa$  for every  $\kappa \geq \aleph_1$ .

Shelah's proved that if  $T$  has one of the following properties

- unsuperstable
- DOP (dimensional order property, defined only for superstable theories)
- OTOP (omitting types order property)

then  $I(\kappa, T) = 2^\kappa$  for every  $\kappa \geq \aleph_1$ .

So if the function  $I(\kappa, T)$  is to have a non-trivial upper bound, then  $T$  must be (stable) superstable, NDOP and NOTOP: such theories are briefly called **classifiable**.

# Classification theory for classifiable theories

# Classification theory for classifiable theories

(John T. Baldwin, *Fundamentals of Stability Theory*)

# Classification theory for classifiable theories

*The solution of the spectrum problem for classifiable theories depends upon a key construction which assigns to each model of size  $\kappa$  a skeleton of submodels. Each submodel has cardinality at most  $2^{\aleph_0}$ , and the skeleton is partially ordered by the natural tree order on a subset of  $\kappa^{<\omega}$ .*

(John T. Baldwin, *Fundamentals of Stability Theory*)

# Classification theory for classifiable theories

*The solution of the spectrum problem for classifiable theories depends upon a key construction which assigns to each model of size  $\kappa$  a skeleton of submodels. Each submodel has cardinality at most  $2^{\aleph_0}$ , and the skeleton is partially ordered by the natural tree order on a subset of  $\kappa^{<\omega}$ . The isomorphism type of the model is determined by the small submodels and this partial ordering.*

(John T. Baldwin, *Fundamentals of Stability Theory*)

# Classification theory for classifiable theories

*The solution of the spectrum problem for classifiable theories depends upon a key construction which assigns to each model of size  $\kappa$  a skeleton of submodels. Each submodel has cardinality at most  $2^{\aleph_0}$ , and the skeleton is partially ordered by the natural tree order on a subset of  $\kappa^{<\omega}$ . The isomorphism type of the model is determined by the small submodels and this partial ordering.*

(John T. Baldwin, *Fundamentals of Stability Theory*)

Using the above decomposition, to each classifiable theory  $T$  one can assign a rank  $\text{dp}(T)$ , called **depth of  $T$** , a priori taking as value either an ordinal  $\leq \kappa^+$  or  $\infty$ ;

# Classification theory for classifiable theories

*The solution of the spectrum problem for classifiable theories depends upon a key construction which assigns to each model of size  $\kappa$  a skeleton of submodels. Each submodel has cardinality at most  $2^{\aleph_0}$ , and the skeleton is partially ordered by the natural tree order on a subset of  $\kappa^{<\omega}$ . The isomorphism type of the model is determined by the small submodels and this partial ordering.*

(John T. Baldwin, *Fundamentals of Stability Theory*)

Using the above decomposition, to each classifiable theory  $T$  one can assign a rank  $\text{dp}(T)$ , called **depth of  $T$** , *a priori* taking as value either an ordinal  $\leq \kappa^+$  or  $\infty$ ; however, Lascar proved that  $\text{dp}(T) \in \omega_1 \cup \{\infty\}$ .



# Classification theory for classifiable theories

*The solution of the spectrum problem for classifiable theories depends upon a key construction which assigns to each model of size  $\kappa$  a skeleton of submodels. Each submodel has cardinality at most  $2^{\aleph_0}$ , and the skeleton is partially ordered by the natural tree order on a subset of  $\kappa^{<\omega}$ . The isomorphism type of the model is determined by the small submodels and this partial ordering.*

(John T. Baldwin, *Fundamentals of Stability Theory*)

Using the above decomposition, to each classifiable theory  $T$  one can assign a rank  $\text{dp}(T)$ , called **depth of  $T$** , *a priori* taking as value either an ordinal  $\leq \kappa^+$  or  $\infty$ ; however, Lascar proved that  $\text{dp}(T) \in \omega_1 \cup \{\infty\}$ .

A classifiable theory  $T$  is called

# Classification theory for classifiable theories

*The solution of the spectrum problem for classifiable theories depends upon a key construction which assigns to each model of size  $\kappa$  a skeleton of submodels. Each submodel has cardinality at most  $2^{\aleph_0}$ , and the skeleton is partially ordered by the natural tree order on a subset of  $\kappa^{<\omega}$ . The isomorphism type of the model is determined by the small submodels and this partial ordering.*

(John T. Baldwin, *Fundamentals of Stability Theory*)

Using the above decomposition, to each classifiable theory  $T$  one can assign a rank  $\text{dp}(T)$ , called **depth of  $T$** , *a priori* taking as value either an ordinal  $\leq \kappa^+$  or  $\infty$ ; however, Lascar proved that  $\text{dp}(T) \in \omega_1 \cup \{\infty\}$ .

A classifiable theory  $T$  is called

- **shallow** if  $\text{dp}(T) \neq \infty$ , in which case  $\text{dp}(T) < \aleph_1$ ;

# Classification theory for classifiable theories

*The solution of the spectrum problem for classifiable theories depends upon a key construction which assigns to each model of size  $\kappa$  a skeleton of submodels. Each submodel has cardinality at most  $2^{\aleph_0}$ , and the skeleton is partially ordered by the natural tree order on a subset of  $\kappa^{<\omega}$ . The isomorphism type of the model is determined by the small submodels and this partial ordering.*

(John T. Baldwin, *Fundamentals of Stability Theory*)

Using the above decomposition, to each classifiable theory  $T$  one can assign a rank  $\text{dp}(T)$ , called **depth of  $T$** , *a priori* taking as value either an ordinal  $\leq \kappa^+$  or  $\infty$ ; however, Lascar proved that  $\text{dp}(T) \in \omega_1 \cup \{\infty\}$ .

A classifiable theory  $T$  is called

- **shallow** if  $\text{dp}(T) \neq \infty$ , in which case  $\text{dp}(T) < \aleph_1$ ;
- **deep** otherwise.

# Shelah's Main Gap Theorem

## Theorem (Shelah)

Let  $\kappa \geq \aleph_1$  be the  $\gamma$ -th cardinal.

# Shelah's Main Gap Theorem

## Theorem (Shelah)

Let  $\kappa \geq \aleph_1$  be the  $\gamma$ -th cardinal.

- ① If  $T$  is classifiable shallow of depth  $\alpha$  , then

$$I(\kappa, T) \leq \beth_\alpha \left( |\gamma|^{2^{\aleph_0}} \right)$$

# Shelah's Main Gap Theorem

## Theorem (Shelah)

Let  $\kappa \geq \aleph_1$  be the  $\gamma$ -th cardinal.

- ① If  $T$  is classifiable shallow of depth  $\alpha$  (necessarily,  $\alpha < \omega_1$ ), then

$$I(\kappa, T) \leq \beth_\alpha(|\gamma|^{2^{\aleph_0}}) < \beth_{\omega_1}(|\gamma|^{2^{\aleph_0}}).$$

# Shelah's Main Gap Theorem

## Theorem (Shelah)

Let  $\kappa \geq \aleph_1$  be the  $\gamma$ -th cardinal.

- ① If  $T$  is classifiable shallow of depth  $\alpha$  (necessarily,  $\alpha < \omega_1$ ), then

$$I(\kappa, T) \leq \beth_\alpha(|\gamma|^{2^{\aleph_0}}) < \beth_{\omega_1}(|\gamma|^{2^{\aleph_0}}).$$

- ② If  $T$  is not classifiable shallow, then

$$I(\kappa, T) = 2^\kappa.$$

# Shelah's Main Gap Theorem

## Theorem (Shelah)

Let  $\kappa \geq \aleph_1$  be the  $\gamma$ -th cardinal.

- ① If  $T$  is classifiable shallow of depth  $\alpha$  (necessarily,  $\alpha < \omega_1$ ), then

$$I(\kappa, T) \leq \beth_\alpha(|\gamma|^{2^{\aleph_0}}) < \beth_{\omega_1}(|\gamma|^{2^{\aleph_0}}).$$

- ② If  $T$  is not classifiable shallow, then

$$I(\kappa, T) = 2^\kappa.$$

**Remark.** The upper bound in ① may trivialize (e.g. when  $\kappa$  is a fixed point of the  $\aleph$ -function),



# Shelah's Main Gap Theorem

## Theorem (Shelah)

Let  $\kappa \geq \aleph_1$  be the  $\gamma$ -th cardinal.

- ① If  $T$  is classifiable shallow of depth  $\alpha$  (necessarily,  $\alpha < \omega_1$ ), then

$$I(\kappa, T) \leq \beth_\alpha(|\gamma|^{2^{\aleph_0}}) < \beth_{\omega_1}(|\gamma|^{2^{\aleph_0}}).$$

- ② If  $T$  is not classifiable shallow, then

$$I(\kappa, T) = 2^\kappa.$$

**Remark.** The upper bound in ① may trivialize (e.g. when  $\kappa$  is a fixed point of the  $\aleph$ -function), but in general it is easy to find cardinals for which this is not the case: for example, under GCH there are unboundedly many  $\kappa$  for which such upper bound is  $< 2^\kappa$ , or even  $\leq \kappa$ .

$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow
$I(\kappa, T) = 1$	$I(\kappa, T) < \beth_\alpha( \gamma ^{2^{\aleph_0}})$	$I(\kappa, T) = 2^\kappa$

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

# Complexity of the isomorphism relation

Let's take another viewpoint.

# Complexity of the isomorphism relation

Let's take another viewpoint.

Intuitively, one may say that the isomorphism relation on  $\kappa$ -sized models of  $T$  is “*simple*” when there is a reasonable way to decide, given any two  $\kappa$ -sized models of  $T$ , whether they are isomorphic or not;

# Complexity of the isomorphism relation

Let's take another viewpoint.

Intuitively, one may say that the isomorphism relation on  $\kappa$ -sized models of  $T$  is “*simple*” when there is a reasonable way to decide, given any two  $\kappa$ -sized models of  $T$ , whether they are isomorphic or not; otherwise it is “*complicated*”.

# Complexity of the isomorphism relation

Let's take another viewpoint.

Intuitively, one may say that the isomorphism relation on  $\kappa$ -sized models of  $T$  is “*simple*” when there is a reasonable way to decide, given any two  $\kappa$ -sized models of  $T$ , whether they are isomorphic or not; otherwise it is “*complicated*”. In the former case, the  $\kappa$ -sized models of  $T$  can be regarded as “easily classifiable” up to isomorphism.

# Complexity of the isomorphism relation

Let's take another viewpoint.

Intuitively, one may say that the isomorphism relation on  $\kappa$ -sized models of  $T$  is “*simple*” when there is a reasonable way to decide, given any two  $\kappa$ -sized models of  $T$ , whether they are isomorphic or not; otherwise it is “*complicated*”. In the former case, the  $\kappa$ -sized models of  $T$  can be regarded as “easily classifiable” up to isomorphism.

But what should “simple” and “reasonable” mean, mathematically?



# Complexity of the isomorphism relation

Let's take another viewpoint.

Intuitively, one may say that the isomorphism relation on  $\kappa$ -sized models of  $T$  is “*simple*” when there is a reasonable way to decide, given any two  $\kappa$ -sized models of  $T$ , whether they are isomorphic or not; otherwise it is “*complicated*”. In the former case, the  $\kappa$ -sized models of  $T$  can be regarded as “easily classifiable” up to isomorphism.

But what should “simple” and “reasonable” mean, mathematically?

Here is where generalized Descriptive Set Theory enters the scene...

# Complexity of the isomorphism relation

Given a countable complete first-order theory  $T$  in the relational countable vocabulary  $\mathcal{L}$ , let

$$\text{Mod}_T^\kappa$$

be the collection of models of  $T$  with *domain*  $\kappa$ .

# Complexity of the isomorphism relation

Given a countable complete first-order theory  $T$  in the relational countable vocabulary  $\mathcal{L}$ , let

$$\text{Mod}_T^\kappa$$

be the collection of models of  $T$  with *domain*  $\kappa$ . By the generalized Lopez-Escobar theorem,  $\text{Mod}_T^\kappa$  can be identified up to a natural coding with a  $(\kappa^+)$ -Borel subset of the generalized Cantor space  ${}^\kappa 2$ ,

# Complexity of the isomorphism relation

Given a countable complete first-order theory  $T$  in the relational countable vocabulary  $\mathcal{L}$ , let

$$\text{Mod}_T^\kappa$$

be the collection of models of  $T$  with *domain*  $\kappa$ . By the generalized Lopez-Escobar theorem,  $\text{Mod}_T^\kappa$  can be identified up to a natural coding with a  $(\kappa^+)$ -Borel subset of the generalized Cantor space  ${}^\kappa 2$ , and the isomorphism relation

$$\cong_T^\kappa$$

on  $\text{Mod}_T^\kappa$  is a  $\kappa$ -analytic equivalence relation on it.

# Complexity of the isomorphism relation

Given a countable complete first-order theory  $T$  in the relational countable vocabulary  $\mathcal{L}$ , let

$$\text{Mod}_T^\kappa$$

be the collection of models of  $T$  with *domain*  $\kappa$ . By the generalized Lopez-Escobar theorem,  $\text{Mod}_T^\kappa$  can be identified up to a natural coding with a  $(\kappa^+)$ -Borel subset of the generalized Cantor space  ${}^\kappa 2$ , and the isomorphism relation

$$\cong_T^\kappa$$

on  $\text{Mod}_T^\kappa$  is a  $\kappa$ -analytic equivalence relation on it.

Within this framework, we can say that

$$\cong_T^\kappa \text{ is “simple” if it is a } (\kappa^+)\text{-Borel subset of } (\text{Mod}_T^\kappa)^2,$$

because in this case we have a “Borel” procedure to decide whether two models are isomorphic or not.

## Main questions

For which *uncountable* cardinals  $\kappa$  and which countable complete first-order theories  $T$  is  $\cong_T^\kappa$  “simple” (= Borel)?

## Main questions

For which *uncountable* cardinals  $\kappa$  and which countable complete first-order theories  $T$  is  $\cong_T^\kappa$  “simple” (= Borel)?

Does the Borelness of  $\cong_T^\kappa$  depend on both parameters, or does it just depend on the theory  $T$ ?

(In the latter case, we can regard  $T$  itself as “simple” if some/any  $\cong_T^\kappa$  is Borel.)

## Main questions

For which *uncountable* cardinals  $\kappa$  and which countable complete first-order theories  $T$  is  $\cong_T^\kappa$  “simple” (= Borel)?

Does the Borelness of  $\cong_T^\kappa$  depend on both parameters, or does it just depend on the theory  $T$ ?

(In the latter case, we can regard  $T$  itself as “simple” if some/any  $\cong_T^\kappa$  is Borel.)

## Theorem (S.D. Friedman-Hyttinen-Kulikov)



## Main questions

For which *uncountable* cardinals  $\kappa$  and which countable complete first-order theories  $T$  is  $\cong_T^\kappa$  “simple” (= Borel)?

Does the Borelness of  $\cong_T^\kappa$  depend on both parameters, or does it just depend on the theory  $T$ ?

(In the latter case, we can regard  $T$  itself as “simple” if some/any  $\cong_T^\kappa$  is Borel.)

## Theorem (S.D. Friedman-Hyttinen-Kulikov)

Let  $T$  be a countable complete first order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

## Main questions

For which *uncountable* cardinals  $\kappa$  and which countable complete first-order theories  $T$  is  $\cong_T^\kappa$  “simple” (= Borel)?

Does the Borelness of  $\cong_T^\kappa$  depend on both parameters, or does it just depend on the theory  $T$ ?

(In the latter case, we can regard  $T$  itself as “simple” if some/any  $\cong_T^\kappa$  is Borel.)

## Theorem (S.D. Friedman-Hyttinen-Kulikov)

Let  $T$  be a countable complete first order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- 1 If  $T$  is classifiable shallow, then  $\cong_T^\kappa$  is Borel.

## Main questions

For which *uncountable* cardinals  $\kappa$  and which countable complete first-order theories  $T$  is  $\cong_T^\kappa$  “simple” (= Borel)?

Does the Borelness of  $\cong_T^\kappa$  depend on both parameters, or does it just depend on the theory  $T$ ?

(In the latter case, we can regard  $T$  itself as “simple” if some/any  $\cong_T^\kappa$  is Borel.)

## Theorem (S.D. Friedman-Hyttinen-Kulikov)

Let  $T$  be a countable complete first order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- 1 If  $T$  is classifiable shallow, then  $\cong_T^\kappa$  is Borel.
- 2 If  $T$  is not classifiable shallow, then  $\cong_T^\kappa$  is not Borel.

$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow
$I(\kappa, T) = 1$	$I(\kappa, T) < \beth_\alpha( \gamma ^{2^{\aleph_0}})$	$I(\kappa, T) = 2^\kappa$

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow
$I(\kappa, T) = 1$	$I(\kappa, T) < \beth_\alpha( \gamma ^{2^{\aleph_0}})$	$I(\kappa, T) = 2^\kappa$
	$(\kappa^{+-})\text{Borel}$	Not $(\kappa^{+-})\text{Borel}$

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

Borel sets can be stratified according to the usual definition:

$\Sigma_1^0$  = open sets

$\Pi_1^0$  = closed sets

$$\Sigma_\alpha^0 = \left\{ \bigcup_{\gamma < \kappa} A_\gamma \mid A_\gamma \in \bigcup_{1 \leq \beta < \alpha} \Pi_\beta^0 \right\}$$

$$\Pi_\alpha^0 = \{ {}^\kappa\kappa \setminus A \mid A \in \Sigma_\alpha^0 \}$$

Borel sets can be stratified according to the usual definition:

$\Sigma_1^0$  = open sets

$\Pi_1^0$  = closed sets

$$\Sigma_\alpha^0 = \left\{ \bigcup_{\gamma < \kappa} A_\gamma \mid A_\gamma \in \bigcup_{1 \leq \beta < \alpha} \Pi_\beta^0 \right\}$$

$$\Pi_\alpha^0 = \{ {}^\kappa \kappa \setminus A \mid A \in \Sigma_\alpha^0 \}$$

If  $A$  is Borel, the smallest ordinal  $\alpha < \kappa^+$  for which  $A \in \Sigma_\alpha^0 \cup \Pi_\alpha^0$  is called the **Borel rank** of  $A$ , and denoted by  $\mathbf{rk}_B(A)$ .

Borel sets can be stratified according to the usual definition:

$\Sigma_1^0$  = open sets

$\Pi_1^0$  = closed sets

$$\Sigma_\alpha^0 = \left\{ \bigcup_{\gamma < \kappa} A_\gamma \mid A_\gamma \in \bigcup_{1 \leq \beta < \alpha} \Pi_\beta^0 \right\}$$

$$\Pi_\alpha^0 = \{ {}^\kappa\kappa \setminus A \mid A \in \Sigma_\alpha^0 \}$$

If  $A$  is Borel, the smallest ordinal  $\alpha < \kappa^+$  for which  $A \in \Sigma_\alpha^0 \cup \Pi_\alpha^0$  is called the **Borel rank** of  $A$ , and denoted by  $\text{rk}_B(A)$ . We stipulate that  $\text{rk}_B(A) = \infty$  when  $A$  is not Borel.



Borel sets can be stratified according to the usual definition:

$\Sigma_1^0$  = open sets

$\Pi_1^0$  = closed sets

$$\Sigma_\alpha^0 = \left\{ \bigcup_{\gamma < \kappa} A_\gamma \mid A_\gamma \in \bigcup_{1 \leq \beta < \alpha} \Pi_\beta^0 \right\}$$

$$\Pi_\alpha^0 = \{ {}^\kappa\kappa \setminus A \mid A \in \Sigma_\alpha^0 \}$$

If  $A$  is Borel, the smallest ordinal  $\alpha < \kappa^+$  for which  $A \in \Sigma_\alpha^0 \cup \Pi_\alpha^0$  is called the **Borel rank** of  $A$ , and denoted by  $\text{rk}_B(A)$ . We stipulate that  $\text{rk}_B(A) = \infty$  when  $A$  is not Borel.

## Question (F-H-K)

If  $T$  is classifiable shallow, what is the Borel rank of  $\cong_T^\kappa$ ? Is it related to the depth of  $T$ ?

# A descriptive version of the Main Gap Theorem

A detailed analysis of Friedman-Hyttinen-Kulikov proof reveals the following refinement of their theorem.

# A descriptive version of the Main Gap Theorem

A detailed analysis of Friedman-Hyttinen-Kulikov proof reveals the following refinement of their theorem. Given  $T$ , define the Borel-rank spectrum function  $B(\kappa, T)$  by setting

$$B(\kappa, T) = \text{rk}_B(\cong_T^\kappa).$$

# A descriptive version of the Main Gap Theorem

A detailed analysis of Friedman-Hyttinen-Kulikov proof reveals the following refinement of their theorem. Given  $T$ , define the Borel-rank spectrum function  $B(\kappa, T)$  by setting

$$B(\kappa, T) = \text{rk}_B(\cong_T^\kappa).$$

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

# A descriptive version of the Main Gap Theorem

A detailed analysis of Friedman-Hyttinen-Kulikov proof reveals the following refinement of their theorem. Given  $T$ , define the Borel-rank spectrum function  $B(\kappa, T)$  by setting

$$B(\kappa, T) = \text{rk}_B(\cong_T^\kappa).$$

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- 1 If  $T$  is classifiable shallow of depth  $\alpha$ , then  $B(\kappa, T) \leq 4\alpha$ .

# A descriptive version of the Main Gap Theorem

A detailed analysis of Friedman-Hyttinen-Kulikov proof reveals the following refinement of their theorem. Given  $T$ , define the Borel-rank spectrum function  $B(\kappa, T)$  by setting

$$B(\kappa, T) = \text{rk}_B(\cong_T^\kappa).$$

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- 1 If  $T$  is classifiable shallow of depth  $\alpha$ , then  $B(\kappa, T) \leq 4\alpha$ .
- 2 If  $T$  is not classifiable shallow, then  $B(\kappa, T) = \infty$ .

# A descriptive version of the Main Gap Theorem

A detailed analysis of Friedman-Hyttinen-Kulikov proof reveals the following refinement of their theorem. Given  $T$ , define the Borel-rank spectrum function  $B(\kappa, T)$  by setting

$$B(\kappa, T) = \text{rk}_B(\cong_T^\kappa).$$

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- 1 If  $T$  is classifiable shallow of depth  $\alpha$ , then  $B(\kappa, T) \leq 4\alpha$ .
- 2 If  $T$  is not classifiable shallow, then  $B(\kappa, T) = \infty$ .

Thus in the “good” case the function  $B(\kappa, T)$  is almost everywhere dominated by a constant function which (unlike Shelah’s upper bound) depends only on  $\text{dp}(T)$  and not on the cardinal  $\kappa$ .

# A descriptive version of the Main Gap Theorem

(To appreciate the descriptive main gap, recall that the Borel hierarchy does not collapse before  $\kappa^+ \geq \aleph_2$ .)



# A descriptive version of the Main Gap Theorem

(To appreciate the descriptive main gap, recall that the Borel hierarchy does not collapse before  $\kappa^+ \geq \aleph_2$ .)

Since the depth of a classifiable shallow theory is always a countable ordinal, we get

# A descriptive version of the Main Gap Theorem

(To appreciate the descriptive main gap, recall that the Borel hierarchy does not collapse before  $\kappa^+ \geq \aleph_2$ .)

Since the depth of a classifiable shallow theory is always a countable ordinal, we get

## Descriptive Main Gap Theorem

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- 1 If  $T$  is classifiable shallow, then  $B(\kappa, T) \leq 4\text{dp}(T) < \aleph_1$ .
- 2 If  $T$  is not classifiable shallow, then  $B(\kappa, T) = \infty$ .

# A descriptive version of the Main Gap Theorem

(To appreciate the descriptive main gap, recall that the Borel hierarchy does not collapse before  $\kappa^+ \geq \aleph_2$ .)

Since the depth of a classifiable shallow theory is always a countable ordinal, we get

## Descriptive Main Gap Theorem

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- 1 If  $T$  is classifiable shallow, then  $B(\kappa, T) \leq 4\text{dp}(T) < \aleph_1$ .
- 2 If  $T$  is not classifiable shallow, then  $B(\kappa, T) = \infty$ .

**Remark.** This gap theorem, unlike Shelah's, is never trivial for the relevant  $\kappa$ 's: in particular, under GCH the descriptive gap is non-trivial for every successor cardinal  $\kappa \geq \aleph_2$ .

$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow
$I(\kappa, T) = 1$	$I(\kappa, T) < \beth_\alpha( \gamma ^{2^{\aleph_0}})$	$I(\kappa, T) = 2^\kappa$
	$(\kappa^{+-})\text{Borel}$	Not $(\kappa^{+-})\text{Borel}$

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow
$I(\kappa, T) = 1$	$I(\kappa, T) < \beth_\alpha( \gamma ^{2^{\aleph_0}})$	$I(\kappa, T) = 2^\kappa$
	$(\kappa^{+-})\text{Borel}$	Not $(\kappa^{+-})\text{Borel}$
	$B(\kappa, T) \leq 4\alpha$	$B(\kappa, T) = \infty$

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

The Descriptive Main Gap Theorem is a consequence of a result connecting  $B(\kappa, T)$  with the Scott height  $S(\kappa, T)$  of the  $\kappa$ -sized models of  $T$ .

The Descriptive Main Gap Theorem is a consequence of a result connecting  $B(\kappa, T)$  with the Scott height  $S(\kappa, T)$  of the  $\kappa$ -sized models of  $T$ .

## Definition

Let  $\kappa$  be an infinite cardinal. Given  $M, N \in \text{Mod}_T^\kappa$  and an ordinal  $\beta$ , set  $M \equiv_\beta N$  if and only if  $M$  and  $N$  satisfy the same  $\mathcal{L}_{\infty\kappa}$ -formulae with quantifier rank  $< \beta$ .

The Descriptive Main Gap Theorem is a consequence of a result connecting  $B(\kappa, T)$  with the Scott height  $S(\kappa, T)$  of the  $\kappa$ -sized models of  $T$ .

## Definition

Let  $\kappa$  be an infinite cardinal. Given  $M, N \in \text{Mod}_T^\kappa$  and an ordinal  $\beta$ , set  $M \equiv_\beta N$  if and only if  $M$  and  $N$  satisfy the same  $\mathcal{L}_{\infty\kappa}$ -formulae with quantifier rank  $< \beta$ .

The  $\mathcal{L}_{\infty\kappa}$ -**Scott height** of  $M \in \text{Mod}_T^\kappa$  is the smallest ordinal  $\beta$ , if it exists, such that  $M \equiv_\beta N \Rightarrow M \cong N$  for every  $N \in \text{Mod}_T^\kappa$ .



The Descriptive Main Gap Theorem is a consequence of a result connecting  $B(\kappa, T)$  with the Scott height  $S(\kappa, T)$  of the  $\kappa$ -sized models of  $T$ .

## Definition

Let  $\kappa$  be an infinite cardinal. Given  $M, N \in \text{Mod}_T^\kappa$  and an ordinal  $\beta$ , set  $M \equiv_\beta N$  if and only if  $M$  and  $N$  satisfy the same  $\mathcal{L}_{\infty\kappa}$ -formulae with quantifier rank  $< \beta$ .

The  $\mathcal{L}_{\infty\kappa}$ -**Scott height** of  $M \in \text{Mod}_T^\kappa$  is the smallest ordinal  $\beta$ , if it exists, such that  $M \equiv_\beta N \Rightarrow M \cong N$  for every  $N \in \text{Mod}_T^\kappa$ . If such a  $\beta$  does not exist, we say that  $M$  has  $\mathcal{L}_{\infty\kappa}$ -Scott height  $\infty$ .

The Descriptive Main Gap Theorem is a consequence of a result connecting  $B(\kappa, T)$  with the Scott height  $S(\kappa, T)$  of the  $\kappa$ -sized models of  $T$ .

## Definition

Let  $\kappa$  be an infinite cardinal. Given  $M, N \in \text{Mod}_T^\kappa$  and an ordinal  $\beta$ , set  $M \equiv_\beta N$  if and only if  $M$  and  $N$  satisfy the same  $\mathcal{L}_{\infty\kappa}$ -formulae with quantifier rank  $< \beta$ .

The  $\mathcal{L}_{\infty\kappa}$ -**Scott height** of  $M \in \text{Mod}_T^\kappa$  is the smallest ordinal  $\beta$ , if it exists, such that  $M \equiv_\beta N \Rightarrow M \cong N$  for every  $N \in \text{Mod}_T^\kappa$ . If such a  $\beta$  does not exist, we say that  $M$  has  $\mathcal{L}_{\infty\kappa}$ -Scott height  $\infty$ .

**Remark.** Since  $M, N \in \text{Mod}_T^\kappa$  are  $\mathcal{L}_{\infty\kappa}$ -equivalent if and only if they are  $\mathcal{L}_{\kappa+\kappa}$ -equivalent, the  $\mathcal{L}_{\infty\kappa}$ -Scott height of  $M$  is either  $\leq \kappa^+$  or  $\infty$ .

The Descriptive Main Gap Theorem is a consequence of a result connecting  $B(\kappa, T)$  with the Scott height  $S(\kappa, T)$  of the  $\kappa$ -sized models of  $T$ .

## Definition

Let  $\kappa$  be an infinite cardinal. Given  $M, N \in \text{Mod}_T^\kappa$  and an ordinal  $\beta$ , set  $M \equiv_\beta N$  if and only if  $M$  and  $N$  satisfy the same  $\mathcal{L}_{\infty\kappa}$ -formulae with quantifier rank  $< \beta$ .

The  $\mathcal{L}_{\infty\kappa}$ -**Scott height** of  $M \in \text{Mod}_T^\kappa$  is the smallest ordinal  $\beta$ , if it exists, such that  $M \equiv_\beta N \Rightarrow M \cong N$  for every  $N \in \text{Mod}_T^\kappa$ . If such a  $\beta$  does not exist, we say that  $M$  has  $\mathcal{L}_{\infty\kappa}$ -Scott height  $\infty$ .

**Remark.** Since  $M, N \in \text{Mod}_T^\kappa$  are  $\mathcal{L}_{\infty\kappa}$ -equivalent if and only if they are  $\mathcal{L}_{\kappa+\kappa}$ -equivalent, the  $\mathcal{L}_{\infty\kappa}$ -Scott height of  $M$  is either  $\leq \kappa^+$  or  $\infty$ .

**Remark.**  $M \equiv_\beta N \iff \text{II} \uparrow \text{EF}_{\mathfrak{T}_\alpha}^\kappa(M, N)$ .

## Definition

The  $\mathcal{L}_{\infty\kappa}$ -**Scott height**  $S(\kappa, T)$  of a theory  $T$  is the sup of all the  $\mathcal{L}_{\infty\kappa}$ -Scott heights of the  $\kappa$ -sized models of  $T$ .

## Definition

The  $\mathcal{L}_{\infty\kappa}$ -**Scott height**  $S(\kappa, T)$  of a theory  $T$  is the sup of all the  $\mathcal{L}_{\infty\kappa}$ -Scott heights of the  $\kappa$ -sized models of  $T$ .

## Theorem (Shelah )

Let  $\kappa > 2^{\aleph_0}$  be regular.

## Definition

The  $\mathcal{L}_{\infty\kappa}$ -**Scott height**  $S(\kappa, T)$  of a theory  $T$  is the sup of all the  $\mathcal{L}_{\infty\kappa}$ -Scott heights of the  $\kappa$ -sized models of  $T$ .

## Theorem (Shelah )

Let  $\kappa > 2^{\aleph_0}$  be regular. Then  $S(\kappa, T) < \infty$  if and only if  $T$  is classifiable.

## Definition

The  $\mathcal{L}_{\infty\kappa}$ -**Scott height**  $S(\kappa, T)$  of a theory  $T$  is the sup of all the  $\mathcal{L}_{\infty\kappa}$ -Scott heights of the  $\kappa$ -sized models of  $T$ .

## Theorem (Shelah)

Let  $\kappa > 2^{\aleph_0}$  be regular. Then  $S(\kappa, T) < \infty$  if and only if  $T$  is classifiable. Furthermore,

- if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $S(\kappa, T) \leq 2\alpha$  ;

## Definition

The  $\mathcal{L}_{\infty\kappa}$ -**Scott height**  $S(\kappa, T)$  of a theory  $T$  is the sup of all the  $\mathcal{L}_{\infty\kappa}$ -Scott heights of the  $\kappa$ -sized models of  $T$ .

## Theorem (Shelah, with a bit of Lascar)

Let  $\kappa > 2^{\aleph_0}$  be regular. Then  $S(\kappa, T) < \infty$  if and only if  $T$  is classifiable. Furthermore,

- if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $S(\kappa, T) \leq 2\alpha < \aleph_1$ ;



## Definition

The  $\mathcal{L}_{\infty\kappa}$ -**Scott height**  $S(\kappa, T)$  of a theory  $T$  is the sup of all the  $\mathcal{L}_{\infty\kappa}$ -Scott heights of the  $\kappa$ -sized models of  $T$ .

## Theorem (Shelah, with a bit of Lascar)

Let  $\kappa > 2^{\aleph_0}$  be regular. Then  $S(\kappa, T) < \infty$  if and only if  $T$  is classifiable. Furthermore,

- if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $S(\kappa, T) \leq 2\alpha < \aleph_1$ ;
- if  $T$  is classifiable deep, then  $S(\kappa, T) = \kappa^+$ .

# Scott height vs Borelness

For countable models, Becker and Kechris proved that

$$B(\omega, T) < \aleph_1 \iff S(\omega, T) < \aleph_1.$$

# Scott height vs Borelness

For countable models, Becker and Kechris proved that

$$B(\omega, T) < \aleph_1 \iff S(\omega, T) < \aleph_1.$$

Their proof heavily uses *effective* descriptive set theory and a boundedness argument,

# Scott height vs Borelness

For countable models, Becker and Kechris proved that

$$B(\omega, T) < \aleph_1 \iff S(\omega, T) < \aleph_1.$$

Their proof heavily uses *effective* descriptive set theory and a boundedness argument, therefore it is not clear how to generalize this to uncountable models.

# Scott height vs Borelness

For countable models, Becker and Kechris proved that

$$B(\omega, T) < \aleph_1 \iff S(\omega, T) < \aleph_1.$$

Their proof heavily uses *effective* descriptive set theory and a boundedness argument, therefore it is not clear how to generalize this to uncountable models.

## Theorem (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  (with possibly  $\kappa = \omega$ ) and  $T$  be a complete first order theory in a signature of size  $< \kappa$ .

# Scott height vs Borelness

For countable models, Becker and Kechris proved that

$$B(\omega, T) < \aleph_1 \iff S(\omega, T) < \aleph_1.$$

Their proof heavily uses *effective* descriptive set theory and a boundedness argument, therefore it is not clear how to generalize this to uncountable models.

## Theorem (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  (with possibly  $\kappa = \omega$ ) and  $T$  be a complete first order theory in a signature of size  $< \kappa$ .

- 1  $S(\kappa, T) \leq B(\kappa, T)$ , thus if  $\cong_T^\kappa$  is Borel then  $S(\kappa, T) < \kappa^+$ ;

# Scott height vs Borelness

For countable models, Becker and Kechris proved that

$$B(\omega, T) < \aleph_1 \iff S(\omega, T) < \aleph_1.$$

Their proof heavily uses *effective* descriptive set theory and a boundedness argument, therefore it is not clear how to generalize this to uncountable models.

## Theorem (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  (with possibly  $\kappa = \omega$ ) and  $T$  be a complete first order theory in a signature of size  $< \kappa$ .

- ①  $S(\kappa, T) \leq B(\kappa, T)$ , thus if  $\cong_T^\kappa$  is Borel then  $S(\kappa, T) < \kappa^+$ ;
- ② if  $S(\kappa, T) < \kappa^+$ , then  $\cong_T^\kappa \in \mathbf{\Pi}_{2S(\kappa, T)}^0$ .

# Scott height vs Borelness

For countable models, Becker and Kechris proved that

$$B(\omega, T) < \aleph_1 \iff S(\omega, T) < \aleph_1.$$

Their proof heavily uses *effective* descriptive set theory and a boundedness argument, therefore it is not clear how to generalize this to uncountable models.

## Theorem (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  (with possibly  $\kappa = \omega$ ) and  $T$  be a complete first order theory in a signature of size  $< \kappa$ .

- ①  $S(\kappa, T) \leq B(\kappa, T)$ , thus if  $\cong_T^\kappa$  is Borel then  $S(\kappa, T) < \kappa^+$ ;
- ② if  $S(\kappa, T) < \kappa^+$ , then  $\cong_T^\kappa \in \mathbf{\Pi}_{2S(\kappa, T)}^0$ .

In particular,

$$B(\kappa, T) < \kappa^+ \iff S(\kappa, T) < \kappa^+$$



# Scott height vs Borelness

For countable models, Becker and Kechris proved that

$$B(\omega, T) < \aleph_1 \iff S(\omega, T) < \aleph_1.$$

Their proof heavily uses *effective* descriptive set theory and a boundedness argument, therefore it is not clear how to generalize this to uncountable models.

## Theorem (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  (with possibly  $\kappa = \omega$ ) and  $T$  be a complete first order theory in a signature of size  $< \kappa$ .

- ①  $S(\kappa, T) \leq B(\kappa, T)$ , thus if  $\cong_T^\kappa$  is Borel then  $S(\kappa, T) < \kappa^+$ ;
- ② if  $S(\kappa, T) < \kappa^+$ , then  $\cong_T^\kappa \in \mathbf{\Pi}_{2S(\kappa, T)}^0$ .

In particular,

$$B(\kappa, T) < \kappa^+ \iff S(\kappa, T) < \kappa^+$$

and in such case  $S(\kappa, T) \leq B(\kappa, T) \leq 2S(\kappa, T)$ .

# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- ① if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $\cong_T^\kappa \in \Pi_{4\alpha}^0$ ;

# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- ① if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $\cong_T^\kappa \in \Pi_{4\alpha}^0$ ;
- ② if  $T$  is not classifiable shallow, then  $\cong_T^\kappa$  is not Borel (i.e.  $B(\kappa, T) = \infty$ ).

# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- ① if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $\cong_T^\kappa \in \mathbf{\Pi}_{4\alpha}^0$ ;
- ② if  $T$  is not classifiable shallow, then  $\cong_T^\kappa$  is not Borel (i.e.  $B(\kappa, T) = \infty$ ).

## Proof.

- ① Let  $T$  be classifiable shallow with  $\text{dp}(T) = \alpha$ .

# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- ① if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $\cong_T^\kappa \in \mathbf{\Pi}_{4\alpha}^0$ ;
- ② if  $T$  is not classifiable shallow, then  $\cong_T^\kappa$  is not Borel (i.e.  $B(\kappa, T) = \infty$ ).

## Proof.

- ① Let  $T$  be classifiable shallow with  $\text{dp}(T) = \alpha$ . Then  $S(\kappa, T) \leq 2\alpha < \aleph_1 < \kappa^+$ ,

# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- ① if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $\cong_T^\kappa \in \mathbf{\Pi}_{4\alpha}^0$ ;
- ② if  $T$  is not classifiable shallow, then  $\cong_T^\kappa$  is not Borel (i.e.  $B(\kappa, T) = \infty$ ).

## Proof.

- ① Let  $T$  be classifiable shallow with  $\text{dp}(T) = \alpha$ . Then  $S(\kappa, T) \leq 2\alpha < \aleph_1 < \kappa^+$ , whence  $\cong_T^\kappa \in \mathbf{\Pi}_{2S(\kappa, T)}^0$

# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- 1 if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $\cong_T^\kappa \in \Pi_{4\alpha}^0$ ;
- 2 if  $T$  is not classifiable shallow, then  $\cong_T^\kappa$  is not Borel (i.e.  $B(\kappa, T) = \infty$ ).

## Proof.

- 1 Let  $T$  be classifiable shallow with  $\text{dp}(T) = \alpha$ . Then  $S(\kappa, T) \leq 2\alpha < \aleph_1 < \kappa^+$ , whence  $\cong_T^\kappa \in \Pi_{2S(\kappa, T)}^0 \subseteq \Pi_{4\alpha}^0$ .



# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- 1 if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $\cong_T^\kappa \in \Pi_{4\alpha}^0$ ;
- 2 if  $T$  is not classifiable shallow, then  $\cong_T^\kappa$  is not Borel (i.e.  $B(\kappa, T) = \infty$ ).

## Proof.

- 1 Let  $T$  be classifiable shallow with  $\text{dp}(T) = \alpha$ . Then  $S(\kappa, T) \leq 2\alpha < \aleph_1 < \kappa^+$ , whence  $\cong_T^\kappa \in \Pi_{2S(\kappa, T)}^0 \subseteq \Pi_{4\alpha}^0$ .
- 2 Suppose that  $B(\kappa, T) < \kappa^+$ .

# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- ① if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $\cong_T^\kappa \in \Pi_{4\alpha}^0$ ;
- ② if  $T$  is not classifiable shallow, then  $\cong_T^\kappa$  is not Borel (i.e.  $B(\kappa, T) = \infty$ ).

## Proof.

- ① Let  $T$  be classifiable shallow with  $\text{dp}(T) = \alpha$ . Then  $S(\kappa, T) \leq 2\alpha < \aleph_1 < \kappa^+$ , whence  $\cong_T^\kappa \in \Pi_{2S(\kappa, T)}^0 \subseteq \Pi_{4\alpha}^0$ .
- ② Suppose that  $B(\kappa, T) < \kappa^+$ . Then since  $S(\kappa, T) \leq B(\kappa, T)$

# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- ① if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $\cong_T^\kappa \in \Pi_{4\alpha}^0$ ;
- ② if  $T$  is not classifiable shallow, then  $\cong_T^\kappa$  is not Borel (i.e.  $B(\kappa, T) = \infty$ ).

## Proof.

- ① Let  $T$  be classifiable shallow with  $\text{dp}(T) = \alpha$ . Then  $S(\kappa, T) \leq 2\alpha < \aleph_1 < \kappa^+$ , whence  $\cong_T^\kappa \in \Pi_{2S(\kappa, T)}^0 \subseteq \Pi_{4\alpha}^0$ .
- ② Suppose that  $B(\kappa, T) < \kappa^+$ . Then since  $S(\kappa, T) \leq B(\kappa, T) < \kappa^+$ ,

# Proof of the Descriptive Main Gap Theorem

## Theorem (Mangraviti-M.)

Let  $T$  be a countable complete first-order theory, and  $\kappa$  be such that  $\kappa^{<\kappa} = \kappa > 2^{\aleph_0}$ .

- ① if  $T$  is classifiable shallow with  $\text{dp}(T) = \alpha$ , then  $\cong_T^\kappa \in \Pi_{4\alpha}^0$ ;
- ② if  $T$  is not classifiable shallow, then  $\cong_T^\kappa$  is not Borel (i.e.  $B(\kappa, T) = \infty$ ).

## Proof.

- ① Let  $T$  be classifiable shallow with  $\text{dp}(T) = \alpha$ . Then  $S(\kappa, T) \leq 2\alpha < \aleph_1 < \kappa^+$ , whence  $\cong_T^\kappa \in \Pi_{2S(\kappa, T)}^0 \subseteq \Pi_{4\alpha}^0$ .
- ② Suppose that  $B(\kappa, T) < \kappa^+$ . Then since  $S(\kappa, T) \leq B(\kappa, T) < \kappa^+$ , the theory  $T$  is classifiable shallow by Shelah's theorem. □

$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow
$I(\kappa, T) = 1$	$I(\kappa, T) < \beth_\alpha( \gamma ^{2^{\aleph_0}})$	$I(\kappa, T) = 2^\kappa$
	$(\kappa^{+-})\text{Borel}$	Not $(\kappa^{+-})\text{Borel}$
	$B(\kappa, T) \leq 4\alpha$	$B(\kappa, T) = \infty$

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow
$I(\kappa, T) = 1$	$I(\kappa, T) < \beth_\alpha( \gamma ^{2^{\aleph_0}})$	$I(\kappa, T) = 2^\kappa$
	$S(\kappa, T) \leq 2\alpha$	$S(\kappa, T) \in \{\kappa^+, \infty\}$
	$(\kappa^+)$ -Borel	Not $(\kappa^+)$ -Borel
	$B(\kappa, T) \leq 4\alpha$	$B(\kappa, T) = \infty$

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow
$I(\kappa, T) = 1$	$I(\kappa, T) < \beth_\alpha( \gamma ^{2^{\aleph_0}})$	$I(\kappa, T) = 2^\kappa$
$S(\kappa, T) = 0$	$S(\kappa, T) \leq 2\alpha$	$S(\kappa, T) \in \{\kappa^+, \infty\}$
	$(\kappa^+)$ -Borel	Not $(\kappa^+)$ -Borel
	$B(\kappa, T) \leq 4\alpha$	$B(\kappa, T) = \infty$

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

# Categoricity from the topological viewpoint

Lemma (completeness  $\rightsquigarrow$  density)

Let  $\kappa$  be any infinite cardinal,



# Categoricity from the topological viewpoint

## Lemma (completeness $\rightsquigarrow$ density)

Let  $\kappa$  be any infinite cardinal,  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ ,

# Categoricity from the topological viewpoint

## Lemma (completeness $\rightsquigarrow$ density)

Let  $\kappa$  be any infinite cardinal,  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ , and let  $M \in \text{Mod}_T^\kappa$ .

# Categoricity from the topological viewpoint

## Lemma (completeness $\rightsquigarrow$ density)

Let  $\kappa$  be any infinite cardinal,  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ , and let  $M \in \text{Mod}_T^\kappa$ . If  $[M]_\cong$  has nonempty interior (with respect to  $\text{Mod}_T^\kappa$ ), then it is dense in  $\text{Mod}_T^\kappa$ .

# Categoricity from the topological viewpoint

## Lemma (completeness $\rightsquigarrow$ density)

Let  $\kappa$  be any infinite cardinal,  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ , and let  $M \in \text{Mod}_T^\kappa$ . If  $[M]_\cong$  has nonempty interior (with respect to  $\text{Mod}_T^\kappa$ ), then it is dense in  $\text{Mod}_T^\kappa$ .

## Theorem (Mangraviti-M.)

Let  $\kappa$  be any infinite cardinal and  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ . The following are equivalent:

# Categoricity from the topological viewpoint

## Lemma (completeness $\rightsquigarrow$ density)

Let  $\kappa$  be any infinite cardinal,  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ , and let  $M \in \text{Mod}_T^\kappa$ . If  $[M]_\cong$  has nonempty interior (with respect to  $\text{Mod}_T^\kappa$ ), then it is dense in  $\text{Mod}_T^\kappa$ .

## Theorem (Mangraviti-M.)

Let  $\kappa$  be any infinite cardinal and  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ . The following are equivalent:

- 1  $T$  is  $\kappa$ -categorical;

# Categoricity from the topological viewpoint

## Lemma (completeness $\rightsquigarrow$ density)

Let  $\kappa$  be any infinite cardinal,  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ , and let  $M \in \text{Mod}_T^\kappa$ . If  $[M]_\cong$  has nonempty interior (with respect to  $\text{Mod}_T^\kappa$ ), then it is dense in  $\text{Mod}_T^\kappa$ .

## Theorem (Mangraviti-M.)

Let  $\kappa$  be any infinite cardinal and  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ . The following are equivalent:

- 1  $T$  is  $\kappa$ -categorical;
- 2  $B(\kappa, T) = 0$ , i.e.  $\cong_T^\kappa$  is clopen;

# Categoricity from the topological viewpoint

## Lemma (completeness $\rightsquigarrow$ density)

Let  $\kappa$  be any infinite cardinal,  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ , and let  $M \in \text{Mod}_T^\kappa$ . If  $[M]_\cong$  has nonempty interior (with respect to  $\text{Mod}_T^\kappa$ ), then it is dense in  $\text{Mod}_T^\kappa$ .

## Theorem (Mangraviti-M.)

Let  $\kappa$  be any infinite cardinal and  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ . The following are equivalent:

- 1  $T$  is  $\kappa$ -categorical;
- 2  $B(\kappa, T) = 0$ , i.e.  $\cong_T^\kappa$  is clopen;
- 3 there is  $M \in \text{Mod}_T^\kappa$  such that  $[M]_\cong$  is clopen.

# Categoricity from the topological viewpoint

## Lemma (completeness $\rightsquigarrow$ density)

Let  $\kappa$  be any infinite cardinal,  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ , and let  $M \in \text{Mod}_T^\kappa$ . If  $[M]_\cong$  has nonempty interior (with respect to  $\text{Mod}_T^\kappa$ ), then it is dense in  $\text{Mod}_T^\kappa$ .

## Theorem (Mangraviti-M.)

Let  $\kappa$  be any infinite cardinal and  $T$  be a (not necessarily countable) complete first-order theory in a language  $\mathcal{L}$  of size  $\leq \kappa$ . The following are equivalent:

- 1  $T$  is  $\kappa$ -categorical;
- 2  $B(\kappa, T) = 0$ , i.e.  $\cong_T^\kappa$  is clopen;
- 3 there is  $M \in \text{Mod}_T^\kappa$  such that  $[M]_\cong$  is clopen.

In particular, there is no complete non- $\kappa$ -categorical first-order theory  $T$  for which  $\cong_T^\kappa$  is true open.



$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow
$I(\kappa, T) = 1$	$I(\kappa, T) < \beth_\alpha( \gamma ^{2^{\aleph_0}})$	$I(\kappa, T) = 2^\kappa$
$S(\kappa, T) = 0$	$S(\kappa, T) \leq 2\alpha$	$S(\kappa, T) \in \{\kappa^+, \infty\}$
	$(\kappa^+)$ -Borel	Not $(\kappa^+)$ -Borel
	$B(\kappa, T) \leq 4\alpha$	$B(\kappa, T) = \infty$

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

$T$  countable complete first-order theory;  $\kappa = \aleph_\gamma$  with  $\gamma \geq 1$ .

Uncountably categorical	Classifiable shallow $\text{dp}(T) = \alpha$	Not classifiable shallow
$I(\kappa, T) = 1$	$I(\kappa, T) < \beth_\alpha( \gamma ^{2^{\aleph_0}})$	$I(\kappa, T) = 2^\kappa$
$S(\kappa, T) = 0$	$S(\kappa, T) \leq 2\alpha$	$S(\kappa, T) \in \{\kappa^+, \infty\}$
(CI)open	$(\kappa^{+-})\text{Borel}$	Not $(\kappa^{+-})\text{Borel}$
$B(\kappa, T) = 0$	$B(\kappa, T) \leq 4\alpha$	$B(\kappa, T) = \infty$

(\*) Recall that  $\alpha = \text{dp}(T) < \aleph_1$ .

# Yet another Descriptive Main Gap Theorem

## Theorem

Let  $\kappa = \aleph_\gamma$  be such that  $\kappa^{<\kappa} = \kappa$  and  $\beth_{\omega_1}(|\gamma|) \leq \kappa$ .

# Yet another Descriptive Main Gap Theorem

## Theorem

Let  $\kappa = \aleph_\gamma$  be such that  $\kappa^{<\kappa} = \kappa$  and  $\beth_{\omega_1}(|\gamma|) \leq \kappa$ . Let  $T, T'$  be arbitrary countable complete first-order theory, and assume that  $T$  is classifiable shallow, while  $T'$  is not.

# Yet another Descriptive Main Gap Theorem

## Theorem

Let  $\kappa = \aleph_\gamma$  be such that  $\kappa^{<\kappa} = \kappa$  and  $\beth_{\omega_1}(|\gamma|) \leq \kappa$ . Let  $T, T'$  be arbitrary countable complete first-order theory, and assume that  $T$  is classifiable shallow, while  $T'$  is not. Then

$$\cong_T^\kappa <_B^\kappa \cong_{T'}^\kappa.$$

# Yet another Descriptive Main Gap Theorem

## Theorem

Let  $\kappa = \aleph_\gamma$  be such that  $\kappa^{<\kappa} = \kappa$  and  $\beth_{\omega_1}(|\gamma|) \leq \kappa$ . Let  $T, T'$  be arbitrary countable complete first-order theory, and assume that  $T$  is classifiable shallow, while  $T'$  is not. Then

$$\cong_T^\kappa <_B^\kappa \cong_{T'}^\kappa.$$

Moreover, the  $\leq_B^\kappa$ -interval between  $\cong_T^\kappa$  and  $\cong_{T'}^\kappa$  is nonempty.

# Yet another Descriptive Main Gap Theorem

## Theorem

Let  $\kappa = \aleph_\gamma$  be such that  $\kappa^{<\kappa} = \kappa$  and  $\beth_{\omega_1}(|\gamma|) \leq \kappa$ . Let  $T, T'$  be arbitrary countable complete first-order theory, and assume that  $T$  is classifiable shallow, while  $T'$  is not. Then

$$\cong_T^\kappa <_B^\kappa \cong_{T'}^\kappa.$$

Moreover, the  $\leq_B^\kappa$ -interval between  $\cong_T^\kappa$  and  $\cong_{T'}^\kappa$  is nonempty.

The non-reducibility direction can be proved either counting the number of isomorphism types, or using the fact that  $\cong_T^\kappa$  is Borel while  $\cong_{T'}^\kappa$  is not.

# Yet another Descriptive Main Gap Theorem

## Theorem

Let  $\kappa = \aleph_\gamma$  be such that  $\kappa^{<\kappa} = \kappa$  and  $\beth_{\omega_1}(|\gamma|) \leq \kappa$ . Let  $T, T'$  be arbitrary countable complete first-order theory, and assume that  $T$  is classifiable shallow, while  $T'$  is not. Then

$$\cong_T^\kappa <_\kappa^B \cong_{T'}^\kappa.$$

Moreover, the  $\leq_\kappa^B$ -interval between  $\cong_T^\kappa$  and  $\cong_{T'}^\kappa$  is nonempty.

The non-reducibility direction can be proved either counting the number of isomorphism types, or using the fact that  $\cong_T^\kappa$  is Borel while  $\cong_{T'}^\kappa$  is not. The reducibility direction uses that  $\cong_T^\kappa$  has  $\leq \kappa$  classes and that all of them are Borel.



# Yet another Descriptive Main Gap Theorem

## Theorem

Let  $\kappa = \aleph_\gamma$  be such that  $\kappa^{<\kappa} = \kappa$  and  $\beth_{\omega_1}(|\gamma|) \leq \kappa$ . Let  $T, T'$  be arbitrary countable complete first-order theory, and assume that  $T$  is classifiable shallow, while  $T'$  is not. Then

$$\cong_T^\kappa <_B^\kappa \cong_{T'}^\kappa.$$

Moreover, the  $\leq_B^\kappa$ -interval between  $\cong_T^\kappa$  and  $\cong_{T'}^\kappa$  is nonempty.

The non-reducibility direction can be proved either counting the number of isomorphism types, or using the fact that  $\cong_T^\kappa$  is Borel while  $\cong_{T'}^\kappa$  is not. The reducibility direction uses that  $\cong_T^\kappa$  has  $\leq \kappa$  classes and that all of them are Borel.

## Question

How large is the  $\leq_B^\kappa$ -gap between  $\cong_T^\kappa$  and  $\cong_{T'}^\kappa$ ?

## Proposition (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  and  $\mathcal{L}$  a signature of size  $< \kappa$ .

# Forbidden complexities for $\cong_T^\kappa$

## Proposition (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  and  $\mathcal{L}$  a signature of size  $< \kappa$ . If  $\alpha < \kappa^+$  is limit, then there is no  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\kappa$  is a true  $\Sigma_\alpha^0$  set.

# Forbidden complexities for $\cong_T^\kappa$

## Proposition (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  and  $\mathcal{L}$  a signature of size  $< \kappa$ . If  $\alpha < \kappa^+$  is limit, then there is no  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\kappa$  is a true  $\Sigma_\alpha^0$  set.

This extends to  $\mathcal{L}_{\kappa^+\kappa}$ -theories and beyond.

# Forbidden complexities for $\cong_T^\kappa$

## Proposition (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  and  $\mathcal{L}$  a signature of size  $< \kappa$ . If  $\alpha < \kappa^+$  is limit, then there is no  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\kappa$  is a true  $\Sigma_\alpha^0$  set.

This extends to  $\mathcal{L}_{\kappa+\kappa}$ -theories and beyond. For successor ordinals, the situation is quite different.

# Forbidden complexities for $\cong_T^\kappa$

## Proposition (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  and  $\mathcal{L}$  a signature of size  $< \kappa$ . If  $\alpha < \kappa^+$  is limit, then there is no  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\kappa$  is a true  $\Sigma_\alpha^0$  set.

This extends to  $\mathcal{L}_{\kappa+\kappa}$ -theories and beyond. For successor ordinals, the situation is quite different.

Let  $\kappa^{<\kappa} = \kappa > \omega$ .

# Forbidden complexities for $\cong_T^\kappa$

## Proposition (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  and  $\mathcal{L}$  a signature of size  $< \kappa$ . If  $\alpha < \kappa^+$  is limit, then there is no  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\kappa$  is a true  $\Sigma_\alpha^0$  set.

This extends to  $\mathcal{L}_{\kappa+\kappa}$ -theories and beyond. For successor ordinals, the situation is quite different.

Let  $\kappa^{<\kappa} = \kappa > \omega$ . There is a countable complete first-order theory  $T$  in the signature  $\mathcal{L} = \{P\}$  with  $P$  unary such that

# Forbidden complexities for $\cong_T^\kappa$

## Proposition (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  and  $\mathcal{L}$  a signature of size  $< \kappa$ . If  $\alpha < \kappa^+$  is limit, then there is no  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\kappa$  is a true  $\Sigma_\alpha^0$  set.

This extends to  $\mathcal{L}_{\kappa^+\kappa}$ -theories and beyond. For successor ordinals, the situation is quite different.

Let  $\kappa^{<\kappa} = \kappa > \omega$ . There is a countable complete first-order theory  $T$  in the signature  $\mathcal{L} = \{P\}$  with  $P$  unary such that

- 1 if  $\kappa = \lambda^+$  is a successor cardinal, then  $S(\kappa, T) = 2$  and  $\cong_T^\kappa$  is a true  $\Delta_3^0$  set (in particular,  $B(\kappa, T) = 3$ );



# Forbidden complexities for $\cong_T^\kappa$

## Proposition (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  and  $\mathcal{L}$  a signature of size  $< \kappa$ . If  $\alpha < \kappa^+$  is limit, then there is no  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\kappa$  is a true  $\Sigma_\alpha^0$  set.

This extends to  $\mathcal{L}_{\kappa^+\kappa}$ -theories and beyond. For successor ordinals, the situation is quite different.

Let  $\kappa^{<\kappa} = \kappa > \omega$ . There is a countable complete first-order theory  $T$  in the signature  $\mathcal{L} = \{P\}$  with  $P$  unary such that

- 1 if  $\kappa = \lambda^+$  is a successor cardinal, then  $S(\kappa, T) = 2$  and  $\cong_T^\kappa$  is a true  $\Delta_3^0$  set (in particular,  $B(\kappa, T) = 3$ );
- 2 if  $\kappa$  is a limit cardinal, then  $S(\kappa, T) = 1$  while  $\cong_T^\kappa$  is a true  $\Pi_2^0$  set (in particular,  $B(\kappa, T) = 2$ ).

# Forbidden complexities for $\cong_T^\kappa$

## Proposition (Mangraviti-M.)

Let  $\kappa^{<\kappa} = \kappa$  and  $\mathcal{L}$  a signature of size  $< \kappa$ . If  $\alpha < \kappa^+$  is limit, then there is no  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\kappa$  is a true  $\Sigma_\alpha^0$  set.

This extends to  $\mathcal{L}_{\kappa+\kappa}$ -theories and beyond. For successor ordinals, the situation is quite different.

Let  $\kappa^{<\kappa} = \kappa > \omega$ . There is a countable complete first-order theory  $T$  in the signature  $\mathcal{L} = \{P\}$  with  $P$  unary such that

- 1 if  $\kappa = \lambda^+$  is a successor cardinal, then  $S(\kappa, T) = 2$  and  $\cong_T^\kappa$  is a true  $\Delta_3^0$  set (in particular,  $B(\kappa, T) = 3$ );
- 2 if  $\kappa$  is a limit cardinal, then  $S(\kappa, T) = 1$  while  $\cong_T^\kappa$  is a true  $\Pi_2^0$  set (in particular,  $B(\kappa, T) = 2$ ).

This also show that both the Scott height and the Borel complexity may vary with the size of the models.

# Back to countable structures

Also in the case of countable structures there are some complexities for  $\cong_T^\omega$  which are forbidden.

# Back to countable structures

Also in the case of countable structures there are some complexities for  $\cong_T^\omega$  which are forbidden.

## Proposition (Mangraviti-M.)

Let  $\mathcal{L}$  be a finite signature and  $\alpha < \omega_1$  be a limit ordinal.

# Back to countable structures

Also in the case of countable structures there are some complexities for  $\cong_T^\omega$  which are forbidden.

## Proposition (Mangraviti-M.)

Let  $\mathcal{L}$  be a finite signature and  $\alpha < \omega_1$  be a limit ordinal. Then there is no (complete or incomplete)  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\omega$  is a true  $\Sigma_\alpha^0$  set.

# Back to countable structures

Also in the case of countable structures there are some complexities for  $\cong_T^\omega$  which are forbidden.

## Proposition (Mangraviti-M.)

Let  $\mathcal{L}$  be a finite signature and  $\alpha < \omega_1$  be a limit ordinal. Then there is no (complete or incomplete)  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\omega$  is a true  $\Sigma_\alpha^0$  set.

Moreover, completeness of a theory turns into an even stronger density condition when stepping back to the countable case.

# Back to countable structures

Also in the case of countable structures there are some complexities for  $\cong_T^\omega$  which are forbidden.

## Proposition (Mangraviti-M.)

Let  $\mathcal{L}$  be a finite signature and  $\alpha < \omega_1$  be a limit ordinal. Then there is no (complete or incomplete)  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\omega$  is a true  $\Sigma_\alpha^0$  set.

Moreover, completeness of a theory turns into an even stronger density condition when stepping back to the countable case.

## Lemma

Let  $T$  be a complete first-order theory in a countable language and  $M \in \text{Mod}_T^\omega$ .

# Back to countable structures

Also in the case of countable structures there are some complexities for  $\cong_T^\omega$  which are forbidden.

## Proposition (Mangraviti-M.)

Let  $\mathcal{L}$  be a finite signature and  $\alpha < \omega_1$  be a limit ordinal. Then there is no (complete or incomplete)  $\mathcal{L}$ -theory  $T$  such that  $\cong_T^\omega$  is a true  $\Sigma_\alpha^0$  set.

Moreover, completeness of a theory turns into an even stronger density condition when stepping back to the countable case.

## Lemma

Let  $T$  be a complete first-order theory in a countable language and  $M \in \text{Mod}_T^\omega$ . Then  $[M]_{\cong}$  is dense in  $\text{Mod}_T^\omega$  (unconditionally!).



Thus we get a purely topological characterization of  $\aleph_0$ -categoricity.

Thus we get a purely topological characterization of  $\aleph_0$ -categoricity.

## Theorem (Mangraviti-M.)

Let  $T$  be a complete first-order theory in a countable language. The following are equivalent:

Thus we get a purely topological characterization of  $\aleph_0$ -categoricity.

## Theorem (Mangraviti-M.)

Let  $T$  be a complete first-order theory in a countable language. The following are equivalent:

- 1  $T$  is  $\aleph_0$ -categorical;

Thus we get a purely topological characterization of  $\aleph_0$ -categoricity.

## Theorem (Mangraviti-M.)

Let  $T$  be a complete first-order theory in a countable language. The following are equivalent:

- 1  $T$  is  $\aleph_0$ -categorical;
- 2  $B(\kappa, T) = \mathbf{1}$ , i.e.  $\cong_T^\kappa$  is open or closed;

Thus we get a purely topological characterization of  $\aleph_0$ -categoricity.

## Theorem (Mangraviti-M.)

Let  $T$  be a complete first-order theory in a countable language. The following are equivalent:

- 1  $T$  is  $\aleph_0$ -categorical;
- 2  $B(\kappa, T) = 1$ , i.e.  $\cong_{\kappa}^T$  is open or closed;
- 3 there is  $M \in \text{Mod}_T^{\omega}$  such that  $[M]_{\cong}$  is open or closed.

Thus we get a purely topological characterization of  $\aleph_0$ -categoricity.

## Theorem (Mangraviti-M.)

Let  $T$  be a complete first-order theory in a countable language. The following are equivalent:

- 1  $T$  is  $\aleph_0$ -categorical;
- 2  $B(\kappa, T) = 1$ , i.e.  $\cong_T^\kappa$  is open or closed;
- 3 there is  $M \in \text{Mod}_T^\omega$  such that  $[M]_\cong$  is open or closed.

In particular, other complexities are forbidden: there is no complete first-order theory  $T$  such that  $\cong_T^\omega$  is true open or true closed.

# Topological smoothness

Recall that an equivalence relation  $E$  on a standard Borel space  $X$  is called **smooth** if  $E \leq_B \text{id}(Y)$  for some *Polish* space  $Y$ .

Recall that an equivalence relation  $E$  on a standard Borel space  $X$  is called **smooth** if  $E \leq_B \text{id}(Y)$  for some *Polish* space  $Y$ .

## Definition

An equivalence relation  $E$  on a topological space  $X$  is called **topologically smooth** if  $E$  is continuously reducible to  $\text{id}(Y)$  for some *Hausdorff* space  $Y$ , that is:



Recall that an equivalence relation  $E$  on a standard Borel space  $X$  is called **smooth** if  $E \leq_B \text{id}(Y)$  for some *Polish* space  $Y$ .

## Definition

An equivalence relation  $E$  on a topological space  $X$  is called **topologically smooth** if  $E$  is continuously reducible to  $\text{id}(Y)$  for some *Hausdorff* space  $Y$ , that is: there is a *continuous* map  $f: X \rightarrow Y$  such that for all  $x, y \in X$

$$x E y \iff f(x) = f(y).$$

# Topological smoothness

Recall that an equivalence relation  $E$  on a standard Borel space  $X$  is called **smooth** if  $E \leq_B \text{id}(Y)$  for some *Polish* space  $Y$ .

## Definition

An equivalence relation  $E$  on a topological space  $X$  is called **topologically smooth** if  $E$  is continuously reducible to  $\text{id}(Y)$  for some *Hausdorff* space  $Y$ , that is: there is a *continuous* map  $f: X \rightarrow Y$  such that for all  $x, y \in X$

$$x E y \iff f(x) = f(y).$$

There are natural examples of non- $\aleph_0$ -categorical complete theories  $T$  for which  $\cong_T^\omega$  is smooth.

# Topological smoothness

Recall that an equivalence relation  $E$  on a standard Borel space  $X$  is called **smooth** if  $E \leq_B \text{id}(Y)$  for some *Polish* space  $Y$ .

## Definition

An equivalence relation  $E$  on a topological space  $X$  is called **topologically smooth** if  $E$  is continuously reducible to  $\text{id}(Y)$  for some *Hausdorff* space  $Y$ , that is: there is a *continuous* map  $f: X \rightarrow Y$  such that for all  $x, y \in X$

$$x E y \iff f(x) = f(y).$$

There are natural examples of non- $\aleph_0$ -categorical complete theories  $T$  for which  $\cong_T^\omega$  is smooth.

What about *topological* smoothness?

# Topological smoothness

Let  $T$  be any complete first-order theory in a countable language.

# Topological smoothness

Let  $T$  be any complete first-order theory in a countable language.

## Theorem (Mangraviti-M.)

$\cong_T^\omega$  is topologically smooth    if and only if     $T$  is  $\aleph_0$ -categorical.

# Topological smoothness

Let  $T$  be any complete first-order theory in a countable language.

## Theorem (Mangraviti-M.)

$\cong_T^\omega$  is topologically smooth if and only if  $T$  is  $\aleph_0$ -categorical.

Thus unless  $T$  has just one countable model

*it is not possible to classify in a continuous way the countable models of  $T$  up to isomorphism using the elements of a Hausdorff space as invariants.*

# Topological smoothness

Let  $T$  be any complete first-order theory in a countable language.

## Theorem (Mangraviti-M.)

$\cong_T^\omega$  is topologically smooth if and only if  $T$  is  $\aleph_0$ -categorical.

Thus unless  $T$  has just one countable model

*it is not possible to classify in a continuous way the countable models of  $T$  up to isomorphism using the elements of a Hausdorff space as invariants.*

In particular, the quotient topology on  $\text{Mod}_T^\omega / \cong$  is never Hausdorff if the space has more than one point.

# Topological smoothness

Let  $T$  be any complete first-order theory in a countable language.

## Theorem (Mangraviti-M.)

$\cong_T^\omega$  is topologically smooth if and only if  $T$  is  $\aleph_0$ -categorical.

Thus unless  $T$  has just one countable model

*it is not possible to classify in a continuous way the countable models of  $T$  up to isomorphism using the elements of a Hausdorff space as invariants.*

In particular, the quotient topology on  $\text{Mod}_T^\omega / \cong$  is never Hausdorff if the space has more than one point.

**Remark.** The result is actually stronger.



# Topological smoothness

Let  $T$  be any complete first-order theory in a countable language.

## Theorem (Mangraviti-M.)

$\cong_T^\omega$  is topologically smooth if and only if  $T$  is  $\aleph_0$ -categorical.

Thus unless  $T$  has just one countable model

*it is not possible to classify in a continuous way the countable models of  $T$  up to isomorphism using the elements of a Hausdorff space as invariants.*

In particular, the quotient topology on  $\text{Mod}_T^\omega / \cong$  is never Hausdorff if the space has more than one point.

**Remark.** The result is actually stronger. If  $f: \text{Mod}_T^\omega \rightarrow Y$  is continuous and invariant with  $Y$  Hausdorff, then the quotient map  $\hat{f}: \text{Mod}_T^\omega / \cong$  is nowhere injective: for every  $M \in \text{Mod}_T^\omega$  we can always find  $N \in \text{Mod}_T^\omega$  such that  $N \not\cong M$ , yet  $f(N) = f(M)$ .

The end

**Thank you for your attention!**