

Philipp von Bachmann,  
Mat.-Nr.: 4116220  
David Ott,  
Mat.-Nr.: 4185646

1	2	3	$\Sigma$

## Assignment Nr. 4

### 4.1 Radiometry

a)

As the point light source emits  $\Phi_S$  Watts in every direction, its radiance is  $L_o(d, \omega) = \frac{\Phi_S}{4\pi}$  in every direction. Because it is the only light source in the scene which emits light to  $x$  from the direction  $\omega_i$ , the integral for the irradiance of  $x$   $I_x = \int_{\omega} L(x, \omega) \cos(\theta) d\omega$  collapses to  $I_x = L_o(d, \omega_i) \cos(\theta)$ . This would assume however that the light source sits on the unit circle around  $x$ . Therefore, we have to normalize by the distance between  $d$  and  $x$ , resulting in  $I_x = \frac{L_o(d, \omega_i) \cos(\theta)}{|x-d|} = \frac{L_o(d, \omega_i) \cos(\theta)}{r^2 + d^2} = \frac{\Phi_S \cos(\theta)}{4\pi(r^2 + d^2)}$

Let  $w'$  be the direction from  $x$  to  $d$

$$\begin{aligned} E &= \int_{\omega} L(x, \omega) \cos(\theta) d\omega \\ &= L(x, w') \cos(\theta) \\ &= \frac{d^2 \Phi \cos(\theta)}{dA \cos(\theta) d\omega} \end{aligned}$$

as a point is perpendicular to another  $\cos = 1$

$$= \frac{d^2 \Phi \cos(\theta)}{dA d\omega}$$

Assume  $dA = 1$ , then  $d\omega = 4\pi r^2$

$$= \frac{d^2 \Phi \cos(\theta)}{4\pi r^2}$$

The radius is given by  $r = \sqrt{r^2 + d^2}$

$$= \frac{d^2 \Phi \cos(\theta)}{4\pi r^2 + d^2}$$

a)

$$\omega = (\theta, \phi)$$

$$\begin{aligned}
E &= \int_0^{2\pi} \int_0^{\frac{\pi}{2}} L(x, \omega) \cos \theta \sin \theta d\theta d\phi \\
&= \int_0^{2\pi} \int_0^{\frac{\pi}{2}} \frac{d^2 \Phi}{dA \cos(\theta) d\omega} \cos \theta \sin \theta d\theta d\phi \\
&= \int_0^{2\pi} \int_0^{\frac{\pi}{2}} \frac{d^2 \Phi}{\sin(\theta) d\theta d\phi \cdot r^2 \cos(\theta) d\omega} \cos \theta \sin \theta d\theta d\phi \\
&= \int_0^{2\pi} \int_0^{\frac{\pi}{2}} \frac{d^2 \Phi}{r^2 d\omega} \\
&= \int_0^{2\pi} \int_0^{\frac{\pi}{2}} \frac{d^2 \Phi}{r^2 d\theta d\phi} \\
&= \\
&= \frac{\Phi_S \cos(\theta)}{4\pi(r^2 + d^2)}
\end{aligned}$$

## 4.2 Analytical solution of the rendering equation in 2D

a)

$$\begin{aligned}
L(x, \omega_o) &= L_e(x, \omega_o) + \int_0^\pi f_r(\omega, x, \omega_o) L(x, \omega) \cos(\phi) d\omega \\
&= 0 + \int_0^\pi \frac{1}{2} L(x, \omega) \cos(\phi) d\omega \\
&= \frac{1}{2} \int_0^\pi L(x, \omega) \cos(\phi) d\omega \\
&= \frac{1}{2} \int_0^\pi L(x, \omega) \cos(\phi) d\omega
\end{aligned}$$

We want to find the first angle, that hits the area light and the last angle that hits the area light. Trigonometric examination shows that those angles can be calculated with

$$\begin{aligned}
\phi_{lower} &= \frac{\pi}{2} - \tan^{-1}(p+1) \\
\phi_{upper} &= \frac{\pi}{2} + \tan^{-1}(1-p)
\end{aligned}$$

This can be seen when examining that the triangle that is formed from the top left point of the lightsource  $(-1, 1)$  with  $(p, 0)$  and  $(-1, 0)$  has an angle  $\alpha'$  at  $(-1, 1)$  with  $\frac{p-(-1)}{1} = \tan(\alpha')$ . Since a triangle has a total inner angle of  $\pi$  and it has a right angle at  $(-1, 0)$ , the angle  $\alpha$  at  $(p, 0)$  with  $\alpha = \pi - \frac{\pi}{2} - \alpha' = \frac{\pi}{2} - \tan^{-1}(p+1)$ . This  $\alpha$  is the desired angle  $\phi_{lower}$ . Note that this even holds for  $p < -1$ . In a similar way  $\phi_{upper}$  is found. Examine the triangle  $((p, 0), (1, 0), (1, 1))$ . The angle  $\alpha'_2$  at  $(1, 1)$  has  $\tan(\alpha'_2) = \frac{1-p}{1}$ . Therefore the angle  $\alpha_2$  at  $(p, 0)$  is  $\alpha_2 = \pi - \frac{\pi}{2} - \tan^{-1}(1-p) = \frac{\pi}{2} - \tan^{-1}(1-p)$ . The desired angle  $\phi_{upper}$  is the remaining part of the half circle with  $\phi_{upper} = \pi - (\frac{\pi}{2} - \tan^{-1}(1-p)) = \frac{\pi}{2} + \tan^{-1}(1-p)$ .

$$= \frac{1}{2} \left( \int_0^{\phi_{lower}} L(x, \omega) \cos(\phi) d\omega + \int_{\phi_{lower}}^{\phi_{upper}} L(x, \omega) \cos(\phi) d\omega + \int_{\phi_{upper}}^{\pi} L(x, \omega) \cos(\phi) d\omega \right)$$

The first and the third integral refer to incident angles that point past the light source, therefore they are 0. The second integral refers to incident angles pointing to the lightsource, where  $L(x, \omega) = 1$

$$\begin{aligned} &= \frac{1}{2} \int_{\phi_{lower}}^{\phi_{upper}} 1 \cdot \cos(\phi) d\omega \\ &= \frac{1}{2} \int_{\phi_{lower}}^{\phi_{upper}} \cos\left(\left|\omega - \frac{\pi}{2}\right|\right) d\omega \\ &= \frac{1}{2} \left( \int_{\frac{\pi}{2} - \tan^{-1}(p+1)}^{\frac{\pi}{2}} \cos\left(\frac{\pi}{2} - \omega\right) d\omega + \int_{\frac{\pi}{2}}^{\frac{\pi}{2} + \tan^{-1}(1-p)} \cos\left(\omega - \frac{\pi}{2}\right) d\omega \right) \\ &= \frac{1}{2} \left( \cos\left(\frac{\pi}{2} - \arctan(p+1)\right) - \cos\left(\arctan(-p+1) + \frac{\pi}{2}\right) \right) \\ &= \frac{1}{2} \left( \frac{1-p}{\sqrt{(p-2)p+2}} + \frac{p+1}{\sqrt{p(p+2)+2}} \right) \end{aligned}$$

**b)**

$$\begin{aligned} L(x, \omega_o) &= L_e(x, \omega_o) + \int_{y \in S} f_r(\omega_i, x, \omega_o) \cdot L(y, -\omega_i(x, y)) \cdot \frac{\cos \phi_i \cos \phi_y}{|x - y|} \cdot dy_x \\ &= L_e(x, \omega_o) + \int_{-1}^1 f_r(\omega_i, x, \omega_o) \cdot L(y, -\omega_i(x, y)) \cdot \frac{\cos \phi_i \cos \phi_y}{|x - y|} \cdot dy_x \end{aligned}$$

If we are not in a light source this is:

$$\begin{aligned} &= \int_{-1}^1 f_r(\omega_i, x, \omega_o) \cdot L(y, -\omega_i(x, y)) \cdot \frac{\cos \phi_i \cos \phi_y}{|x - y|} \cdot dy_x \\ &= \int_{-1}^1 f_r(\omega_i, x, \omega_o) \cdot L(y, -\omega_i(x, y)) \cdot \frac{\cos \phi_i \cos \phi_y}{\sqrt{(x_x - y_x)^2 + (x_y - y_y)^2}} \cdot dy_x \\ &= \int_{-1}^1 f_r(\omega_i, x, \omega_o) \cdot 1 \cdot \frac{\cos \phi_i \cos \phi_y}{\sqrt{(x_x - y_x)^2 + (x_y - y_y)^2}} \cdot dy_x \\ &= \end{aligned}$$

### 4.3 Simple Path Tracer