





Oncology

In randomized head-to-head comparisons with either
CAP (cyclophosphamide, Adriamycin*, prednisone)¹ or chlorambucil²
FLUDARA® induced:

SUPERIOR RESPONSE RATES



LONGER REMISSIONS



- Fludara has been shown to be better tolerated than CAP¹...
(NAUSEA/VOMITING 5% WITH FLUDARA VS. 25% WITH CAP, p<0.001;
ALOPECIA/COMPLETE HAIR LOSS 2% WITH FLUDARA VS. 65% WITH CAP, p<0.001)
- ...and as well tolerated** as chlorambucil.²

**FLUDARA "SHOULD BE USED AS SOON AS A LACK OF RESPONSE
TO STANDARD THERAPY IS OBSERVED." ^{3§}**

* Adriamycin is a registered trade mark of Pharmacia & Upjohn.


** Patients on Fludara had a higher incidence of grade III/IV leukopenia than patients on chlorambucil.

§ Single agent Fludara is generally well tolerated at recommended doses. The most common side-effect observed is myelosuppression (anemia, leukopenia, neutropenia, and thrombocytopenia). Please see Prescribing Information for additional warnings.

References:

1. The French Cooperative Group on CLL, *et al.* Multicentre prospective randomised trial of fludarabine versus cyclophosphamide, doxorubicin, and prednisone (CAP) for treatment of advanced-stage chronic lymphocytic leukaemia. *Lancet* 1996; **347**:1432-8.
2. Rai KR, *et al.* Cancer and Leukemia Group B, Chicago, IL. A Randomized Comparison of Fludarabine and Chlorambucil for Patients With Previously Untreated Chronic Lymphocytic Leukemia. A CALGB, SWOG, CTG/NCI-C and ECOG Inter-Group Study. *Blood* 1996; **88** (Suppl 1):141a.
3. Montserrat E, *et al.* Fludarabine in Resistant or Relapsing B-Cell Chronic Lymphocytic Leukemia. The Spanish Group Experience. *Leukemia and Lymphoma* 1995; **21**:467-72.

Fludara®
(fludarabine phosphate)



FOR INJECTION

CHOOSE IT FIRST, IN SECOND LINE

PAAB
CCPP

MEMBER
PMAC