Exploring Lagrangian Optimization

Aaron @philosolog
Brennan @Brensum
Jordan @Jadams06
Kerem @Ottoerm7
Oliver @aureliusandreas

Contents

Section 1: The Extreme Value Theorem in $\mathbb{R}^2$

Chapter 1

Hungry Joe

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Chapter 2

Utilmaxxing

Theorem 1 († The Extreme Value Theorem in $\mathbb{R}^2$)**.** Suppose that $f(x)$ is continuous on the interval $[a, b]$ then there are two numbers $a \leq c, d \leq b$ so that $f(c)$ is an absolute maximum for the function and $f(d)$ is an absolute minimum for the function.

Section 2: The Extreme Value Theorem in $\mathbb{R}^3$

Chapter 3

Hangry Joe

Chapter 4

He's a Nerd!

Section 3: The Method of Lagrange Multipliers

Chapter 5

Poor Joe

Chapter 6

Joe's Math

Chapter 7

A Brief Generalization

Section 4: The Cobb-Douglas Production Function

Chapter 8

Rich Joe

Chapter 9

Business Joe

Chapter 10

Another Brief Generalization

Section 5: The Stuff I Thought I Should Put at the End but Wasn't Sure if It Was Necessary

Chapter 11

Concluding Remarks