

Calibration of colour gradient bias in shear measurement using CANDELS

X. Er¹ ^{*}, H. Hoekstra², T. Schrabback³, V. F. Cardone¹, R. Scaramella¹, R. Maoli⁴, M. Vicinanza^{1,4,5}, B. Gillis⁶ L. Miller⁷, J. Rhodes^{8,9}

¹ I.N.A.F. - Osservatorio Astronomico di Roma, via Frascati 33, 00040 - Monte Porzio Catone, Roma, Italy

² Leiden Observatory, Leiden University, PO Box 9513, NL-230 RA, Leiden, the Netherlands

³ Argelander Institute fuer Astronomie, Auf dem Huegel 71, D-53121 Bonn, Germany

⁴ Dipartimento di Fisica, Universita di Roma "La Sapienza", Piazzale Aldo Moro, 00185 - Roma, Italy

⁵ Dipartimento di Fisica, Universita di Roma "Tor Vergata", via della Ricerca Scientifica 1, 00133 - Roma, Italy

⁶ Royal Observatory, University of Edinburgh, Blackford Hill, Edinburgh EH9 3HJ, UK

⁷ Department of Physics, Oxford University, keble Road, Oxford OX1 3RH, UK

⁸ Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA

⁹ California Institute of Technology, 1200 East California Blvd, Pasadena, CA 91125, USA

Accepted –; received –; in original from 17 April 2017

ABSTRACT

Euclid will image about two billion galaxies that can be used to infer cosmological parameters using weak gravitational lensing. Exploiting the precision afforded by these data relies critically on our ability to correct for instrumental effects, such as the convolution by the point spread function (PSF). A complication is the fact that the optical data are obtained using a broad bandpass (550–920 nm) while the diffraction-limited PSF depends on wavelength. This leads to biases in the recovered galaxy shapes because the colours of galaxies vary spatially. We show that the colour-gradient bias can be determined with high accuracy in simulated noisy data. We also find that higher order image distortions, such as flexion, enhance the bias, which may be relevant for the study of lensing in high density regions. We estimate the size of this colour-gradient biases using multi-band observations from the *Hubble* Space Telescope and find correlations with the colours and sizes of the galaxies, but do not observe a significant dependence with redshift. **We need some concluding remark about the impact for Euclid.**

Key words: cosmology, weak lensing, systematics

1 INTRODUCTION

The images of distant galaxies are distorted, or sheared, by the tidal effect of the gravitational potential generated by intervening matter, an effect commonly referred to as weak gravitational lensing (e.g. Bartelmann & Schneider 2001, for a detailed introduction). The measurement of the resulting correlations in the shapes can be used to directly infer the statistical properties of the mass distribution in the Universe, which in turn provide constraints on cosmological parameters. Hence weak gravitational lensing by large-scale structure, or cosmic shear, has been identified as a powerful tool for cosmology. The measurement of the signal as a function of cosmological time is sensitive to the expansion history and the growth rate of large-scale structures, and can be used to constrain models for dark energy and modified gravity.

A useful measurement of the cosmic shear signal requires averaging over large numbers of galaxies to reduce the uncertainty caused by the intrinsic ellipticities of galaxies. The result is, how-

ever, only meaningful if biases in the shape estimates are negligible. Various instrumental effects change the observed ellipticities by more than the typical lensing signal, which is of order one per cent. The most dominant source of bias is the smearing of the images by the point spread function (PSF), driving the desire for space-based observations (Paulin-Henriksson et al. 2008; Massey et al. 2013). Despite these observational challenges, the most recent cosmic shear studies are starting to yield competitive constraints on cosmological parameters (Heymans et al. 2013; Jarvis et al. 2016; Jee et al. 2016; Hildebrandt et al. 2017). These results are based on surveys of modest areas of the sky, which limits their ability to study the nature of dark energy, which requires more than an order of magnitude improvement in precision.

Such a measurement is the objective of *Euclid* (Laureijs et al. 2011), the dark energy mission of the European Space Agency (ESA) scheduled for a launch in 2020, which will survey the 15 000 deg² of extragalactic sky that has both low extinction and zodiacal light. To reduce the detrimental effects of noise on the shape measurements, the images used for the lensing analysis are observed using a wide bandpass (550–920 nm). The much smaller PSF in

^{*} er.xinzhong@oa-roma.inaf.it

space-based observations is a major advantage, but the diffraction-limited PSF leads to new complications.

The most prominent one is that the correction for the smearing by the chromatic PSF depends on the spectral energy distribution (SED) of the galaxy of interest (Cypriano et al. 2010; ?) and ignoring this would lead to significant biases in the case of *Euclid*. Fortunately this can be accounted for using the supporting broadband observations that are used to derive photometric redshifts for the sources: the correction employs an effective PSF which is derived from the estimate of the observed SED of the galaxy. This correction is sufficient if the SED does not vary spatially (?). If this is not the case, the underlying brightness distribution, which is needed for an unbiased estimate of the shear, cannot be unambiguously recovered from the observed images. This results in a higher order systematic bias, which we call colour-gradient (hereafter CG) bias. As shown by Semboloni et al. (2013) (S13 in the rest of this paper) the amplitude depends on several factors: the SED of the galaxy, the relative size of the galaxy to the PSF, and the width of the bandpass, $\Delta\lambda$. For instance, the bias scales as $\Delta\lambda^2$, and thus is particularly relevant in the case of *Euclid*.

Galaxies show a wide variety in colour gradients, with elliptical galaxies typically showing negative colour gradients (redder in the centre and bluer in the outskirts), with steeper gradients more commonly found in bluer or more luminous early type galaxies (e.g. ??). ?? also suggest correlations between colour gradients and the overall colours and luminosities of the galaxies. Hence the relation between galaxy morphology and density may cause the CG bias to vary across the sky and lead to correlations with the lensing signal itself.

It is important that all systematic sources of biases are accounted for to a level that is smaller than the statistical uncertainties. In the case of *Euclid* this leads to tight requirements, as detailed in (e.g. Massey et al. 2013; Cropper et al. 2013). Initial studies by Voigt et al. (2012) and S13 used simulated images to show that the CG bias could be substantial, exceeding nominal requirements for the multiplicative bias in the shear. They also argued that it should be possible to calibrate the bias using *Hubble* Space Telescope (HST) observations of a large sample of galaxies in the F606W and F814W filters. However, their conclusion is based on the analysis of simulated noiseless data. In this work, we revisit the issue of the calibration of CG bias, with a particular focus on determining the bias from data with realistic noise levels.

In Section 2, we describe the main concepts and introduce the notation. We present the results from the analysis of simulated images in Section 3. In particular we explore the impact of having to use noisy data to measure the CG bias in Section 3.2. In Section 4 we estimate the CG bias using HST observations from the Cosmic Assembly Near-infrared Deep Extragalactic Legacy Survey (CANDELS; ?).

2 THE ORIGIN OF COLOUR GRADIENT BIAS

Following the notation of S13, we consider an image of a galaxy, and denote the photon brightness distribution of the image at each position θ and wavelength λ by $I(\theta; \lambda)$, which is related to the intensity $S(\theta; \lambda)$ by $I^0(\theta; \lambda) = \lambda S(\theta; \lambda) T(\lambda)$, where $T(\lambda)$ is the normalised transmission. We take this to be a top-hat with a width $\Delta\lambda$ around a central wavelength λ_{cen} . The resulting image of the galaxy, observed using a telescope with a PSF $P(\theta; \lambda)$ is given by:

$$I^{\text{obs}}(\theta) = \int_{\Delta\lambda} I^0(\theta; \lambda) * P(\theta, \lambda) d\lambda, \quad (1)$$

where $*$ denotes a convolution.

A measurement of the ellipticity of a galaxy provides an unbiased (but noisy) estimate of the weak gravitational lensing signal, quantified by the complex shear $\gamma = \gamma_1 + i\gamma_2$. The ellipticity ϵ in turn can be determined from the second order brightness moments Q_{ij}^0 of the PSF-corrected image $I^0(\theta)$ (?):

$$\epsilon_1 + i\epsilon_2 \approx \frac{Q_{11}^0 - Q_{22}^0 + 2iQ_{12}^0}{Q_{11}^0 + Q_{22}^0 + 2(Q_{11}^0 Q_{22}^0 - (Q_{12}^0)^2)^{1/2}} \quad (2)$$

where the second order brightness moments are given by¹

$$Q_{ij}^0 = \frac{1}{F} \int I^0(\theta) \theta_i \theta_j d^2\theta \quad (i, j = 1, 2), \quad (3)$$

where $F = \int d^2\theta I^0(\theta)$ is the total observed photon flux.

In practice, however, the observed moments are measured from the PSF-convolved image given by Eqn. (1). Moreover, the moments are evaluated using a weight function $W(\theta)$ to reduce the effect of noise in the images. Hence, the observed quadrupole moments are given by

$$Q_{ij}^{\text{obs}} = \frac{1}{F_w} \int_{\Delta\lambda} d\lambda \int d^2\theta I^0(\theta; \lambda) * P(\theta, \lambda) \theta_i \theta_j W(\theta), \quad (4)$$

where F_w is the weighted flux. The use of a weight function biases the observed moments, and the aim of moment-based shape measurement algorithms is to correct for this using estimates of the higher order moment (e.g. ?Melchior et al. 2011). An alternative approach is to fit sheared, PSF-convolved models to the observed images (e.g. ?Miller et al. 2013); in these fitting methods the profile itself acts as a weight.

S13 showed that the inevitable use of a weight function gives rise to the CG bias, which scales as $\Delta\lambda^2$. The bias also depends on the choice of the weight function, and vanishes in the case of *unweighted* moments. In the latter case it is possible to determine the PSF-corrected moments Q_{ij}^0 from the observed quadrupole moments because

$$Q_{ij}^{\text{obs}} = Q_{ij}^0 + P_{ij}^{\text{eff}} \quad (5)$$

where P_{ij}^{eff} are the quadrupole moments of the effective PSF, defined as

$$P_{\text{eff}}(\theta) = \frac{1}{F} \int d\lambda P(\theta, \lambda) F(\lambda), \quad (6)$$

where $F(\lambda)$ is the photon flux as a function of wavelength, which is directly related to the spectral energy distribution (SED) of the galaxy. Hence the correction for the chromatic PSF requires an estimate of the SED. ? have shown that the broadband observations that are used to determine photometric redshifts for *Euclid* can also be used to estimate the effective PSF with sufficient accuracy to meet the stringent requirements presented in Cropper et al. (2013).

Our approach to evaluate the multiplicative bias in the lensing signal that is introduced by colour gradient is similar to the one in S13. Figure 1 shows the flowchart of the steps that enable us to evaluate the CG bias. In both cases we start with the same wavelength dependent image $I^0(\theta; \lambda)$, but the bottom flow resembles what happens in the actual observations: the original image is

¹ We implicitly assume that the moments are evaluated around the position where the dipole moments vanish.




Figure 1. Flowchart describing how the colour-gradient bias is determined. The initial image is the same in both flows, but in the top flow an image without a colour gradient is created to which a shear is applied. In the bottom flow, the image is sheared before the PSF steps are applied. The ellipticities of the resulting images differ slightly, and can be used to quantify the bias that is introduced.

sheared² before the convolution with the PSF. The deconvolution with the effective PSF then yields the PSF-corrected shape. In the top flow the PSF steps are applied first, resulting in an image without a colour gradient that is subsequently sheared.

We measure the ellipticities of the resulting images to estimate the CG bias. To reduce noise in our estimate of the multiplicative bias m we use the ring-test method (Nakajima & Bernstein 2007) where we create eight copies of the original galaxy but with different orientations. The ensemble averaged ellipticities then provide an estimate of the multiplicative CG bias, m (we do not explore additive bias here), via

$$m = \frac{\epsilon_i^{\text{CG}}}{\epsilon_i^{\text{NCG}}} - 1, \quad (7)$$

where ‘CG’ indicates the case where the galaxy has a colour gradient, and ‘NCG’ is the galaxy with a uniform colour. Note that our approach differs slightly from that in S13, who quantify the response of the observed ellipticity to an applied shear. Hence, they do not apply the last step in the bottom flow (the deconvolution), but rather convolve the final image in the top flow. The steps presented in Fig. 1 yield a more symmetric result, highlighting the fact that the CG bias is the consequence of the fact that the shearing of the image does not commute with the convolution with the PSF. However, we verify in Sect. 3 that we recover the results of S13 (see Fig. 3).

3 COLOUR GRADIENT BIAS IN SIMULATED DATA

The CG bias is a higher order systematic bias, and thus the changes in the measured ellipticities are small. It is therefore important to verify that numerical errors in the calculations are subdominant compared to the small effects we aim to measure. To do so, we compare results from two independent codes that are used to generate the simulated images: one is written in C/C++ and the other uses the python-based GalSim package (Rowe et al. 2015), which

is widely used to create simulated images (e.g. Fenech Conti et al. 2016; ?). In the C code we compute the image by multiplying the surface brightness at the centre of each pixel using a sheared Sérsic profile with the pixel area. In the case of GalSim we use the Shear() function (which convolves the image by the pixel). Since we are interested in small differences in the shapes of deconvolved images, we checked for numerical errors. We convolved and subsequently deconvolved elliptical images. Comparison of the recovered ellipticities revealed small differences between the codes that ranged from 10^{-7} to 10^{-6} , two orders of magnitude smaller than the CG biases we are concerned with. Hence can safely neglect this numerical artefacts here.

As a further test we compare directly to the results obtained by S13 for two reference galaxy models. The reference galaxies are modeled as the sum of a bulge and disk component. To describe the wavelength dependence of the images we use the galaxy SED templates from ?: we use the SED for an elliptical galaxy for the bulge and take the SED of an irregular galaxy for the disk. This choice ensures that the resulting colour gradients are large. The two components are described by a circular Sérsic profile:

$$I_S(\theta) = I_0 e^{-\kappa (\frac{\theta}{a})^{1/n}}, \quad (8)$$

where I_0 is the central intensity, and $\kappa = 1.9992 n - 0.3271$. For the bulge component we adopt $n = 1.5$ and for the disk we use $n = 1$. The profiles are normalised such that the bulge contains 25% of the flux at a wavelength of 550 nm. The galaxies are circular and the sizes for the bulge and disk for galaxy ‘B’ are $0''.17$ and $1''.2$, respectively. The second galaxy ‘S’ is smaller with sizes of $0''.09$ and $0''.6$ for the bulge and disk, respectively (also see Table 3 in S13). We create images with a size of 256×256 pixels, and resolution 0.05 arcsec/pixel at wavelengths 1 nm apart and sum these in the range [550 : 920] nm to mimic the Euclid pass-band.

To create the PSF-convolved images we consider several PSF profiles. For a direct comparison with S13 use their reference PSF (PSF1), which is a Gaussian with a wavelength dependent dispersion

$$\sigma_{\text{PSF}}(\lambda) = 0''.102 \left(\frac{\lambda}{800\text{nm}} \right), \quad (9)$$

² We use $\gamma_1 = 0.05$ and $\gamma_2 = 0.02$ as reference, but we verified that other values yield similar results.




Figure 2. Comparison of the obscured Airy profile (solid), which is a good approximation to the *Euclid* PSF, to PSF1 (Gaussian; dashed) and PSF3 (compact Gaussian and top-hat; dotted) from S13. The profiles for 550 nm are indicated by the blue lines and the results for 920 nm are shown in red.

As discussed in S13 this PSF has a similar size as the nominal *Euclid* PSF, but a steeper wavelength-dependence. To approximate the *Euclid* PSF they consider a model that consists of a compact Gaussian core and an appropriately scaled top-hat (PSF3 in S13). Instead we use here a more realistic obscured Airy profile, which is actually close to the *Euclid* design profile (Laureijs et al. 2011):

$$P(\theta) = \frac{I_0}{(1 - \epsilon^2)^2} \left(\frac{2J_1(x)}{x} - \frac{2\epsilon J_1(\epsilon x)}{x} \right)^2, \quad (10)$$

where I_0 is the maximum intensity at the center, ϵ is the aperture obscuration ratio, and $J_1(x)$ is the first kind of Bessel function of order one; x is defined as $x = \pi\theta/\lambda D$. In the case of *Euclid* $D = 1.2\text{m}$ and $\epsilon = 1/3$. We compare this model to the Gaussian case and PSF3 from S13 in Fig. 2 at the 550 nm and 920 nm.

As a test of our implementation of the pipeline to determine the CG bias we compare our results to those from S13. We reproduced all their results, but here we only show the comparison when θ_w , the width of the weight function that is used to compute the quadrupole moments, is varied. As discussed in Sect. 2 the amplitude of the noise bias depends on the width of the weight function that is used to compute the (weighted) quadrupole moments.

The results of this test are presented in Fig. 3. The solid lines indicate the results from S13 for the two reference galaxies B (red) and S (blue). We compare these to the outcome of the C code (dashed lines) and the GalSim code (dotted lines). The agreement is excellent for the large galaxy (B) for the full range in θ_w . Our two codes agree also well for the small galaxy (S), but the difference with S13 is noticeable. However, given the consistent results between the C and GalSim code we conclude that numerical errors are negligible in our implementation. In the remainder, we limit the simulations to those generated with GalSim.

Do we understand why there is a difference for the blue lines? I see there is a comment in the tex file that the parameters for the galaxies were slightly different? Can this be fixed?

Figure 3 shows that the CG bias decreases rapidly when the width of the weight function is increased. This allows for an interesting trade-off between CG bias and noise bias. The latter increases with increasing θ_w but relatively slowly (see Fig. 4 in S13). As a proxy for the optimal weight function (which maximizes the signal-to-noise ratio) we adopt the value of the half-light radius in




Figure 3. The CG bias in shear versus width of the weight function (in units of the half-light radius w_h) used to compute the quadrupole moments. The solid lines correspond to the results from S13 for the large ('B'; red) and small ('S'; blue) reference galaxy. The dashed (dash-dotted) lines are our results for images simulated using the C (GalSim) code. In the top panel, we use the same PSF as in S13, and in the bottom panel, we use the Airy profile as PSF.

the remainder of this paper. This yield $m = 0.8 \times 10^{-3}$ for galaxy B and $m = 2 \times 10^{-3}$ for galaxy S, demonstrating that the CG bias is a strong function of galaxy size.

3.1 Impact for large distortions

Besides shear, there are higher order image distortions in weak lensing, such as flexion (e.g. ??). In previous analysis, we only consider the shear effect in the lensing process, while the higher order image distortions also suffer from CG bias as well. In additional tests, we perform the complete lens ray-tracing instead of solely shearing the images. (We only use our C code in this part, since the current version of GalSim only provides shear effect.) We adopt an Singular Isothermal Sphere halo model as the lens model, and vary the separation between the lens and the source image. The results are presented in Fig. 4. The solid (dotted) lines indicate the results from complete lens process (sole shear process). The solid lines approach to the dotted lines with the decreasing of the shear. When the shear becomes greater than 0.1, one needs to take account of the complete lens process. However, the typical shear value is about a




Figure 4. The CG bias versus tangential shear in complete lensing process (solid lines). The dotted lines correspond to the results from only shear process.

few per cents. The complete lens process are necessary only in the strong lensing region or in some galaxy cluster lensing.

3.2 Calibration of CG bias using simulated HST images

The *Euclid* observations lack high-resolution multi-band images to measure the CG bias directly for each source galaxy. However, the cosmological lensing signal is typically inferred from the ellipticity correlation function (REF), which involves averaging the shapes of large ensembles of galaxies. Provided the average bias that is caused by colour gradients is known for a selection of sources, it is possible in principle to obtain unbiased estimates of the ellipticity correlation function. Here it is particularly important that the correction for the average CG bias accounts for the variation in redshift and colour. The former is relevant for tomographic cosmic shear studies, whereas the latter avoids significant spatial variation in the bias because of the correlation between galaxy colour, or morphology, and density.

S13 showed that HST observations in both the F606W and F814W filters can be used to determine the CG bias to meet *Euclid* requirements. However, S13 did not consider the complicating factor that the HST images themselves are noisy. Although the HST data are typically deeper than the nominal *Euclid* data, and the HST is considerably smaller, it is nonetheless necessary to investigate the impact of noise in more detail. We therefore address this particular question here, before determine the CG bias from actual HST data in Sect. 4.

The method to calibrate the CG bias using observations in two bands is described in detail in S13, but here we outline the main steps for completeness. To model the wavelength dependence of the image we use two narrow-band³ images, each of which is given by:

$$I_i(\theta) = \int_{\Delta\lambda_i} T_i(\lambda) I(\theta, \lambda) d\lambda, \quad (11)$$

³ To distinguish these filters from the broad VIS pass-band we refer to the F606W and F814W as narrow bands, but acknowledge that these are commonly referred to broad-band filters and that genuine narrow-band filters are significantly narrower.

where $T_i(\lambda)$ is the transmission of the i th narrow filter. We assume that for each pixel the wavelength dependence of the image can be interpolated linearly:

$$I(\theta, \lambda) \approx \tilde{I}(\theta, \lambda) = a(\theta)\lambda + b(\theta). \quad (12)$$

Eqs.11 and 12 yield a linear set of equations on each pixel, which can be used to solved for the coefficients (a, b) :

$$T_{ai}\lambda a(\theta) + T_{bi}b(\theta) = I_i(\theta), \quad i = 1, 2, \quad (13)$$

where $T_{ai, bi}$ is the integrated transmission function at two filters. With (a, b) and Eq.12, one can obtain approximated galaxy images of each wavelength $I(\theta, \lambda)$. Then we will follow the same procedure as in previous section to estimate the CG bias. **need to rephrase and think about the notation.**

We first consider the recovery of the CG bias for noiseless observations of the two reference galaxies. We simulate the images in the F606W and F814W filters at different redshifts, ignoring evolution in the SEDs. We adopt the native sampling of the Advanced Camera for Surveys (ACS) on HST of 0.05 arcsec/pixel. As shown in S13, we cannot ignore the blurring of the observed images by the HST PSF, and to mimic this we assume an obscured Airy function for a mirror with diameter $D = 2.5$ and obscuration 0.33 as a proxy for the HST PSF. We deconvolve our synthetic HST images and create the images at different wavelengths as the starting point for the flow presented in Fig. 1.

Following S13, we show the CG bias as a function of redshift for galaxy ‘B’ (left panels) and ‘S’ (right panels) in Fig. 5. The results for the actual CG bias are indicated by the solid black lines, whereas the dashed black lines indicate the recovered values from the noiseless synthetic HST observations in the F606W and F814W filters. These results show that the CG bias varies significantly with redshift. The bottom panels show the residuals between the recovered and the true bias. **we should also show the residual for the noiseless case!** The residual bias is within the target tolerance for *Euclid*, indicated by the grey band, for all redshifts.

This comparison, however, does not account for the fact that the HST images will contain noise. Although the *Euclid* data are shallower, it is important to quantify the impact of noise on our ability to recover the CG bias using archival HST data. To do so, we add Gaussian noise to the simulated HST images, where the r.m.s. noise level σ is determined by the signal-to-noise ratio of the galaxy, SNR, the total flux within an aperture of radius $1.5 r_h$, F_{tot} , and the number of pixels within this aperture, N_{tot} , such that

$$\sigma = \frac{F_{\text{tot}}}{\text{SNR} \sqrt{N_{\text{tot}}}}. \quad (14)$$

For reference, we compared the input SNR for the two reference galaxies to that estimated by SExtractor (?). We find good agreement for galaxy ‘B’ for SNR values ranging from 5 to 50 in both HST filters. The agreement is also good for the ‘S’ galaxy, but SExtractor returns lower values if the input SNR is larger than 30. We consider two noise levels: a SNR=50 corresponds roughly to the typical galaxy used in the *Euclid* weak lensing analysis; a SNR=15 corresponds to the faintest galaxies that might be used. **we should see what magnitude SNR=50 and SNR=15 correspond to in the actual HST data.**

The deconvolution of noisy images is problematic, because the presence of noise will lead to biased estimates of the underlying galaxy. Instead we regulate the problem by assuming that galaxies can be fit by a bulge and disk component, each described by a Sersic profile. We fit this model, convolved with the PSF, to the noisy images in each band and use the best fit model to compute




Figure 5. The CG bias in shear measurement as a function of redshift using simulated images. The black solid lines are the true CG bias; the dashed lines are the estimation using two band HST images without noise; the colour lines are the estimation using noisy images of input $SNR = 50$, the value are the average over 40 realizations each redshift. In the bottom, Δm is the residual with respect to the true CG bias, and the error bars show the standard variations (Eq.15). Up(Down)-figure are the results for B-(S-)galaxy. **the legend states that this is for SNR=15?**

	S-606W	S-814W	B-606W	B-814W
n_1	0.5-2.5	0.5-2.5	0.5-2.5	0.5-2.5
n_2	0.5-2.5	0.5-2.5	0.5-2.5	0.5-2.5
R_{bulge}	1-10	1-10	3-30	3-30
R_{disk}	5-30	5-30	10-60	10-60
q	0.6-1	0.6-1	0.6-1	0.6-1

Table 1. Constraints for the fitting parameters in Galfit. The first two columns are for two images of the S-galaxy, the other two are the image of B-galaxy. n_1 is the Sersic index for bulge, and n_2 is the Sersic index for disk. The effect radius is given in unit of pixel (0.05 arcsec).

the CG bias. To perform the fit, we use GALFIT (?) with the prior constraints on the galaxy parameters (Sersic index, effective radius, and axis ratio) listed in Table 1.

We use combine the images in the two filters and use SEXTRACTOR to estimate the center and some initial galaxy parameters that are used as the starting point by GALFIT. The resulting best fit images depend somewhat on these initial values, and thus could affect the estimate for the CG bias. This will be more prominent as the SNR of the images decreases. To explore this we perform the fits using two sets of initial parameters: in the first we leave all parameters free, while in the other case we fix the Sersic index to its simulated value, but leave the other parameters free.

We use the best fit models to compute the CG bias, following the same algorithm as was used to compute the signal in the noiseless case. We show the results in Fig. 5 for $SNR=50$. We find that fixing the Sersic index (blue line) or leaving all parameters free (red line) results in a similar CG bias as a function of redshift. Moreover, the results closely resemble the noiseless case (dashed lines). The residuals presented in the bottom panel of Fig. 5 shows that for the $SNR=50$ case, the average CG bias can be determined with an accuracy that meets the adopted *Euclid* tolerance, indicated by the grey band.

This needs to be clarified: For the orange lines we also free all the parameters, but we perform another convolution with the effective PSF (see appendix for more detail). In Fig.1 one can see that the convolution is canceled with the last deconvolution for the CG images. The reason for that is the deconvolution may cause some numerical errors, especially for the small size noisy images. Therefore, in the following result for noise images, we will perform the convolution to the NCG image instead of deconvolution to the CG one.

Moreover, although we use the circular source image in the simulation, the fitted noisy images will become slightly elliptical. In order to get rid of the error due to the “intrinsic shape”, we rotate the source image 6 times, and use the average value as our estimate for the galaxy ellipticity (?). At each redshift, we use one noise free image and 40(200, 300) noisy images for $SNR = 50$ ($SNR = 15$). In the bottom panel of each figure, we also show the residues with respect to the true CG bias, and the error bars, which are given by the standard deviation from e_{cg} and e_{ncg}

$$\sigma_m = |m| \sqrt{\left(\frac{\sigma_{cg}}{\langle e_{cg} \rangle}\right)^2 + \left(\frac{\sigma_{ncg}}{\langle e_{ncg} \rangle}\right)^2}. \quad (15)$$

One can see that in general our estimate can reproduce the properties of CG bias in ellipticity measurement using both ideal or noisy images. In the high SNR cases (Fig.5), all the estimates with different initial parameters basically agree with each other. The estimate with all fitting parameters free gives lower bias than that fixed the Sersic index. In the low SNR case (Fig.6), the fitting using all free parameters show better estimates for the CG bias, and the residual are within the requirement for most cases. The variance due to the image fitting with different initial parameters are larger than high SNR cases, but can still be reduced by large sample of images and reach the requirement. It means that for each single type of galaxy, one need about 300 galaxy images in order to calibrate the bias. The scatters in the estimation are proportional to the magnitude of the bias, i.e. we have large uncertainty for S-galaxy at low redshift. We also perform tests using images with lower SNR, and calculate the average residue with respect to true bias. When the SNR of images drop to 12, the average residue increases by about 70%(60%) for small(big) galaxy than that of $SNR = 15$. This test may not reflect the real dependence on the SNR of images, since in general the lower SNR, the smaller the image size. Therefore, image simulation with specific configuration of the survey is necessary, e.g. the limit magnitude, the population of the galaxies and the relation between the size and SNR etc. In additional tests using




Figure 6. Same as Fig. 5 but for images of SNR=15. 1000 realizations are used at each redshift bin for both B- and S-galaxy.

other measurement methods for PSF correction, such as the image fitting methods (e.g. ???), and KSB+ (e.g. ??), we find that the bias has similar dependence on the SED of the source galaxy in the two HST filter. In Fig. 8, we compare the SED of the galaxy in redshift $z = 0, 0.5$. One can see that at $z = 0.5$, besides the strong emission lines, the linear approximation cannot reflect the properties of the source galaxies. Thus for some galaxies with complex properties, two band image may not be sufficient to calibrate the CG bias.

In both two figures, we notice that at $z = 0.5$ and 0.9 , the estimates are significantly smaller than the true bias for both galaxy models. This is due to the uneven SED of the source galaxy in the two HST filter. In Fig. 8, we compare the SED of the galaxy in redshift $z = 0, 0.5$. One can see that at $z = 0.5$, besides the strong emission lines, the linear approximation cannot reflect the properties of the source galaxies. Thus for some galaxies with complex properties, two band image may not be sufficient to calibrate the CG bias.

3.3 PSF variations in calibration data from HST

So far we implicitly assumed that the simple axisymmetric PSF used to mimic the HST data is perfectly known. In reality, however, the HST PSF is more complex and varies spatially and as a function of time. The small field-of-view of ACS implies that typically a small number of stars can be used to model the PSF. We therefore examine how well the HST PSF properties need to be determined so that they do not affect the CG bias measurement significantly.

To do so, we generate models where we slightly increase the PSF size in the two bands. (**by how much?**) These models are used only in the step where we deconvolve the simulated HST images




Figure 7. The bias histogram from all the simulated images. The top (bottom) panel shows that from images of SNR=50 (SNR=15).




Figure 8. Spectral energy distributions used to create the disc and bulge components of our mock galaxies. They are normalized at $\lambda = 5500\text{\AA}$ by ratio $F_{\text{bulge}}/F_{\text{disk}} = 1/3$, and they are convolved with HST filter function in F606W and F814W. *left-* redshift 0; *right-* redshift 0.5.

in the absence of noise. The change in CG bias, Δm as a function of redshift is shown in Fig. 9 for the ‘B’ galaxy (top panel) and ‘S’ galaxy (bottom panel). The results for an increase in the PSF size in the F606W band are indicated by the solid, dashed and dotted lines, respectively; the red, green and blue lines indicate the impact of increasing the size of the PSF in the F814W band. The sensitivity to the PSF errors is typically larger for low redshift galaxies, but the change in CG bias is much smaller than the bias itself. As expected, small galaxies are more sensitive to errors in the estimate of the PSF size.

To test the impact of imperfect PSF modelling, we make two comparisons: Comparing a control set of “actual” PSFs to a mock set of stars using these PSFs, including some unresolved binaries;




Figure 9. The PSF in the deconvolution are varied to see the effect on the CG bias. From red, green to blue lines, we increase the size of PSF for the F814W; from the solid, dashed to dotted lines we increase the size of PSF for the F606W. **here we only need to show the residuals. Hence the top plots can be removed.**

and comparing the stars in actual images to the best-fit models. In the former case, we generate mock star fields using PSFs generated with the TinyTim tool (?) and allow 30 per cent of stars to be unresolved binaries, with another star laid on top of them within a radius of one pixel. In the latter case, we use actual star fields observed with HST and try two methods of fitting the model to the stars. In both cases, we perform this for both the F606W and F814W filters.

The model PSFs are generated using the default parameters for TinyTim where possible, and the appropriate camera, detector, and filter passband settings for each image. For the rest of the options, we use:

- **Position on Detector:** Chosen for each star based on its detected position.
- **Spectrum:** 1; 15 (Use the K7V spectrum, which is chosen to represent a typical star in the sample. The choice of a fixed spectrum for stars was found to have a negligible impact on the models.)
- **PSF Diameter:** 2.0 arcsec
- **Focus-Secondary Mirror Despace:** Fit per image. As HST “breathes” due to its varying angle relative to the Sun, the position of the focus relative to the secondary mirror changes over time and cannot be perfectly predicted for any given observation. We thus




Figure 10. Comparison of PSF between TinyTim model and stars. Top (Bottom) is for the B-galaxy (S-galaxy). The solid (dashed) lines are the result using undrizzled (drizzled) PSF. The dotted lines are that if we also include binary PSF effect.

have to fit the best focus position by simulating multiple sets of PSFs for each image.

- **Zernike Polynomials:** In the first test, “focus only”, set to default values. In the second test, “all parameters”, these are fit in addition to the focus despace.

- **Subsampling Factor:** $8\times$

Stars are selected such that they have a signal-to-noise ratio of at least 50, there are no detected objects within 1 arcsecond (20 pixels), and outlier rejection is performed in the fitting procedure. The stars are normalized and then stacked using inverse-variance weighting. The corresponding model PSFs are stacked using the same weights.

In Fig. 10, one can see that the inaccuracy of PSF can cause error on the estimation of CG bias. For the small size galaxy, the error can reach about 15% for undrizzled PSF, while for the big galaxy, the error is only a few percent, can be neglect. In the result using drizzled PSF, the difference becomes smaller. The most problematic PSF will be those from binary stars. One can see for both big and small galaxies, we have a significant error.

Field	Area (arcmin ²)	N_1	N_2	N_3
AEGIS	180	2094	2112	3460
COSMOS	139	1593	1656	2449
UDS	146	1455	1497	2341
Total	465	5142	5265	8250

Table 2. Size of the HST CANDELS data sample in F606W and F814W bands. The number of galaxies are shown in three selection methods to match the Euclid survey, $N_1: m_V < 25$ and $m_I < 24.5$, $N_2: m_{VIS} < 24.5$, $N_3: m_V < 25.5$ and $m_I < 25$.

4 CANDELS DATA

In the previous section we demonstrated that it should be possible to determine the CG bias from HST observations in the F606W and F814W filters, and that the effects of noise in the data do not bias the results. In this section we use actual HST observations to determine the expected CG bias in *Euclid* weak lensing measurements.

4.1 CANDELS data

To investigate the impact of colour gradients using realistic galaxies populations, we employ HST/ACS data taken in the F606W and F814W filters in the three CANDELS fields (AEGIS, COSMOS, and UDS), which have a roughly homogeneous coverage in both bands (see ???). We base our analysis on a tile-wise reduction of the ACS data, incorporating pointings that have at least four exposures to facilitate good cosmic ray removal, yielding combined exposure times of 1.3-2.3ks in F606W and 2.1-3.0ks in F814W. We employ the updated correction for charge-transfer inefficiency from ?, MultiDrizzle (?) for the cosmic ray removal and stacking, as well as careful shift refinement, optimised weighting, and masking for stars and image artefacts as detailed in ?. Schrabback et al. (in prep.) describe the generation of weak lensing catalogues for these images. We base our analysis on the galaxies passing their source selection and apply additional magnitude cuts as detailed below. To investigate the dependence of the colour gradient influence on galaxy colour and redshift, we match this galaxy catalogue to the photometric redshift catalogue from ?. We list the total non-masked areas in which these catalogues overlap in Table 2.

We match the galaxy from F606W and F814W bands by selecting that the difference of galaxy coordinates in two bands is smaller than 1 pixel (0.05 arcsec). Moreover, in order to resemble the *Euclid* wide survey, we apply a selection based on the magnitude of two bands. In the first selection, the galaxy must be brighter than magnitude 25 in the F606W band and 24.5 in the F814W band. In the second selection (VIS, N2 sample), we apply the linear interpolation from F606W and F814W band using the effective wavelength to approximate the Euclid VIS magnitude, and select the galaxy brighter than 24.5 in VIS. In the last one (N3 sample), we enlarge the sample by using lower threshold in two bands: 25.5 in V and 25.0 in I band. The amount of galaxies is listed in Table 2. The galaxy VIS magnitudes will determined by several factors, such as exposure time and filter transmission, etc. Thus, the actual number of galaxies can be different from what we have shown in this work. The approximation that we use rely on the flatness of the source spectrum. In the VIS sample, the number of galaxies is similar to that in the first selection in all three fields, which suggest that the spectra of our galaxies are relatively smooth. In the N3 sample the galaxy number is much higher. The total galaxy number and number density ($\sim 12/\text{arcmin}^2$) is lower than that in S13 for several rea-

sons: we only selected galaxies for which the 3D-HST photometric redshift are available, and those suitable for weak lensing analysis, e.g. the very big or bright galaxies are not include. Moreover, in S13 the galaxies are selected by the magnitude of F814W. However we select galaxies by the estimated VIS magnitude and well match in F606W and F814W.

There is no significant difference found in the distribution of galaxy parameters, such as effective radius, axis ratio, photometric redshift and colour (we use $m_{F606} - m_{F814}$ as colour in this work, and will write as m_{V-I} later). The SNR of most images in VIS sample are larger than 15, thus they will be able to provide relatively stable estimate for CG bias.

We also compare the galaxies from 3 catalogues (AEIGS, COSMOS, UDS). In the distribution of galaxy parameters (Fig. 11), we find no significant difference in half light radius (r_h), galaxy axis ratio or redshift distribution. However, in the colour distribution, there are more blue galaxies (small m_{V-I}) in AEGIS, which is more significant in N3 sample. We will see later that the colour of galaxy is also related to the CG bias in shape measurement.

4.2 CG bias from CANDELS

In this section, we use the real galaxy images to estimate the CG bias which will appear in the Euclid weak lensing survey. For the reference PSF of Euclid, we use the Airy model (Eq. 10), and we use the PSF model from TinyTim for the HST data.

We estimate the CG bias following the same procedure for the simulated galaxy:

- fit the galaxy using one Sersic component for both band images. Some constraints such as Sersic index ($0.5 < n < 5.0$), effective radius ($1 < r_e < 50$ pixel) and axis ratio ($0.6 < q < 1.0$) are adopted in Galfit.
- interpolate the SED on each pixel of the galaxy image, and generate the galaxy at each wavelength (Eq. 11 - 13), and then integrate over the wavelength to simulate the CG and NCG galaxy image.
- measure the shape of two images and calculate the bias m . We also apply 6 different orientation of the images in order to reduce the intrinsic shape noise.

In Fig. 12, we show the histogram of the CG bias. The CG bias in most of the galaxies is smaller than 0.01 (94%). We did not perform averaging as we did for the simulated noisy images, which can further reduce the scatter of the bias. Thus the actual bias, especially the scatter in the sample, is smaller. In the bottom panel, we estimate the CG bias using a larger weight function ($2r_h$). As one expected, the bias decreases by about one order of magnitude.

The colour gradients as a tracer of galaxy evolution have been found to be correlate with some aspects. We try to explore relations between CG bias and the properties of galaxy. First we show the relation between CG bias with two tracers of the colour gradients: the ratio of Sersic index from two bands, and the ratio of effective radius from two bands (Fig. 13). One can see that there is a linear relation between the bias and the radius, but for that of Sersic index it is not obvious. The reason is that the Sersic index is mainly account for the type of the galaxy, the radius correlates with the colour gradient directly. Moreover, the CG bias depends on several factors of the galaxy, e.g. the total size of the galaxy. It is not surprise to see large bias scatters for the whole sample. In principle, one expect that when $r_{e606} = r_{e814}$ and $n_{606} = n_{814}$, the CG bias in principle will be vanish, since the identical images from two bands will not have a colour gradient. This is confirmed




Figure 11. The histogram for basic galaxy properties in our sample. The lines with different colours represent galaxies from different catalogues (AEGIS, COSMOS, UDS). From left to right: half light radius, axis ratio, photometric redshift, and colour (m_{V-I}).

in our result: the blue line (bin average) in Fig. 13 meets zero at $r_{e814}/r_{e606} = 1$. However, those colour gradients information will not be available in Euclid. We also need the relation between the bias with other parameters.

In Fig. 12, the galaxies in AEGIS field have more positive CG bias than the galaxies in the other two. The colour distribution in AEGIS is different from the other two as well, which suggests the correlation between the CG bias and the colour of the galaxy. In Fig. 14, we show the relation between the CG bias and the colour of the galaxies. The bias is inversely proportional to the colour of the galaxies. This is consistent with the trend of total colour (e.g. ?): the bluer galaxies have positive colour gradients, while the redder ones have negative gradients. Since this marks a possible transition of two types of galaxies, we split the galaxy sample into two groups according to their colour: the red galaxies ($m_{V-I} > 0.5$) and the blue ones ($m_{V-I} < 0.5$). They are shown in the bottom panel of Fig. 14 as a function of redshift. Most of the red galaxies are located at moderate redshifts, mainly between redshift [0.5, 1.0], while the blue galaxies are either at the lower redshift ($z < 0.5$) or higher redshift ($z > 1.0$). The CG bias in red galaxies are obviously more negative than that of the blue ones. It again confirms that the colour/colour gradients is an important tracer of the galaxy evolution, since apparently galaxies at different stages of the evolution.

In addition, we stack the images from two bands as our mock VIS band images, together with the mock VIS magnitude (m_{VIS}).

In Fig. 15, we show the bias as a function of m_{VIS} . There is no obvious dependence on m_{VIS} , which seems conflict with some study of colour gradients, e.g. ? find tight relation between colour gradients and r -band magnitude. However, there are two points one needs to notice: 1) the actual m_{VIS} is different from our linear approximation, also the filter transmission are different of two telescope. 2) the more important thing is that the wide band magnitude may not contain sufficient information about the type or colour of galaxy, thus may not be a good tracer for CG bias.

The VIS image, on the other hand, contains more information. Fig. 16 shows the CG bias with the Sersic index and the effective radius fitted from the VIS images. In the top panel, we divide the sample into two groups by the fitted effective radius, either larger or smaller than 0.35 arcsec. The small (large) galaxies are shown by the blue (orange) points, and the blue (red) line is the bin average. The large galaxies cover a large range of Sersic index, have negative average CG bias. Most of small galaxies have small Sersic index (< 2.5). The bias of small galaxies are positive and approximately proportional to the Sersic index. The scatters of the bias for both large and small galaxies increase with the Sersic index. In the bottom panel, the galaxies are divided into three groups: the first is red galaxy whose color is large ($m_{V-I} > 1.0$); the second and third group are the rest galaxies either with large Sersic index ($n > 2.25$, elliptical galaxy) or small Sersic index (disk galaxy). The solid lines show the bin average over effective radius. We can see that the VIS image alone can also provide an rough estima-




Figure 12. CG bias histogram from CANDELS: different colours show the result from three catalogues. In the bottom panel we show the CG bias using different weight function ($r_w = 2r_h$).

tion for CG bias, but classification of the galaxies is necessary. As shown in the figure, the disk galaxies have small bias, small radius (< 1 arcsec), and also small bias scatters. The elliptical galaxies cover large radius range, and the bias is larger than the disk ones. The bias in red galaxies are significant, and mainly negative. The scatters of red galaxies are larger than the other two kinds of galaxies. Extra photometry can definitely provide more constraints on the CG bias, as it has been shown the correlations between colour gradients and other properties of galaxy. Moreover, the multi-band information is required for the photometric redshift study. One can obtain that for free to calibrate the CG bias. Although the dependence on the multi-band is different between photometric redshift and CG bias, the experience from photometric redshift can be used for CG bias, such as some machine learning algorithms.

We calculate the average bias and the dispersions over the redshift bins for both red ($m_{V-I} > 0.5$) and blue galaxies (Table 3). The red galaxies are mainly located between redshift (0.4, 1.2), while the blue galaxies are low density in redshift (0.8, 1.2). The bias from red galaxies are significantly smaller than that of blue galaxies, as one expected, the colour gradient in the elliptical galaxies are smaller. The dispersions of the bias in each bin are large, which probably indicate that in each bin there are several kinds and sizes of galaxies. Therefore, in order to calibrate the bias with high precision, one need bigger galaxy samples. From our simulation, we need about 200 galaxies for one type of galaxy in every redshift bin. If we make rough bins, for instance, 2 types of colour: red and blue; 5 different sizes from about 0.1 arcsec to 1.0 arcsec (Fig. 16), and 5 redshift bins, at least 20000 galaxies are required. For more




Figure 13. CG bias as a function of galaxy properties: ratio of Sersic index between two band (top) and effective radius between two bands (bottom). The blue lines are the average CG bias over the parameter bins.

photo-z	Number	\bar{m}	σ_m
0 – 0.4	187	-1.3×10^{-3}	0.012
0.4 – 0.8	1415	-7.6×10^{-4}	0.011
0.8 – 1.2	1116	-1.0×10^{-3}	0.017
> 1.2	245	-2.1×10^{-3}	0.015
0 – 0.4	667	6.6×10^{-4}	0.0026
0.4 – 0.8	513	2.8×10^{-4}	0.0028
0.8 – 1.2	187	1.2×10^{-3}	0.0041
> 1.2	935	5.4×10^{-4}	0.0046

Table 3. The number, average CG bias and dispersion in redshift bins for blue (bottom half) and red (top half) galaxies.

realistic SED classifications and redshift bins, several times larger sample are also necessary.

5 SUMMARY AND DISCUSSION

In the image survey for weak gravitational lensing, the wide band filter can provide high signal-to-noise images and large coverage of redshift range. There is however a shape bias due to the chromatic shape of galaxy and the PSF, which is named as colour gradient bias. For very wide band surveys, such as Euclid, this effect can cause a non-negligible bias. In this work, we exam such a kind of bias in measuring the shape of a galaxy using both simulated images and real data taken from the HST ACS CANDELS survey. In the simulated galaxy images, we confirm the bias behaviour from




Figure 14. CG bias as a function of galaxy properties, top: color (m_{V-I}), bottom: photo-z. In the bottom panel, the red and blue points are the bias for red ($m_{V-I} > 0.5$) and blue ($m_{V-I} < 0.5$) galaxies respectively. The lines are the average CG bias in the redshift bins.




Figure 15. CG bias as a function of mock VIS magnitude.

previous results (S13). We further apply the calibration method to the noisy images in the simulations, and find that with reasonable signal-to-noise ratio (SNR= 15) and sufficient numbers of galaxies (300 images for one type of galaxy in one redshift bin), we can estimate the CG bias to a high precision. However, the underestimate cannot be avoided due to strong emission lines, or the uneven SED of source galaxies. Moreover, the simulations are performed with only two galaxy models, and the SNR of the simulated images in two bands are assigned with equal value. In reality, the relation between the SNR with the size and SED of the galaxies has to be taken into account. We also perform comparison of TinyTim and




Figure 16. CG bias with Sersic index (top) and effective radius (bottom) from the mock VIS images. The unit of radius is a pixel (= 0.05 arcsec). In the top panel, the blue (red) is the average of small (big) galaxies. In the bottom panel, the red line is average bias of red galaxy ($m_{V-I} > 1$); the green line is that of elliptical galaxy ($n_{\text{Sersic}} > 2.75$); the blue line is for the disk galaxy.

star PSF models. The inaccuracy, especially that due to the binary stars, will cause errors in estimating the CG bias.

In the estimation using CANDELS data, we select the images from two filters (F606W, F814W). For most of the galaxy, the bias ($|m|$) is smaller than 0.01. As we find from simulated images, the estimation using noisy image has a large scatter, thus the CG bias in reality may be even smaller than that shown here. In our sample of galaxies, the CG bias shows a correlation with the colour of galaxies, and a linear relation with the ratio of two band images. We also generate the mock VIS band images. From the parameters of the image (Sersic index and effective radius), one can classify the galaxy in order to obtain tighter constraints on the bias. For example, the galaxies with small Sersic index, i.e. disk-like, have smaller CG bias. On the other hand, those with large Sersic index have large bias and also bias variation. The relations show consistent results about the colour gradient dependence on the properties of galaxy (e.g. ?). However, since the CG bias also depends on the relative size with respect to the PSF, the dependence of CG bias is certainly more complicate. We did not provide any fitting formula for the bias at current sample, since the scatters are too large. More importantly, it has to be performed according to the types or morphology of the galaxies, which require larger sample of data.

The multi-band photometry from several bands, which can be used to estimate the redshift, can be also use for the CG bias analysis. Although the redshift dependence is not significant in our sample of galaxies, this may not be the case for a larger survey, or if we look at the bias according to the type of galaxy. Moreover, the

colour of galaxy also indicates the evolution history, or the large scale tide force. It can be also used for the study of intrinsic alignment analysis in weak lensing (e.g. ?). Therefore, the dependence of the CG bias on the colour of the galaxies will further increase the sysmatics of the intrinsic alignment. The detail behaviour will require large cosmological simulations, which is beyond the scope of this paper, but definitely needs further studies for the project such as Euclid.

The role of environment on the colour gradients is not clear. On the one hand, it has been shown that colour gradients depend on the environment where galaxies reside, with steeper colour gradients in poor rather than rich clusters (e.g. ?), which is possible due to the different processes during galaxy formation. On the other hand, in recent study using integral field spectra from SDSS-IV, the metallicity gradients show weak or no correlation with density environment (?). In any cases, close galaxy pairs or nearby bright star(s) may also cause weak brightness/colour gradient, which will affect our estimate for CG bias as well. Moreover, in the calibration we linearly interpolate the SED on pixels from two bands. Some advanced method to estimate the SED (e.g. ?) may help to improve the calibration of the bias.

In this work, we use the brightness moments to estimate the ellipticity of the galaxy. The PSF correction is not taken into account. The bias thus will appear in every method of measurement. However, the bias using the measurement method for real data will be different, since every method has its own property and weight function. The CG bias will have method-dependent properties as well, although they will in principle have same dependence on the colour gradient of the galaxy images. As the first step of the CG bias analysis, we did not adopt any specific method in order to obtain general properties of the CG bias. Before the real analysis of Euclid data, one needs to study the bias with specific methods and simulated images with real properties in the Euclid weak lensing survey.

ACKNOWLEDGMENTS

We would like to thank Emiliano Merlin, Marco Castellano for helping on SExtractor and Galfit, Gary Bernstein, Adam Rogers and also in general, the members of the Euclid Consortium for useful discussions. XE and VFC are funded by Italian Space Agency (ASI) through contract Euclid -IC (I/031/10/0) and acknowledge financial contribution from the agreement ASI/INAF/I/023/12/0. XE is also partly support by NSFC Grant No. 11473032. JR is supported by JPL, which is run by Caltech under a contract for NASA, and is supported by grant NASA ROSES 12-EUCLID12- 0004. The fast Fourier transforms are supplied by the FFTW library (?). We use CFITSIO (?) for the FITS file.

REFERENCES

- Bartelmann, M. & Schneider, P. 2001, Phys. Rep., 340, 291
 Cropper, M., Hoekstra, H., Kitching, T., et al. 2013, MNRAS, 431, 3103
 Cypriano, E. S., Amara, A., Voigt, L. M., et al. 2010, MNRAS, 405, 494
 Fenech Conti, I., Herbonnet, R., Hoekstra, H., et al. 2016, ArXiv e-prints
 Heymans, C., Grocott, E., Heavens, A., et al. 2013, MNRAS, 432, 2433
 Hildebrandt, H., Viola, M., Heymans, C., et al. 2017, MNRAS, 465, 1454
 Jarvis, M., Sheldon, E., Zuntz, J., et al. 2016, MNRAS, 460, 2245
 Jee, M. J., Tyson, J. A., Hilbert, S., et al. 2016, ApJ, 824, 77
 Laureijs, R., Amiaux, J., Arduini, S., et al. 2011, ArXiv e-prints
 Massey, R., Hoekstra, H., Kitching, T., et al. 2013, MNRAS, 429, 661
 Melchior, P., Viola, M., Schäfer, B. M., & Bartelmann, M. 2011, MNRAS, 412, 1552
 Miller, L., Heymans, C., Kitching, T. D., et al. 2013, MNRAS, 429, 2858
 Nakajima, R. & Bernstein, G. 2007, AJ, 133, 1763
 Paulin-Henriksson, S., Amara, A., Voigt, L., Refregier, A., & Bridle, S. L. 2008, A&A, 484, 67
 Rowe, B. T. P., Jarvis, M., Mandelbaum, R., et al. 2015, Astronomy and Computing, 10, 121
 Semboloni, E., Hoekstra, H., Huang, Z., et al. 2013, MNRAS, 432, 2385
 Voigt, L. M., Bridle, S. L., Amara, A., et al. 2012, MNRAS, 421, 1385