




التكنولوجيا التطبيقية
APPLIED TECHNOLOGY
وزارة التربية والتعليم والتعليم الفني



الرياضيات البحتة

الصف الثاني

التخصصات الصناعية (لغة انجليزية)

PURE MATHEMATICS

Second Grade

For industrial (English)

2nd.

2023 - 2024

Unit 1 : Differentiation and Its Applications–Integration

	Lesson	Title	Page
Second Term	one	Rules of differentiation	5
	two	Derivative of the product of two functions	9
	three	Derivative of the quotient of two functions	11
	four	Derivative of composite functions	14
	five	Derivative of trigonometric functions	17
	six	Derivative of trigonometric functions (follow)	20
	seven	Equation of tangent	23
	eight	Equation of normal	26
	nine	Indefinite integration	29
	ten	Indefinite integration (follow)	33
	eleven	Test	37
	twelve	Evaluation	38

Unit 1 : Differentiation and Its Applications

Dear student by the end of this unit you will recognize the following:

- The derivative of the constant function.
- The derivative of the function $f(x) = x^n$.
- The derivative of the function $f(x) = x$.
- The derivative of the function $f(x) = ax^n$.
- The derivative of the product of two functions.
- The derivative of the quotient of two functions.
- The derivative of the function $y = (f(x))^n$.
- The derivative of trig. functions
- To find the equation of the tangent, the equation of the normal to a curve at a point on it.
- Some rules of indefinite integration.

Lesson One

Rules of differentiation (derivative)

Let $y = f(x)$, the first derivative of this function is symbolized as :

$$y', \quad f'(x), \quad \frac{dy}{dx}, \quad \frac{d}{dx}(f(x))$$

$$1. \quad f(x) = a \in R \quad \text{where } a \text{ is constant (independent on } x) \Rightarrow f'(x) = 0$$

$$2. \quad f(x) = x^n, \quad n \in R \quad \Rightarrow \quad f'(x) = nx^{n-1}$$

$$3. \quad f(x) = x \quad \Rightarrow \quad f'(x) = 1$$

Important note:

First derivative of $f(x)$ is equal to rate of change.

$$\text{i.e.} \quad f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \text{slope of tangent} = \tan \theta$$

(where θ is inclination angle of the tangent with +ve X-axis)

Example (1)

Find the first derivative of of :

$$\bullet \quad y = 2x^7 \quad \Rightarrow \quad \frac{dy}{dx} = 14x^6$$

$$\bullet \quad y = \frac{3}{x^4} = 3x^{-4} \quad \Rightarrow \quad y' = -12x^{-5}$$

$$\bullet \quad f(x) = \sqrt[4]{x} = (x)^{\frac{1}{4}} \quad \Rightarrow \quad f'(x) = \frac{1}{4} 12x^{-\frac{3}{4}}$$

Example (2)

Find the slope of the tangent to the curve of the function f where $f(x) = x^2 - x + 1$ at the point $(-2, 7)$

Solution

$$f'(x) = 2x - 1$$

$$\text{at } x = -2$$

$$f'(-2) = -5 \quad \therefore \text{slope of the tangent} = -5$$

Example (3)

Prove that the tangent to the curve of the function $y = 3x^2 - 5x$ at the point $(1, -2)$ makes an angle of measure 45° with the positive direction of X-axis

Solution

$$\therefore y = 3x^2 - 5x \quad \therefore y' = 6x - 5$$

$$\text{at } x = 1 \quad \therefore y' = 1$$

$$\text{i.e. } \tan \theta = 1 \quad \therefore \theta = \tan^{-1} 1 = 45^\circ$$

Example (4)

Find the points on the curve $y = x^3 - 3x^2$ at which the slope of the tangent equals zero (the tangent // X-axis)

Solution

$$\frac{dy}{dx} = 3x^2 - 6x = 0 \quad \Rightarrow \quad 3x(x - 2) = 0$$

$$3x = 0$$

$$x = 0$$

$$\therefore y = (0)^3 - 3(0)^2$$

$$y = 0$$

$$\text{Points : } (0, 0)$$

or

$$x - 2 = 0$$

$$x = 2$$

$$y = (2)^3 - 3(2)^2$$

$$y = -4$$

$$\text{and } (2, -4)$$

Example (5) If $y = mx^2 - 3x$ and $\frac{dy}{dx} = 7$ at $x = 1$ find m .

Solution:

$$\frac{dy}{dx} = 2mx - 3 = 7$$

$$2mx = 10$$

$$2m(1) = 10 \Rightarrow m = 5$$

Example (6)

Find the slope of the tangent to the curve $y = x - \frac{2}{x}$ at $x = 1$

Solution

$$y = x - 2x^{-1}$$

$$\therefore \frac{dy}{dx} = 1 + 2x^{-2}$$

$$\text{at } x = 1 \Rightarrow \frac{dy}{dx} = 1 + 2 = 3$$

\therefore the slope of the tangent = 3

EXERCISE (1)

1. Find the points on the curve $y = x^3 - 4x + 2$ at which the slope of the tangent $= -1$
2. Find the points on the curve $y = 2x^3 - x + 3$ at which the slope of the tangent $= 5$
3. Find the first derivative of :
 - a) $y = x^5 - 3x^2 + 5x - 9$
 - b) $y = 2x^2 + 5$
4. Find the slope of the tangent to the curve $y = x^2 - \frac{1}{x}$ at $x = 1$
5. Find the measure of the angle which the tangent to the curve $y = 2x^3 + 3x$ makes with the positive direction of X-axis at the point $(-1, 0)$.
6. Find the points on the curve $y = x^3 - 4x + 1$ at which the tangent makes an angle of measure 135° with the positive direction of X-axis.
7. If the slope of the tangent to the curve $y = x^2 + ax$ at the point $(1, 3)$ is equal to 6 then, find the value of a .

Lesson Two

Derivative Of The Product Of Two Functions

If $y = f(x) \cdot g(x)$ then, $\frac{dy}{dx} = f(x) \cdot g'(x) + g(x) \cdot f'(x)$

i.e. $\frac{dy}{dx} = \text{first} \times (\text{second})' + \text{second} \times (\text{first})'$

Example (1)

Find the first derivative of the function $y = (3x^2 - 2x + 1)(x^4 - 5x^2 + 2)$ when $x = 0$

Solution:

$$\frac{dy}{dx} = (3x^2 - 2x + 1)(4x^3 - 10x) + (x^4 - 5x^2 + 2)(6x - 2)$$

when $x = 0$

$$\frac{dy}{dx} = (0 - 0 + 1)(0 - 0) + (0 - 0 + 2)(0 - 2)$$

$$\frac{dy}{dx} = -4$$

Example (2)

Find the slope of the tangent to the curve of the function
 $y = (2x + 5)(3x^2 - 1)$ at $x = -1$

Solution:

$$\frac{dy}{dx} = (2x + 5)(6x) + (3x^2 - 1)(2)$$

At $x = -1$

$$\frac{dy}{dx} = (-2 + 5)(-6) + (3 - 1)(2) = -14$$

\therefore Slope of the tangent = -14

EXERCISE (2)

1. Find the points on the curve $y = (x^2 + 5)(x - 1)$ at which the slope of the tangent is equal to -2 .
2. Find the first derivative of :
 - a) $y = (x^5 - 3x)(5x - 9)$
 - b) $y = (x^2 + 5)(x^2 - 3)$
3. Find the slope of the tangent to the curve $y = x^2(5x - 1)$ at $x = 1$
4. Find the measure of the angle which the tangent to the curve $y = (x^3 + 3x)(x - 5)$ makes with the positive direction of X-axis at the point $(-1, 0)$.

Lesson Three

Derivative Of The Quotient Of Two Functions

If $y = \frac{f(x)}{g(x)}$ where $g(x) \neq 0$

$$\text{then, } \frac{dy}{dx} = \frac{g(x) \cdot f'(x) - f(x) \cdot g'(x)}{[g(x)]^2}$$

$$\text{i.e. } \frac{dy}{dx} = \frac{\text{denominator} \cdot (\text{numerator})' - \text{numerator} \cdot (\text{denominator})'}{[\text{denominator}]^2}$$

Example (1)

Find the first derivative of the function $y = \frac{3x-2}{2x+3}$ when $x = 1$

Solution

$$\frac{dy}{dx} = \frac{(2x+3)(3) - (3x-2)(2)}{(2x+3)^2}$$

$$\text{when } x = 1 \Rightarrow \frac{dy}{dx} = \frac{(2+3)(3) - (3-2)(2)}{(2+3)^2} = \frac{13}{25}$$

Example (2)

Find the first derivative of the function $y = \frac{5x+11}{x+2}$ when $x = 0$

Solution

$$\frac{dy}{dx} = \frac{(x+2)(5) - (5x+11)(1)}{(x+2)^2}$$

$$\text{when } x = 0 \quad \frac{dy}{dx} = \frac{(0+2)(5) - (0+11)(1)}{(0+2)^2} = \frac{-1}{4}$$

Example (3)

If $f(x) = \frac{x^2}{3x+2}$ find $f'(-1)$

Solution:

$$f'(x) = \frac{(3x+2)(2x) - x^2(3)}{(3x+2)^2}$$

$$f'(-1) = \frac{(-3+2)(-2) - 1(3)}{(-3+2)^2} = -1$$

Example (4)

Find the measure of the angle which the tangent to the curve $y = \frac{x-1}{x^2-x+1}$ makes with the positive direction of X-axis at the point (1, 0).

Solution:

$$y' = \frac{(x^2 - x + 1)(1) - (x - 1)(2x - 1)}{(x^2 - x + 1)^2}$$

$$\text{At } x = 1 \quad y' = \frac{(1-1+1)(1) - (1-1)(2-1)}{(1-1+1)^2} = 1$$

$$\tan \theta = 1$$

$$\theta = 45^\circ$$

EXERCISE (3)

1. Find the first derivative of the function $y = \frac{2x+3}{x-2}$ when $x = 3$
2. Find the slope of the tangent to the curve $y = x - \frac{2}{x}$ at $x = 1$
3. Find the measure of the angle which the tangent to the curve $y = \frac{1+x}{1-x}$ makes with the positive direction of X-axis at the point $(-1, 0)$.
4. Find the points on the curve $y = \frac{x-2}{x-1}$ at which the slope of the tangent is equal to 1 .

Lesson Four

Derivative Of Composite Functions

Corollary:

If $y = (f(x))^n$, then:

$$\frac{dy}{dx} = n(f(x))^{n-1} \cdot f'(x)$$

Example (1)

If $y = (x^3 + 3x^2 - 1)^7$ then, find $\frac{dy}{dx}$ at $x = 0$

Solution:

$$\frac{dy}{dx} = 7(x^3 + 3x^2 - 1)^6 \cdot (3x^2 + 6x)$$

At $x = 0$

$$\frac{dy}{dx} = 7(0 + 0 - 1)^6 \cdot (0 + 0) = 0$$

Example (2)

Find the slope of the tangent to the curve of the function
 $y = x(2x - 1)^5 + 3$ at the point $(0, 3)$.

Solution:

$$\frac{dy}{dx} = (x)[5(2x - 1)^4 \cdot (2)] + 1(2x - 1)^5$$

at $x = 0$

$$\frac{dy}{dx} = (0)[5(0 - 1)^4 \cdot (2)] + 1(0 - 1)^5 = -1$$

\therefore slope of the tangent = -1

Example (3)

If $y = z^3$, $z = 2x^2 - 3x + 1$ find $\frac{dy}{dx}$ at $x = 2$

Solution

$$y = (2x^2 - 3x + 1)^3$$

$$\therefore \frac{dy}{dx} = 3(2x^2 - 3x + 1)^2 \cdot (4x - 3)$$

$$\text{At } x = 2$$

$$\frac{dy}{dx} = 3(8 - 6 + 1)^2 \cdot (8 - 3) = 135$$

Example (4)

If $y = 2z^3$, $z = \sqrt{x^2 - 2x}$ find $\frac{dy}{dx}$ at $x = 3$

Solution:

$$y = 2(\sqrt{x^2 - 2x})^3 = 2(x^2 - 2x)^{\frac{3}{2}}$$

$$\therefore \frac{dy}{dx} = 3(x^2 - 2x)^{\frac{1}{2}} \cdot (2x - 2)$$

$$\text{At } x = 3 \Rightarrow \frac{dy}{dx} = 3(9 - 6)^{\frac{1}{2}} \cdot (6 - 2) = 12\sqrt{3}$$

Example (5)

If $y = (3z - 1)^4$, $z = x(3x - 2)$ find $\frac{dy}{dx}$ at $x = 1$

Solution:

$$y = [3(3x^2 - 2x) - 1]^4 = [9x^2 - 6x - 1]^4$$

$$\therefore \frac{dy}{dx} = 4[9x^2 - 6x - 1]^3 \cdot (18x - 6)$$

$$\text{At } x = 1 \Rightarrow \frac{dy}{dx} = 4[9 - 6 - 1]^3 \cdot (18 - 6) = 384$$

EXERCISE (4)

1. Find $\frac{dy}{dx}$ if $y = (2x^2 - 3x - 5)^5$ at $x = 2$
2. If $f(x) = \frac{3x^2+1}{x^2+2}$, find $f'(x)$ at $x = 1$
3. If $y = z^2$, $z = x^2 + 1$, then find $\frac{dy}{dx}$ at $x = -1$
4. If $y = z^3$, $z = \frac{x}{x+1}$ then, find $\frac{dy}{dx}$ at $x = 1$
5. If $y = z + 2$, $z = x^2 + 2x$, then find $\frac{dy}{dx}$ at $x = 2$
6. If $y = z^3 + z$, $z = x^2 - x$ then, find $\frac{dy}{dx}$ at $x = 2$

Lesson Five

Derivative Of Trigonometric Functions

$$1. \text{ If } y = \sin x \quad \Rightarrow \quad \frac{dy}{dx} = \cos x$$

$$2. \text{ If } y = \cos x \quad \Rightarrow \quad \frac{dy}{dx} = -\sin x$$

$$3. \text{ If } y = \tan x \quad \Rightarrow \quad \frac{dy}{dx} = \sec^2 x$$

Corollaries:

$$1. \text{ If } y = \sin(ax + b) \quad \Rightarrow \quad \frac{dy}{dx} = a \cos(ax + b)$$

$$2. \text{ If } y = \cos(ax + b) \quad \Rightarrow \quad \frac{dy}{dx} = -a \sin(ax + b)$$

$$3. \text{ If } y = \tan(ax + b) \quad \Rightarrow \quad \frac{dy}{dx} = a \sec^2(ax + b)$$

Example (1)

Find the first derivative of the function $y = \sin 4x$

Solution:

$$\frac{dy}{dx} = 4 \cos 4x$$

Example (2)

Find the first derivative of the function $y = \cos (2x + 5)$

Solution:

$$\frac{dy}{dx} = -2 \sin(2x + 5)$$

Example (3)

Find the first derivative of the function $y = x \sin 2x$

Solution:

$$\begin{aligned} \frac{dy}{dx} &= x (2 \cos 2x) + \sin 2x \\ &= 2x \cos 2x + \sin 2x \end{aligned}$$

Example (4)

Find the first derivative of the function $y = \frac{\sin 2x}{\cos 3x}$ at $x = \pi$

Solution:

$$\frac{dy}{dx} = \frac{\cos 3x (2 \cos 2x) - \sin 2x (-3 \sin 3x)}{(\cos 3x)^2}$$

at $x = \pi$

$$\frac{dy}{dx} = \frac{\cos 3\pi (2 \cos 2\pi) - \sin 2\pi (-3 \sin 3\pi)}{(\cos 3\pi)^2} = \frac{(-1)(2) - 0(0)}{(-1)^2} = -2$$

Example (5)

Find the first derivative of the function $y = 2\cos x - \tan 5x$

Solution:

$$\frac{dy}{dx} = -2\sin x - 5\sec^2 5x$$

EXERCISE (5)

Find the first derivative of each of the following functions:

1. $y = \tan 2x$

2. $y = \sin 2x + \sin 3x$

3. $y = x \sin x - x^2$

4. $y = \sin 2x + \cos 3x$

5. $y = 2x + \sin 2x$ at $x = 0$

6. $y = \cos 2x + \sin 5x$

7. $y = \tan (3x + 2)$

8. $y = \cos x \cdot \sin x$ at $x = \pi$

9. $y = (3x - \cos 2x)^3$

10. $y = x^3 \cos 2x$

Lesson six

Derivative Of Trigonometric Functions (follow)

Example (1)

If $y = \sin\left(2x + \frac{\pi}{3}\right)$ find y' at $x = \pi$

Solution:

$$y' = 2\cos\left(2x + \frac{\pi}{3}\right) = 2\cos\left(2\pi + \frac{\pi}{3}\right) = 2 \times \frac{1}{2} = 1$$

Example (2) If $y = 2x \sin x \cos x$, prove that: $y' = \sin 2x + 2x \cos 2x$

Solution

$$y = x \sin 2x$$

$$y' = x(2\cos 2x) + \sin 2x = 2x \cos 2x + \sin 2x$$

Note:

$$\sin 2x = 2\sin x \cdot \cos x$$

Example (3)

If $y = \frac{\sin x}{\sin x + \cos x}$, prove that: $y' = \frac{1}{1 + \sin 2x}$

Solution

$$\begin{aligned} y' &= \frac{\cos x (\sin x + \cos x) - \sin x (\cos x - \sin x)}{(\sin x + \cos x)^2} \\ &= \frac{\cos x \sin x + \cos^2 x - \sin x \cos x + \sin^2 x}{\sin^2 x + \cos^2 x + 2\sin x \cos x} \\ &= \frac{1}{1 + \sin 2x} \end{aligned}$$

Example (4)

If $y = \sqrt{1 + \cos 2x}$, find $\frac{dy}{dx}$ at $x = \frac{\pi}{4}$

Solution:

$$y = (1 + \cos 2x)^{\frac{1}{2}}$$

$$\therefore y' = \frac{1}{2}(1 + \cos 2x)^{-\frac{1}{2}} (-2 \sin 2x)$$

$$\text{When } x = \frac{\pi}{4} \Rightarrow y' = -1$$

EXERCISE (6)

1. If $y = \cos x \cdot \sin x$, find $\frac{dy}{dx}$ at $x = \frac{\pi}{4}$
2. If $y = (\sin x + \cos x)^2$, prove that: $y' = 2\cos 2x$
3. Find $\frac{d}{dx}(2x^2 \sin x \cos x)$ at $x = \pi$
4. If $y = \sin x - \frac{1}{3}\sin^3 x$, prove that: $y' = \cos^3 x$
5. If $y = \frac{x}{\sin x}$, find $\frac{dy}{dx}$ at $x = \frac{\pi}{2}$
6. If $y = x \cos 2x$, find $\frac{dy}{dx}$ at $x = \frac{\pi}{2}$
7. Find $\frac{d}{dx}\left(\frac{x}{\cos 3x}\right)$ at $x = \frac{\pi}{3}$
8. Find the slope of the tangent to the curve $y = \sin 2x \cdot \cos x$ at $x = \pi$

Lesson seven

Equation of Tangent

Slope of the tangent to a curve:

Remember: Let m_1, m_2 be the slopes of two straight lines L_1, L_2

1. If $L_1 // L_2$ then, $m_1 = m_2$
2. If $L_1 \perp L_2$ then, $m_1 m_2 = -1$
3. Equation of the tangent to a curve at any point (x_1, y_1) is

$$(y - y_1) = m(x - x_1)$$

Example (1) Find the equation of the tangent to the curve

$y = 2x^3 - 4x^2 + 3$ at the point $(2, 3)$ which lies on the curve.

Solution:

$$y' = 6x^2 - 8x$$

$$\text{When } x = 2, \quad y' = 8$$

$$\text{Equation of the tangent : } y - 3 = 8(x - 2)$$

$$y - 8x + 13 = 0$$

Example (2) Find the equation of the tangent to the curve $y = \frac{x-1}{x^2-x+1}$
at the point $(1, 0)$ which lies on the curve.

Solution:

$$y' = \frac{1(x^2 - x + 1) - (x - 1)(2x - 1)}{(x^2 - x + 1)^2}$$

$$\text{When } x = 1,$$

$$y' = \frac{1(1 - 1 + 1) - (1 - 1)(2 - 1)}{(1 - 1 + 1)^2} \quad m = 1$$

$$\text{Equation of the tangent : } y - 0 = 1(x - 1)$$

$$y = x - 1$$

Example (3)

Find the equation of the tangent to the curve $y = (2x + 5)(3x^2 - 1)$ at $x = 0$.

Solution:

$$y' = (2x + 5)(6x) + (3x^2 - 1)(2)$$

at $x = 0$,

$\therefore y' = -2 =$ the slop of the tangent

and $y = (0 + 5)(0 - 1) = -5$

\Rightarrow point $(0, -5)$

Equation of the tangent : $y + 5 = -2(x - 0)$

$$y + 2x + 5 = 0$$

EXERCISE (7)

1. Find the equation of the tangent to the curve $y = x - \frac{2}{x}$ at $x = 1$
2. Find the equation of the tangent to the curve $y = \frac{x-2}{x-3}$ at $x = 2$
3. Find the equation of the tangent to the curve
 $y = (x^2 + 5)(x - 1)$ at $x = 2$
4. Find the equation of the tangent to the curve
 $y = x^2(5x - 1)$ at $x = 3$

Lesson eight

Equation Of Normal

Equation of the normal to a curve at any point (x_1, y_1) is :

$$y - y_1 = \frac{-1}{m}(x - x_1)$$

Example (1) Find the equation of the normal to the curve

$y = 2x^3 - 4x^2$ at the point $(2,0)$ which lies on the curve

Solution:

$$y' = 6x^2 - 8x$$

$$\text{When } x = 2, \quad y' = 8$$

$$\text{Equation of the normal : } y - 0 = \frac{-1}{8}(x - 2)$$

$$8y + x - 2 = 0$$

Example (2)

Find the equation of the normal to the curve $y = \frac{x-1}{x^2-x+1}$
at the point $(1,0)$ which lies on the curve.

Solution:

$$y' = \frac{1(x^2 - x + 1) - (x - 1)(2x - 1)}{(x^2 - x + 1)^2}$$

$$\text{When } x = 1,$$

$$y' = \frac{1(1 - 1 + 1) - (1 - 1)(2 - 1)}{(1 - 1 + 1)^2}$$

$$, m = 1$$

$$\text{Equation of the normal : } y - 0 = -1(x - 1)$$

$$y + x - 1 = 0$$

Example (3)

Find the equation of the normal to the curve $y = (2x + 5)(3x^2 - 1)$
at $x = 0$.

Solution:

$$y' = (2x + 5)(6x) + (3x^2 - 1)(2)$$

When $x = 0$,

$$\therefore y' = -2 = m \text{ (slope of tangent)}$$

$$\text{and } y = (0 + 5)(0 - 1) = -5$$

\Rightarrow point $(0, -5)$

$$\text{Equation of the normal : } y + 5 = \frac{1}{2}(x - 0)$$

$$2y - x + 10 = 0$$

EXERCISE (8)

1. Find the equation of the normal to the curve $y = x - \frac{2}{x}$ at $x = 2$
2. Find the equation of the normal to the curve $y = \frac{x-2}{x-3}$ at $x = 2$
3. Find the equation of the normal to the curve
 $y = (x^2 + 5)(x - 1)$ at $x = 1$
4. Find the equation of the normal to the curve
 $y = x^2(5x - 1)$ at $x = 1$

Lesson nine

Indefinite Integration

Rules:

$$1. \int x^n dx = \frac{x^{n+1}}{n+1} + c$$

$$2. \int a dx = ax + c$$

$$3. \int (ax + b)^n dx = \frac{(ax + b)^{n+1}}{a(n+1)} + c$$

$$4. \int [f(x)]^n \cdot [f'(x)] dx = \frac{[f(x)]^{n+1}}{n+1} + c$$

Example (1)

Find: (a) $\int 5x dx$ (b) $\int (3x^2 - x + 1) dx$

Solution:

$$\underline{a)} \int 5x dx = \frac{5}{2}x^2 + c$$

$$\underline{b)} \int (3x^2 - x + 1) dx = x^3 - \frac{1}{2}x^2 + x + c$$

Example (2)

Find: $\int (x - 2)(x - 1)(x + 1) \, dx$

Solution:

$$\begin{aligned}\int (x - 2)(x - 1)(x + 1) \, dx &= \int (x - 2)(x^2 - 1) \, dx \\ &= \int (x^3 - 2x^2 - x + 2) \, dx \\ &= \frac{x^4}{4} - \frac{2x^3}{3} - \frac{x^2}{2} + 2x + c\end{aligned}$$

Example (3) Find: $\int \frac{x^2 + m^2 x}{x} \, dx$, m is constant

Solution:

$$\int \frac{x^2 + m^2 x}{x} \, dx = \int (x + m^2) \, dx = \frac{1}{2}x^2 + m^2 x + c$$

Example (4) Find: $\int (x + \frac{1}{x})^2 \, dx$

Solution:

$$\begin{aligned}\int (x + \frac{1}{x})^2 \, dx &= \int (x^2 + 2 + \frac{1}{x^2}) \, dx \\ &= \int (x^2 + 2 + x^{-2}) \, dx = \frac{1}{3}x^3 + 2x - x^{-1} + c\end{aligned}$$

Example (5) Find: $\int (7x - 3)^8 \, dx$

Solution:

$$\begin{aligned}\int (7x - 3)^8 \, dx &= \frac{(7x-3)^9}{9 \times 7} + c \\ &= \frac{(7x-3)^9}{63} + c = \frac{1}{63} (7x - 3)^9 + c\end{aligned}$$

Example (6) Find : $\int (2x + 11)(\sqrt{2x + 11})dx$

Solution:

$$\begin{aligned} & \int (2x + 11)(2x + 11)^{\frac{1}{2}} dx \\ &= \int (2x + 11)^{\frac{3}{2}} dx = \frac{(2x+11)^{\frac{5}{2}}}{\frac{5}{2} \times 2} + c = \frac{1}{5} (2x + 11)^{\frac{5}{2}} + c \end{aligned}$$

Example (7) Find: $\int 2x (x^2 + 2)^5 dx$

Solution:

$$\int 2x (x^2 + 2)^5 dx = \frac{(x^2 + 2)^6}{6} + c$$

EXERCISE (9)

Find

1. $\int x(x + 3) dx$

2. $\int x^2(4x + 2) dx$

3. $\int (x - 4)(x + 4) dx$

4. $\int \frac{x^2 + 2x - 3}{x - 1} dx$

5. $\int \frac{x^3 + 27}{x + 3} dx$

6. $\int (2x - 4)^5 dx$

7. $\int x^2(4x^3 - 1)^7 dx$

Lesson ten

Indefinite Integration(follow)

Integrals of some trig. functions:

$$1. \int \sin x \, dx = -\cos x + c$$

$$2. \int \cos x \, dx = \sin x + c$$

$$3. \int \sec^2 x \, dx = \tan x + c$$

Corollaries:

$$1. \int \sin(ax + b) \, dx = -\frac{1}{a} \cos(ax + b) + c$$

$$2. \int \cos(ax + b) \, dx = \frac{1}{a} \sin(ax + b) + c$$

$$3. \int \sec^2(ax + b) \, dx = \frac{1}{a} \tan(ax + b) + c$$

Remember:

$$\sin^2 x + \cos^2 x = 1$$

$$1 + \tan^2 x = \sec^2 x$$

$$1 + \cot^2 x = \csc^2 x$$

$$\sin 2x = 2 \sin x \cdot \cos x$$

$$\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$$

$$\cos 2x = \cos^2 x - \sin^2 x$$

$$= 2 \cos^2 x - 1$$

$$= 1 - 2 \sin^2 x$$

$$\cos^2 x = \frac{1}{2} + \frac{1}{2} \cos 2x$$

$$\sin^2 x = \frac{1}{2} - \frac{1}{2} \cos 2x$$

Example (1): $\int (\cos x - \sin x) dx = \sin x + \cos x + c$

Example (2): $\int (\sec^2 x + \cos x) dx = \tan x + \sin x + c$

Example (3): $\int (1 + \sin x)^2 dx$

$$\begin{aligned} &= \int (1 + 2\sin x + \sin^2 x) dx \\ &= \int \left(1 + 2\sin x + \frac{1}{2} - \frac{1}{2}\cos 2x\right) dx = \int \left(\frac{3}{2} + 2\sin x - \frac{1}{2}\cos 2x\right) dx \\ &= \frac{3}{2}x - 2\cos x - \frac{1}{4}\sin 2x + c \end{aligned}$$

Example (4): $\int (\sin x + \cos x)^2 dx$

$$\begin{aligned} &= \int (\sin^2 x + \cos^2 x + 2\sin x \cos x) dx \\ &= \int (1 + \sin 2x) dx = x - \frac{1}{2}\cos 2x + c \end{aligned}$$

Example (5): $\int \frac{\sin^3 x + \cos^3 x}{\sin x + \cos x} dx$

$$= \int \frac{(\sin x + \cos x) (\sin^2 x - \sin x \cos x + \cos^2 x)}{\sin x + \cos x} dx$$

$$= \int (1 - \sin x \cos x) . dx$$

$$= \int (1 - \frac{1}{2} \sin 2x) dx = x + \frac{1}{4} \cos 2x + c$$

EXERCISE (10)

Find

1. $\int (7\sin x - 2\cos x) dx$

2. $\int (\sin 2x - 5\cos x) dx$

3. $\int (\sin x + \sec^2 x) dx$

4. $\int \sin (7x + 3) dx$

5. $\int \cos (5 - 2x) dx$

6. $\int 2\cos^2 x dx$

7. $\int (\cos^2 x - \sin^2 x) dx$

Unit TEST

FIRST QUESTION:

1. Find the measure of the angle which the tangent to the curve $y = 2x^3 + 3x$ makes with the positive direction of X-axis at the point $(-1, 0)$.
2. Find the slope of the tangent to the curve of the function $y = x^2(5x - 1)$ at $x = 1$.

SECOND QUESTION:

1. Find the first derivative of the function $y = \frac{2x+3}{x-2}$ when $x = 3$
2. If $y = z + 2$, $z = x^2 + 2x$ then, find $\frac{dy}{dx}$ at $x = 2$

THIRD QUESTION:

1. Find the first derivative of the function $y = x \sin x - x^2$
2. Find the equation of the tangent to the curve $y = 2x^3 - 4x^2 + 3$ at the point $(2,3)$ which lies on the curve.

FOURTH QUESTION:

Find

1. $\int x^2 (4x + 2) dx$
2. $\int \cos (5 - 2x) dx$

EVALUATION

Applied Technology School