

If the limit does not exist or has an infinite limit, you should point it out. In addition, do not use the L'Hôpital's rule to solve the limit problem.

1. (16%) Find the following limit

(a) $\lim_{x \rightarrow 1} \frac{x^3 - 3x + 2}{x^3 - x^2 - x + 1}$

(b) $\lim_{x \rightarrow \infty} x(\sqrt{x^2 + 1} - x)$

(c) $\lim_{\theta \rightarrow 0} \frac{1 - \cos(\theta)}{\theta^2}$

(d) $\lim_{x \rightarrow 3} \frac{\sqrt{3x+1}}{x-3}$

2. (8%) Considering the following function.

$$f(x) = \begin{cases} |x| \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

(a) Is $f(x)$ continuous at $x = 0$? Explain your answer.

(b) Is $f(x)$ differentiable at $x = 0$? Explain your answer.

3. (8%) Proof that there is only one intersect point between $f(x) = 2x - 2$ and $g(x) = \cos x$. (Hint: use the intermediate value theorem and mean value theorem/Rolle's theorem)

4. (15%) Remember that you can solve the derivative using the definition or the differentiation rule for the following question.

(a) Find the following limit. $\lim_{x \rightarrow 0} \frac{\cos(\pi+x) + 1}{x}$

(b) Find the derivative of $f(x) = \frac{x^3 + 3x - 1}{x + 1}$

(c) Let $f(x) = x \cos(x) - \tan(x) + 2\pi$, find $f''(x)$

5. (8%) Given $x^2 + \frac{y^2}{4} = 1$, find all the tangent lines of the graph that pass the point (3,0) (Note (3,0) is not on the graph).

6. (15%) Let $f(x) = \frac{x^3}{(x+2)^2}$

- (a) Find the critical numbers and the possible points of inflection of $f(x)$
- (b) Find the open intervals on which f is increasing or decreasing
- (c) Find the open intervals of concavity
- (d) Find all the asymptotes (Vertical/horizontal/Slant)
- (e) Sketch the graph of $f(x)$ (Label any intercepts, relative extrema, points of inflection, and asymptotes)

7. (16%) Remember the meaning and the definition of definite integral when solving the following question

(a) $\int \frac{2+t+t^3}{\sqrt{t}} dt$

(b) $\int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} (t^3 + t^6 \tan(t)) dt$

(c) $\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} \left(\frac{1}{\sqrt{n+1}} + \frac{1}{\sqrt{n+2}} + \dots + \frac{1}{\sqrt{n+n}} \right)$

8. (8%) Considering the function $f(x) = \cos(x) + 2\cos(2x) + \dots + n\cos(nx)$.

Proof that there exists at least one root between $(0, \pi)$ (Hint: Let $F(x) =$

$\int_0^x f(t) dt$ and use the fundamental theorem of calculus as well as Rolle's

theorem.)

9. (5%) Evaluate $\int_{\frac{1}{4}}^1 \frac{\sqrt{1-\sqrt{x}}}{\sqrt{x}} dx$