

## Stats PS4 Solution

1. (DGS 9.2.8)

- (a) The PDF's  $f_0(x)$  and  $f_1(x)$  are as sketched in Fig. ?? . Under  $H_0$  it is impossible to obtain a value of  $X$  greater than 1, but such values are possible under  $H_1$ . Therefore, if a test procedure rejects  $H_0$  only if  $x > 1$ , then it is impossible to make an error of type I, and  $\alpha(\delta) = 0$ . Also,  $\beta(\delta) = Pr(X < 1|H_1) = 1/2$ .
- (b) To have  $\alpha(\delta) = 0$ , we can include in the critical region only a set of points having probability 0 under  $H_0$ . Therefore, only points  $x > 1$  can be considered. To minimize  $\beta(\delta) = 0$  we should choose this set to have maximum probability under  $H_1$ . Therefore, all points  $x > 1$  should be used in the critical region.




Figure 1.1: Figure for ??.

2. (DGS 9.5.10)

$$\frac{S^2}{\sigma^2} \sim \chi_{n-1}^2$$

$$\text{Thus, } P(S_n^2 \leq c_1) = P\left(\chi^2 \leq \frac{c_1}{4}\right) = P(S_n^2 \geq c_2) = P\left(\chi^2 \geq \frac{c_2}{4}\right) = 0.025$$

$$\text{Therefore, } \frac{c_1}{4} = \chi_{0.975,9}^2 = 2.7 \text{ and } \frac{c_2}{4} = \chi_{0.025,9}^2 = 19.023$$

$$\Rightarrow c_1 = 10.8, c_2 = 76.095$$

3. (DGS 9.6.4)

$$\text{Let } Z = \bar{X} - \bar{Y} = \frac{\sigma_1^2}{m} + \frac{\sigma_2^2}{n} = \left(\frac{1}{m} + \frac{k}{n}\right) \sigma_1^2, \text{ thus,}$$

$$\frac{\bar{X}_m - \bar{Y}_n}{\sqrt{\left(\frac{1}{m} + \frac{k}{n}\right)\sigma_1^2}} = Z \sim N(0, 1)$$

$$\text{Also } \sum \frac{(X - \mu_1)^2}{\sigma_1^2} + \sum \frac{(Y - \mu_2)^2}{\sigma_2^2} = \left(S_X^2 + \frac{S_Y^2}{k}\right) \frac{1}{\sigma_1^2} \sim \chi_{m+n-2}^2.$$

As we know  $\frac{Z}{\sqrt{\chi^2/V}} \sim t$ , we have:

$$\frac{(\bar{X}_m - \bar{Y}_n) / \sqrt{\left[\left(\frac{1}{m} + \frac{k}{n}\right)\sigma_1^2\right]}}{\sqrt{\left(S_X^2 + \frac{S_Y^2}{k}\right) \frac{1}{\sigma_1^2} / (m+n-2)}} = \frac{(m+n-2)^{1/2}(\bar{X}_m - \bar{Y}_n)}{\left(\frac{1}{m} + \frac{k}{n}\right)^{1/2} \left(S_X^2 + \frac{S_Y^2}{k}\right)^{1/2}} \sim t_{m+n-2}$$

4. (DGS 9.6.6)

Similarly to the solution to the above Question 9.6.4, we can construct a test statistics:

$$\frac{(m+n-2)^{1/2}(\bar{X}_m - \bar{Y}_n - \lambda)}{\left(\frac{1}{m} + \frac{1}{n}\right)^{1/2} (S_X^2 + S_Y^2)^{1/2}} \sim t_{m+n-2}$$

5. DGS ch. 7.2 problem 10

¶10. In this exercise

$$f(x|\theta) = \begin{cases} 1 & \text{for } \theta - \frac{1}{2} < x < \theta + \frac{1}{2}, \\ 0 & \text{otherwise,} \end{cases}$$

and

$$\xi(\theta) = \begin{cases} \frac{1}{10} & \text{for } 10 < \theta < 20, \\ 0 & \text{otherwise.} \end{cases}$$

The condition that  $\theta - 1/2 < x < \theta + 1/2$  is the same as the condition that  $x - 1/2 < \theta < x + 1/2$ . Therefore,  $f(x | \theta)\xi(\theta)$  will be positive only for values of  $\theta$  which satisfy both the requirement that  $x - 1/2 < \theta < x + 1/2$  and the requirement that  $10 < \theta < 20$ . Since  $X = 12$  in this exercise,  $f(x | \theta)\xi(\theta)$  is positive only for  $11.5 < \theta < 12.5$ . Furthermore, since  $f(x | \theta)\xi(\theta)$  is constant over this interval, the posterior p.d.f.  $\xi(\theta | x)$  will also be constant over this interval. In other words, the posterior distribution of  $\theta$  must be a uniform distribution on this interval.

6. DGS ch. 7.4 problem 18

18. Suppose that the prior distribution of  $\theta$  is the Pareto distribution with parameters  $x_0$  and  $\alpha$  ( $x_0 > 0$  and  $\alpha > 0$ ). Then the prior p.d.f.  $\xi(\theta)$  has the form

$$\xi(\theta) \propto 1/\theta^{\alpha+1} \quad \text{for } \theta \geq x_0.$$

If  $X_1, \dots, X_n$  form a random sample from a uniform distribution on the interval  $[0, \theta]$ , then

$$f_n(\mathbf{x} \mid \theta) \propto 1/\theta^n \quad \text{for } \theta > \max\{x_1, \dots, x_n\}.$$

Hence, the posterior p.d.f. of  $\theta$  has the form

$$\xi(\theta \mid \mathbf{x}) \propto \xi(\theta) f_n(\mathbf{x} \mid \theta) \propto 1/\theta^{\alpha+n+1},$$

for  $\theta > \max\{x_0, x_1, \dots, x_n\}$ , and  $\xi(\theta \mid \mathbf{x}) = 0$  for  $\theta \leq \max\{x_0, x_1, \dots, x_n\}$ . This posterior p.d.f. can now be recognized as also being the Pareto distribution with parameters  $\alpha + n$  and  $\max\{x_0, x_1, \dots, x_n\}$ .

7. DGS ch. 7.4 problem 6

6. Suppose that the parameters of the prior gamma distribution of  $\theta$  are  $\alpha$  and  $\beta$ . Then  $\mu_0 = \alpha/\beta$ . The posterior distribution of  $\theta$  was given in Theorem 7.3.2. The mean of this posterior distribution is

$$\frac{\alpha + \sum_{i=1}^n X_i}{\beta + n} = \frac{\beta}{\beta + n} \mu_0 + \frac{n}{\beta + n} \bar{X}_n.$$

Hence,  $\gamma_n = n/(\beta + n)$  and  $\gamma_n \rightarrow 1$  as  $n \rightarrow \infty$ .

$$\gamma_n \sim \frac{n}{n}$$

8. DGS ch. 7.4 problem 12

12. (a)  $A$ 's prior distribution for  $\theta$  is the beta distribution with parameters  $\alpha = 2$  and  $\beta = 1$ . Therefore,  $A$ 's posterior distribution for  $\theta$  is the beta distribution with parameters  $2+710 = 712$  and  $1+290 = 291$ .  $B$ 's prior distribution for  $\theta$  is a beta distribution with parameters  $\alpha = 4$  and  $\beta = 1$ . Therefore,  $B$ 's posterior distribution for  $\theta$  is the beta distribution with parameters  $4 + 710 = 714$  and  $1 + 290 = 291$ .

(b)  $A$ 's Bayes estimate of  $\theta$  is  $712/(712+291) = 712/1003$ .  $B$ 's Bayes estimate of  $\theta$  is  $714/(714+291) = 714/1005$ .

(c) If  $y$  denotes the number in the sample who were in favor of the proposition, then  $A$ 's posterior distribution for  $\theta$  will be the beta distribution with parameters  $2 + y$  and  $1 + 1000 - y = 1001 - y$ , and  $B$ 's posterior distribution will be a beta distribution with parameters  $4 + y$  and  $1 + 1000 - y = 1001 - y$ . Therefore,  $A$ 's Bayes estimate of  $\theta$  will be  $(2 + y)/1003$  and  $B$ 's Bayes estimate of  $\theta$  will be  $(4 + y)/1005$ . But

$$\left| \frac{4 + y}{1005} - \frac{2 + y}{1003} \right| = \frac{2(1001 - y)}{(1005)(1003)}.$$

This difference is a maximum when  $y = 0$ , but even then its value is only

$$\frac{2(1001)}{(1005)(1003)} < \frac{2}{1000}.$$