



EMBO
Practical Course


Plant microbiota

26 March – 07 April 2017 | Cologne, Germany


Alpha- and beta-diversity
Ruben Garrido-Oter

A

Beta-diversity
(unweighted UniFrac distance)

**B**

Alpha-diversity
(phylogenetic diversity)



Host description

- Human
- Wild animal
- Hydra
- Fish
- Arabidopsis and relatives
- Barley
- Maize
- Grapevine
- Rice

Indices of alpha-diversity (within sample diversity)

Chao index (Chao, 1984) is based on the number of low abundance OTUs (rare biota) found in a sample.

The **Shannon** index is based on the concept of entropy. It takes into account simultaneously how many species and how evenly distributed they are in a community.

Observed species consists on simply counting the number of detected OTUs in a sample.

Faith's Phylogenetic Diversity is related to the phylogenetic diversity of a community, assuming that communities of consisting of highly unrelated species are more complex.

Indices of beta-diversity (between samples diversity)

beta-diversity metrics provide a quantitative evaluation of how much samples differ from one another. They can be classified in multiple ways:

	quantitative (take into account relative abundances)	qualitative (considers only OTU presence / absence)
phylogenetic (considers OTU relatedness)	weighted UniFrac	unweighted UniFrac
non phylogenetic (assumes all species are equally related)	Bray-Curtis	Jaccard

Indices of beta-diversity (between samples diversity) (cont.)

