

Literatur

- [1] H. A. Abro und M. M. Shaikh, „A new time-efficient and convergent nonlinear solver“, *Applied Mathematics and Computation*, Jg. 355, S. 516–536, 2019, doi: 10.1016/j.amc.2019.03.012.
- [2] K. Amini und F. Rostami, „A modified two steps Levenberg–Marquardt method for nonlinear equations“, *Journal of Computational and Applied Mathematics*, Jg. 288, Nr. 288, S. 341–350, 2015, doi: 10.1016/j.cam.2015.04.040.
- [3] K. Amini, F. Rostami und G. Caristi, „An efficient Levenberg–Marquardt method with a new LM parameter for systems of nonlinear equations“, *Optimization*, Jg. 67, Nr. 5, S. 637–650, 2018, doi: 10.1080/02331934.2018.1435655.
- [4] J. Fan, „The modified Levenberg-Marquardt method for nonlinear equations with cubic convergence“, *Math. Comp.*, Jg. 81, Nr. 277, S. 447–466, 2012, doi: 10.1090/S0025-5718-2011-02496-8.
- [5] J. Fan, „Accelerating the modified Levenberg-Marquardt method for nonlinear equations“, *Math. Comp.*, Jg. 83, Nr. 287, S. 1173–1187, 2014, doi: 10.1090/S0025-5718-2013-02752-4.
- [6] J. Fan und J. Pan, „An improved trust region algorithm for nonlinear equations“, *Comput Optim Appl*, Jg. 48, Nr. 1, S. 59–70, 2011, doi: 10.1007/s10589-009-9236-7.
- [7] J. M. Martínez, „Practical quasi-Newton methods for solving nonlinear systems“, *Journal of Computational and Applied Mathematics*, Jg. 124, 1-2, S. 97–121, 2000, doi: 10.1016/S0377-0427(00)00434-9.
- [8] J. Nocedal und S. J. Wright, *Numerical optimization*, 2. Aufl. New York: Springer, 2006.
- [9] F. Rahpeymaii, „An efficient conjugate gradient trust-region approach for systems of nonlinear equation“, *Afr. Mat.*, Jg. 30, 3-4, S. 597–609, 2019, doi: 10.1007/s13370-019-00669-0.
- [10] Roger Behling, „Local convergence analysis of the Levenberg-Marquardt framework for nonzero-residue nonlinear least-squares problems under an error bound condition“, 2018.
- [11] X. Wang und Y. Li, „An Efficient Sixth-Order Newton-Type Method for Solving Nonlinear Systems“, *Algorithms*, Jg. 10, Nr. 2, S. 45, 2017, doi: 10.3390/a10020045.
- [12] M. Waseem, M. A. Noor und K. I. Noor, „Efficient method for solving a system of nonlinear equations“, *Applied Mathematics and Computation*, Jg. 275, S. 134–146, 2016, doi: 10.1016/j.amc.2015.11.061.
- [13] G. Yuan, Z. Wei und X. Lu, „A BFGS trust-region method for nonlinear equations“, *Computing*, Jg. 92, Nr. 4, S. 317–333, 2011, doi: 10.1007/s00607-011-0146-z.
- [14] M. Zeng und H. Fu, „A Broyden Trust Region Quasi-Newton Method for Nonlinear Equations“, *IAENG International Journal of Computer Science*, 46:3, 2019.
- [15] L. Zheng, L. Chen und Y. Ma, „A variant of the Levenberg-Marquardt method with adaptive parameters for systems of nonlinear equations“, *MATH*, Jg. 7, Nr. 1, S. 1241–1256, 2021, doi: 10.3934/math.2022073.