

# Linear Regression


# Acknowledgement

- Most of these slides were either created by Prof. Andrew Ng or else are modifications of his slides

# Model Representation

## Housing Prices (Portland, OR)

Price  
(in 1000s  
of dollars)



# Model Representation

Training set of housing prices (Portland, OR)	Size in feet <sup>2</sup> (x)	Price (\$) in 1000's (y)
	2104	460
	1416	232
	1534	315
	852	178
	...	...


Notation:

**m** = Number of training examples

**x**'s = "input" variable / features

**y**'s = "output" variable / "target" variable

# Model Representation



# Cost function

Training Set	Size in feet <sup>2</sup> (x)	Price (\$) in 1000's (y)
	2104	460
	1416	232
	1534	315
	852	178
	...	...


Hypothesis:  $h_{\theta}(x) = \theta_0 + \theta_1 x$

$\theta_i$ 's: Parameters


How to choose  $\theta_i$ 's ?

# Cost function


$$h_{\theta}(x) = \theta_0 + \theta_1 x$$



$$\theta_0 = 1.5$$
$$\theta_1 = 0$$




$$\theta_0 = 0$$
$$\theta_1 = 0.5$$



$$\theta_0 = 1$$
$$\theta_1 = 0.5$$

# Cost function



Idea: Choose  $\theta_0, \theta_1$  so that  $h_{\theta}(x)$  is close to  $y$  for our training examples  $(x, y)$

Hypothesis:

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

Parameters:

$$\theta_0, \theta_1$$



Cost Function:

$$J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$



Goal: minimize  $J(\theta_0, \theta_1)$   
 $\theta_0, \theta_1$





# Cost function - intuition





# Cost function - intuition





# Cost function - intuition





# Cost function - intuition





# Cost function - intuition



# Cost function - intuition



# Cost function - intuition






# Cost function - intuition


$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )



$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )





# Gradient Descent


Have some function  $J(\theta_0, \theta_1)$

Want  $\min_{\theta_0, \theta_1} J(\theta_0, \theta_1)$

## Outline:

- Start with some  $\theta_0, \theta_1$
- Keep changing  $\theta_0, \theta_1$  to reduce  $J(\theta_0, \theta_1)$   
until we hopefully end up at a minimum

# Gradient Descent



# Gradient Descent

## Gradient descent algorithm



repeat until convergence {  
     $\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$     (for  $j = 0$  and  $j = 1$ )  
}

---

Correct: Simultaneous update

$\text{temp0} := \theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$   
 $\text{temp1} := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$   
 $\theta_0 := \text{temp0}$   
 $\theta_1 := \text{temp1}$

# Gradient Descent Intuition




# Gradient Descent Intuition

$$\theta_1 := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_1)$$

If  $\alpha$  is too small, gradient descent can be slow.

If  $\alpha$  is too large, gradient descent can overshoot the minimum. It may fail to converge, or even diverge.



# Gradient Descent for Linear Regression

Gradient descent algorithm

repeat until convergence {  
     $\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$   
    (for  $j = 1$  and  $j = 0$ )  
}

Linear Regression Model

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

$$J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

# Gradient Descent for Linear Regression

$$\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) =$$

$$j = 0 : \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1) =$$

$$j = 1 : \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1) =$$

# Gradient Descent for Linear Regression

## Gradient descent algorithm

repeat until convergence {



$$\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$$

$$\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x^{(i)}$$

}




# Gradient Descent for Linear Regression



# Gradient Descent for Linear Regression


$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )





$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )



# Gradient Descent for Linear Regression



# Multivariate Linear Regression (MLR)

# Multiple Features (variables)

Size (feet <sup>2</sup> )	Price (\$1000)
$x$	$y$
2104	460
1416	232
1534	315
852	178
...	...

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

Size (feet <sup>2</sup> )	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
2104	5	1	45	460
1416	3	2	40	232
1534	3	2	30	315
852	2	1	36	178
...	...	...	...	...

# Multiple Features (variables)

Size (feet <sup>2</sup> )	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
2104	5	1	45	460
1416	3	2	40	232
1534	3	2	30	315
852	2	1	36	178
...	...	...	...	...

Notation:

$n$  = number of features

$x^{(i)}$  = input (features) of  $i^{th}$  training example.

$x_j^{(i)}$  = value of feature  $j$  in  $i^{th}$  training example.

# Hypothesis

$$h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3 + \cdots + \theta_n x_n$$

$$h_{\theta}(x) = \begin{bmatrix} \theta_0 & \theta_1 & \cdots & \theta_n \end{bmatrix} \begin{bmatrix} x_0 \\ x_1 \\ \vdots \\ x_n \end{bmatrix} = \theta^T x$$

# Gradient Descent for Multiple Variables

Hypothesis:  $h_{\theta}(x) = \theta^T x = \theta_0 x_0 + \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_n x_n$

Parameters:  $\theta_0, \theta_1, \dots, \theta_n$

Cost function:

$$J(\theta_0, \theta_1, \dots, \theta_n) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

repeat until convergence: {

$$\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x_j^{(i)} \quad \text{for } j := 0 \dots n$$

}



## Gradient Descent

Previously (n=1):

Repeat {

$$\theta_0 := \theta_0 - \underbrace{\alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})}_{\frac{\partial}{\partial \theta_0} J(\theta)}$$

$$\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x^{(i)}$$

(simultaneously update  $\theta_0, \theta_1$ )

}

New algorithm ( $n \geq 1$ ):

Repeat {

$$\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

(simultaneously update  $\theta_j$  for  
 $j = 0, \dots, n$ )

}

---

$$\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)}$$

$$\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_1^{(i)}$$

$$\theta_2 := \theta_2 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_2^{(i)}$$


...

# Trick 1: Feature Scaling

Idea: Make sure features are on a similar scale.


E.g.  $x_1$  = size (0-2000 feet<sup>2</sup>)

$x_2$  = number of bedrooms (1-5)



$$x_1 = \frac{\text{size (feet}^2\text{)}}{2000}$$

$$x_2 = \frac{\text{number of bedrooms}}{5}$$



# Mean normalization

$$x_i := \frac{x_i - \mu_i}{s_i}$$


Replace  $x_i$  with  $x_i - \mu_i$  to make features have approximately zero mean  
(Do not apply to  $x_0 = 1$ ).

E.g.  $x_1 = \frac{size - 1000}{2000}$

$$x_2 = \frac{\#bedrooms - 2}{5}$$

$$-0.5 \leq x_1 \leq 0.5, -0.5 \leq x_2 \leq 0.5$$

## Trick 2: learning rate



# Features and Polynomial Regression

## Housing prices prediction

$$h_{\theta}(x) = \theta_0 + \theta_1 \times \textit{frontage} + \theta_2 \times \textit{depth}$$



# Features and Polynomial Regression

