

Mean-Force Hamiltonians from Influence Functionals

Gerard McCaul

(Dated: April 27, 2026)

The Hamiltonian of mean force (HMF) provides the standard starting point for strong-coupling thermodynamics, yet explicit operator forms are known only in restricted settings. We present a “quenched density” framework that uses the Hubbard-Stratonovich transformation to rewrite the reduced equilibrium state as an average over local propagators in imaginary time. This approach rigorously separates the statistical definition of the environment from the algebraic structure of the system response. We apply this framework to the minimal case of a harmonic environment with a coupling commuting with the system Hamiltonian. In this scenario the correction to the HMF has an exact, closed-form expression. We validate this result against finite-bath trace-out calculations and stochastic imaginary-time sampling in a five-level projector-coupled model.

I. INTRODUCTION

What governs equilibrium? The canonical answer is the thermal Gibbs state, where the environment is characterised by a single parameter: the temperature. Yet, this canonical answer is puzzling. It implies that neither the specific character of the environment nor the form of its coupling to the system alters the system’s equilibrium. This of course is a consequence of the assumptions used to derive the answer - the weak-coupling or Markovian regimes open-systems theory has historically relied upon [1]. At strong coupling however, the reduced state inherits explicit temperature dependence and interaction-induced operator content that definitionally cannot be captured by these assumptions [2, 3]. These issues have motivated a broad investigation into strong-coupling thermodynamics, including exact fluctuation relations [4–6], consistent heat and work definitions [7–9], and operational measurability [10–12]. This representability question sits alongside operational ambiguities in thermodynamic bookkeeping beyond weak coupling, where heat/work/state assignments can depend on the chosen framework [13].

While formally one can write the reduced state in canonical exponential form, the effective Hamiltonian governing this state is generally not the bare Hamiltonian of the system. Instead, it must account for the non-trivial influence of the environment in a manner that goes beyond the specification of temperature. If an effective operator capturing this influence can be constructed, the standard description of equilibrium holds generically across all coupling strengths. *The Hamiltonian of mean force* (HMF) is the operator that reproduces this reduced equilibrium object as a Gibbs-like state, and provides the standard starting point for strong-coupling thermodynamics [4, 5, 14–16]. At finite coupling, the HMF can become temperature dependent and encode interaction-induced terms that are absent in H_Q , including effective many-body or nonlocal operator content [2, 3, 7, 15–17]. This is the conceptual tension: the reduced equilibrium state is well defined, but its generator is not generally simple. In the strong-

coupling regime one usually characterises equilibrium through the mean-force Gibbs state, whose structure and limiting forms (including ultrastrong-coupling projector dominance) have been analysed from a ‘state-first’ perspective [18–21].

The target of thermodynamic descriptions is therefore unambiguous, and in fact a formal definition of HMF is equally unambiguous - it is simply the logarithm of the total state after tracing out its environment. It is the *representation* of this operator that remains a challenge. Is the HMF a local operator? Does it admit a compact description in terms of few-body interactions? To address this, a veritable zoo of approximations - polaron transforms [22], weak-coupling expansions [18], reaction-coordinate mappings [23–27] - have proliferated strong-coupling thermodynamics. These approximations often disagree in intermediate regimes however, and while powerful numerical techniques have been derived from path-integrals [28–33], they typically do not yield compact operator forms for the HMF.

Ultimately, the source of these difficulties lie in the trace operation, which mixes the bath and system degrees of freedom in a way that is difficult to invert. For this reason, a general framework for representing the HMF has proved elusive. The purpose of this paper is to lay the groundwork for just such a framework. To do so, we construct a representation of the equilibrium density matrix that disentangles the statistical properties of the environment from the algebraic response of the system. We term this object the *quenched density*, and to obtain it we employ the *influence functional* approach [34–36]. This formalism provides a natural language for the problem. For a recent unifying overview that links hierarchical, chain-mapping, pseudomode, stochastic, and extended-state-space approaches to non-Markovian open-system dynamics, see Ref. [37]. By employing the influence functional in concert with the Hubbard-Stratonovich decoupling [38–40], we are able to rewrite the reduced equilibrium density not as a partial trace, but as a classical average over an ensemble of “quenched” imaginary-time propagators.

$$\bar{\rho}_S(\beta) = \mathbb{E}_\xi \left[\hat{\mathcal{T}} e^{-\int_0^\beta d\tau H_{\text{eff}}[\xi(\tau)]} \right]. \quad (1)$$

In this picture, the bath is replaced by a stochastic field $\xi(\tau)$ [41–46], and the system evolves under a time-dependent effective Hamiltonian for each realisation of the noise.

This formulation is obtained in two steps. The path integral sets the system in a frame where it can be expressed in terms of classical trajectories for the bath degrees of freedom, which can be integrated out. The resulting action is then a non-local kernel in imaginary time. The Hubbard-Stratonovich transformation remedies this by representing the bath information locally. This then allows the path integral to be reversed and the reduced density to be formulated at the operator level. The framework thus provides a rigorous bridge between the nonlocal path-integral description (where the bath is exact but the system operator structure is opaque) and the local operator description required for thermodynamics.

We apply this framework to establishing a rigorous benchmark: the linearly coupled Gaussian bath in the commuting (pure-dephasing) sector [1, 47]. In this limit, relevant for quantum nondemolition measurements and qubit dephasing [48, 49], the technical obstruction of imaginary time ordering vanishes, and the quenched density’s average can be performed analytically. We use this to recover the known result that the bath induces a static potential renormalisation $-\lambda f^2$ (the reorganisation energy) [50]. While this specific result is standard, its derivation within the quenched density framework serves two critical purposes. First, it demonstrates how the framework naturally recovers thermodynamic identities from path-integral primitives. Second, and most importantly, it provides a foundation for the generalisation of the framework to both non-commuting couplings and anharmonic environments.

The paper is structured as follows. In Sec. II we define the total Hamiltonian and reduced equilibrium objects. Sec. III develops the core of the framework: the Quenched Density representation of the imaginary-time propagator. Sec. IV connects this to the exact commuting Gaussian benchmark. Sec. V presents the numerical validation, confirming that the quenched density sampling, analytic prediction, and full finite-bath trace-out agree to machine precision. Sec. VI closes the paper with a summary of results, and highlights the essential contribution of this framework as a foundation for the future extension of the theory to non-Gaussian and non-commuting environments.

II. THE TOTAL HAMILTONIAN AND THE HAMILTONIAN OF MEAN FORCE

The Hamiltonian of mean force is defined by tracing the global equilibrium operator, and therefore presupposes a concrete model for the composite Hamiltonian.

We consider a system with Hamiltonian H_Q (system coordinates q), coupled to a bath with Hamiltonian H_X (bath coordinates x) through an interaction H_I , so that

$$H_{\text{tot}} = H_Q + H_X + H_I. \quad (2)$$

To obtain an explicit influence-functional representation we specialise to a Gaussian environment with linear coupling, i.e. a Caldeira-Leggett bath [35]. This choice is appropriate when the bath is large and near equilibrium, so that its fluctuations are well captured by a quadratic Hamiltonian and the leading (system-local) term in the interaction dominates.

The bath is modelled as a collection of harmonic oscillators,

$$H_X = \sum_k \left(\frac{p_k^2}{2m_k} + \frac{1}{2} m_k \omega_k^2 x_k^2 \right), \quad (3)$$

and we take a linear coupling to a system operator $f \equiv f(q)$,

$$H_I = \sum_k c_k f(q) x_k, \quad (4)$$

so that

$$H_{\text{tot}} = H_Q + \sum_k \left(\frac{p_k^2}{2m_k} + \frac{1}{2} m_k \omega_k^2 x_k^2 + c_k f(q) x_k \right). \quad (5)$$

As is standard in Caldeira-Leggett constructions, one may include a counter-term proportional to $f(q)^2$ to enforce translational invariance of the bath coordinates and to compensate the static potential renormalisation induced by the coupling [35]. In what follows we absorb any such counterterm into the definition of H_Q .

Although the microscopic bath is parametrised by $\{m_k, \omega_k, c_k\}$, its influence on the system enters only through the spectral density [1, 51],

$$J(\omega) = \frac{\pi}{2} \sum_k \frac{c_k^2}{m_k \omega_k} \delta(\omega - \omega_k), \quad (6)$$

which compactly encodes the bath two-point structure and fixes the imaginary-time influence kernel used below [34].

Having specified the composite dynamics, we define the Hamiltonian of mean force by tracing the global Hamiltonian of mean force $H_{\text{MF}}(\beta)$ as the operator whose Gibbs form reproduces the reduced equilibrium object up to a chosen normalisation. If the total system is at inverse temperature β , the unnormalised global equilibrium operator is $e^{-\beta H_{\text{tot}}}$, then the reduced (unnormalised) equilibrium operator is

$$\bar{\rho}_Q(\beta) \equiv \text{Tr}_X e^{-\beta H_{\text{tot}}}, \quad (7)$$

where Tr_X traces out the bath degrees of freedom. The Hamiltonian of mean force is defined by

$$e^{-\beta H_{\text{MF}}(\beta)} \equiv \frac{\text{Tr}_X e^{-\beta H_{\text{tot}}}}{Z_X(\beta)}, \quad Z_X(\beta) \equiv \text{Tr}_X e^{-\beta H_X}, \quad (8)$$

or equivalently

$$H_{\text{MF}}(\beta) = -(1/\beta) \log[\text{Tr}_X e^{-\beta H_{\text{tot}}}] \quad (9)$$

up to an additive scalar fixed by the choice of Z_X .

For a harmonic bath linearly coupled through a collective coordinate, the bath can be eliminated exactly, yielding the Feynman-Vernon influence functional [2, 3, 34, 36]. In equilibrium this produces a Euclidean (imaginary-time) influence functional which is bilocal in τ and whose kernel is fixed by the bath spectral density $J(\omega)$ [28, 42, 52, 53]. While this gives an exact representation of the reduced equilibrium operator, it does not generically yield a compact local expression for H_{MF} . The reduced equilibrium object is a path integral over the system degrees of freedom, rather than a simple exponential $e^{-\beta H_{\text{eff}}}$ of a time-independent operator. Extracting H_{MF} - the logarithm of this ordered object - is therefore an inverse representability problem [54–57], not merely a question of coupling strength or temperature [15, 58].

There are two logically distinct aspects to this problem. First, the bath produces an *imaginary-time nonlocal* self-interaction with a memory kernel determined by $J(\omega)$. This can be resolved with a local representation of the path integral, which can be achieved via a Hubbard-Stratonovich (HS) transformation [38, 39]. This mapping allows for a return to the operator picture by replacing the nonlocal self-interaction with a local coupling to a stochastic auxiliary field. However, this transformation introduces stochasticity that must be averaged over, and it is in this averaging process that we encounter the second obstruction: the algebraic properties of the system and its coupling. While the HS transformation allows one to return to the operator picture, to obtain the exact state now requires an averaging procedure over the stochastic auxiliary field. When $[H_Q, f] \neq 0$, evaluating this average analytically presents an (apparently) [59] insuperable difficulty. Nevertheless, in the commuting sector $[H_Q, f] = 0$, the algebraic obstruction disappears, and the averaging can be performed. Under these conditions, the influence functional contains exactly the ingredients required to construct $H_{\text{MF}}(\beta)$ in closed form.

Summarising, our strategy to find $H_{\text{MF}}(\beta)$ is to rewrite the reduced state such that the contributions from the bath statistics (entering only through $J(\omega)$) and the system-coupling algebra may be cleanly separated. To achieve this, it is necessary to first develop a formal treatment connecting the quenched density representation of the reduced system to a path integral representation of the influence functional.

III. THE QUENCHED DENSITY REPRESENTATION

While many representations of reduced equilibrium exist [1, 36, 51], tracing out environmental degrees of freedom is most naturally achieved in a path integral formulation. Indeed, one of the principal advantages of this approach is that it requires no assumptions on the spectral character of the bath or the strength of its coupling to the system [51]. In this framework, tracing out the bath degrees of freedom leads to the following representation of the reduced density matrix $\bar{\rho}_Q(\beta)$ [60, 61]:

$$\langle q_f | \bar{\rho}_Q(\beta) | q_i \rangle = \frac{1}{Z_{\text{tot}}} \int \mathcal{D}q \mathcal{D}\mathbf{x} e^{-S_E[q, \mathbf{x}]}. \quad (10)$$

This is an imaginary-time path integral [60, 61], with a Euclidean action $S_E[q, \mathbf{x}]$. This contains both the bare system action, and the *influence functional* $\Gamma_E[q, \mathbf{x}]$ that remains after tracing the bath degrees of freedom. While this representation makes exact treatment of the bath possible, it comes at a conceptual cost: the path integral trades operator structure for functional structure. In particular, questions about the *operator algebra* underlying the reduced equilibrium state are obscured once the dynamics is expressed solely in terms of paths and kernels.

If one wishes to understand when a local or otherwise restricted Hamiltonian of mean force can exist, it is therefore useful to reconnect the path-integral description to an operator-based notion of imaginary-time propagation. The key observation is that the reduced equilibrium operator may be regarded as an imaginary-time evolution over an interval of length β , but with a generator that need not be time independent. Allowing for explicit τ -dependence in the generator provides a natural bridge between the nonlocal Euclidean influence functional and a time-ordered exponential acting on the system Hilbert space.

Motivated by this, we introduce a class of imaginary-time propagators generated by a τ -dependent Hamiltonian $\bar{H}(\tau)$ acting on the system Hilbert space. With this, we define the *quenched* imaginary-time propagator $U(\beta)$ and the *quenched density* $\rho_{\bar{H}}(\beta)$ as its normalised counterpart:

$$\begin{aligned} U(\beta) &:= \hat{\tau} \exp \left[- \int_0^\beta d\tau \bar{H}(\tau) \right], \\ \rho_{\bar{H}}(\beta) &:= \frac{1}{Z_{\bar{H}}} U(\beta), \\ Z_{\bar{H}} &:= \text{Tr} U(\beta), \end{aligned} \quad (11)$$

where $\hat{\tau}$ orders increasing imaginary times (inverse temperatures) to the left [60, 61]. We refer to this construction as *quenched* as the τ -dependence is treated as fixed

during the construction of the propagator - we may consider it as a quasistatic quench, where the Hamiltonian is modulated as temperature is lowered.

Having defined this object, we must make contact with the path integral representation. To this end, we assume the standard kinematic structure

$$\bar{H}(\tau) = \bar{T} + \bar{V}(\tau), \quad \bar{T} = \frac{\hat{p}^2}{2m}, \quad \bar{V}(\tau) \equiv \bar{V}(\hat{q}, \tau), \quad (12)$$

i.e. the τ -dependence enters through a potential term that is multiplicative in the $|q\rangle$ basis. (This is the case relevant to the influence-functional construction in Sec. IV.)

We now discretise $[0, \beta]$ into N slices of width $\Delta = \beta/N$ and define $\tau_j = j\Delta$, $\bar{H}_j := \bar{H}(\tau_j)$. The τ -ordered exponential is then given by the Trotter product [61, 62]

$$\hat{\tau} \exp \left[- \int_0^\beta d\tau \bar{H}(\tau) \right] = \lim_{N \rightarrow \infty} e^{-\Delta \bar{H}_N} \dots e^{-\Delta \bar{H}_1} e^{-\Delta \bar{H}_0}. \quad (13)$$

Inserting N resolutions of the identity [61] in the $|q\rangle$ basis between factors, with $q_0 = q_i$ and $q_{N+1} = q_f$, we obtain

Substituting Eq. (17) and Eq. (16) into Eq. (14) yields

$$\bar{\rho}(q_f, q_i; \beta) = \lim_{\Delta \rightarrow 0} \int \prod_{j=1}^N dq_j \prod_{j=0}^N \left(\sqrt{\frac{m}{2\pi\Delta}} \right) \exp \left[- \sum_{j=0}^N \Delta \left(\frac{m}{2} \left(\frac{q_{j+1} - q_j}{\Delta} \right)^2 + \bar{V}(q_j, \tau_j) \right) \right]. \quad (18)$$

In the limit $N \rightarrow \infty$ (i.e. $\Delta \rightarrow 0$), Eq. (18) becomes the Euclidean path integral [60, 61, 63], such that the reduced density (normalised by $Z_{\bar{H}}$) is given by

$$\rho_{\bar{H}}(q_f, q_i; \beta) = \frac{1}{Z_{\bar{H}}} \int_{q(0)=q_i}^{q(\beta)=q_f} \mathcal{D}q(\tau) \exp \left[- \int_0^\beta d\tau \left(\frac{m}{2} \dot{q}(\tau)^2 + \bar{V}(q(\tau), \tau) \right) \right]. \quad (19)$$

The Euclidean action $S[\bar{q}] = \int_0^\beta d\tau \left[\frac{m}{2} \dot{q}^2 + \bar{V}(q, \tau) \right]$ appearing in the exponent defines the system action referred to in subsequent sections. We note that for finite-dimensional systems, while the specific form of the path integral measure changes (e.g., to a sum over discrete trajectories or coherent states), the quenched density representation retains precisely the same form [51, 64]. The separation into a bare system action and a bath-induced stochastic driving is a consequence of the bath properties alone, and does not depend on the dimensionality of the system Hilbert space [65].

$$\bar{\rho}_Q(q_f, q_i; \beta) = \int_{q(0)=q_i}^{q(\beta)=q_f} \mathcal{D}q \exp \left[- S_Q[q] + \frac{1}{2} \int_0^\beta d\tau \int_0^\beta d\tau' f(q(\tau)) K(\tau - \tau') f(q(\tau')) \right]. \quad (20)$$

Here $Z_X(\beta) = \text{Tr}_X e^{-\beta H_X}$ is the free bath partition function, S_Q is the Euclidean system action defined and $K(\tau)$

the kernel

$$\bar{\rho}(q_f, q_i; \beta) = \lim_{N \rightarrow \infty} \int \prod_{j=1}^N dq_j \prod_{j=0}^N \langle q_{j+1} | e^{-\Delta \bar{H}_j} | q_j \rangle. \quad (14)$$

For small Δ we use a first-order Trotter splitting (sufficient in the continuum limit) [61]

$$e^{-\Delta(\bar{T} + \bar{V}_j)} = e^{-\Delta \bar{T}} e^{-\Delta \bar{V}_j} + \mathcal{O}(\Delta^2), \quad \bar{V}_j := \bar{V}(\hat{q}, \tau_j), \quad (15)$$

so that

$$\langle q_{j+1} | e^{-\Delta \bar{H}_j} | q_j \rangle = \langle q_{j+1} | e^{-\Delta \bar{T}} | q_j \rangle e^{-\Delta \bar{V}(q_j, \tau_j)} + \mathcal{O}(\Delta^2). \quad (16)$$

The kinetic matrix element is evaluated by inserting a momentum resolution of the identity [60, 62]:

$$\begin{aligned} \langle q_{j+1} | e^{-\Delta \bar{T}} | q_j \rangle &= \int \frac{dp}{2\pi} \langle q_{j+1} | p \rangle \langle p | q_j \rangle e^{-\Delta p^2/(2m)} \\ &= \sqrt{\frac{m}{2\pi\Delta}} \exp \left[-\frac{m}{2\Delta} (q_{j+1} - q_j)^2 \right]. \end{aligned} \quad (17)$$

IV. FROM INFLUENCE FUNCTIONALS TO THE HAMILTONIAN OF MEAN FORCE

Having established the equivalence between the quenched density and the standard Euclidean path integral in the previous section, we now outline the specific form of the reduced system when coupled to a harmonic bath. Following the influence-functional formalism [34–36] and specifically its stochastic representation for Gaussian baths [42, 52, 53]. The resulting reduced density matrix admits the imaginary-time path integral representation:

is the imaginary-time influence kernel. In terms of the

bath spectral density $J(\omega)$, the kernel is given by [34, 51]

$$K(\tau) = \frac{1}{\pi} \int_0^\infty d\omega J(\omega) \frac{\cosh\left[\omega\left(\frac{\beta}{2} - |\tau|\right)\right]}{\sinh\left(\frac{\beta\omega}{2}\right)}. \quad (21)$$

While the representation in Eq. (20) is exact, it is non-local (bilocal) in imaginary time. This nonlocality can be resolved by a Hubbard-Stratonovich (HS) transformation [38, 39, 66], which decouples the bilocal interaction by introducing an auxiliary Gaussian field $\xi(\tau)$. This step provides the bridge to an operator-level representation of $\bar{\rho}_Q$ as a Gaussian average over the local quenched canonical densities discussed in Sec. III:

$$\bar{\rho}_Q(\beta) = \mathbb{E}_\xi \left[\hat{T} \exp \left(- \int_0^\beta d\tau \bar{H}(\tau; \xi) \right) \right], \quad (22)$$

where $\bar{H}(\tau; \xi) = H_Q - \xi(\tau)f$ is the time-dependent quenched Hamiltonian. The statistical properties of the auxiliary field $\xi(\tau)$ are set by the influence kernel [42, 52, 53]

$$\mathbb{E}_\xi[\xi(\tau)] = 0, \quad \mathbb{E}_\xi[\xi(\tau)\xi(\tau')] = K(\tau - \tau'). \quad (23)$$

This Hubbard-Stratonovich unravelling is the same as that underlying partition-free stochastic Liouville formulations of Gaussian environments and equilibrated initial conditions [42]. Recent work also develops a time-local minimally extended-state-space embedding (QD-MESS) via influence-functional decomposition and Hubbard-Stratonovich unravelling, recovering Caldeira-Leggett and Smoluchowski-type limits as special cases [66].

Having performed this transformation, the operator representation maps directly to the quenched density framework. The challenge now lies in performing the average over ξ . To do so, we must isolate the stochastic term. Moving to the interaction picture with respect to H_Q , the propagator factorises as

$$\hat{T} e^{-\int_0^\beta d\tau [H_Q - \xi(\tau)f]} = e^{-\beta H_Q} \hat{T} e^{\int_0^\beta d\tau \xi(\tau)f(\tau)}, \quad (24)$$

where $f(\tau) = e^{\tau H_Q} f e^{-\tau H_Q}$. While we have successfully isolated the stochastic field $\xi(\tau)$, the coupling operator $f(\tau)$ has acquired a nontrivial imaginary-time dependence. This greatly complicates the averaging out of the stochastic term, and we defer its explicit evaluation to future work.

Here we focus instead on the commuting (QND) sector $[H_Q, f] = 0$, where the algebraic obstruction disappears: $f(\tau) = f$ is τ -independent, the $\hat{T}$ ordering becomes redundant and the expression factorises:

$$\hat{T} \exp \left[- \int_0^\beta d\tau \bar{H}(\tau; \xi) \right] = e^{-\beta H_Q} \exp[f X], \quad (25)$$

where $X := \int_0^\beta d\tau \xi(\tau)$ is a Gaussian variable with mean zero and variance

$$C(\beta) = \mathbb{E}_\xi[X^2] = \int_0^\beta d\tau \int_0^\beta d\tau' K(\tau - \tau'). \quad (26)$$

Substituting the kernel Eq. (21) and performing the double imaginary-time integral gives

$$C(\beta) = \frac{2\beta}{\pi} \int_0^\infty d\omega \frac{J(\omega)}{\omega} \equiv 2\beta \lambda, \quad \lambda := \frac{1}{\pi} \int_0^\infty d\omega \frac{J(\omega)}{\omega}, \quad (27)$$

where λ is the reorganisation energy [51, 65]. Crucially, $C(\beta)$ is *linear* in β , so the ratio $C(\beta)/(2\beta) = \lambda$ is temperature independent. The closed commuting-sector result depends on the bath only through this spectral integral; the Ohmic form used below is therefore a numerical benchmark choice rather than a restriction of the derivation.

With this the Gaussian average $\mathbb{E}_\xi[e^{fX}] = \exp[\frac{1}{2}C(\beta)f^2]$ can be performed exactly, and using Eq. (22), we obtain:

$$\bar{\rho}_Q(\beta) = e^{-\beta H_Q} \exp[\beta \lambda f^2]. \quad (28)$$

This is (up to normalisation) the canonical exponential form for a thermal state, and the Hamiltonian of mean force is identically

$$H_{\text{MF}}(\beta) = H_Q - \lambda f^2 \quad (29)$$

The only nontrivial operator content is the correction $-\lambda f^2$, which is temperature independent.

An immediate consequence that can be inferred from this result is that it is identical to the result in the classical mean-force construction [15, 50, 67]. This equivalence is most easily understood through the link to Koopman von-Neumann dynamics. In real time, classical evolution is generated by the Liouville/Koopman operator $\mathcal{L}_H = \{\cdot, H\}$, where $\{\cdot, H\}$ denotes the classical Poisson bracket, with propagator $e^{t\mathcal{L}_H}$. To obtain Gibbs states from imaginary time however, the relevant generator becomes the Hamiltonian. This is true in both the quantum and classical scenarios, but in the latter case all observable operators (and hence the Hamiltonian) commute identically. The classical equilibrium construction will therefore always lie within the commuting sector of the fully quantum problem. For a fuller discussion of the dynamical distinctions between quantum and classical systems in both real and imaginary time, see ref. [68].

A slight (but interesting) corollary of this result concerns integrable systems where the coupling operator f is a conserved quantity ($[H_Q, f] = 0$). In this case, the Hamiltonian of mean force H_{MF} shares the same eigenbasis as the bare system Hamiltonian. The strong coupling to the bath therefore does not mix the system energy

levels, but rather induces a level-dependent energy shift $-\lambda f_n^2$, where f_n are the eigenvalues of the coupling operator. The integrability of the open system is thus preserved in the sense that the equilibrium state remains diagonal in the energy basis of the isolated system, but depending on the specific structure of f , the bath can induce a non-trivial operator-level effect on the reduced equilibrium state.

A final important consequence concerns the relationship between the mean force correction, and the system counterterm that is often added to enforce translational invariance of the bath in the Caldeira-Leggett model. Explicitly, we see that λ coincides with the static potential renormalisation $\frac{1}{2} \sum_k c_k^2 / (m_k \omega_k^2)$. If the standard counterterm $+\lambda f^2$ has been added to H_Q (as stated in Sec. II), the bath-induced shift $-\lambda f^2$ cancels it exactly:

$$H_Q = H_Q^{\text{bare}} + \lambda f^2 \implies H_{\text{MF}}(\beta) = H_Q^{\text{bare}} + \text{const.} \quad (30)$$

Thus, in the commuting sector with the standard counterterm convention, the normalised mean-force Gibbs state coincides with the bare-system Gibbs state: the bath has no residual operator-level effect on the reduced equilibrium state. Conversely, if no counterterm is included, $-\lambda f^2$ represents a genuine bath-induced static renormalisation of the system potential. Amusingly, this result closely aligns to the master equation folk wisdom where superior results can be obtained when one neglects *both* counterterms and Lamb-shifts [17].

There is perhaps a deeper lesson hiding within this observation. Consider that in the commuting sector, the system operator f does not drive the bath dynamically, but merely labels a family of displaced harmonic equilibria. Tracing out the bath then asks a simple thermodynamic question: what is the free-energy change of a harmonic environment whose equilibrium position has been shifted by an amount proportional to f ? Because the harmonic trace is invariant under this translation, the only remnant is the displacement free energy, $-\lambda f^2$. The counterterm is precisely the same object inserted with the opposite sign before the bath is traced out. Another equivalent perspective regards the counterterm as subtracting the static susceptibility (i.e. a Matsubara zero-frequency mode) contribution of the influence kernel. In the commuting case only this zero mode contributes to the HMF, as within the interaction picture f is definitionally τ -independent. From these perspectives, the cancellation between counterterm and commuting HMF correction may be regarded as definitionally necessary, rather than merely coincidental.

Having established the quenched density representation exactly captures the reorganisation-energy result [50], we now apply it to stochastic numerical evaluation of the equilibrium state.

V. NUMERICAL EVALUATION

We now verify the predicted H_{MF} can be obtained numerically through stochastic averaging. To do so, we consider a toy five-level system $H_S = \sum_{n=0}^4 E_n |n\rangle\langle n|$ coupled to the bath via the projector $f = |4\rangle\langle 4|$. This coupling is manifestly commuting ($[H_S, f] = 0$), ensuring that the exact HMF should be

$$H_{\text{MF}} = H_S - \lambda |4\rangle\langle 4| + \text{const.} \quad (31)$$

Although the benchmark uses the projector $P = |4\rangle\langle 4|$, this is not a special assumption in the derivation. For any commuting Hermitian coupling $f = \sum_n f_n |n\rangle\langle n|$, Eq. (29) gives level shifts $-\lambda f_n^2$. As such, the choice to project to a single level is purely one of convenience and interpretational clarity, as in the commuting system-interaction regime all bath-induced operator content is localised to a single sector, whose effect can be resolved directly in the populations and fitted HMF coefficients.

The precise level spacings are selected randomly and kept fixed across all (β, g) simulations. For the bath, we use $J_g(\omega) = 2g^2 \omega e^{-\omega/\omega_c}$, incorporating system-bath coupling strength directly into the parameter g . We choose $\omega_c = 4$ for all simulations. This Ohmic spectrum is used only to provide a concrete numerical test case: the analytic commuting result depends on the bath through λ alone, so any spectral density for which the reorganisation integral is well defined gives the same operator form. As a point of comparison, we further implement an explicit bath of K harmonic modes a_k , coupling to the system through $\sum_k a_k + a_k^\dagger$. The bath spectrum is then mimicked by discretising the interval $\omega \in [0, \omega_{\text{max}}]$ into mode bins with centres ω_k and widths $\Delta\omega_k$, setting $c_k^2 = \frac{2}{\pi} J_g(\omega_k) \omega_k \Delta\omega_k$. This ensures that the discretised reorganisation scale $\lambda_{\text{disc}} = \sum_k \frac{c_k^2}{2\omega_k}$ tracks the continuum expression $\lambda_{\text{cont}} = \frac{1}{\pi} \int_0^\infty \frac{J_g(\omega)}{\omega} d\omega$ [51]. This provides an exact reference for the reduced state $\rho_Q^{\text{ED}} = \text{Tr}_B[e^{-\beta H_{\text{tot}}}] / \text{Tr}[e^{-\beta H_{\text{tot}}}]$.

Finally, we implement the stochastic formulation of the quenched density given by Eq. (22). The Gaussian noise $\xi(\tau)$ is generated by convolution of white noise with the covariance matrix of the discretised kernel $K(\tau)$, a standard technique for sampling coloured noise (see e.g. [52]). Each stochastic estimate uses 50,000 path samples on a 32-point imaginary-time grid. Accuracy is assessed both by the resulting standard errors and the convergence of the noise of $X = \int_0^\beta d\tau \xi(\tau)$ to its predicted variance $2\beta\lambda_{\text{disc}}$. Fig. 1 confirms this scaling, plotting the sample variance extracted from the stochastic simulations against β (panel a) and g^2 (panel b).

With this setup the core predictions of the HMF framework can be verified. Fig. 2 confirms the thermodynamic consistency of the quenched density framework, showing that the effective Hamiltonian correctly captures the




FIG. 1. Variance scaling. The sample variance σ_ξ^2 of the stochastic field grows linearly with inverse temperature β (a) and linearly with the reorganisation energy $\lambda \propto g^2$ (b). Symbols are simulation results from 50,000 path samples on a 32-point imaginary-time grid; dotted lines are the theoretical prediction $2\beta\lambda_{\text{disc}}$.

temperature dependence of the open system, with the extracted reorganisation energy matching the theoretical value.

Fig. 3 demonstrates robustness in the strong-coupling regime, where the interaction strength rivals the system energy scales. To verify the operator structure of the HMF, we fit the numerically extracted H_{MF} to the ansatz $c_0\mathbb{I} + a_{H_S}H_S + a_P P$. As shown in panel (b), the coefficient a_P tracks the theoretical prediction $-\lambda$ perfectly, while $a_{H_S} \approx 1$ confirms that the bare system Hamiltonian is not renormalised. Finally, Fig. 4 provides the microscopic resolution: the HMF modifies only the coupled subspace, leaving the orthogonal manifold untouched, as predicted by the operator structure of Eq. 29.




FIG. 2. Temperature dependence and scaling. (a) Population of the coupled state ($|4\rangle$) vs β for various couplings g . Circles (ED) match solid lines (analytic). (b) The extracted reorganisation energy λ_{est} collapsed by g^2 vs β . All curves hover near the theoretical thermodynamic limit (dashed lines), verifying that the bath's effect reduces to a single reorganisation energy λ .

VI. DISCUSSION AND OUTLOOK

We have presented the quenched density framework as a rigorous bridge between path-integral influence functionals and operator-level thermodynamics. By mapping the bath trace to a stochastic average over quenched propagators, we isolate the algebraic structure of the system's response from the statistical properties of the environment.

In the case of harmonic baths where H_Q and f commute, the algebraic and statistical contributions to the HMF can be cleanly partitioned: the bath statistics collapse to a scalar reorganisation energy λ , and the system algebra reduces to a static potential shift $-\lambda f^2$. Our numerical benchmarks (Sec. V) confirm that this framework reproduces the exact finite-bath physics, regardless




FIG. 3. Strong-coupling benchmark. (a) Population p_4 vs g^2 at fixed β , showing excellent agreement over the full range. (b) Coefficients in the system basis vs g^2 . The projector component $P_4 = |4\rangle\langle 4|$ scales as $-\lambda$, while off-diagonal coherences are exponentially suppressed. This strong-coupling projector dominance is consistent with ultrastrong-coupling analyses of the mean-force Gibbs state [18]. The bare system Hamiltonian H_S remains unrenormalised (dotted), confirming $H_{MF} = H_S - \lambda f^2$.

of coupling strength.

The exact result derived here is hardly novel, and in isolation can hardly justify the formal machinery used to derive it. In this regard, our treatment is incomplete. The power of the quenched density lies in its modularity, which naturally accommodates both more general environments and system-bath couplings. While we defer the full treatment to future work, it is possible to sketch the route to generalisation.

For non-commuting couplings, the object to control is the interaction-picture operator $f(\tau) = e^{\tau H_Q} f e^{-\tau H_Q}$. A natural extension is to expand it in the adjoint chain $f_n = \text{ad}_{H_Q}^n(f)$, so that the Gaussian bath enters through scalar kernel moments multiplying products $f_n f_m$. This turns the problem of finding the HMF into a question




FIG. 4. Microscopic validation. (a) Detailed population snapshot at fixed β, g : Analytic HMF (blue) corrects the bare Gibbs (grey) to match ED (orange). (b) Effective energy shifts Δ_i for each level; only the coupled level ($|4\rangle$) shifts, by exactly $-\lambda$. Non-coupled levels remain unperturbed.

of closure on the Lie Algebra generated by products of adjoint operators f_n and H_Q . When this algebra closes, there exists a finite representation of the HMF. In the sequel this result will be demonstrated explicitly, but we see its shadow even in the present commuting case, where the algebra closes trivially and the HMF reduces to a static potential renormalisation. In this sense, we can already see the shape of the answer to the question with which we began. What governs equilibrium? Algebra.

-
- [1] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2002).
 - [2] P. Hänggi, G.-L. Ingold, and P. Talkner, Finite quantum dissipation: The challenge of obtaining specific heat, *New Journal of Physics* **10**, 115008 (2008).

- [3] G.-L. Ingold, P. Hänggi, and P. Talkner, Specific heat anomalies of open quantum systems, *Physical Review E* **79**, 061105 (2009).
- [4] M. Campisi, P. Talkner, and P. Hänggi, Fluctuation Theorem for Arbitrary Open Quantum Systems, *Physical Review Letters* **102**, 210401 (2009).
- [5] C. Jarzynski, Nonequilibrium work theorem for a system strongly coupled to a thermal environment, *Journal of Statistical Mechanics: Theory and Experiment* **2004**, P09005 (2004).
- [6] F. C. Binder, J. Goold, and K. Modi, Energetic fluctuations in an open quantum process, *Phys. Rev. E* **100**, 062120 (2019).
- [7] M. Esposito, M. A. Ochoa, and M. Galperin, Nature of heat in strongly coupled open quantum systems, *Physical Review B* **92**, 235440 (2015).
- [8] Á. Rivas, Strong Coupling Thermodynamics of Open Quantum Systems, *Physical Review Letters* **124**, 160601 (2020).
- [9] A. M. Lacerda, A. Purkayastha, M. Kewming, G. T. Landi, and J. Goold, Quantum thermodynamics with fast driving and strong coupling via the mesoscopic leads approach, *Phys. Rev. B* **107**, 195117 (2023).
- [10] P. Strasberg and M. Esposito, Measurability of nonequilibrium thermodynamics in terms of the Hamiltonian of mean force, *Physical Review E* **101**, 050101 (2020).
- [11] H. J. D. Miller and J. Anders, Entropy production and time asymmetry in the presence of strong interactions, *Phys. Rev. E* **95**, 032123 (2017).
- [12] F. Binder, R. Auccaise, J. Goold, and K. Modi, Operational thermodynamics of open quantum systems, *Physical Review E* **99**, 032102 (2019).
- [13] F. Binder, S. Vinjanampathy, K. Modi, and J. Goold, Quantum thermodynamics of general quantum processes, *Physical Review E* **91**, 032119 (2015).
- [14] C. Jarzynski, Stochastic and Macroscopic Thermodynamics of Strongly Coupled Systems, *Physical Review X* **7**, 011008 (2017).
- [15] P. Talkner and P. Hänggi, *Colloquium* : Statistical mechanics and thermodynamics at strong coupling: Quantum and classical, *Reviews of Modern Physics* **92**, 041002 (2020).
- [16] U. Seifert, First and Second Law of Thermodynamics at Strong Coupling, *Physical Review Letters* **116**, 020601 (2016).
- [17] L. A. Correa and J. Glatthard, Potential renormalisation, Lamb shift and mean-force Gibbs state – to shift or not to shift?, *Quantum* **9**, 1868 (2025).
- [18] J. D. Cresser and J. Anders, Weak and Ultrastrong Coupling Limits of the Quantum Mean Force Gibbs State, *Physical Review Letters* **127**, 250601 (2021).
- [19] A. S. Trushechkin, M. Merkli, J. D. Cresser, and J. Anders, Open quantum system dynamics and the mean force Gibbs state, *AVS Quantum Science* **4**, 012301 (2022).
- [20] C. R. Hogg, J. Glatthard, F. Cerisola, and J. Anders, Tutorial on the stochastic simulation of dissipative quantum oscillators, *The Journal of Chemical Physics* **161**, 071501 (2024).
- [21] C. R. Hogg, F. Cerisola, J. D. Cresser, S. A. R. Horsley, and J. Anders, Enhanced entanglement in multi-bath spin-boson models, *Quantum* **8**, 1357 (2024).
- [22] S. Jang, Y.-C. Cheng, D. R. Reichman, and R. J. Silbey, Nonequilibrium theory of electronic excitation energy transfer: The small polaron quantum master equation, *The Journal of Chemical Physics* **129**, 101104 (2008).
- [23] M. Shubbrook, J. Iles-Smith, and A. Nazir, Non-markovian quantum heat statistics with the reaction coordinate mapping, *Quantum Science and Technology* **10**, 025063 (2025).
- [24] X. Du, J. Wang, and Y. Ma, Generalized Hamiltonian of mean-force approach to open quantum systems coupled to finite baths in thermoequilibrium, *Physical Review E* **111**, 024139 (2025).
- [25] A. Nazir and G. Schaller, The reaction coordinate mapping in quantum thermodynamics, *Thermodynamics in the Quantum Regime* , 551 (2018).
- [26] N. Anto-Sztrikacs, A. Nazir, and D. Segal, Effective-hamiltonian theory of open quantum systems at strong coupling, *PRX Quantum* **4**, 020307 (2023).
- [27] J. Iles-Smith, A. G. Dijkstra, N. Lambert, and A. Nazir, Energy transfer in structured and unstructured environments: Master equations beyond the born–markov approximations, *The Journal of Chemical Physics* **144**, 044110 (2016).
- [28] J. M. Moix, Y. Zhao, and J. Cao, Equilibrium-reduced density matrix formulation: Influence of noise, disorder, and temperature on localization in excitonic systems, *Physical Review B* **85**, 115412 (2012).
- [29] X. Chen, The rigorous stochastic matrix multiplication scheme for the calculations of reduced equilibrium density matrices of open multilevel quantum systems, *The Journal of Chemical Physics* **140**, 154101 (2014).
- [30] N. Makri, Exploiting classical decoherence in dissipative quantum dynamics: Memory, phonon emission, and the blip sum, *Chemical Physics Letters* **593**, 93 (2014).
- [31] Y. Tanimura, Reduced hierarchical equations of motion in real and imaginary time: Correlated initial states and thermodynamic quantities, *The Journal of Chemical Physics* **141**, 044114 (2014).
- [32] L. Song and Q. Shi, Calculation of correlated initial state in the hierarchical equations of motion method using an imaginary time path integral approach, *The Journal of Chemical Physics* **143**, 194106 (2015).
- [33] P. L. Walters and F. Wang, Path integral quantum algorithm for simulating non-markovian quantum dynamics in open quantum systems, *Phys. Rev. Res.* **6**, 013135 (2024).
- [34] R. P. Feynman and F. L. Vernon, The theory of a general quantum system interacting with a linear dissipative system, *Annals of Physics* **24**, 118 (1963).
- [35] A. O. Caldeira and A. J. Leggett, Quantum tunnelling in a dissipative system, *Annals of Physics* **149**, 374 (1983).
- [36] H. Grabert, P. Schramm, and G.-L. Ingold, Quantum Brownian motion: The functional integral approach, *Physics Reports* **168**, 115 (1988).
- [37] M. Xu, V. Vadimov, J. T. Stockburger, and J. Ankerhold, Colloquium: Simulating non-markovian dynamics in open quantum systems, *Rev. Mod. Phys.* , (2026).
- [38] J. Hubbard, Calculation of Partition Functions, *Physical Review Letters* **3**, 77 (1959).
- [39] R. L. Stratonovich, On a method of calculating quantum distribution functions, *Soviet Physics Doklady* **2**, 416 (1957).
- [40] J. T. Stockburger and H. Grabert, Exact c -Number Representation of Non-Markovian Quantum Dissipation, *Physical Review Letters* **88**, 170407 (2002).
- [41] J. T. Stockburger, Simulating spin-boson dynamics with

- stochastic Liouville–von Neumann equations, *Chemical Physics* **296**, 159 (2004).
- [42] G. M. G. McCaul, C. D. Lorenz, and L. Kantorovich, Partition-free approach to open quantum systems in harmonic environments: An exact stochastic Liouville equation, *Physical Review B* **95**, 125124 (2017).
- [43] M. Wiedmann, J. T. Stockburger, and J. Ankerhold, Non-Markovian dynamics of a quantum heat engine: Out-of-equilibrium operation and thermal coupling control, *New Journal of Physics* **22**, 033007 (2020).
- [44] M. A. Lane, D. Matos, I. J. Ford, and L. Kantorovich, Exactly thermalized quantum dynamics of the spin-boson model coupled to a dissipative environment, *Physical Review B* **101**, 224306 (2020).
- [45] D. Matos, M. A. Lane, I. J. Ford, and L. Kantorovich, Efficient Choice of Colored Noise in the Stochastic Dynamics of Open Quantum Systems, *Physical Review E* **102**, 062134 (2020).
- [46] D. Matos, L. Kantorovich, and I. J. Ford, Stochastic Entropy Production for Continuous Measurements of an Open Quantum System, *Journal of Physics Communications* **6**, 125003 (2022).
- [47] A. A. Clerk, M. H. Devoret, S. M. Girvin, F. Marquardt, and R. J. Schoelkopf, Introduction to quantum noise, measurement, and amplification, *Rev. Mod. Phys.* **82**, 1155 (2010).
- [48] Y. Makhlin, G. Schön, and A. Shnirman, Quantum-state engineering with josephson-junction devices, *Rev. Mod. Phys.* **73**, 357 (2001).
- [49] G. M. Palma, K.-A. Suominen, and A. K. Ekert, Quantum computers and dissipation, *Proc. Roy. Soc. Lond. A* **452**, 567 (1996).
- [50] M. Campisi, P. Talkner, and P. Hänggi, Thermodynamics and fluctuation theorems for a strongly coupled open quantum system: An exactly solvable case, *Journal of Physics A: Mathematical and Theoretical* **42**, 392002 (2009).
- [51] U. Weiss, *Quantum Dissipative Systems*, 4th ed. (World Scientific, Singapore, 2012).
- [52] G. M. G. McCaul, C. D. Lorenz, and L. Kantorovich, Driving spin-boson models from equilibrium using exact quantum dynamics, *Physical Review B* **97**, 224310 (2018).
- [53] G. McCaul and Denys. I. Bondar, How to win friends and influence functionals: Deducing stochasticity from deterministic dynamics, *The European Physical Journal Special Topics* **230**, 733 (2021).
- [54] A. J. Coleman and V. I. Yukalov, *Reduced Density Matrices: Coulson's Challenge*, Lecture Notes in Chemistry, Vol. 72 (Springer, 2000).
- [55] D. A. Mazziotti, ed., *Reduced-Density-Matrix Mechanics: With Application to Many-Electron Atoms and Molecules*, Advances in Chemical Physics, Vol. 134 (John Wiley & Sons, 2007).
- [56] Y.-K. Liu, M. Christandl, and F. Verstraete, Quantum computational complexity of the N -representability problem: QMA complete, *Physical Review Letters* **98**, 110503 (2007).
- [57] A. A. Klyachko, Quantum marginal problem and N -representability, *Journal of Physics: Conference Series* **36**, 72 (2006).
- [58] K. V. Hovhannisyanyan, S. Nemati, C. Henkel, and J. Anders, Hamiltonian of mean force for strongly-coupled systems, *Physical Review E* **102**, 012104 (2020).
- [59] In the sequel we shall revisit and resolve this issue.
- [60] R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York, 1965).
- [61] H. Kleinert, *Path Integrals in Quantum Mechanics, Statistics, Polymer Physics, and Financial Markets*, 5th ed. (World Scientific, Singapore, 2009).
- [62] L. S. Schulman, *Techniques and Applications of Path Integration* (Wiley, New York, 1981).
- [63] C. Grosche and F. Steiner, *Handbook of Feynman Path Integrals*, Springer Tracts in Modern Physics, Vol. 145 (Springer, Berlin, 1998).
- [64] A. Altland and B. D. Simons, *Condensed Matter Field Theory*, 2nd ed. (Cambridge University Press, 2010).
- [65] A. J. Leggett, S. Chakravarty, A. T. Dorsey, M. P. A. Fisher, A. Garg, and W. Zwerger, Dynamics of the dissipative two-state system, *Reviews of Modern Physics* **59**, 1 (1987).
- [66] M. Xu and J. Ankerhold, Quantum brownian motion for embedded system, *The European Physical Journal Special Topics* 10.1140/epjs/s11734-025-01925-0 (2025), published 19 September 2025.
- [67] F. Cerisola, M. Berritta, S. Scali, S. A. R. Horsley, J. D. Cresser, and J. Anders, Quantum–classical correspondence in spin–boson equilibrium states at arbitrary coupling, *New Journal of Physics* **26**, 053032 (2024).
- [68] G. McCaul, D. V. Zhdanov, and D. I. Bondar, Wave operator representation of quantum and classical dynamics, *Physical Review A* **108**, 052208 (2023).