




Girls aged 0-5 years: Head Circumference

THAI SOCIETY
for PEDIATRIC
ENDOCRINOLOGY

Name _____ DOB _____ HN _____



Tanner Stage			Breast	Pubic Hair
I			Elevation of nipple only, as in early childhood	No pubic hair
II			Breast bud Slight elevation of breast and nipple Enlargement of areola diameter	Sparse, slightly pigmented, straight or only slightly curled Spreads along labia
III			Further enlarged breast and areola with contour development	Darker, coarser, and more curled Spreads sparsely over the junction of pubes
IV			Projection of areola and nipple to form a secondary mound above breast contour level	Adult type Area covered smaller than in adults, no spread to medial thighs
V			Mature stage Protruding of nipple only	Adult in quantity and type Distributed as an inverse triangle, spread to medial surface of thighs but not up the linea alba or above base of inverse triangle

References: Marshall WA, Tanner JM. Variations in pattern of pubertal changes in girls. Arch Dis Child 1969;44:291-303.

permadit ภูมิศาสตร์ การเป็นหนุ่มสาวก่อนวัย พิมพ์ครั้งที่ 1 กรุงเทพฯ: ไฮลิสติก พับลิชิ่ง; 2558.

Designed by Thai Society for Pediatric Endocrinology, 2022