

Introduction to Algorithms

Lecture 13 Number Theoretic Algorithm

Xue Chen

xuechen1989@ustc.edu.cn

2025 spring in



Outline

1 Introduction

2 Prime Number Theorem


3 Basic Tools

4 Primality Testing

Introduction

Many algorithms uses number theory

- 1 Hash functions
- 2 Coding theory
- 3 Cryptography





■ $y^2 = x^3 - 2x + 15$

Introduction

Many algorithms uses number theory

- 1 Hash functions
- 2 Coding theory
- 3 Cryptography



Primes are the backbone of Number theory

Most Fundamental Problem in Number Theory

How to find a **large prime** number p ?

Set up

Our Focus

Given n say 10^3 or 10^4 , find a prime number of n digits (in binary).

Set up

Our Focus

Given n say 10^3 or 10^4 , find a prime number of n digits (in binary).

- 1 Basic idea: Sieve algorithm?

Set up

Our Focus

Given n say 10^3 or 10^4 , find a prime number of n digits (in binary).


- 1 Basic idea: Sieve algorithm?
- 2 But we want a number between 2^{n-1} and $2^n - 1$!

Set up

Our Focus

Given n say 10^3 or 10^4 , find a prime number of n digits (in binary).

- 1 Basic idea: Sieve algorithm?
- 2 But we want a number between 2^{n-1} and $2^n - 1$!
- 3 Lots of interesting math: prime number THM, Euclid's ALGO, Fermat's THM, chinese remainder THM, primality testing



Basic Algorithm

```
procedure GENERATE-PRIME( $n$ )  
  repeat  
    Generate a random number  $q$  of  $n$  digits  
  until  $q$  is a prime
```

Basic Algorithm

```
procedure GENERATE-PRIME( $n$ )  
  repeat  
    Generate a random number  $q$  of  $n$  digits  
  until  $q$  is a prime
```

To finish this algorithm and analyze it, lots of questions:

- 1 Density of primes in n -digit numbers?

Basic Algorithm

```
procedure GENERATE-PRIME( $n$ )  
  repeat  
    Generate a random number  $q$  of  $n$  digits  
  until  $q$  is a prime
```

To finish this algorithm and analyze it, lots of questions:

- 1 Density of primes in n -digit numbers?
- 2 How to test whether q is a prime or not efficiently?
- 3 Many tools ...

Basic Algorithm

```
procedure GENERATE-PRIME( $n$ )  
  repeat  
    Generate a random number  $q$  of  $n$  digits  
  until  $q$  is a prime
```

To finish this algorithm and analyze it, lots of questions:

- 1 Density of primes in n -digit numbers?
- 2 How to test whether q is a prime or not efficiently?
- 3 Many tools ...
- 4 To the best of my knowledge, fastest algorithm in time $\approx n^3$


Outline

1 Introduction

2 Prime Number Theorem


3 Basic Tools

4 Primality Testing



Prime Number Theorem — Most fundamental theorem of primes

$\pi(X)$ denotes # primes less than X — $\pi(N) \approx \frac{N}{\ln N}$.




Prime Number Theorem — Most fundamental theorem of primes

$\pi(X)$ denotes # primes less than X — $\pi(N) \approx \frac{N}{\ln N}$.



- Legendre, Gauss and others discovered the curve around 1800s, the THM was proved around 1890 by Hadamard and Poussin




Prime Number Theorem — Most fundamental theorem of primes

$\pi(X)$ denotes # primes less than X — $\pi(N) \approx \frac{N}{\ln N}$.




- ① Legendre, Gauss and others discovered the curve around 1800s, the THM was proved around 1890 by Hadamard and Poussin
- ② Riemann made a bolder claim 1852 — called Riemann hypothesis



Prime Number Theorem — Most fundamental theorem of primes

$\pi(X)$ denotes # primes less than X — $\pi(N) \approx \frac{N}{\ln N}$.



- ① Legendre, Gauss and others discovered the curve around 1800s, the THM was proved around 1890 by Hadamard and Poussin
- ② Riemann made a bolder claim 1852 — called Riemann hypothesis
- ③ Twin prime number THM by Yitang Zhang 2013

Legendre



Weak Proof by Chebyshev

Let us show $\pi(N) = \Theta(N/\ln N)$

Weak Proof by Chebyshev

Let us show $\pi(N) = \Theta(N/\ln N)$

- 1 Consider $\binom{2N}{N}$ — all primes in $[N+1, 2N]$ are its factors

Weak Proof by Chebyshev

Let us show $\pi(N) = \Theta(N/\ln N)$

- 1 Consider $\binom{2N}{N}$ — all primes in $[N+1, 2N]$ are its factors
- 2 On the 1st hand, $\binom{2N}{N} = \Theta(2^{2N}/\sqrt{N})$

Weak Proof by Chebyshev

Let us show $\pi(N) = \Theta(N/\ln N)$

- 1 Consider $\binom{2N}{N}$ — all primes in $[N+1, 2N]$ are its factors
- 2 On the 1st hand, $\binom{2N}{N} = \Theta(2^{2N}/\sqrt{N})$
- 3 On the 2nd hand, let us consider the contribution of $p \leq N$ in $\binom{2N}{N}$
— Question: Can we bound k s.t. at most k factors of p in $\binom{2N}{N}$?

Outline

1 Introduction

2 Prime Number Theorem

3 Basic Tools

4 Primality Testing

Tools from number theory

Algorithms test whether a large integer is a prime or not — called primality testing. Before designing testers, what properties do prime number have?

Tools from number theory

Algorithms test whether a large integer is a prime or not — called primality testing. Before designing testers, what properties do prime number have?

- 1 Fix a large prime p of n digits and consider all operations in $\text{mod } p$

Tools from number theory

Algorithms test whether a large integer is a prime or not — called primality testing. Before designing testers, what properties do prime number have?

- 1 Fix a large prime p of n digits and consider all operations in $\text{mod } p$
- 2 In group theory, all elements are in $Z_p = \{0, 1, \dots, p-1\}$
- 3 Let $Z_p^* = \{1, \dots, p-1\}$

Tools from number theory

Algorithms test whether a large integer is a prime or not — called primality testing. Before designing testers, what properties do prime number have?

- 1 Fix a large prime p of n digits and consider all operations in $\text{mod } p$
- 2 In group theory, all elements are in $Z_p = \{0, 1, \dots, p-1\}$
- 3 Let $Z_p^* = \{1, \dots, p-1\}$

Basic Property

For any $a \in \{1, \dots, p-1\}$, $\exists b$ such that $ab \equiv 1 \pmod{p}$, called a^{-1} .

In fact, we could find a^{-1} in time $O(\log p) = O(n) \odot$

Revisit Euclid's Algorithm

Recall that Euclid's algorithm computes $\gcd(a, b)$

Algorithm Euclid's algorithm for GCD

```
function EUCLID( $a, b$ )  
  if  $b=0$  then  
    return  $a$   
  else  
    return Euclid( $b, a \bmod b$ )
```

Revisit Euclid's Algorithm

Recall that Euclid's algorithm computes $\gcd(a, b)$

Algorithm Extended Euclid's algorithm

```
function EUCLID( $a, b$ )  
  if  $b=0$  then  
    return ( $x = 1, y = 0$ )  
  else  
    ( $x_0, y_0$ ) = Euclid( $b, a \bmod b$ )  
    Return ( $y_0, x_0 - [a/b] \cdot y_0$ )
```

Extension

For any a, b , it finds $x, y \in \mathbb{Z}$ such that $ax + by = \gcd(a, b)$

Properties of Prime

Fermat's little Theorem

Fix a prime p , for any $a \in \mathbb{Z}_p^*$, $a^{p-1} \equiv 1 \pmod{p}$.

① Proof?

Properties of Prime

Fermat's little Theorem

Fix a prime p , for any $a \in \mathbb{Z}_p^*$, $a^{p-1} \equiv 1 \pmod{p}$.

- 1 Proof?
- 2 It gives an alternate way to find a^{-1} , how?

Properties of Prime

Fermat's little Theorem

Fix a prime p , for any $a \in \mathbb{Z}_p^*$, $a^{p-1} \equiv 1 \pmod{p}$.

- 1 Proof?
- 2 It gives an alternate way to find a^{-1} , how?
- 3 Can we use it to test primes? Say randomly pick a and compute a^{p-1}
- 4 Does it reject composites?

Properties of Prime

Fermat's little Theorem

Fix a prime p , for any $a \in \mathbb{Z}_p^*$, $a^{p-1} \equiv 1 \pmod{p}$.

- 1 Proof?
- 2 It gives an alternate way to find a^{-1} , how?
- 3 Can we use it to test primes? Say randomly pick a and compute a^{p-1}
- 4 Does it reject composites?
- 5 ☹ 561 is a counter-example: for any $\gcd(a, 561) = 1$, $a^{560} \equiv 1 \pmod{p}$

Properties of Prime

Fermat's little Theorem

Fix a prime p , for any $a \in \mathbb{Z}_p^*$, $a^{p-1} \equiv 1 \pmod{p}$.

- 1 Proof?
- 2 It gives an alternate way to find a^{-1} , how?
- 3 Can we use it to test primes? Say randomly pick a and compute a^{p-1}
- 4 Does it reject composites?
- 5 ☹ 561 is a counter-example: for any $\gcd(a, 561) = 1$, $a^{560} \equiv 1 \pmod{p}$

Properties of Prime

Fermat's little Theorem

Fix a prime p , for any $a \in \mathbb{Z}_p^*$, $a^{p-1} \equiv 1 \pmod{p}$.

- 1 Proof?
- 2 It gives an alternate way to find a^{-1} , how?
- 3 Can we use it to test primes? Say randomly pick a and compute a^{p-1}
- 4 Does it reject composites?
- 5 ☹ 561 is a counter-example: for any $\gcd(a, 561) = 1$, $a^{560} \equiv 1 \pmod{p}$
- 6 Wait! How many a that has $\gcd(a, 561) = 1$?

Properties of Prime

Fermat's little Theorem

Fix a prime p , for any $a \in \mathbb{Z}_p^*$, $a^{p-1} \equiv 1 \pmod{p}$.

- 1 Proof?
- 2 It gives an alternate way to find a^{-1} , how?
- 3 Can we use it to test primes? Say randomly pick a and compute a^{p-1}
- 4 Does it reject composites?
- 5 ☹ 561 is a counter-example: for any $\gcd(a, 561) = 1$, $a^{560} \equiv 1 \pmod{p}$
- 6 Wait! How many a that has $\gcd(a, 561) = 1$?
- 7 Let us introduce Euler function $\phi(N)$

Property: Prime Factorization

Unique Factorization

For any integer N , there is an **unique** factorization of primes

$$N = p_1^{e_1} \cdot p_2^{e_2} \cdots p_r^{e_r}.$$

- 1 Definition of Euler Function: $\phi(N) = (1 - 1/p_1) \cdots (1 - 1/p_r) \cdot N$ if N 's distinct prime factors are p_1, \dots, p_r

Property: Prime Factorization

Unique Factorization

For any integer N , there is an **unique** factorization of primes

$$N = p_1^{e_1} \cdot p_2^{e_2} \cdots p_r^{e_r}.$$

- 1 Definition of Euler Function: $\phi(N) = (1 - 1/p_1) \cdots (1 - 1/p_r) \cdot N$ if N 's distinct prime factors are p_1, \dots, p_r
- 2 Fact: Number of $a \in Z_N$ with $(a, N) = 1$ is $\phi(N)$

Property: Prime Factorization

Unique Factorization

For any integer N , there is an **unique** factorization of primes

$$N = p_1^{e_1} \cdot p_2^{e_2} \cdots p_r^{e_r}.$$

- 1 Definition of Euler Function: $\phi(N) = (1 - 1/p_1) \cdots (1 - 1/p_r) \cdot N$ if N 's distinct prime factors are p_1, \dots, p_r
- 2 Fact: Number of $a \in Z_N$ with $(a, N) = 1$ is $\phi(N)$
- 3 Extension of Fermat's little THM: $a^{\phi(N)} \equiv 1 \pmod{N}$ for any $(a, N) = 1$.

Composite Numbers

To distinguish prime numbers from composites, what properties does a composite have?

Chinese Remainder THM


Given distinct primes p_1, \dots, p_r , consider $(a_1, \dots, a_r) \in \mathbb{Z}_{p_1} \times \dots \times \mathbb{Z}_{p_r}$.
There is a 1-1 map between $a \in \mathbb{Z}_{p_1 \cdot p_2 \dots p_r}$ and (a_1, \dots, a_r)

Composite Numbers

To distinguish prime numbers from composites, what properties does a composite have?

Chinese Remainder THM

Given distinct primes p_1, \dots, p_r , consider $(a_1, \dots, a_r) \in \mathbb{Z}_{p_1} \times \dots \times \mathbb{Z}_{p_r}$.
There is a 1-1 map between $a \in \mathbb{Z}_{p_1 \cdot p_2 \dots p_r}$ and (a_1, \dots, a_r)



- 1 Most fundamental THM in group/ring theory and number theory
- 2 Both directions are useful in algorithm design and analysis

Outline

1 Introduction

2 Prime Number Theorem

3 Basic Tools

4 Primality Testing

Basic Idea

Problem Description

Given q , test whether it is a prime or not.

Basic Idea

Problem Description

Given q , test whether it is a prime or not.

- 1 The issue of Fermat's little THM is that for some composites like $q = 561$, **most** $a \in \mathbb{Z}_q$ has $a^{q-1} \equiv 1$

Basic Idea

Problem Description

Given q , test whether it is a prime or not.

- 1 The issue of Fermat's little THM is that for some composites like $q = 561$, **most** $a \in Z_q$ has $a^{q-1} \equiv 1$
- 2 Let us try Chinese Remainder THM.

Basic Idea

Problem Description

Given q , test whether it is a prime or not.

- 1 The issue of Fermat's little THM is that for some composites like $q = 561$, **most** $a \in \mathbb{Z}_q$ has $a^{q-1} \equiv 1$
- 2 Let us try Chinese Remainder THM.

Key Observation

- 1 If q is a prime, $x^2 \equiv 1$ has at most 2 roots. What are they?
- 2 If q is not a prime, $x^2 \equiv 1$ has at least 4 roots. Why?

New Idea

Taking Square Root

If we can generate a **random** root of $x^2 \equiv 1 \pmod{q}$, it tells whether q is a prime or not — prime q has at most 2 roots and composite q has ≥ 4 roots

New Idea

Taking Square Root

If we can generate a **random** root of $x^2 \equiv 1 \pmod{q}$, it tells whether q is a prime or not — prime q has at most 2 roots and composite q has ≥ 4 roots

- 1 Unfortunately, finding square roots is computational hard — equivalent to factoring and break RSA system

New Idea

Taking Square Root

If we can generate a **random** root of $x^2 \equiv 1 \pmod{q}$, it tells whether q is a prime or not — prime q has at most 2 roots and composite q has ≥ 4 roots

- 1 Unfortunately, finding square roots is computational hard — equivalent to factoring and break RSA system
- 2 Wait ... we can use Fermat's little THM to help with it

New Idea

Taking Square Root

If we can generate a **random** root of $x^2 \equiv 1 \pmod{q}$, it tells whether q is a prime or not — prime q has at most 2 roots and composite q has ≥ 4 roots

- 1 Unfortunately, finding square roots is computational hard — equivalent to factoring and break RSA system
- 2 Wait ... we can use Fermat's little THM to help with it
- 3 Since $a^{q-1} \equiv 1$ for any a , $a^{\frac{q-1}{2}}$ might be a random root for a random a

New Idea

Taking Square Root

If we can generate a **random** root of $x^2 \equiv 1 \pmod{q}$, it tells whether q is a prime or not — prime q has at most 2 roots and composite q has ≥ 4 roots

- ① Unfortunately, finding square roots is computational hard — equivalent to factoring and break RSA system
- ② Wait ... we can use Fermat's little THM to help with it
- ③ Since $a^{q-1} \equiv 1$ for any a , $a^{\frac{q-1}{2}}$ might be a random root for a random a
 - ▶ If $a^{\frac{q-1}{2}} \neq \pm 1$, claim q is composite
 - ▶ If $a^{\frac{q-1}{2}} = -1$, give up this a

New Idea

Taking Square Root

If we can generate a **random** root of $x^2 \equiv 1 \pmod{q}$, it tells whether q is a prime or not — prime q has at most 2 roots and composite q has ≥ 4 roots

- ① Unfortunately, finding square roots is computational hard — equivalent to factoring and break RSA system
- ② Wait ... we can use Fermat's little THM to help with it
- ③ Since $a^{q-1} \equiv 1$ for any a , $a^{\frac{q-1}{2}}$ might be a random root for a random a
 - ▶ If $a^{\frac{q-1}{2}} \neq \pm 1$, claim q is composite
 - ▶ If $a^{\frac{q-1}{2}} = -1$, give up this a
 - ▶ Else if $a^{\frac{q-1}{2}} = 1$, keep trying $a^{\frac{q-1}{4}}$, $a^{\frac{q-1}{8}}$, ...

Formal Description

Pick many random $a \sim Z_q$ and call $\text{WITNESS}(a)$ for each one — output “composite” if any call does so

Algorithm Miller-Rabin Tester

function $\text{WITNESS}(a, q)$

Decompose $q - 1 = 2^t \cdot u$

$x_0 \equiv a^u$

for $i = 1, \dots, t$ **do**

$x_i \equiv x_{i-1}^2$

// x_{i-1} is a square root of x_i

if $x_i \equiv 1$ and $x_{i-1} \not\equiv \pm 1$ **then**

Return Composite

Return Composite if $x_t \neq 1$ o.w. return Prime

Formal Description

Pick many random $a \sim Z_q$ and call $\text{WITNESS}(a)$ for each one — output “composite” if any call does so

Algorithm Miller-Rabin Tester

function $\text{WITNESS}(a, q)$

Decompose $q - 1 = 2^t \cdot u$

$x_0 \equiv a^u$

for $i = 1, \dots, t$ **do**

$x_i \equiv x_{i-1}^2$

// x_{i-1} is a square root of x_i

if $x_i \equiv 1$ and $x_{i-1} \not\equiv \pm 1$ **then**

Return **Composite**

Return **Composite** if $x_t \neq 1$ o.w. return **Prime**

Running Time

$O(\log^3 q)$: $t = O(\log q)$ — but $x_i^2 \bmod q$ takes $O(\log^2 q)$ time

Analysis

Correctness

If q is composite, at least half a make $\text{WITNESS}(a, q) = \text{Composite}$.

The proof is similar to the proof $x^2 \equiv 1$ has at least 4 roots but uses group theory

Analysis

Correctness

If q is composite, at least half a make $\text{WITNESS}(a, q) = \text{Composite}$.

The proof is similar to the proof $x^2 \equiv 1$ has at least 4 roots but uses **group theory**

- 1 Suppose $q = p_1 \cdot p_2$ for two distinct primes

Analysis

Correctness

If q is composite, at least half a make $\text{WITNESS}(a, q) = \text{Composite}$.

The proof is similar to the proof $x^2 \equiv 1$ has at least 4 roots but uses **group theory**

- 1 Suppose $q = p_1 \cdot p_2$ for two distinct primes
- 2 Define (a, j) is acceptable if $a^{2^j \cdot u} \equiv -1$
- 3 Find the largest j with acceptable pairs (existence from $j = 0$ and $a = -1$) such that $x^{2^{j+1}u} \equiv 1$ for all x

Analysis

Correctness

If q is composite, at least half a make $\text{WITNESS}(a, q) = \text{Composite}$.

The proof is similar to the proof $x^2 \equiv 1$ has at least 4 roots but uses **group theory**

- 1 Suppose $q = p_1 \cdot p_2$ for two distinct primes
- 2 Define (a, j) is acceptable if $a^{2^j \cdot u} \equiv -1$
- 3 Find the largest j with acceptable pairs (existence from $j = 0$ and $a = -1$) such that $x^{2^{j+1}u} \equiv 1$ for all x
- 4 But $B = \{x : x^{2^j u} \equiv \pm 1\} \subsetneq Z_q^*$ is a proper subgroup
- 5 By group theory, only need to show $\exists b \notin B$, which is implied by Chinese Remainder THM

Analysis

Correctness

If q is composite, at least half a make $\text{WITNESS}(a, q) = \text{Composite}$.

The proof is similar to the proof $x^2 \equiv 1$ has at least 4 roots but uses **group theory**

- 1 Suppose $q = p_1 \cdot p_2$ for two distinct primes
- 2 Define (a, j) is acceptable if $a^{2^j \cdot u} \equiv -1$
- 3 Find the largest j with acceptable pairs (existence from $j = 0$ and $a = -1$) such that $x^{2^{j+1}u} \equiv 1$ for all x
- 4 But $B = \{x : x^{2^j u} \equiv \pm 1\} \subsetneq Z_q^*$ is a proper subgroup
- 5 By group theory, only need to show $\exists b \notin B$, which is implied by Chinese Remainder THM
- 6 When $i = j + 1$, half elements will be caught

Summary

- ① Primality tester faster than $O(n^3)$?
- ② How to generate primes efficiently?
- ③ Many other problem: factoring, ...

Questions?