

Instantons from Blow-up

Joonho Kim,^a Sung-Soo Kim,^b Ki-Hong Lee,^c Kimyeong Lee,^a and Jaewon Song^a

^a*School of Physics, Korea Institute for Advanced Study, Seoul 02455, Korea*

^b*School of Physics, University of Electronic Science and Technology of China,
No.4, Section 2, North Jianshe Road, Chengdu, Sichuan 610054, China*

^c*Department of Physics and Astronomy & Center for Theoretical Physics
Seoul National University, Seoul 08826, Korea*

E-mail: joonhokim@kias.re.kr, sungsoo.kim@uestc.edu.cn,
khlee11812@gmail.com, klee@kias.re.kr, jsong@kias.re.kr

ABSTRACT: The Nekrasov partition function for 4d $\mathcal{N} = 2$ or 5d $\mathcal{N} = 1$ gauge theory on the blow up of a point $\hat{\mathbb{C}}^2$ can be written in terms of the partition function on the flat space \mathbb{C}^2 . At the same time, the partition function on the blow up is identical to the partition function on a flat space for sufficiently large class of examples. This relation enables us to compute the instanton partition functions for 4d $\mathcal{N} = 2$ and 5d $\mathcal{N} = 1$ gauge theories for arbitrary gauge theory with large class of matter representations without knowing explicit construction of the instanton moduli space. Remarkably, the instanton partition function is completely determined by the perturbative part. We obtain the partition function for the previously unknown theories: exceptional gauge groups EFG with fundamental/spinor hypermultiplets and more. We also compute the case with $SU(6)$ with rank-3 antisymmetric tensor and compare with the topological vertex computation using the recently found 5-brane web configuration.

Contents

1	Introduction	1
2	Instanton Counting from Blow-up	3
2.1	Blowup equation	3
2.2	Recursion formula for 5d instanton partition function	8
2.3	Number of independent blowup equations	12
3	Examples	16
3.1	Theories with the known ADHM description	16
3.2	Theories with spinor hypermultiplets	19
3.3	Theories with an exceptional gauge group	24
3.4	Theories with a rank-3 antisymmetric hypermultiplet	25
4	Conclusion	31
A	One-instanton partition functions	32

1 Introduction

The Seiberg-Witten prepotential provides a complete description for the low energy dynamics of 4d $\mathcal{N} = 2$ or 5d $\mathcal{N} = 1$ gauge theory in its Coulomb branch [1, 2]. It is a function of the vacuum expectation value (VEV) of the scalar in the vector multiplet that parameterizes the Coulomb branch moduli space. Quantum correction to the prepotential is known to be one-loop exact, while there also exist non-perturbative corrections coming from Yang-Mills instantons.

An efficient way to compute the fully quantum corrected prepotential \mathcal{F} is to study the Nekrasov partition function \mathcal{Z} on Ω -deformed \mathbb{C}^2 or $\mathbb{C}^2 \times S^1$. It can be written as the product of the classical, one-loop, and instanton contributions,

$$\mathcal{Z}(\vec{a}, \vec{m}, \epsilon_1, \epsilon_2, \mathbf{q}) = \mathcal{Z}_{\text{class}}(\vec{a}, \epsilon_1, \epsilon_2, \mathbf{q}) \mathcal{Z}_{\text{1-loop}}(\vec{a}, \vec{m}, \epsilon_1, \epsilon_2) \mathcal{Z}_{\text{inst}}(\vec{a}, \vec{m}, \epsilon_1, \epsilon_2, \mathbf{q}), \quad (1.1)$$

where the instanton piece is the fugacity sum over all multi-instanton contributions:

$$\mathcal{Z}_{\text{inst}}(\vec{a}, \vec{m}, \epsilon_1, \epsilon_2, \mathbf{q}) = 1 + \sum_{k=1}^{\infty} \mathbf{q}^k \mathcal{Z}_k(\vec{a}, \vec{m}, \epsilon_1, \epsilon_2). \quad (1.2)$$

Once the Nekrasov partition function is known, one can extract the prepotential via taking $\epsilon_i \rightarrow 0$ limit as $\mathcal{F} = \lim_{\epsilon_i \rightarrow 0} \epsilon_1 \epsilon_2 \log \mathcal{Z}$ [3–6].

The instanton part of the partition function in the Ω -background can be computed once we know appropriate instanton moduli space. For the classical gauge group, the ADHM construction of the moduli space provides a direct way to compute the instanton partition function. The ADHM construction can be understood as the quantum mechanics described by the Dp - $D(p+4)$ system. The Higgs branch moduli space of the Dp system gives the desired moduli space. Matter fields can be also introduced by considering the world-volume theory on the D0-branes of the D0-D4-D8 system. By using the localization on the 1d system on the D0-branes or its dimensional reduction [7, 8], the partition function has been obtained for variety of cases: classical gauge groups [9–13], exceptional gauge groups [14–17]. 3d $N=4$ Coulomb branch realization of the instanton moduli space: [18] The precise choice of the Contour of the ADHM integral has been derived in [19–21] following the Jeffrey-Kirwan residue formula in 2d elliptic genus [22, 23].

But there is no ADHM type construction for the exceptional gauge groups or generic type of matter fields. String-theoretic picture implies that they require strong-coupling dynamics or non-Lagrangian field theories to realize instanton moduli space of exceptional group as a vacuum moduli space. There has been a few results regarding the exceptional instantons.

In this paper, we generalize the approach of Nakajima-Yoshioka (NY) [5, 24, 25] for the pure YM theory. In [16], the NY blow-up formula were used to compute the instanton partition function for exceptional gauge group and tested against the superconformal index of 4d SCFT where the Higgs branch is given by the instanton moduli space [26]. We propose a general blow-up formula for general gauge theory with arbitrary representations, under the condition that the matter representation is not too large.¹ This enables us to compute the instanton partition functions for numerous gauge theories that have not been known before, without relying on to the explicit construction of the moduli space.

The basic idea is as follows: Let us consider a one-point blow up $\hat{\mathbb{C}}^2$ of the flat space \mathbb{C}^2 . The *full* partition function on $\hat{\mathbb{C}}^2$ can be written in terms of the products of the *full* partition function of \mathbb{C}^2 . But at the same time, the partition function on the blowup is identical to that of the flat space since we can smoothly blow-down $\hat{\mathbb{C}}^2$ to \mathbb{C}^2 as long as the matter representation is not ‘too large’. This provides us a functional relation for the partition function, which turns out to be sufficient to determine the instanton partition function itself. Remarkably, this relation is completely determined by the perturbative part of the partition function. Therefore we arrive at a surprising conclusion: The *perturbative* physics determine the *non-perturbative* physics!

For example, we find the following universal expression for the 1-instanton partition function with arbitrary gauge group and matters:

$$Z_1 = \frac{(p_1 p_2)^{\frac{b}{2}} \prod_l (y_l^{\text{tw}})^{-\frac{I_2(\mathbf{R}_l)}{2}}}{(1-p_1)(1-p_2)} \sum_{\vec{k} \in \Delta_\ell} \frac{e^{\frac{\kappa_{\text{eff}}}{2}(\vec{a} \cdot \vec{k} - d_{ijk} a^i k^j k^k)} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_l^{\text{tw}})}{(1-p_1 p_2 e^{-\vec{a} \cdot \vec{k}})(1-e^{\vec{a} \cdot \vec{k}}) \prod_{\vec{\alpha} \cdot \vec{k} = -1} (1-e^{-\vec{a} \cdot \vec{\alpha}})} \quad (1.3)$$

where ...

¹This has to do with the 1-loop beta function coefficient as we will discuss in detail.

2 Instanton Counting from Blow-up

The essential idea of using the blow-up of \mathbb{C}^2 for instanton counting is that the gauge theory partition function for a 4d $\mathcal{N} = 2$ (or 5d $\mathcal{N} = 1$) theory on the blow-up of a point $\hat{\mathbb{C}}^2$ (or $S^1 \times \hat{\mathbb{C}}^2$) can be written in two different ways. This will allow us to write a recursion relation for the instanton partition function that can be solved rather easily [5, 16, 24, 25].

2.1 Blowup equation

Localization on the blow-up $\hat{\mathbb{C}}^2$ One of the expressions for the partition function $\hat{\mathcal{Z}}$ on the blow-up $\hat{\mathbb{C}}^2$ comes from the Coulomb branch localization, which results that $\hat{\mathcal{Z}}$ can be obtained by patching together the flat-space partition function \mathcal{Z} [27].

The blow-up $\hat{\mathbb{C}}^2$ is constructed from \mathbb{C}^2 by replacing the origin with a compact 2-cycle. In particular, the geometry is identical to the total space of the line bundle of degree (-1) over \mathbb{P}^1 . One can parametrize $\mathcal{O}(-1) \rightarrow \mathbb{P}^1$ using the homogeneous coordinates (z_0, z_1, z_2) , satisfying the projective condition $(z_0, z_1, z_2) \sim (\lambda^{-1}z_0, \lambda^1z_1, \lambda^1z_2)$ for any $\lambda \in \mathbb{C}^*$, where the two-cycle $\mathbb{P}^1 \subset \hat{\mathbb{C}}^2$ corresponds to the locus $z_0 = 0$. We are interested in the $U(1)^2$ equivariant partition function, with the $U(1)^2$ action V rotating the complex coordinates (z_0, z_1, z_2) as follows:

$$(z_0, z_1, z_2) \mapsto (z_0, e^{\epsilon_1} z_1, e^{\epsilon_2} z_2). \quad (2.1)$$

Instantons are located at two fixed points of the $U(1)^2$ action, i.e., the north/south poles of the \mathbb{P}^1 , whose coordinates are $(z_0, z_1, z_2) = (0, 1, 0)$ and $(0, 0, 1)$. Around these fixed points, the local weights under the $U(1)^2$ action V are:

$$(z_0 z_1, z_2/z_1) \mapsto (e^{\epsilon_1} z_0 z_1, e^{\epsilon_2 - \epsilon_1} z_2/z_1) \quad (\text{near the north pole}) \quad (2.2)$$

$$(z_0 z_2, z_1/z_2) \mapsto (e^{\epsilon_2} z_0 z_2, e^{\epsilon_1 - \epsilon_2} z_1/z_2) \quad (\text{near the south pole}) \quad (2.3)$$

The full partition function $\hat{\mathcal{Z}}$ on $\hat{\mathbb{C}}^2$, which includes both the perturbative and instanton contributions, can be obtained by performing the localization on the Coulomb branch. On the Coulomb branch, the gauge group is generically broken to $U(1)^r$ where r is the rank of the gauge group. The $U(1)^r$ equivariant parameters \vec{a} naturally appear in the partition function. One needs to sum over all distinct field configurations with zero-sized instantons located at the north and south poles. All the inequivalent configurations are labeled by the r -dimensional vector \vec{k} of the first Chern numbers, corresponding to different flux configurations on the two-cycle \mathbb{P}^1 . When the gauge group has $U(1)$ factor, we can turn on the external flux that can be supported on the \mathbb{P}^1 . We assume there is no such a factor in the gauge group. Summing up, $\hat{\mathcal{Z}}$ can be expressed in terms of the partition function \mathcal{Z} on \mathbb{C}^2 as [27–31]

$$\hat{\mathcal{Z}}(\vec{a}, \epsilon_1, \epsilon_2, q, \vec{m}) = \sum_{\vec{k} \in \Lambda} \mathcal{Z}^{(N)}(\vec{k}) \mathcal{Z}^{(S)}(\vec{k}), \quad (2.4)$$

where the flux sum is taken over the co-root lattice Λ of the gauge algebra. Each factor represents the partition function localized at the $U(1)^2$ fixed points (north/south-poles of the $\mathbb{P}^1 \subset \hat{\mathbb{C}}^2$ given as

$$\begin{aligned}\mathcal{Z}^{(N)}(\vec{k}) &\equiv \mathcal{Z}(\vec{a} + \vec{k}\epsilon_1, \epsilon_1, \epsilon_2 - \epsilon_1, q, \vec{m} + \frac{1}{2}\epsilon_1) , \\ \mathcal{Z}^{(S)}(\vec{k}) &\equiv \mathcal{Z}(\vec{a} + \vec{k}\epsilon_2, \epsilon_1 - \epsilon_2, \epsilon_2, q, \vec{m} + \frac{1}{2}\epsilon_2) .\end{aligned}\tag{2.5}$$

In addition to the Coulomb branch parametres, the partition function depends on the Omega deformation parameters ϵ_1, ϵ_2 and also mass parameters \vec{m} . The parameter q is the instanton parameter. For a 4d theory, it is given as $q = e^{2\pi i\tau} = \Lambda^{b_0}$ where τ is the complexified gauge coupling and Λ being the dynamical scale of the gauge theory. The exponent b_0 is the 1-loop beta function coefficient. For a 5d theory, it is also given by a exponentiated gauge coupling as $q = e^{-\frac{1}{g^2}} \equiv e^{-m_0}$. Notice that the Coulomb parameter \vec{a} gets an appropriate shift at each fixed point p , induced by the non-trivial magnetic flux \vec{k} on the blown-up \mathbb{P}^1 , with the proportionality constant $H|_p$. The values of the moment map H for the $U(1)^2$ action V , *i.e.*, $dH = \iota_V \omega$, at the north and south poles are given as

$$H|_{\text{NP}} = \epsilon_1 \text{ and } H|_{\text{SP}} = \epsilon_2.\tag{2.6}$$

The mass parameters also get shifted since the hypermultiplet mass is twisted by $SU(2)_R$, which makes the combination $m + \frac{\epsilon_1 + \epsilon_2}{2}$ invariant at the fixed points.²

Partition function on $\hat{\mathbb{C}}^2$ vs \mathbb{C}^2 An alternative fact for the partition function $\hat{\mathcal{Z}}$ on the blow-up $\hat{\mathbb{C}}^2$ is that $\hat{\mathcal{Z}}$ is actually identical to the flat-space partition function \mathcal{Z} [5, 24, 25]. The blow-up $\hat{\mathbb{C}}^2$ is identical to \mathbb{C}^2 except for the origin, which is replaced by the blown-up sphere \mathbb{P}^1 . Since the Nekrasov partition function gets contributions only from the small instantons localized at the fixed points of the $U(1)^2$ equivariant action V , the size of the divisor should not affect the partition function as we smoothly shrink it. So we expect that $\hat{\mathcal{Z}} = \mathcal{Z}$. This implies the following relation: [5, 16, 24, 25, 32, 33]

$$\mathcal{Z} = \hat{\mathcal{Z}} = \sum_{\vec{k} \in \Lambda} \mathcal{Z}^{(N)}(\vec{k}) \mathcal{Z}^{(S)}(\vec{k})\tag{2.7}$$

This blow-up identity can be thought of as a special case of more generalized to orbifold partition functions [31, 34, 35]. For example, the Nekrasov partition function on the orbifold $\mathbb{C}^2/\mathbb{Z}_2$ can be computed in two different ways, one is via formula analogous to (2.4) by combining the contributions from two fixed points of the blown-up geometry $\mathcal{O}(-2) \rightarrow \mathbb{P}^1$. The other way is to compute the partition function at the orbifold point using the ADHM construction for the orbifolds. The Nekrasov partition function still remains the same as we

²One can instead use the shifted mass to simplify the formula involving mass. We use unshifted mass to match with the existing formulae in the literature.

blow up or down the singular point.³ The only difference in our case is that we blow-up or down a non-singular point instead of a singular point.

Correlation functions in 4d In fact, (2.7) is not enough to fix the partition function completely, since there are 3 unknown functions and only one relation. It turns out the necessary additional relations can be found from the insertion of non-trivial \mathcal{Q} -closed operators [5, 25] associated to the two-cycle on the blow-up.

In the 4d Donaldson-twisted theory, the \mathcal{Q} -invariant observable \mathcal{O}_2 associated to a two-cycle can be constructed by applying the topological descent procedure twice to the Casimir invariant $\mathcal{O}_0 = \text{Tr}(\Phi^2)$ as [36]

$$\begin{aligned} 0 &= \{\mathcal{Q}, \mathcal{O}_0\}, \quad d\mathcal{O}_0 = \{\mathcal{Q}, \mathcal{O}_1\}, \quad d\mathcal{O}_1 = \{\mathcal{Q}, \mathcal{O}_2\}, \\ d\mathcal{O}_2 &= \{\mathcal{Q}, \mathcal{O}_3\}, \quad d\mathcal{O}_3 = \{\mathcal{Q}, \mathcal{O}_4\}, \quad d\mathcal{O}_4 = 0. \end{aligned} \quad (2.8)$$

In our case, we consider a $U(1)^2$ -equivariant version of the topological descent procedure, that is to choose $\mathcal{Q} = Q_{\text{top}} + H$ to obtain the operator associated to the two-cycle. In terms of the component fields, it can be written as [37]

$$\mathcal{O}_{\mathbb{P}^1} = \int_{\mathbb{P}^1} \mathcal{O}_2 = \int_{M_4} \left\{ \omega \wedge \text{Tr} \left(\Phi F + \frac{1}{2} \psi \wedge \psi \right) + H \text{Tr} (F \wedge F) \right\}. \quad (2.9)$$

Here ω and H are the Kähler two-form on the \mathbb{P}^1 and the moment map $\iota_V \omega = dH$, respectively. The first part of (2.9) without H is the non-equivariant version of the topological operator associated to two-cycle. It is convenient to study the generating function $\langle e^{t\mathcal{O}_{\mathbb{P}^1}} \rangle$ of the correlators $\langle \mathcal{O}_{\mathbb{P}^1} \dots \mathcal{O}_{\mathbb{P}^1} \rangle$. This cause a shift of instanton parameter by $q \rightarrow q \exp(tH)$ at the fixed points of the blow-up $\hat{\mathbb{C}}^2$ [5, 24, 25]. The expectation value of the generating function can be written as

$$\hat{\mathcal{Z}}^t \equiv \langle e^{t\mathcal{O}_{\mathbb{P}^1}} \rangle = \sum_{\vec{k} \in \Lambda} \mathcal{Z}^{(N),t}(\vec{k}) \cdot \mathcal{Z}^{(S),t}(\vec{k}), \quad (2.10)$$

where

$$\begin{aligned} \mathcal{Z}^{(N),t}(\vec{k}) &\equiv \mathcal{Z}(\vec{a} + \vec{k}\epsilon_1, \epsilon_1, \epsilon_2 - \epsilon_1, q \exp(t\epsilon_1), \vec{m} + \frac{1}{2}\epsilon_1), \\ \mathcal{Z}^{(S),t}(\vec{k}) &\equiv \mathcal{Z}(\vec{a} + \vec{k}\epsilon_2, \epsilon_1 - \epsilon_2, \epsilon_2, q \exp(t\epsilon_2), \vec{m} + \frac{1}{2}\epsilon_2). \end{aligned} \quad (2.11)$$

Now, as we shrink the two-cycle \mathbb{P}^1 to recover the flat \mathbb{C}^2 , the effect of inserting $(\mathcal{O}_{\mathbb{P}^1})^d$ turns out to give a vanishing contribution for small d due to the selection rule. We recall that the instanton breaks the $U(1)_R$ symmetry to the discrete subgroup \mathbb{Z}_{2b_0} with $b_0 = 2h^\vee - \sum_i I_2(\mathbf{R}_i)$ where the sum is over all hypermultiplets and h^\vee is the dual Coxeter number of the gauge group and \mathbf{R}_i denotes the representation of the i -th hypermultiplet and $I_2(\mathbf{R})$

³This simple picture does not necessarily hold when there are too many hypermultiplets, due to some subtle scheme dependence related to the wall-crossing [35].

being the quadratic Dynkin index.⁴ The first term of the operator $\mathcal{O}_{\mathbb{P}^1}$ (the two-form piece) carries R -charge $+2$, which is the familiar non-equivariant version. This discrete R -charge is sometimes called as a ghost number. The correlation functions vanish unless the R -charges add up to zero, modulo $2b_0 = 4h^\vee - 2\sum_i I_2(\mathbf{R}_i)$. Therefore, expanding (2.10) in powers of t , we find

$$\langle e^{t\mathcal{O}_{\mathbb{P}^1}} \rangle = \mathcal{Z} + \mathcal{O}(t^{2h^\vee - \sum_i I_2(\mathbf{R}_i)}) . \quad (2.12)$$

This is our blowup equation. Each term at order t^n carries pieces with R -charge between 0 and $2n$. When $n < 2b_0$, the only possible non-trivial contribution comes from the $R = 0$ piece $\int HF \wedge F$ at zero instanton sector. For n -instanton sector, one should have $R = 2b_0n$, which is the condition to absorb the fermionic zero modes. This piece vanishes for zero instanton sector (at north/south poles). For $n \geq 2b_0$, we always have a term that absorb all the fermionic zero modes (or the term that has $R = 0 \bmod (2b_0n)$ so they do not vanish).

We see that as long as the hypermultiplet representation is not too large, *i.e.*, when $b_0 = 2h^\vee - \sum_i I_2(\mathbf{R}_i) > 2$, this allows us to write 3 independent relations for the 3 unknown variables. One can expand $\langle e^{t\mathcal{O}_{\mathbb{P}^1}} \rangle$ until $\mathcal{O}(t^2)$ order and recursively solve for \mathcal{Z} at each instanton number. So the instanton part of the partition function will be completely determined from the perturbative partition function. An explicit form of the recursion relation will be studied in Section 2.2.

Correlation functions in 5d We now turn to 5d $\mathcal{N} = 1$ gauge theory wrapped on S^1 . The Casimir invariant $\text{Tr}(\Phi^2)$ and its descendants are no longer considered as well-defined observables. Instead, there are two types of \mathcal{Q} -invariant observables [38]. The first type of observables are constructed from the 5d Wilson loop on the S^1 by applying the descent procedure. The second type of observables introduce the 3d (Kähler) Chern-Simons term, which can be written as [38, 39]

$$\begin{aligned} \mathcal{O}_{\mathbb{P}^1} = \exp \left[\int_{S^1 \times M_4} \left(\omega \wedge \text{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) \right. \right. \\ \left. \left. + \omega \wedge \left(\phi F + \frac{1}{2} \psi \wedge \psi \right) \wedge dt + H \text{Tr} \left(F \wedge F \right) \wedge dt \right) \right] . \end{aligned} \quad (2.13)$$

It can be viewed as the natural S^1 uplift of (2.9) via exponentiation. The correlation function is now given by

$$\hat{\mathcal{Z}}^d \equiv \langle (\mathcal{O}_{\mathbb{P}^1})^d \rangle = \sum_{\vec{k} \in \Lambda} \mathcal{Z}^{(N),d}(\vec{k}) \cdot \mathcal{Z}^{(S),d}(\vec{k}) , \quad (2.14)$$

where

$$\begin{aligned} \mathcal{Z}^{(N),d}(\vec{k}) &\equiv \mathcal{Z}(\vec{a} + \vec{k}\epsilon_1, \epsilon_1, \epsilon_2 - \epsilon_1, q \exp((d - \frac{b}{2})\epsilon_1), \vec{m} + \frac{1}{2}\epsilon_1) , \\ \mathcal{Z}^{(S),d}(\vec{k}) &\equiv \mathcal{Z}(\vec{a} + \vec{k}\epsilon_2, \epsilon_1 - \epsilon_2, \epsilon_2, q \exp((d - \frac{b}{2})\epsilon_2), \vec{m} + \frac{1}{2}\epsilon_2) . \end{aligned} \quad (2.15)$$

⁴We normalize it so that $I_2(\mathbf{F}) = 1$ for the fundamental representation \mathbf{F} .

Here the quantity b is given as

$$b \equiv h^\vee - \frac{1}{2} \sum_i I_2(\mathbf{R}_i) - \kappa_{\text{eff}} , \quad \kappa_{\text{eff}} = \kappa - \frac{1}{2} \sum_i I_3(\mathbf{R}_i) , \quad (2.16)$$

where $I_2(\mathbf{R})$ and $I_3(\mathbf{R})$ are quadratic and cubic Casimir invariants respectively. Notice that d has to be an integer to be gauge-invariant.

The reason that the instanton parameter is further shifted by $\exp(\frac{b}{2}H|_p)$ is because the instanton mass parameter is twisted by $SU(2)_R$ as in the case of the hypermultiplet mass. Moreover, notice that the instanton soliton carries the $SU(2)_R$ twisted effective mass, given by $m_{\text{inst}} \equiv m_{0,\text{eff}} - \kappa_{\text{eff}} \epsilon_+$. In addition, the effective Chern-Simons coupling κ_{eff} induces an electric charge to the instanton, contributing to its ground state energy as $E_0 = m_{\text{inst}} - \vec{a} \cdot \vec{\Pi}$, where $\vec{\Pi}$ is the $U(1)^r \subset G$ electric charge.⁵ To keep it invariant the effective instanton mass m_{inst} at a fixed point p of the blow-up $\hat{\mathbb{C}}^2$, we require the shifted gauge coupling $m_0|_p$ to be

$$m_0|_p = m_0 + \frac{b}{2}H|_p \quad \text{with} \quad b \equiv h^\vee - \sum_i \frac{I_2(\mathbf{R}_i)}{2} - \kappa_{\text{eff}}. \quad (2.17)$$

For the case of 5d pure $\mathcal{N} = 1$ SYM, the correlation function turns out to be

$$\boxed{\langle (\mathcal{O}_{\mathbb{P}^1})^d \rangle = \mathcal{Z} \quad \text{for} \quad 0 \leq d \leq d_{\text{max}}} , \quad (2.18)$$

where $d_{\text{max}} = h^\vee$.⁶ We call (2.18) as the blowup equation. The value of d_{max} depends on the matter content and gauge group. For $d_{\text{max}} \geq 2$, there are sufficient number of algebraic relations to determine the instanton partition function recursively in increasing order of instantons. This fact was utilized in [16] to compute instanton partition function for the gauge theories with exceptional gauge groups, for which the ADHM construction of instanton moduli space is unknown.

In this paper, we aim at developing the relation (2.18) for various 5d $\mathcal{N} = 1$ gauge theories with hypermultiplets in various representations, and compute the instanton partition function. We will identify a certain bound on d in Section 2.3 as the *necessary* condition for (2.18) for a large number of theories. We conjecture that the bound on d we obtain is actually sufficient to obtain the blowup equation (2.18). While we do not attempt to prove this sufficiency, we compute n -instanton partition function Z_n , based on the recursion formula that will be derived shortly from (2.18), and confirm the agreement with the known result obtained from an alternative method.

We find a universal expression for the bound on d when the gauge group is not of SU or Sp type:

$$d_{\text{max}} = h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) \quad \text{for } G \neq SU(N) \text{ or } Sp(N) \quad (2.19)$$

⁵This agrees with the supersymmetric Casimir energy of the ADHM quantum mechanics.

⁶This was shown in [25] for the case of $G = SU(N)$.

This is essentially identical condition as in $4d$ $\mathcal{N} = 2$ gauge theory. But in $5d$, some new effects come into play. For the $SU(N)$ case, we can have a Chern-Simons term generated at 1-loop, which alters the bound on d . When there is neither bare nor effective Chern-Simons coupling, the same bound holds for the $SU(N)$ case as well. The detailed condition will be given in section 2.3. For the case of $Sp(N)$, one can turn on the discrete θ -parameter and it turns out the bound on d depends on this parameter.

2.2 Recursion formula for 5d instanton partition function

The blowup equation (2.18) can be translated to a recursion formula on the (5d) n -instanton contribution Z_n to the full partition function \mathcal{Z} . To derive this, we decompose the partition function \mathcal{Z} in terms of the classical, one-loop, and instanton pieces:

$$\mathcal{Z}(\vec{a}, \epsilon_1, \epsilon_2, q, \vec{m}) = Z_{\text{class}}(\vec{a}, \epsilon_1, \epsilon_2, q, \vec{m}) \cdot Z_{1\text{-loop}}(\vec{a}, \epsilon_1, \epsilon_2, \vec{m}) \cdot Z_{\text{inst}}(\vec{a}, \epsilon_1, \epsilon_2, q, \vec{m}) \quad (2.20)$$

where Z_{inst} can be further expanded in terms of the instanton fugacity q as⁷

$$Z_{\text{inst}}(\vec{a}, \epsilon_1, \epsilon_2, q, \vec{m}) = \sum_{n \geq 0} q^n Z_n(\vec{a}, \epsilon_1, \epsilon_2, \vec{m}) . \quad (2.21)$$

Then the blowup equation (2.18) can be written as

$$\begin{aligned} Z_{\text{inst}} &= \sum_{\vec{k}} \left[\frac{Z_{\text{class}}^{(N),d}(\vec{k}) Z_{\text{class}}^{(S),d}(\vec{k})}{Z_{\text{class}}} \frac{Z_{1\text{-loop}}^{(N),d}(\vec{k}) Z_{1\text{-loop}}^{(S),d}(\vec{k})}{Z_{1\text{-loop}}} \right] Z_{\text{inst}}^{(N),d}(\vec{k}) Z_{\text{inst}}^{(S),d}(\vec{k}) \\ &\equiv \sum_{\vec{k}} f_d(\vec{k}) Z_{\text{inst}}^{(N),d}(\vec{k}) Z_{\text{inst}}^{(S),d}(\vec{k}) , \end{aligned} \quad (2.22)$$

where the superscript $(N/S), d$ implies the appropriate shift of parameters, specified in (2.11). The function $f_d(\vec{k})$ is determined only via perturbative part of the partition function.

We recall the known expressions for the classical and 1-loop partition function (in 5d)

⁷Sometimes the instanton partition function is expanded in powers of the shifted instanton mass $q \exp(-b \frac{\epsilon_1 + \epsilon_2}{2})$ instead of q [16, 25, 28]. We expand it with the true instanton fugacity, which makes the symmetry property $\epsilon_{1,2} \rightarrow -\epsilon_{1,2}$ of Z_n manifest. This is the one that we obtain using the ADHM quantum mechanics.

[3, 40, 41]:⁸

$$Z_{\text{class}} = \exp \left[\frac{1}{\epsilon_1 \epsilon_2} \left(\frac{1}{2} m_0 h_{ij} a_i a_j + \frac{\kappa}{6} d_{ijk} a^i a^j a^k \right) \right], \quad (2.23)$$

$$Z_{1\text{-loop}}^{\text{vec}} = \exp \left[\frac{1}{\epsilon_1 \epsilon_2} \sum_{\vec{\alpha} \in \Delta} \left(\frac{(\vec{a} \cdot \vec{\alpha} + \epsilon_+)^3}{12} - \frac{\epsilon_1^2 + \epsilon_2^2 + 24}{48} (\vec{a} \cdot \vec{\alpha} + \epsilon_+) + 1 \right) \right] \times \text{PE} \left[- \frac{p_1 p_2}{(1-p_1)(1-p_2)} \sum_{\vec{\alpha} \in \Delta} e^{-\vec{a} \cdot \vec{\alpha}} \right] \quad \text{for the vector multiplet} \quad (2.24)$$

$$Z_{1\text{-loop}}^{\text{hyp}, l} = \exp \left[- \frac{1}{\epsilon_1 \epsilon_2} \sum_{\vec{\omega} \in \mathbf{R}_l} \left(\frac{(\vec{a} \cdot \vec{\omega} + m_l)^3}{12} x - \frac{\epsilon_1^2 + \epsilon_2^2 + 24}{48} (\vec{a} \cdot \vec{\omega} + m_l) + 1 \right) \right] \times \text{PE} \left[+ \frac{(p_1 p_2)^{\frac{1}{2}} \cdot y_l}{(1-p_1)(1-p_2)} \sum_{\vec{\omega} \in \mathbf{R}_l} e^{-\vec{a} \cdot \vec{\omega}} \right] \quad \text{for the } \ell\text{'th hypermultiplet} \quad (2.25)$$

where $p_1 \equiv e^{-\epsilon_1}$, $p_2 \equiv e^{-\epsilon_2}$, $y_l \equiv e^{-m_l}$, $q \equiv e^{-m_0}$.⁹ Also Δ is the set of all roots and \mathbf{R}_ℓ denotes the set of all weight vectors in representation \mathbf{R}_ℓ . Here, PE represents the Plethystic exponential

$$\text{PE} [f(\vec{a}, \epsilon_1, \epsilon_2, m_0, \vec{m})] \equiv \exp \left(\sum_{n=1}^{\infty} \frac{1}{n} f(n\vec{a}, n\epsilon_1, n\epsilon_2, nm_0, n\vec{m}) \right). \quad (2.26)$$

We also set the radius of S^1 as $\beta = 1$. Also, the symbols h_{ij} and d_{ijk} are defined as

$$h_{ij} = \text{Tr}(T_i T_j), \quad d_{ijk} = \frac{1}{2} \text{Tr} T_i \{T_j, T_k\}, \quad (2.27)$$

where T_i are the generators of the gauge algebra. They satisfy the relations

$$\begin{aligned} \sum_{\vec{\omega} \in \mathbf{R}} (\vec{a} \cdot \vec{\omega})(\vec{b} \cdot \vec{\omega})(\vec{c} \cdot \vec{\omega}) &= I_3(\mathbf{R}) d_{ijk} a^i b^j c^k, \\ \sum_{\vec{\omega} \in \mathbf{R}} (\vec{a} \cdot \vec{\omega})(\vec{b} \cdot \vec{\omega}) &= I_2(\mathbf{R}) h_{ij} a^i b^j, \\ \sum_{\vec{\omega} \in \mathbf{R}} (\vec{a} \cdot \vec{\omega}) &= 0, \end{aligned} \quad (2.28)$$

where $I_2(\mathbf{R})$ and $I_3(\mathbf{R})$ are the quadratic and cubic Dynkin indices.

⁸The ‘Casimir part’ of the 1-loop free energy (the ones inside the exp) depends on the choice of ...

⁹We assume a particular Weyl chamber in the Coulomb branch, i.e., $0 < a_i < \epsilon_+ < m$ for all $i \in \{1, \dots, r\}$.

Substituting them to (2.22), we obtain the ratio of three different Z 's given as

$$f_d(\vec{k})_{\text{class}} = q^{\frac{\vec{k} \cdot \vec{k}}{2}} (p_1 p_2)^{(\frac{b}{2}-d)(\frac{\vec{k} \cdot \vec{k}}{2}) + \frac{\kappa}{6} d_{ijk} k^i k^j k^k} \times e^{-(\frac{b}{2}-d)(\vec{a} \cdot \vec{k})} e^{-\frac{\kappa}{2} d_{ijk} a^i k^j k^k}, \quad (2.29)$$

$$f_d(\vec{k})_{1\text{-loop}}^{\text{vec}} = e^{\frac{h^\vee}{2}(\vec{a} \cdot \vec{k})} \prod_{\alpha \in \Delta} \mathcal{L}_{\vec{k} \cdot \vec{\alpha}}(\vec{a} \cdot \vec{\alpha}, \epsilon_1, \epsilon_2)^{-1}, \quad (2.30)$$

$$f_d(\vec{k})_{1\text{-loop}}^{\text{hyp}} = e^{-\frac{I_2(\mathbf{R}_l)}{4}(\vec{a} \cdot \vec{k}) + \frac{I_3(\mathbf{R}_l)}{4} d_{ijk} a^i k^j k^k} (p_1 p_2)^{\frac{I_2(\mathbf{R}_l)}{8}(\vec{k} \cdot \vec{k}) - \frac{I_3(\mathbf{R}_l)}{12} d_{ijk} k^i k^j k^k} \\ \times y_\ell^{-\frac{I_2(\mathbf{R}_l)}{4}(\vec{k} \cdot \vec{k})} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_{\text{tw},l}, \epsilon_1, \epsilon_2), \quad (2.31)$$

where we split the $f_d(\vec{k})$ into classical and 1-loop pieces for vector and hypermultiplet. Here we used $I_2(\mathbf{adj}) = 2h^\vee$, $I_3(\mathbf{adj}) = 0$, and also the fact h_{ij} and d_{ijk} are totally symmetric. We also define $m_{\text{tw}} \equiv m + \epsilon_+$. The function $\mathcal{L}_k(x, \epsilon_1, \epsilon_2)$ is introduced to denote concisely the combination of the PE parts:

$$\mathcal{L}_k(x, \epsilon_1, \epsilon_2) \equiv \text{PE} \left[e^{-x} \left(\frac{p_1^k p_2}{(1-p_1)(1-\frac{p_2}{p_1})} + \frac{p_1 p_2^k}{(1-\frac{p_1}{p_2})(1-p_2)} - \frac{p_1 p_2}{(1-p_1)(1-p_2)} \right) \right] \quad (2.32)$$

One can easily check that the expression inside the PE vanishes at $k = 0, 1$. After some work, it is not difficult to find that

$$\mathcal{L}_k(x, \epsilon_1, \epsilon_2) = \begin{cases} \prod_{m+n \leq k-2} (1 - p_1^{m+1} p_2^{n+1} e^{-x}) & \text{for } k \geq +2 \\ \prod_{m+n \leq -k-1} (1 - p_1^{-m} p_2^{-n} e^{-x}) & \text{for } k \leq -1 \\ 1 & \text{for } k = 0, 1. \end{cases} \quad (2.33)$$

Combining them all, the resursion formula on the n -instanton piece Z_n can be written as

$$Z_n = \sum_{\frac{1}{2}\vec{k} \cdot \vec{k} + \ell + m = n} \left((p_1 p_2)^{(\frac{h^\vee}{2} - \frac{\kappa_{\text{eff}}}{2} - d)(\frac{\vec{k} \cdot \vec{k}}{2}) + \frac{\kappa_{\text{eff}}}{6} d_{ijk} k^i k^j k^k} e^{(d + \frac{\kappa_{\text{eff}}}{2})(\vec{a} \cdot \vec{k})} e^{-\frac{\kappa_{\text{eff}}}{2} d_{ijk} a^i k^j k^k} \right. \\ \left. \times \frac{\prod_l y_{\text{tw},l}^{-I_2(\mathbf{R}_l)(\frac{\vec{k} \cdot \vec{k}}{4})} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_{\text{tw},l}, \epsilon_1, \epsilon_2)}{\prod_{\alpha \in \Delta} \mathcal{L}_{\vec{k} \cdot \vec{\alpha}}(\vec{a} \cdot \vec{\alpha}, \epsilon_1, \epsilon_2)} \cdot p_1^{(\frac{b}{2}-d)\ell} p_2^{(\frac{b}{2}-d)m} Z_\ell^{(N)}(\vec{k}) Z_m^{(S)}(\vec{k}) \right), \quad (2.34)$$

where $y_{\text{tw},l} \equiv e^{-m_{\text{tw},l}} = y_l / \sqrt{p_1 p_2}$. This is a generalization of the recursion formula found for the pure SYM case [25, 28]. Here l runs over all hypermultiplets in the theory.

Solving the recursion formulae The recursion relation (2.34) can be rewritten as

$$Z_n = p_1^{n(\frac{b}{2}-d)} Z_n^{(N)} + p_2^{n(\frac{b}{2}-d)} Z_n^{(S)} + I_n^{(d)} \quad \text{with an allowed range of } d, \quad (2.35)$$

where $I_n^{(d)}$ is defined as

$$I_n^{(d)} = \sum_{\substack{\frac{1}{2}\vec{k}\cdot\vec{k}+\ell+m=n \\ \ell, m \neq n}} \left((p_1 p_2)^{(\frac{b}{2}-d)} e^{(d+\frac{\kappa_{\text{eff}}}{2})(\vec{a}\cdot\vec{k})} e^{-\frac{\kappa_{\text{eff}}}{2} d_{ijk} a^i k^j k^k} \right. \\ \left. \times \frac{\prod_l y_{\text{tw},l}^{-I_2(\mathbf{R}_l)(\vec{k}\cdot\vec{k}/4)} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k},\vec{\omega}}(\vec{a}\cdot\vec{\omega} + m_{\text{tw},l}, \epsilon_1, \epsilon_2)}{\prod_{\alpha \in \Delta} \mathcal{L}_{\vec{k},\vec{\alpha}}(\vec{a}\cdot\vec{\alpha}, \epsilon_1, \epsilon_2)} \cdot p_1^{(\frac{b}{2}-d)\ell} p_2^{(\frac{b}{2}-d)m} Z_\ell^{(N)}(\vec{k}) Z_m^{(S)}(\vec{k}) \right). \quad (2.36)$$

Notice that we have a set of equations labelled by the parameter d . If the blowup equation holds for at least 3 values of d , we can solve it for Z_n . The n -instanton partition function Z_n is given as the solution to the three linear equations (2.35) with consecutive integers $\{d_0, d_0 + 1, d_0 + 2\}$.

$$Z_n(\vec{a}, \epsilon_1, \epsilon_2, \vec{m}) = \frac{p_1^n p_2^n I_n^{(d_0+2)} - (p_1^n + p_2^n) I_n^{(d_0+1)} + I_n^{(d_0)}}{(1 - p_1^n)(1 - p_2^n)} \quad (2.37)$$

Since $I_n^{(d)}$ only involves low-order instanton corrections, the n -instanton partition function Z_n can be constructed from $Z_{m < n}$, allowing us to obtain the full non-perturbative part Z_{inst} in a recursive manner starting from $Z_0 = 1$.

Therefore we arrive at a remarkable conclusion. The non-perturbative partition function Z_{inst} is completely fixed by the perturbative partition function! Note that we do not obtain this conclusion by requiring the perturbative series to be well-defined, as is often done in the resurgence analysis. Instead, we demand consistency upon smooth deformation of the spacetime \mathbb{C}^2 or $\mathbb{C}^2 \times S^1$. Such consistency condition requires non-perturbative parts to exist and even enough to fix the instanton partition function (at least for the large number of examples).

Now, let us write the solution for 1-instanton explicitly. At one instanton level, the formula (2.36) can be written as

$$I_1^{(d)} = \sum_{\vec{k} \in \Delta_\ell} \left((p_1 p_2)^{(\frac{b}{2}-d)} e^{(d+\frac{\kappa_{\text{eff}}}{2})(\vec{a}\cdot\vec{k})} e^{-\frac{\kappa_{\text{eff}}}{2} d_{ijk} a^i k^j k^k} \right. \\ \left. \times \frac{\prod_l y_{\text{tw},l}^{-I_2(\mathbf{R}_l)/2} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k},\vec{\omega}}(\vec{a}\cdot\vec{\omega} + m_{\text{tw},l}, \epsilon_1, \epsilon_2)}{(1 - p_1 p_2 e^{-\vec{a}\cdot\vec{k}})(1 - p_1^{-1} e^{\vec{a}\cdot\vec{k}})(1 - p_2^{-1} e^{\vec{a}\cdot\vec{k}})(1 - e^{\vec{a}\cdot\vec{k}}) \prod_{\vec{\alpha}\cdot\vec{k}=-1} (1 - e^{-\vec{a}\cdot\vec{\alpha}})} \right), \quad (2.38)$$

where Δ_ℓ is the set of long roots ($\vec{k} \cdot \vec{k} = 2$) and we used $Z_0 = 1$. It turns out to be more convenient to express Z_1 by decomposing $I_1^{(d)}$ into the flux sum, *i.e.*, $I_1^{(d)} \equiv \sum_{\vec{k} \in \Delta_\ell} i_1^{(d)}(\vec{k})$, where

$$i_1^{(d)}(\vec{k}) \equiv (p_1 p_2)^{(\frac{b}{2}-d)} e^{(d+\frac{\kappa_{\text{eff}}}{2})(\vec{a}\cdot\vec{k})} e^{-\frac{\kappa_{\text{eff}}}{2} d_{ijk} a^i k^j k^k} \\ \times \frac{\prod_l (y_l^{\text{tw}})^{-I_2(\mathbf{R}_l)/2} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k},\vec{\omega}}(\vec{a}\cdot\vec{\omega} + m_{\text{tw},l}, \epsilon_1, \epsilon_2)}{(1 - p_1 p_2 e^{-\vec{a}\cdot\vec{k}})(1 - e^{\vec{a}\cdot\vec{k}})(1 - p_1^{-1} e^{\vec{a}\cdot\vec{k}})(1 - p_2^{-1} e^{\vec{a}\cdot\vec{k}}) \prod_{\vec{\alpha}\cdot\vec{k}=-1} (1 - e^{-\vec{a}\cdot\vec{\alpha}})}. \quad (2.39)$$

Using the property $i_1^{(d_0+\aleph)}(\vec{k})/i_1^{(d_0)}(\vec{k}) = (p_1 p_2)^{-\aleph} e^{\aleph(\vec{a} \cdot \vec{k})}$, the one-instanton partition function Z_1 can be written as

$$\begin{aligned} Z_1 &= \sum_{\vec{k} \in \Delta_\ell} \frac{(1 - p_1^{-1} e^{\vec{a} \cdot \vec{k}})(1 - p_2^{-1} e^{\vec{a} \cdot \vec{k}})}{(1 - p_1)(1 - p_2)} \cdot i_1^{(d_0)}(\vec{k}) \\ &= \frac{(p_1 p_2)^{(\frac{b}{2} - d_0)} \prod_l (y_l^{\text{tw}})^{-\frac{I_2(\mathbf{R}_l)}{2}}}{(1 - p_1)(1 - p_2)} \sum_{\vec{k} \in \Delta_l} \frac{e^{(d_0 + \frac{\kappa_{\text{eff}}}{2})(\vec{a} \cdot \vec{k}) - \frac{\kappa_{\text{eff}}}{2} d_{ijk} a^i k^j k^k} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_l^{\text{tw}})}{(1 - p_1 p_2 e^{-\vec{a} \cdot \vec{k}})(1 - e^{\vec{a} \cdot \vec{k}}) \prod_{\vec{\alpha} \cdot \vec{k} = -1} (1 - e^{-\vec{a} \cdot \vec{\alpha}})}. \end{aligned} \quad (2.40)$$

Notice that there are multiple options for choosing d_0 . However, we find that (2.40) is independent of a specific choice of d_0 . Once we choose $d_0 = 0$, for instance, which works in most cases,¹⁰ (2.40) becomes

$$\boxed{Z_1 = \frac{(p_1 p_2)^{\frac{b}{2}} \prod_l (y_l^{\text{tw}})^{-\frac{I_2(\mathbf{R}_l)}{2}}}{(1 - p_1)(1 - p_2)} \sum_{\vec{k} \in \Delta_\ell} \frac{e^{\frac{\kappa_{\text{eff}}}{2}(\vec{a} \cdot \vec{k} - d_{ijk} a^i k^j k^k)} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_l^{\text{tw}})}{(1 - p_1 p_2 e^{-\vec{a} \cdot \vec{k}})(1 - e^{\vec{a} \cdot \vec{k}}) \prod_{\vec{\alpha} \cdot \vec{k} = -1} (1 - e^{-\vec{a} \cdot \vec{\alpha}})}}. \quad (2.41)$$

When the hypermultiplets are in the representations with $|\vec{k} \cdot \vec{w}| \leq 1$ for all $\vec{w} \in \mathbf{R}$, we have

$$\prod_{\omega \in \mathbf{R}} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_{\text{tw}}, \epsilon_1, \epsilon_2) = \prod_{\vec{k} \cdot \vec{\omega} = -1} (1 - y_{\text{tw}} e^{-\vec{a} \cdot \vec{\omega}}). \quad (2.42)$$

The formula (2.40) indeed reduces to the pure YM partition function derived in [15, 16] upon removing hypermultiplets and Chern-Simons levels up to the overall factor $(p_1 p_2)^{\frac{b}{2}} = e^{-\frac{\hbar^V}{2}(\epsilon_1 + \epsilon_2)}$ that accounts for the shift of instanton fugacity.

We claim that (2.40) is the closed-form expression for the one-instanton partition function, which holds *universally for any gauge theory* with $d_{\text{max}} > 2$. In section 2.3, we study the structure of the blowup equations to bound the number of possible independent equations.

2.3 Number of independent blowup equations

We are mainly interested in 4d $\mathcal{N} = 2$ and 5d $\mathcal{N} = 1$ gauge theories which are UV-complete. UV complete 4d $\mathcal{N} = 2$ gauge theories are classified in [42]. In the case of 5d SCFTs that are UV complete as 5d SCFTs, the possible matter representations are restricted to [43]

- fundamental representation for $SU(N)$, $SO(N)$, $Sp(N)$, G_2 , F_4 , E_6 , E_7
- antisymmetric representation for $SU(N)$, $Sp(N)$
- spinor representation for $SO(N)$ with $7 \leq N \leq 14$
- rank-3 antisymmetric representation for $Sp(3)$, $Sp(4)$, $SU(6)$, $SU(7)$
- symmetric representation for $SU(N)$.

¹⁰A numerical value of d_0 should be a half-integer for theories with $G = Sp(N)_{\theta=\pi}$.

when the gauge group is simple. In the case of 4d, we can also have following additional cases:

- adjoint representation for arbitrary group
- rank-3 antisymmetric for $SU(8)$
- **16** for $Sp(2)$ (half-hypermultiplet)

We are able to obtain the partition function for a large number of cases in 5d using our formula. But we are not able to apply our formula for some cases including the one with adjoint hypermultiplets since the number of independent blowup equations is smaller than 3.

The formula (2.34) is valid only for a certain range of d , for which $\langle (\mathcal{O}_{\mathbb{P}^1})^d \rangle = \mathcal{Z}$. We want to narrow down the valid range of d by performing a simple sanity check on the blowup equation for the one-instanton partition function:

$$Z_1 = p_1^{\frac{b}{2}-d} Z_1^{(N)} + p_2^{\frac{b}{2}-d} Z_1^{(S)} + I_1^{(d)} \quad \text{with an allowed range of } d. \quad (2.43)$$

Specifically, we want to examine the expansion of each term in (2.43) with in powers of $p_1 p_2 \ll 1$. The leading exponent behave as

$$\begin{aligned} I_1^{(d)} &\sim \begin{cases} g_0(\vec{a}, \vec{m}_{\text{tw}}) \cdot (p_1 p_2)^{\frac{b}{2}-d+1} + \dots & \text{for } N_{\text{sym}} = 0 \\ g_0(\vec{a}, \vec{m}_{\text{tw}}) \cdot (p_1 p_2)^{\frac{b}{2}-d} + \dots & \text{for } N_{\text{sym}} = 1 \end{cases} \\ Z_1 &\sim g_1(\vec{a}, \vec{m}_{\text{tw}}) \cdot (p_1 p_2)^{\frac{s}{2}} + \dots \\ p_1^{\frac{b}{2}-d} Z_1^{(N)} &\sim p_2^{\frac{b}{2}-d} Z_1^{(S)} \sim g_2(\vec{a}, \vec{m}_{\text{tw}}) \cdot (p_1 p_2)^{\frac{b}{4}-\frac{d}{2}+\frac{s}{4}} + \dots \end{aligned} \quad (2.44)$$

where $g_{0,1,2}(\vec{a}, \vec{m}_{\text{tw}})$ are functions independent of $p_{1,2}$ and N_{sym} denotes the number of symmetric representation. The numerical value of s will be obtained shortly for a variety of gauge theories for the cases when ADHM-like construction is available. Notice that for the equation (2.43) to be true, some terms on the right-hand side should have the leading exponent less than or equal to that of Z_1 . Therefore, the condition $d - \frac{b}{2} \geq -\frac{s}{2}$ is naturally imposed, setting a lower bound on d .

Similarly, an upper bound on d can be found from an expansion of (2.43) with respect to $1/p_1 p_2 \ll 1$.¹¹ Each term in (2.43) can be written as

$$\begin{aligned} I_1^{(d)} &\sim \begin{cases} h_0(\vec{a}, \vec{m}_{\text{tw}}) \cdot (1/p_1 p_2)^{d-\frac{b}{2}+1} + \dots & \text{for } N_{\text{sym}} = 0 \\ h_0(\vec{a}, \vec{m}_{\text{tw}}) \cdot (1/p_1 p_2)^{d-\frac{b}{2}} + \dots & \text{for } N_{\text{sym}} = 1 \end{cases} \\ Z_1 &\sim h_1(\vec{a}, \vec{m}_{\text{tw}}) \cdot (1/p_1 p_2)^{\frac{s'}{2}} + \dots \\ p_1^{\frac{b}{2}-d} Z_1^{(N)} &\sim p_2^{\frac{b}{2}-d} Z_1^{(S)} \sim h_2(\vec{a}, \vec{m}_{\text{tw}}) \cdot (1/p_1 p_2)^{\frac{d}{2}-\frac{b}{4}+\frac{s'}{4}} + \dots \end{aligned} \quad (2.45)$$

¹¹This is equivalent to assuming a different parameter regime $0 < a_i < -\epsilon_+ < m$ for all $1 \leq i \leq r$. In general, an explicit form of the 1-loop partition function (2.24)–(2.25) can change depending on a parameter regime, thus affecting (2.34). However, all the above expressions remain valid under flipping a sign of ϵ_+ , such that we can simply study the expansion of the single terms in (2.43) with respect to $1/p_1 p_2 \ll 1$.

Again, for (2.43) to be consistent, the leading exponent of Z_1 should be greater than or equal to those of the terms on the right-hand side. Such requirement imposes an upper bound on d , namely $\frac{s'}{2} \geq d - \frac{b}{2}$. Combining the two inequalities, one can identify the following range

$$-\frac{s}{2} + \frac{b}{2} \leq d \leq \frac{s'}{2} + \frac{b}{2}, \quad (2.46)$$

as a necessary condition for (2.43). We explicitly checked that the n -instanton partition function Z_n actually satisfies all the $(\frac{s+s'}{2})$ recursion relations up to a certain value of $n > 1$ for numerous examples whose Z_n is already known from alternative methods. This is true even though the bound (2.46) itself is merely a *necessary* condition found from one-instanton analysis. Based on this empirical observations, we claim that the 5d recursion formulae (2.34) within the above range of d is true at all instanton orders.

Another remarkable thing is that a numerical value of (s, s') exhibits the very simple pattern across a broad range of theories whose gauge group is not $SU(N)_\kappa$. (See Table 1.)

$$\begin{aligned} s = s' &= h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) & \text{for } G \neq SU(N)_\kappa \text{ nor } Sp(N) \\ s = s' - 2 \left\{ \frac{N_f}{2} \right\} &= h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) & \text{for } G = Sp(N)_{\theta=0} \\ s = s' + 2 \left\{ \frac{N_f}{2} \right\} &= h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + 1 & \text{for } G = Sp(N)_{\theta=\pi} \end{aligned} \quad (2.47)$$

where $\{x\} \equiv x - [x]$ denote the non-integer part of x . As the above numerical pattern (2.47) emerges for all $G \neq SU(N)_\kappa$ examples that we studied, we conjecture that (2.47) is generally true, thereby taking the recursion formulae (2.34) with

$$\begin{aligned} 0 \leq d &\leq h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) & \text{for } G \neq SU(N)_\kappa \text{ nor } Sp(N) \\ 0 \leq d &\leq h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + \left\{ \frac{N_f}{2} \right\} & \text{for } G = Sp(N)_{\theta=0} \\ -\frac{1}{2} \leq d &\leq h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + \frac{1}{2} - \left\{ \frac{N_f}{2} \right\} & \text{for } G = Sp(N)_{\theta=\pi} \end{aligned} \quad (2.48)$$

as a basic assumption to obtain the partition function \mathcal{Z} for any $G \neq SU(N)_\kappa$ gauge theory. It would be desirable to understand from the first principle the range (2.48) of d for which (2.34) holds true.

It turns out to be more difficult to characterize a general pattern behind (s, s') for $SU(N)_\kappa$ gauge theories, due to extra complication caused by the 5d Chern-Simons level κ . Here we consider two particular classes of $SU(N)_\kappa$ gauge theories for illustration. For $SU(N)_\kappa + N_f \mathbf{F}$

gauge theory (N_f fundamental hypermultiplets) with $N_f + 2|\kappa| \leq 2N$, we find that

$$s = \begin{cases} \frac{N_f}{2} & \text{if } \kappa_{\text{eff}} = N - N_f, \\ N - \frac{1}{2} \sum_l I_2(\mathbf{F}) + |\kappa_{\text{eff}}| & \text{otherwise.} \end{cases} \quad (2.49)$$

$$s' = \begin{cases} \frac{N_f}{2} & \text{if } \kappa_{\text{eff}} = -N, \\ N - \frac{1}{2} \sum_l I_2(\mathbf{F}) + |\bar{\kappa}_{\text{eff}}| & \text{otherwise.} \end{cases} \quad (2.50)$$

where $\bar{\kappa}_{\text{eff}} \equiv \kappa + \frac{1}{2} \sum_l I_3(\mathbf{F})$. Plugging in these values to (2.46), we find the range of d to be

$$\begin{aligned} 0 \leq d \leq N & \quad \text{if } \kappa = -N + \frac{N_f}{2} \\ 0 \leq d \leq N - \frac{N_f}{2} - \kappa & \quad \text{if } \kappa \in \left(-N + \frac{N_f}{2}, -\frac{N_f}{2} \right] \\ 0 \leq d \leq N & \quad \text{if } \kappa \in \left[-\frac{N_f}{2}, +\frac{N_f}{2} \right] \\ \frac{N_f}{2} - \kappa \leq d \leq N & \quad \text{if } \kappa \in \left[\frac{N_f}{2}, N - \frac{N_f}{2} \right) \\ 0 \leq d \leq N & \quad \text{if } \kappa = N - \frac{N_f}{2} \end{aligned} \quad (2.51)$$

which always includes the range $0 \leq d \leq N$. Thus the recursion formula (2.34) holds for at least 3 values of d , which is enough to determine the partition function Z_{inst} completely.

For the $SU(N)_\kappa + N_f \mathbf{F} + 1 \mathbf{AS}$ theory (N_f fundamentals and 1 anti-symmetric tensor) with $N_f + 2|\kappa| \leq N + 4$, we find

$$\begin{aligned} s &= \min \left(N - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) - (\kappa_{\text{eff}} - 2), N - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + 2 \left\{ \frac{\kappa_{\text{eff}}}{2} \right\} \right) \\ s' &= \min \left(N - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + \left(\kappa_{\text{eff}} + \sum_l I_3(\mathbf{R}_l) + 2 \right), \right. \\ &\quad \left. N - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + 2 \left\{ -\frac{\kappa_{\text{eff}}}{2} + \frac{1}{2} \sum_l I_3(\mathbf{R}_l) \right\} \right), \end{aligned} \quad (2.52)$$

from which one can identify the valid range of d via (2.46). So as long as there exist at least three distinct allowed values for d for given (N, κ) , the corresponding partition function Z_{inst} can be solved from the recursion formula (2.34).

We also consider $SU(6)_\kappa + 1 \mathbf{TAS}$ theory (one rank-3 antisymmetric tensor) with $|\kappa| \leq 3$ in Section 3. This model can be Higgsed to two disjoint copies of $SU(3)_\kappa$ theory without a bifundamental hypermultiplet [44]. At the level of the partition function, Higgsing is realized by turning off $m_{\text{tw}} = 0$ and imposing the $SU(3)$ traceless conditions. As neither of them

modify s and s' , the numerical value of (s, s') must be identical to that of $SU(3)_\kappa$ gauge theory, which always allow the range $0 \leq d \leq 3$. Therefore, the recursion formula (2.34) is enough to determine the instanton partition function Z_{inst} for $SU(6)_\kappa + 1\text{TAS}$ theory as well.

3 Examples

Recall that the recursion formula (2.34) for the k -instanton partition function, and especially the general expression (2.40) at one-instanton order, are widely applicable to 5d $\mathcal{N} = 1$ gauge theory whose (n, n') satisfies $\frac{n+n'}{2} \geq 2$. Combined with the observation that (n, n') follows (2.47) in most cases, they actually become a very efficient approach at obtaining the BPS partition function on $\mathbb{C}^2 \times S^1$, unless a matter representation is too large.

Conventionally, the instanton partition function has been computed by employing the ADHM construction of the instanton moduli space [3, 4, 45] or by applying the topological vertex formalism to the 5-brane web [46, 47]. Both are based on a certain UV realization of 5d $\mathcal{N} = 1$ gauge theory, motivated by its string theory engineering. Sometimes the UV completion of the gauge theory can be achieved only with specific string theory realizations, but not with the others. For instance, $SU(2) + N_f \mathbf{F}$ gauge theory with $N_f \geq 5$ must be embedded into D4-D8-O8 brane system, to be UV-completed as 5d E_{N_f+1} Minahan-Nemeschansky SCFT, instead of a typical (p, q) 5-brane web with colliding branes which indicate UV inconsistency. A sensible UV QFT observable can thus be obtained only through a proper embedding of the gauge theory into string theory. At some occasions, an extra factor entangled with a true QFT observable may appear during the above instanton computation, which is sensitive to the choice of a string theory embedding.

Our blow-up formula (2.34) implicitly assumes a particular string theory embedding of the gauge theory. There are a wide variety of ‘exceptional’ gauge theories, whose UV completion is found as M-theory wrapped on a singular Calabi-Yau 3-fold [48–51], but for which neither the ADHM quantum mechanics nor the topological vertex formalism has been developed. We will illustrate that bootstrapping the partition function \mathcal{Z} based on the recursion formula (2.34) works well for those ‘exceptional’ theories, providing their BPS spectrum efficiently.

3.1 Theories with the known ADHM description

Let us first consider the ‘standard’ gauge theories with classical gauge groups, whose hypermultiplet admits UV realization as a perturbative string ending on non-perturbative branes. In these cases, the ADHM construction of the instanton moduli space is well-known [3, 41, 45]. As for the k -instanton partition function Z_k , the Witten index of the relevant ADHM quantum mechanics can be computed by SUSY localization [19, 52–54], ending up collecting all Jeffrey-Kirwan residues of a multi-dimensional contour integral. We will examine whether the recursion formula (2.34) actually produces the same result as the localization computation.

SU(N) The ADHM computation of the k -instanton partition function, for $SU(N)_\kappa + N_f \mathbf{F}$ theories with $N_f + 2|\kappa| \leq 2N$, can be summarized as the Young tableaux sum expression.

$$Z_k^{\text{ADHM}} = \sum_{|\vec{Y}|=k} \prod_{i=1}^N \prod_{s \in Y_i} \frac{e^{-\kappa \phi(s)} \prod_{l=1}^{N_f} 2 \sinh \frac{\phi(s) + m_l}{2}}{\prod_{j=1}^N 2 \sinh \frac{E_{ij}}{2} 2 \sinh \frac{E_{ij} - 2\epsilon_+}{2}} \quad (3.1)$$

where

$$\begin{aligned} E_{ij}(s) &= a_i - a_j - \epsilon_1 h_i(s) + \epsilon_2 (v_j(s) + 1) \\ \varphi(s) &= a_i - \epsilon_+ - (n-1)\epsilon_1 - (m-1)\epsilon_2 \quad \text{for } s = (m, n) \in Y_i. \end{aligned}$$

$h_i(s)$ denotes the distance from s to the right end of the diagram Y_i by moving right. $v_j(s)$ denotes the distance from s to the bottom of the diagram Y_j by moving down.

We checked that the instanton corrections Z_1 and Z_2 obtained from the recursion formulae (2.34) with (2.51) and the 1-instanton expression (2.40) precisely agree with the above $Z_{k=1,2}^{\text{ADHM}}$ for $N = 2, 3, 4$. Recall that Z_k^{ADHM} often contains an additional factor Z_{extra} that captures the contribution from an extra branch of vacua. It is sensitive to the string theory embedding of gauge theory and regarded as spurious from the 5d QFT perspective. It is usually factorized from the true QFT partition function as

$$1 + \sum_{k=1}^{\infty} q^k Z_k^{\text{ADHM}}(\vec{a}, \epsilon_1, \epsilon_2, \vec{m}) = Z_{\text{QFT}}(\vec{a}, \epsilon_1, \epsilon_2, \vec{m}, \Lambda) \cdot Z_{\text{extra}}(\epsilon_1, \epsilon_2, \vec{m}, \Lambda). \quad (3.2)$$

A non-trivial $Z_{\text{extra}} \neq 1$ appears in the above expression (3.1) if and only if $N_f + 2|\kappa| = 2N$. This factor can be identified as the contribution of D1-branes escaping from D5-branes which engineer the $SU(N)_\kappa + N_f \mathbf{F}$ gauge theory. Since $Z_k = Z_k^{\text{ADHM}}$, the same factor Z_{extra} emerges from the recursion formula (2.34) as well. The 5-brane web construction of the gauge theory is thus indirectly reflected in the recursion formula.

A similar observation is that the 1-instanton expression (2.40) applied to $SU(2)_\kappa + N_f \mathbf{F}$ with $N_f \geq 5$ does not match the Witten index of the D0-D4-D8-O8⁻ quantum mechanics, which is the correct 1-instanton partition function.¹² Instead, it coincides with the topological vertex computation applied to the 5-brane web with a colliding pair of branes, which engineers the $SU(2)$ gauge theory with $N_f \geq 5$ in IR, but behaves badly in UV. Again, this suggests that the recursion formula (2.34) is rooted in a specific string theory construction of the gauge theory, i.e., the web of (p, q) 5-branes. It would be more satisfactory if one can trace at which step in our formulation of (2.34) the particular UV embedding of the gauge theory has been chosen. In fact, there must be a broader set of blow-up relations for the 5d Nekrasov partition function \mathcal{Z} , similar to those recently found for 6d SCFTs [32, 33, 56, 57]. We expect that some recursion formulae, derived from the generalized blow-up equations, may realize a different string theory embedding of the gauge theory. It would be very interesting if one can reveal the connection between the choice of UV embedding and the blow-up equations.

¹²The case with $SU(2) \simeq Sp(1)$ is an exception, which allows $N_f \leq 7$ fundamental hypermultiplets [55].

In $SU(N)_\kappa + N_f \mathbf{F} + 1 \mathbf{AS}$ theory with $N_f + 2|\kappa| \leq N + 4$, the ADHM quantum mechanics is the worldvolume theory of D1-branes, probing the D5-NS5-D7-O7⁻ brane configuration that realizes the gauge theory. Let us compute the Witten index for 1 and 2 D1-branes, then compare with the blow-up computation based on the recursion formula (2.34). For instance, the Witten index for the single D1-brane can be written as

$$Z_1^{\text{ADHM}} = - \sum_{i=1}^N \frac{e^{-\kappa(a_i - \epsilon_+)}}{2 \sinh \frac{\epsilon_1}{2} 2 \sinh \frac{\epsilon_2}{2}} \frac{\prod_{l=1}^{N_f} 2 \sinh \frac{-\epsilon_+ + a_i + m_l}{2}}{2 \sinh \frac{-3\epsilon_+ + 2a_i + m_a}{2}} \prod_{j \neq i} \frac{2 \sinh \frac{a_i + a_j + m_a - \epsilon_+}{2}}{2 \sinh \frac{a_i - a_j}{2} 2 \sinh \frac{2\epsilon_+ - a_i + a_j}{2}} \\ - \frac{1}{2} \frac{e^{-\frac{\kappa}{2}(\epsilon_+ - m_a)}}{2 \sinh \frac{\epsilon_1}{2} 2 \sinh \frac{\epsilon_2}{2}} \left(\frac{\prod_{l=1}^{N_f} 2 \sinh \frac{\epsilon_+ + 2m_l - m_a}{4}}{\prod_{i=1}^N 2 \sinh \frac{3\epsilon_+ - M - 2a_i}{4}} - (-1)^{\kappa + (N - N_f)/2} \frac{\prod_{l=1}^{N_f} 2 \cosh \frac{\epsilon_+ + 2m_l - m_a}{4}}{\prod_{i=1}^N 2 \cosh \frac{3\epsilon_+ - M - 2a_i}{4}} \right). \quad (3.3)$$

Note that Z_k^{ADHM} captures an extra factor $Z_{\text{extra}} \neq 1$ if $N_f + 2|\kappa| = N + 4$, displaying the BPS spectrum of D1-branes escaping from the D5-branes on which the gauge theory is supported. The appearance of $Z_{\text{extra}} \neq 1$ is an artifact of the string theory embedding, spurious from the 5d QFT perspective. We checked the agreement between Z_1^{ADHM} and the 1-instanton formula (2.40) for the $SU(3)$, $SU(4)$, $SU(5)$ theories whose (n, n') satisfies $\frac{n+n'}{2} \geq 2$. We confirmed $Z_2 = Z_2^{\text{ADHM}}$ as well, where Z_2 is the solution of the recursion formulae (2.34) with (2.52). The same spurious factor Z_{extra} arises from the recursion formula, implying that our blowup equations are implicitly based on the D5-NS5-D7-O7⁻ brane realization of the gauge theory.¹³

Sp(N) The k -instanton partition function for $Sp(N)_\theta + N_f \mathbf{F}$ theory with $N_f \leq 2N + 4$ can be computed from the ADHM quantum mechanics of D1-D5-NS5-O5 branes, which engineers the gauge theory and its instantons. The Witten index for a D1-brane can be written as

$$Z_1^{\text{ADHM}} = \frac{1}{2} \frac{1}{2 \sinh \frac{\epsilon_1}{2} 2 \sinh \frac{\epsilon_2}{2}} \left(\frac{\prod_{l=1}^{N_f} 2 \sinh \frac{m_l}{2}}{\prod_{i=1}^N 2 \sinh \frac{\epsilon_+ \pm a_i}{2}} + e^{i\theta} \frac{\prod_{l=1}^{N_f} 2 \cosh \frac{m_l}{2}}{\prod_{i=1}^N 2 \cosh \frac{\epsilon_+ \pm a_i}{2}} \right). \quad (3.4)$$

Our 1-instanton formula (2.40) was checked to be in agreement with Z_1^{ADHM} for the $Sp(2)$ and $Sp(3)$ gauge theories satisfying $N - 1 \geq \lfloor \frac{N_f}{2} \rfloor$ (at $\theta = 0$) and $N \geq \lceil \frac{N_f}{2} \rceil$ (at $\theta = \pi$). It was also confirmed that $Z_2^{\text{ADHM}} = Z_2$, where Z_2 is the solution of the recursion formulae (2.34) with (2.48). Note that Z_{extra} does not appear in $Z_k^{\text{ADHM}} = Z_k$ for the above theories.

In $Sp(N)_\theta + N_f \mathbf{F} + 1 \mathbf{AS}$ theory with $N_f \leq 7$, the relevant ADHM quantum mechanics is the worldvolume gauge theory of D0-branes which probe the D4-D8-O8 brane configuration. The QFT on D4-branes sees an enhanced E_{N_f+1} flavor symmetry at the UV fixed point [55].

¹³An exceptional case is the $SU(2)$ gauge theory, in which the antisymmetric hypermultiplet decouples and never affects the recursion formula. The corresponding Z_k is the same as the Young tableau formula (3.1).

Let us consider the Witten index for one and two D0-branes [19, 58]. For a single D0-brane,

$$Z_1^{\text{ADHM}} = \frac{1}{2} \frac{1}{2 \sinh \frac{\epsilon_1}{2} 2 \sinh \frac{\epsilon_2}{2} 2 \sinh \frac{m_a + \epsilon_+}{2} 2 \sinh \frac{m_a - \epsilon_+}{2}} \times \left(\frac{\prod_{i=1}^N 2 \sinh \frac{m_a \pm a_i}{2} \prod_{l=1}^{N_f} 2 \sinh \frac{m_l}{2}}{\prod_{i=1}^N 2 \sinh \frac{\epsilon_+ \pm a_i}{2}} + \frac{e^{i\theta} \prod_{i=1}^N 2 \cosh \frac{m_a \pm a_i}{2} \prod_{l=1}^{N_f} 2 \cosh \frac{m_l}{2}}{\prod_{i=1}^N 2 \cosh \frac{\epsilon_+ \pm a_i}{2}} \right). \quad (3.5)$$

We checked Z_1^{ADHM} itself is *not* the same as the 1-instanton expression (2.40) for the $Sp(2)_\theta$, $Sp(3)_\theta$ theories with $N_f \leq 1$ (at $\theta = 0$) and $N_f \leq 2$ (at $\theta = \pi$). Instead, the difference between Z_1 and Z_1^{ADHM} can be identified as the BPS index of D0-branes moving away from the D4-D8-O8 brane system [19, 58]. Similarly, it was confirmed that the 2-instanton correction Z_2 captures the same 5d QFT spectrum as in Z_2^{ADHM} , while excluding the spurious contribution of escaping D0-branes. Our blow-up formulae seem to reflect a different UV embedding of the gauge theory, which does *not* allow an extra branch of vacua leading to $Z_{\text{extra}} \neq 1$.

SO(N) One can compute the instanton partition function of $SO(N) + N_v \mathbf{V}$ theory with $N_v \leq N - 4$ using the ADHM quantum mechanics of the D1-D5-NS5-O5 brane system. At even N , the Witten index for a D1-brane can be written as

$$Z_1^{\text{ADHM}} = \sum_{i=1}^{N/2} \left(\frac{2 \sinh(2\epsilon_+ - a_i) 2 \sinh(a_i - \epsilon_+) \prod_{l=1}^{N_v} 2 \sinh \frac{m_l \pm (a_i - \epsilon_+)}{2}}{2 \cdot 2 \sinh \frac{\epsilon_1}{2} 2 \sinh \frac{\epsilon_2}{2} \prod_{j \neq i} 2 \sinh \frac{a_i \pm a_j}{2} 2 \sinh \frac{2\epsilon_+ - a_i \pm a_j}{2}} + (a_i \rightarrow -a_i) \right). \quad (3.6)$$

At odd N ,

$$Z_1^{\text{ADHM}} = \sum_{i=1}^{\lfloor N/2 \rfloor} \left(\frac{2 \cosh \frac{2\epsilon_+ - a_i}{2} 2 \sinh(a_i - \epsilon_+) \prod_{l=1}^{N_f} 2 \sinh \frac{m_l \pm (a_i - \epsilon_+)}{2}}{2 \cdot 2 \sinh \frac{\epsilon_1}{2} 2 \sinh \frac{\epsilon_2}{2} 2 \sinh \frac{a_i}{2} \prod_{j \neq i} 2 \sinh \frac{a_i \pm a_j}{2} 2 \sinh \frac{2\epsilon_+ - a_i \pm a_j}{2}} + (a_i \rightarrow -a_i) \right). \quad (3.7)$$

The general 1-instanton expression (2.40) and the recursion formula (2.34) are applicable for all $N_v \leq N - 4$. We explicitly confirmed that $Z_k^{\text{ADHM}} = Z_k$, where Z_1 was written in (2.40) and Z_2 was obtained as the solution of the recursion formula (2.34), for $4 \leq N \leq 9$. Notice that $Z_k^{\text{ADHM}} = Z_k$ involves a non-trivial extra factor $Z_{\text{extra}} \neq 1$ when $N_v = N - 4$. This extra factor is associated to D1-branes moving away from the D5-NS5-O5 brane system, where the 5d QFT lives. It implies that the specific UV realization of the gauge theory, i.e., IIB string theory with D1-D5-NS5-O5, is implicit in our recursion formulae (2.34) with (2.48).

3.2 Theories with spinor hypermultiplets

So far, we have investigated the ‘standard’ gauge theories which have certain D-brane set-ups in IIA/IIB string theory to realize themselves and their non-perturbative solitons. Once there are a sufficient number of the blowup equations, the k -instanton partition function Z_k can be obtained as the solution. Solving specifically at 1-instanton level, Z_1 is expressed as (2.40).

All these results were found to be in agreement with the ADHM instanton calculus, sometimes modulo an extra factor Z_{extra} sensitive to the string theory embedding of the gauge theory.

There are also ‘exceptional’ gauge theories, which involve exceptional gauge groups or matters that cannot be engineered from perturbative open strings ending on non-perturbative branes. Instead, their UV embedding is geometrically realized as M-theory wrapped on local Calabi-Yau threefolds. Since the exceptional theories lack for the ADHM formalism, their instanton partition function \mathcal{Z} has been studied on a case-by-case basis; Once the 5-brane web configuration engineering an exceptional theory is identified, the topological vertex formalism can be applied to compute the relevant partition function \mathcal{Z} . Alternatively, one can first construct the $\mathbb{C}^2 \times T^2$ partition function for a related 6d gauge theory, based on its modularity and anomaly, then take the circle reduction to obtain the 5d partition function \mathcal{Z} . Several interesting exceptional theories have been studied so far, based on the above two approaches.

We can take advantage of the universality of the blowup equation. Recall that the blowup recursion formula (2.34) holds for a certain range of d , i.e., the set of all integers between $0 \leq d \leq h^\vee - \frac{1}{2} \sum_l I(\mathbf{R}_l)$, when the gauge group G is neither $SU(N)_\kappa$ nor $Sp(N)_\theta$ so that an extra complication due to the Chern-Simons level κ or the theta angle θ cannot occur. One can solve the recursion formulae for the k -instanton correction Z_k to the partition function, as long as $h^\vee - \frac{1}{2} \sum_l I(\mathbf{R}_l) \geq 2$, even for the exceptional gauge theories. We assume that Z_k solved from the recursion formulae would be the correct BPS data for UV-consistent 5d SCFTs, modulo an extra factor Z_{extra} independent of the Coulomb VEV \vec{a} . This conjecture will be tested via comparison with [44, 59, 60] which compute \mathcal{Z} for some exceptional cases.

In this section, we will focus on the $SO(N)$ gauge theories with spinor hypermultiplets. There are a sufficient number of recursion formulae (2.34) to fix the k -instanton partition function Z_k of the $SO(N)$ gauge theory, if and only if

$$\begin{aligned} N - 4 &\geq N_{\mathbf{v}} + 2^{\frac{N-7}{2}} \cdot N_{\mathbf{s}} && \text{for odd } N \\ N - 4 &\geq N_{\mathbf{v}} + 2^{\frac{N-8}{2}} \cdot (N_{\mathbf{s}} + N_{\mathbf{c}}) && \text{for even } N \end{aligned} \quad (3.8)$$

where $N_{\mathbf{v}}$, $N_{\mathbf{s}}$, and $N_{\mathbf{c}}$ denote the number of hypermultiplets in the vector, spinor, and conjugate spinor representations, respectively. Our 1-instanton expression (2.40) is also applicable to the cases satisfying (3.8). Let us consider the special cases whose Z_k is already known.

SO(7) The k -instanton contribution Z_k of $SO(7) + N_{\mathbf{s}} \mathbf{S}$ theory can be obtained from the SUSY quantum mechanics proposed in [59], which can be summarized as the following $SU(4)$ Young tableau expression:

$$Z_k^{\text{YD}} = \sum_{|\vec{Y}|=k} \prod_{i=1}^4 \prod_{s \in Y_i} \frac{2 \sinh(\phi(s))}{\prod_{j=1}^4 2 \sinh \frac{E_{ij}}{2}} \frac{2 \sinh(\phi(s) - \epsilon_+)}{2 \sinh \frac{E_{ij} - 2\epsilon_+}{2}} \frac{\prod_{l=1}^{N_{\mathbf{s}}} 2 \sinh(\frac{m_l \pm \phi(s)}{2})}{2 \sinh \frac{\epsilon_+ - \phi(s) - a_j}{2}} \quad (3.9)$$

$$\times \prod_{i \leq j}^4 \prod_{\substack{s_i, j \in Y_{i,j} \\ s_i < s_j}} \frac{2 \sinh \frac{\phi(s_i) + \phi(s_j)}{2}}{2 \sinh \frac{\epsilon_1 - \phi(s_i) - \phi(s_j)}{2}} \frac{2 \sinh \frac{\phi(s_i) + \phi(s_j) - 2\epsilon_+}{2}}{2 \sinh \frac{\epsilon_2 - \phi(s_i) - \phi(s_j)}{2}} \quad (3.10)$$

We checked the agreement between Z_1^{YD} and the 1-instanton formula Z_1 in (2.40) for $N_s \leq 3$. At two instantons, it was also confirmed for $N_s \leq 3$ that $Z_2^{\text{YD}} = Z_2$, where Z_2 is the solution of the recursion formula (2.34) with (2.48). Such explicit comparison implies that the blow-up recursion formula (2.34) actually works for the $SO(7) + N_s \mathbf{S}$ theory.

The 1-instanton partition function of $SO(7) + 4\mathbf{S} + 1\mathbf{V}$ theory is written in (H.15) of [60]. We deduced the 1-instanton correction of $SO(7) + N_s \mathbf{S} + N_v \mathbf{V}$ theory with $(N_s, N_v) \leq (2, 1)$ from that, by taking several flavor chemical potentials to infinity. All of those were confirmed to be in agreement with our general 1-instanton expression (2.40) up to $(p_1 p_2)^{13/2}$.

SO(8) The 1-instanton result of $SO(8) + 1\mathbf{S} + 1\mathbf{C} + 1\mathbf{V}$ theory is found in (H.28) of [60]. That is expressed in terms of irreducible characters $\chi_{\mathbf{R}}^S$, whose superscript $S \in \{G, v, s, c\}$ means either the gauge symmetry (G) or the flavor symmetry acting on the vector (v), spinor (s), or conjugate spinor (c) hypermultiplets. Their representation \mathbf{R} is specified by the Dynkin label in the subscript. All irreducible characters for the flavor symmetry are assumed to be written in the orthogonal basis, to be compatible with our convention of mass parameters in (2.25), (2.34), (2.40). The mass parameters will be often distinguished by the superscript $S \in \{s, c, v\}$ according to the matter representation. The flavor symmetry is $Sp(N_v)_v \times Sp(N_s)_s \times Sp(N_c)_c$.

We obtained the 1-instanton partition function of $SO(8) + N_s \mathbf{S} + N_c \mathbf{C} + N_v \mathbf{V}$ theory with $(N_s, N_c, N_v) \leq (1, 1, 1)$ from (H.28) by sending appropriate mass parameters to infinity. All of those were consistent with our general 1-instanton expression (2.40) up to t^{20} order, where $t \equiv \sqrt{p_1 p_2}$. Furthermore, we could determine the unknown part of (H.28) of [60] as

$$\begin{aligned} \hat{Z}_1 = t^4 &+ \sum_{n=0}^{\infty} t^{5+2n} (\chi_{(0n00)}^G \chi_{(1)}^v \chi_{(1)}^s \chi_{(1)}^c + \sum_{n=0}^{\infty} t^{6+2n} (\chi_{(1n00)}^G \chi_{(1)}^s \chi_{(1)}^c + \chi_{(0n10)}^G \chi_{(1)}^s \chi_{(1)}^v) \\ &+ \chi_{(0n01)}^G \chi_{(1)}^c \chi_{(1)}^v) + \sum_{n=0}^{\infty} t^{7+2n} (\chi_{(1n10)}^G \chi_{(1)}^s + \chi_{(1n01)}^G \chi_{(1)}^c + \chi_{(0n11)}^G \chi_{(1)}^v) - \sum_{n=0}^{\infty} t^{8+2n} \chi_{(1n11)}^G, \end{aligned} \quad (3.11)$$

where $\hat{Z}_1 \equiv (2 \sinh \frac{\epsilon_{1,2}}{2}) \cdot Z_1$ is the 1-instanton result after removing the center-of-mass factor.

Now we compare (2.40) with the 1-instanton partition function of $SO(8) + 2\mathbf{S} + 2\mathbf{C} + 2\mathbf{V}$ theory, written in (H.19) of [60]. Our 1-instanton formula (2.40) applied to the $SO(8)$ theories having $(N_s, N_c, N_v) \leq (2, 1, 1), (1, 2, 2), (1, 1, 2), (2, 2, 0), (2, 0, 2), (0, 2, 2)$ was consistent with (H.19) up to t^{20} order, after suitably setting some mass parameters in (H.19) to infinity. By comparison, we could further determine the unknown part of (H.19) of [60] as

$$\begin{aligned} \hat{Z}_1 = t^{-1} &- t^3 (\chi_{(01)}^v + \chi_{(01)}^s + \chi_{(01)}^c) + t^5 (\chi_{(1000)}^G \chi_{(10)}^s \chi_{(10)}^c + \chi_{(0010)}^G \chi_{(10)}^s \chi_{(10)}^v + \chi_{(0001)}^G \chi_{(10)}^c \chi_{(10)}^v) \\ &- t^6 (\chi_{(1010)}^G \chi_{(10)}^s + \chi_{(1001)}^G \chi_{(10)}^c + \chi_{(0011)}^G \chi_{(10)}^v) + t^7 \chi_{(1011)}^G - \sum_{n=0}^{\infty} \left(t^{5+2n} \chi_{(0n00)}^G \chi_{(10)}^s \chi_{(10)}^c \chi_{(10)}^v \right. \\ &+ t^{6+2n} (\chi_{(1n00)}^G \chi_{(01)}^s \chi_{(01)}^c \chi_{(10)}^v + \chi_{(0n10)}^G \chi_{(01)}^s \chi_{(10)}^c \chi_{(01)}^v + \chi_{(0n01)}^G \chi_{(10)}^s \chi_{(01)}^c \chi_{(01)}^v) \\ &- t^{7+2n} (\chi_{(1n10)}^G \chi_{(01)}^s \chi_{(10)}^c \chi_{(10)}^v + \chi_{(1n01)}^G \chi_{(10)}^s \chi_{(01)}^c \chi_{(10)}^v + \chi_{(0n11)}^G \chi_{(10)}^s \chi_{(10)}^c \chi_{(01)}^v) \\ &\left. + t^{8+2n} (\chi_{(2n10)}^G \chi_{(01)}^s \chi_{(10)}^c + \chi_{(2n01)}^G \chi_{(10)}^s \chi_{(01)}^c + \chi_{(1n20)}^G \chi_{(01)}^s \chi_{(10)}^v + \chi_{(1n02)}^G \chi_{(01)}^c \chi_{(10)}^v) \right) \end{aligned} \quad (3.12)$$

$$\begin{aligned}
& + \chi_{(0n21)}^G \chi_{(10)}^s \chi_{(01)}^v + \chi_{(0n12)}^G \chi_{(10)}^c \chi_{(01)}^v - t^{9+2n} (\chi_{(2n11)}^G \chi_{(10)}^s \chi_{(10)}^c + \chi_{(1n21)}^G \chi_{(10)}^s \chi_{(10)}^v \\
& + \chi_{(1n12)}^G \chi_{(10)}^c \chi_{(10)}^v) + t^{10+2n} (\chi_{(2n21)}^G \chi_{(10)}^s + \chi_{(2n12)}^G \chi_{(10)}^c + \chi_{(1n22)}^G \chi_{(10)}^v) - t^{11+2n} \chi_{(2n22)}^G.
\end{aligned}$$

Notice that (3.11) and (3.12) are manifestly invariant under the $SO(8)$ triality, transforming the $SO(8)$ representations as $(n_v n_a n_c n_s) \rightarrow (n_s n_a n_v n_c)$ along with $\chi_{\mathbf{R}}^v \rightarrow \chi_{\mathbf{R}}^s \rightarrow \chi_{\mathbf{R}}^c \rightarrow \chi_{\mathbf{R}}^v$. It can be done by shuffling the Coulomb VEVs and renaming the flavor chemical potentials. We rearranged Z_1 in terms of the new variables \vec{a}' or \vec{a}'' ,

$$\begin{aligned}
(a'_1, a'_2, a'_3, a'_4) &= \left(\frac{-a_1+a_2+a_3-a_4}{2}, \frac{-a_1+a_2+a_3-a_4}{2}, \frac{-a_1+a_2+a_3-a_4}{2}, \frac{-a_1+a_2+a_3-a_4}{2} \right) \\
(a''_1, a''_2, a''_3, a''_4) &= \left(\frac{+a_1-a_2-a_3-a_4}{2}, \frac{-a_1+a_2-a_3-a_4}{2}, \frac{-a_1-a_2+a_3-a_4}{2}, \frac{+a_1+a_2+a_3-a_4}{2} \right),
\end{aligned} \tag{3.13}$$

which exchanges the $SO(8)$ irreducible characters as

$$\chi_{(n_c n_a n_s n_v)}(\vec{a}) = \chi_{(n_v n_a n_c n_s)}(\vec{a}')|_{\vec{a}' \rightarrow \vec{a}}, \quad \chi_{(n_s n_a n_v n_c)}(\vec{a}) = \chi_{(n_v n_a n_c n_s)}(\vec{a}'')|_{\vec{a}'' \rightarrow \vec{a}}. \tag{3.14}$$

Dropping off primes from $Z_1(\vec{a}', \epsilon_1, \epsilon_2; \vec{m}^s, \vec{m}^c, \vec{m}^v)$ or $Z_1(\vec{a}'', \epsilon_1, \epsilon_2; \vec{m}^s, \vec{m}^c, \vec{m}^v)$, we confirmed

$$\begin{aligned}
Z_1^{N_s=N_c=N_v}(\vec{a}, \epsilon_1, \epsilon_2; \vec{m}^s, \vec{m}^c, \vec{m}^v) &= Z_1^{N_s=N_c=N_v}(\vec{a}', \epsilon_1, \epsilon_2; \vec{m}^v, \vec{m}^s, \vec{m}^c)|_{\vec{a}' \rightarrow \vec{a}} \\
Z_1^{N_s=N_c=N_v}(\vec{a}, \epsilon_1, \epsilon_2; \vec{m}^s, \vec{m}^c, \vec{m}^v) &= Z_1^{N_s=N_c=N_v}(\vec{a}'', \epsilon_1, \epsilon_2; \vec{m}^c, \vec{m}^v, \vec{m}^s)|_{\vec{a}'' \rightarrow \vec{a}}.
\end{aligned} \tag{3.15}$$

Similarly, we also found the 1-instanton formula (2.40) applied to $SO(8)$ theories with $(N_s, N_c, N_v) \leq (4, 0, 0)$ or $(0, 4, 0)$ is compatible with the $SO(8)$ triality. Starting with the 1-instanton result $Z_1^{\text{ADHM}} = Z_1^{\text{ADHM}}(\vec{a}, \epsilon_1, \epsilon_2, \vec{m})$ obtained from the relevant ADHM quantum mechanics for $SO(8) + N_v \mathbf{V}$ theory with $N_v \leq 4$, we checked that

$$\begin{aligned}
Z_1^{N_c, N_c=N_v=0}(\vec{a}, \epsilon_1, \epsilon_2, \vec{m}) &= Z_1^{\text{ADHM}}(\vec{a}', \epsilon_1, \epsilon_2, \vec{m})|_{\vec{a}' \rightarrow \vec{a}} \\
Z_1^{N_s, N_s=N_v=0}(\vec{a}, \epsilon_1, \epsilon_2, \vec{m}) &= Z_1^{\text{ADHM}}(\vec{a}'', \epsilon_1, \epsilon_2, \vec{m})|_{\vec{a}'' \rightarrow \vec{a}}.
\end{aligned} \tag{3.16}$$

SO(9) The 1-instanton formula (2.40) can be applied to $SO(9) + N_s \mathbf{S} + N_v \mathbf{V}$ theory with $(N_s, N_v) \leq (1, 3)$ or $(2, 1)$, which has $Sp(N_s)_s \times Sp(N_v)_v$ flavor symmetry. It can be compared with the 1-instanton partition function of $SO(9) + 2\mathbf{S} + 3\mathbf{V}$ theory, which is written in (H.20) of [60] up to t^7 order, after appropriately taking some mass parameters to infinity. We checked all their consistency up to the given order. For example, the character expansion of \hat{Z}_1 for $SO(9) + 2\mathbf{S} + 1\mathbf{V}$ can be written as

$$\begin{aligned}
\hat{Z}_1 &= t^4 \chi_{(1)}^v + t^5 \chi_{(20)}^s - t^6 \chi_{(0001)}^G \chi_{(10)}^s + \sum_{n=0}^{\infty} \left(t^{6+2n} \chi_{(0n00)}^G \chi_{(02)}^s \chi_{(1)}^v \right. \\
&\quad - t^{7+2n} (\chi_{(1n00)}^G \chi_{(02)}^s + \chi_{(0n01)}^G \chi_{(11)}^s \chi_{(1)}^v) \\
&\quad + t^{8+2n} (\chi_{(1n01)}^G \chi_{(11)}^s + \chi_{(0n10)}^G \chi_{(20)}^s \chi_{(1)}^v + \chi_{(0n02)}^G \chi_{(01)}^s \chi_{(1)}^v) \\
&\quad - t^{9+2n} (\chi_{(1n10)}^G \chi_{(20)}^s + \chi_{(1n02)}^G \chi_{(01)}^s + \chi_{(0n11)}^G \chi_{(10)}^s \chi_{(1)}^v) \\
&\quad \left. + t^{10+2n} (\chi_{(1n11)}^G \chi_{(10)}^s + \chi_{(0n20)}^G \chi_{(1)}^v) - t^{11+2n} \chi_{(1n20)}^G \right),
\end{aligned} \tag{3.17}$$

which was tested against the general formula (2.40) up to t^{20} order. It is the same as (H.20) of [60] after reducing the $Sp(3)_v$ characters

$$\chi_{(001)}^v \rightarrow \chi_{(1)}^v, \quad \chi_{(010)}^v \rightarrow 1, \quad \chi_{(100)}^v \rightarrow 0, \quad \chi_{(000)}^v \rightarrow 0. \tag{3.18}$$

SO(10) We apply our 1-instanton expression (2.40) to $SO(10) + N_s \mathbf{S} + N_c \mathbf{C} + N_v \mathbf{V}$ theory with $(N_s, N_c, N_v) \leq (2, 0, 2), (1, 1, 2), (0, 2, 2), (1, 0, 4), (0, 1, 4)$. The relevant flavor symmetry is $U(N_s + N_c) \times Sp(N_v)$ because the $SO(10)$ (conjugate) spinor is a complex representation. Since the $SO(10)$ charge conjugation exchanges the spinor and conjugate spinor representations, i.e., $\chi_{(00001)}^G = (\chi_{(00010)}^G)^*$, the instanton partition function for $SO(10) + (N_s \mp 1) \mathbf{S} + (N_c \pm 1) \mathbf{C} + N_v \mathbf{V}$ must be identified with that of $SO(10) + N_s \mathbf{S} + N_c \mathbf{C} + N_v \mathbf{V}$ simply by flipping the sign of mass parameters for (conjugate) spinor hypermultiplets.

$$\begin{aligned} \hat{Z}_1^{N_s, N_c, N_v}(m_{1, \dots, N_s}^s; m_{1, \dots, N_c}^c) &= \hat{Z}_1^{N_s-1, N_c+1, N_v}(m_{1, \dots, N_s-1}^s; m_{1, \dots, N_c+1}^c) \Big|_{m_{N_c+1}^c = -m_{N_s}^s} \\ &= \hat{Z}_1^{N_s+1, N_c-1, N_v}(m_{1, \dots, N_s+1}^s; m_{1, \dots, N_c-1}^c) \Big|_{m_{N_s+1}^s = -m_{N_c}^c} \end{aligned} \quad (3.19)$$

This relation was explicitly confirmed in all above cases at 1-instanton order. We may want to compare (2.40) with the known 1-instanton partition function of $SO(10) + 1\mathbf{S} + 1\mathbf{C} + 4\mathbf{V}$ theory, written in (H.21) of [60], after taking relevant mass parameters to infinity. However, (H.21) specifies \hat{Z}_1 only up to $\mathcal{O}(t^5)$, which leaves nothing for comparison once we reduce the mass parameters. Thus the consistency between two expressions could be only weakly tested. For instance, \hat{Z}_1 obtained from (2.40) for $SO(10) + N_s \mathbf{S} + N_c \mathbf{C} + 4\mathbf{V}$ theory with $N_s + N_c = 2$ is displayed in (A.5), which turns out to be void until t^4 order.

SO(12) The 1-instanton partition function of $SO(12) + 1\mathbf{S} + 6\mathbf{V}$ theory is written in (H.22) of [60], up to t^8 order. It can be compared with our 1-instanton formula (2.40) applied to $SO(12) + N_s \mathbf{S} + N_c \mathbf{C} + N_v \mathbf{V}$ theory with $(N_s, N_c, N_v) \leq (1, 0, 4)$ or $(0, 1, 4)$, whose flavor symmetry acting on matter multiplets is $SO(2N_s)_s \times SO(2N_c)_c \times Sp(N_v)_v$. For comparison, we need to appropriately decouple some mass parameters in (H.22) to infinity. It reduces the $Sp(6)_v$ characters in (H.22) to, e.g., the $Sp(4)_v$ irreducible characters as follows:

$$\begin{aligned} \chi_{(000000)}^v &\rightarrow 0, & \chi_{(100000)}^v &\rightarrow 0, & \chi_{(010000)}^v &\rightarrow 1, \\ \chi_{(001000)}^v &\rightarrow \chi_{(1000)}^v, & \chi_{(000100)}^v &\rightarrow \chi_{(0100)}^v, & \chi_{(000001)}^v &\rightarrow \chi_{(0001)}^v \end{aligned} \quad (3.20)$$

The agreement between (H.22) and (2.40) was explicitly confirmed up to the given order, for $(N_s, N_c, N_v) = (1, 0, 4)$. Moreover, we checked that the 1-instanton results Z_1 from (2.40) for $(N_s, N_c, N_v) = (1, 0, N_v)$ and $(0, 1, N_v)$ could be interchanged as follows:

$$\hat{Z}_1^{N_s=1, N_c=0, N_v}(a_1, a_2, a_3, a_4, a_5, a_6) = \hat{Z}_1^{N_s=0, N_c=1, N_v}(a_1, a_2, a_3, a_4, a_5, -a_6). \quad (3.21)$$

We have compared so far the solution Z_1 of the recursion formulae (2.34) with the known 1-instanton partition function for various $SO(N)$ theories with spinor hypermultiplets. The comparison showed consistency for all the examples whose Z_1 had been computed [59, 60]. We also collect the character expansion of the 1-instanton partition function (2.40) in Appendix A for novel $SO(N)$ theories with spinor matters. See Table 3 for the list of character expansions.

3.3 Theories with an exceptional gauge group

Let us continue to apply the recursion formulae (2.34) and the general 1-instanton expression (2.40) to study the instanton partition function of exceptional gauge theories. One can find a sufficient number of recursion formulae (2.34) to fix the k -instanton partition function Z_k , if and only if the gauge theory has the following number of fundamental hypermultiplets. Notice that other representations do not appear in the recent classification of 5d SCFTs [43].

$$\begin{aligned}
N_f &\leq 2 & \text{if } G &= G_2 \\
N_f &\leq 2 & \text{if } G &= F_4 \\
N_f + N_{\bar{f}} &\leq 3 & \text{if } G &= E_6 \\
N_f &\leq 2 & \text{if } G &= E_7 \\
\emptyset & & \text{if } G &= E_8
\end{aligned} \tag{3.22}$$

G₂ A supersymmetric quantum mechanical model was proposed in [59], whose Witten index corresponds to the k -instanton partition function of $G_2 + N_f \mathbf{F}$ theory with $N_f \leq 3$. Its index can be written as the following sum over $SU(3)$ colored Young diagrams:

$$Z_k^{\text{YD}} = \sum_{|\vec{Y}|=k} \prod_{i=1}^3 \prod_{s \in Y_i} \frac{2 \sinh(\phi(s))}{2 \sinh \frac{\epsilon_+ - \phi(s)}{2}} \frac{2 \sinh(\epsilon_+ - \phi(s))}{\prod_{j=1}^3 2 \sinh \frac{E_{ij}}{2}} \frac{\prod_{l=1}^{N_f} 2 \sinh(\frac{m_l \pm \phi(s)}{2})}{2 \sinh \frac{E_{ij} - 2\epsilon_+}{2}} \frac{2 \sinh \frac{\epsilon_+ - \phi(s) - a_j}{2}}{2 \sinh \frac{\epsilon_+ - \phi(s)}{2}} \tag{3.23}$$

$$\begin{aligned}
&\times \prod_{i \leq j}^3 \prod_{\substack{s_{i,j} \in Y_{i,j} \\ s_i < s_j}} \frac{2 \sinh \frac{\phi(s_i) + \phi(s_j)}{2}}{2 \sinh \frac{\epsilon_1 - \phi(s_i) - \phi(s_j)}{2}} \frac{2 \sinh \frac{\phi(s_i) + \phi(s_j) - 2\epsilon_+}{2}}{2 \sinh \frac{\epsilon_2 - \phi(s_i) - \phi(s_j)}{2}} \\
&\tag{3.24}
\end{aligned}$$

Our 1-instanton formula (2.40) was confirmed to agree with the above Z_1^{YD} for all $N_f \leq 2$. Also at two instantons, we explicitly checked that $Z_2^{\text{YD}} = Z_2$, where Z_2 is the solution of the recursion formulae (2.34) with (2.48).

F₄ The 1-instanton partition function of $F_4 + 2\mathbf{F}$ gauge theory was already given in (H.31) of [60], which has $Sp(2)_f$ flavor symmetry. In terms of F_4 and $Sp(2)_f$ characters,

$$\begin{aligned}
\hat{Z}_1 &= t^6 \chi_{(01)}^f + t^7 \chi_{(30)}^f - t^8 (\chi_{(0001)}^G \chi_{(20)}^f + \chi_{(1000)}^G) + t^9 \chi_{(0010)}^G \chi_{(10)}^f - t^{10} \chi_{(0100)}^G \\
&+ \sum_{n=0}^{\infty} \left(t^{8+2n} \chi_{(n000)}^G \chi_{(03)}^f - t^{9+2n} \chi_{(n001)}^G \chi_{(12)}^f + t^{10+2n} (\chi_{(n010)}^G \chi_{(21)}^f + \chi_{(n002)}^G \chi_{(02)}^f) \right. \\
&\quad - t^{11+2n} (\chi_{(n100)}^G \chi_{(30)}^f + \chi_{(n011)}^G \chi_{(11)}^f) + t^{12+2n} (\chi_{(n101)}^G \chi_{(20)}^f + \chi_{(n020)}^G \chi_{(01)}^f) \\
&\quad \left. - t^{13+2n} \chi_{(n110)}^G \chi_{(10)}^f + t^{14+2n} \chi_{(n200)}^G \right). \tag{3.25}
\end{aligned}$$

We compared it with our 1-instanton formula (2.40) and found the agreement up to t^{15} order.

E₆ Let us apply our general 1-instanton expression (2.40) to $E_6 + N_f \mathbf{F} + N_{\bar{f}} \bar{\mathbf{F}}$ gauge theory with $(N_f, N_{\bar{f}}) \leq (3, 0), (2, 1), (1, 2), (0, 3)$ whose flavor symmetry is $U(N_f + N_{\bar{f}})$. Since the fundamental and anti-fundamental representations are interchanged by the E_6 charge conjugation, their instanton partition functions should be identified by inverting the sign of relevant mass parameters. We explicitly confirmed that (2.40) satisfies the relations

$$\begin{aligned} Z_1^{N_f, N_{\bar{f}}}(m_{1, \dots, N_f}^f; m_{1, \dots, N_{\bar{f}}}^{\bar{f}}) &= Z_1^{N_f-1, N_{\bar{f}}+1}(m_{1, \dots, N_f-1}^f; m_{1, \dots, N_{\bar{f}}+1}^{\bar{f}}) \Big|_{m_{N_{\bar{f}}+1}^{\bar{f}} = -m_{N_f}^f} \\ &= Z_1^{N_f+1, N_{\bar{f}}-1}(m_{1, \dots, N_f+1}^f; m_{1, \dots, N_{\bar{f}}-1}^{\bar{f}}) \Big|_{m_{N_f+1}^f = -m_{N_{\bar{f}}}^{\bar{f}}} \end{aligned} \quad (3.26)$$

in all above cases. Furthermore, Z_1 at $(N_f, N_{\bar{f}}) = (3, 0)$ can be compared with (H.35) of [60] which displays the character expansion up to t^{11} order. We checked their consistency except a sign mistake in the second term of (H.35). The full character expansion of Z_1 at $N_f = 3$ and $N_{\bar{f}} = 0$ is written in (A.17), after turning off the E_6 Coulomb VEV $\vec{a} = 0$ for simplicity.

E₇ Our 1-instanton expression (2.40) is applicable to $E_7 + N_f \mathbf{F}$ gauge theory with $N_f \leq 2$, which has $SO(2N_f)_f$ flavor symmetry. We show the full character expansion of Z_1 at $N_f = 2$ in (A.20) after setting $\vec{a} = 0$ to shorten the expression. We also compared our (2.40) applied to the $N_f = 1$ case with (H.40) of [60] and found the agreement up to t^{280} order.

E₈ The 1-instanton partition function of E_8 gauge theory can be written as [14]

$$\hat{Z}_1 = \sum_{n=0}^{\infty} t^{29+2n} \chi_{(000000n0)}^G. \quad (3.27)$$

We confirmed that it agrees with our 1-instanton expression (2.40) up to t^{520} order.

3.4 Theories with a rank-3 antisymmetric hypermultiplet

Another non-trivial test of our blow-up recursion formulae (2.34) is the partition function for 5d $SU(6)$ theory with an hypermultiplet in the rank-3 antisymmetric representation (**TAS**). This theory allows an intriguing Higgs branch with $SU(3) \times SU(3)$ gauge symmetry that can be explicitly checked via the partition function.

To have a UV fixed point, 5d $SU(6)$ theories can have up to 2 hypermultiplets in the rank-3 antisymmetric representation [43]. Their type IIB 5-brane configurations were constructed in [44] with/without O5-planes. In particular, 5-brane web diagrams for $SU(6) + \frac{1}{2} \mathbf{TAS}$ and $SU(6) + 1 \mathbf{TAS}$ do not contain orientifold planes, so that topological vertex method [46, 47] can be straightforwardly applied to compute their partition functions. In [44], for instance, the partition function of $SU(6)_{\frac{5}{2}} + \frac{1}{2} \mathbf{TAS}$ theory was computed up to two instantons using the topological vertex formalism.

Our blow-up equation (2.18) demands all mass parameters to be generically turned on. In particular, we need a mass parameter for the rank-3 antisymmetric hypermultiplet. As one cannot introduce mass for a half-hypermultiplet, let us consider $SU(6)_3$ theory with one full




Figure 1: A 5-brane web for $SU(6)_3$ theory with one massless hypermultiplet in the rank-3 antisymmetric representation.

hypermultiplet in the rank-3 antisymmetric representation ($SU(6) + 1\mathbf{TAS}$). An example for 5-brane web for $SU(6)_3 + 1\mathbf{TAS}$ is depicted in Figure 1. It is instructive to see if Figure 1 is consistent with the expected prepotential. The effective prepotential on the Coulomb branch of a 5d gauge theory with a gauge group G and matter f in a representation R_f is [40]

$$\mathcal{F}(\phi) = \frac{m_0}{2} h_{ij} \phi_i \phi_j + \frac{\kappa}{6} d_{ijk} \phi_i \phi_j \phi_k + \frac{1}{12} \left(\sum_{r \in \text{roots}} |r \cdot \phi|^3 - \sum_f \sum_{w \in R_f} |w \cdot \phi - m_f|^3 \right). \quad (3.28)$$

Here, m_0 is the inverse of the gauge coupling squared, κ is the Chern-Simons level and m_f is a mass parameter for the matter f . r is a root of the Lie algebra \mathfrak{g} associated to G and w is a weight of the representation R_f of \mathfrak{g} . We also defined $h_{ij} = \text{Tr}(T_i T_j)$, $d_{ijk} = \frac{1}{2} \text{Tr}(T_i \{T_j, T_k\})$ where T_i are the Cartan generators of the Lie algebra \mathfrak{g} . With Coulomb branch moduli assigned in Figure 1 and the identification of Weyl chamber for the Coulomb VEV ($a_1 \geq a_2 \geq \dots \geq a_6$, $\sum_{i=1}^6 a_i = 0$),

$$a_1 = \phi_1, \quad a_2 = \phi_2 - \phi_1, \quad a_3 = \phi_3 - \phi_2, \quad a_4 = \phi_4 - \phi_3, \quad a_5 = \phi_5 - \phi_4, \quad a_6 = -\phi_5, \quad (3.29)$$

one finds that the prepotential for $SU(6)_3$ with one massless rank-3 antisymmetry hyper takes the form

$$\begin{aligned} \mathcal{F}_{SU(6)_3+1\mathbf{TAS}} = & m_0 (\phi_1^2 + \phi_2^2 + \phi_3^2 + \phi_4^2 + \phi_5^2 - \phi_1 \phi_2 - \phi_2 \phi_3 - \phi_3 \phi_4 - \phi_4 \phi_5) \\ & + \frac{\phi_1^3}{3} + \frac{4\phi_2^3}{3} + \frac{4\phi_3^3}{3} + \frac{4\phi_4^3}{3} + \frac{4\phi_5^3}{3} + 4\phi_1^2 \phi_2 - 5\phi_1 \phi_2^2 - 2\phi_1 (\phi_3^2 + \phi_4^2 + \phi_5^2) \\ & + \phi_2^2 \phi_3 - 2\phi_2 \phi_3^2 - \phi_3 \phi_4^2 - \phi_4^2 \phi_5 + 2\phi_1 \phi_2 \phi_3 + 2\phi_1 \phi_3 \phi_4 + 2\phi_1 \phi_4 \phi_5. \end{aligned} \quad (3.30)$$

From which, one can easily obtain the monopole string tensions $T_i = \partial \mathcal{F} / \partial \phi_i$ as the areas of the compact faces of the 5-brane web

$$T_1 = \textcircled{1} + 2 \times \textcircled{2}, \quad T_2 = \textcircled{3}, \quad T_3 = \textcircled{4}, \quad T_4 = \textcircled{5}, \quad T_5 = \textcircled{6} + 2 \times \textcircled{7}, \quad (3.31)$$




Figure 2: (a) A Higgsing of $SU(6)_3 + 1\mathbf{TAS}$ into two $SU(3)_3$ theories by aligning internal D5-branes in red. (a) Two different $SU(3)_3$ theories are painted in blue and red, respectively.

where the encircled numbers represent the area of apparent faces in Figure 1. One can then see that the 5-brane web given in Figure 1 is consistent with the prepotential for $SU(6)_3$ with one massless rank-3 antisymmetric hyper.

Notice that this 5-brane web for $SU(6)_3 + 1\mathbf{TAS}$ suggests an intriguing Higgsing of the theory, that is the Higgsing of $SU(6)$ theory with one rank-3 antisymmetric hyper into two pure $SU(3)$ theories. It can be achieved by setting the Coulomb branch parameters as

$$a_5 = -a_1 - a_6, \quad \text{or equivalently} \quad \phi_4 = \phi_1. \quad (3.32)$$

This tuning of the parameters, of course, reduces the dimension of Coulomb brach by one and also opens up a Higgs branch in such a way that the 5-brane web in Figure 1 becomes 5-brane web in Figure 2(a) where the D5-branes in the upper parts of ⑥ and ⑦ are aligned and joint to become a single D5-brane that is paint in red in Figure 2(a). The resulting configuration is then a 5-brane configuration for two pure $SU(3)_3$ theories that are on top of each other, as shown in Figure 2(b). This Higgsing procedure is a 5-brane realization of the Higgsing of $SU(6)_3$ theory with a rank-3 antisymmetric hyper into two pure $SU(3)_3$ theories. It follows that under this Higgsing, the prepotential for $SU(6)_3 + 1\mathbf{TAS}$ (3.30) becomes a sum of prepotentials for two independent pure $SU(3)_3$ theories (without bi-fundamentals):

$$\mathcal{F}_{SU(6)_3+1\mathbf{TAS}} \Big|_{\phi_4=\phi_1} \rightarrow \mathcal{F}_{SU(3)_3}(m_0, \phi_1, \phi_5 - \phi_1, -\phi_5) + \mathcal{F}_{SU(3)_3}(m_0, \phi_2 - \phi_1, \phi_3 - \phi_2, \phi_1 - \phi_3),$$

or,

$$\mathcal{F}_{SU(6)_3+1\mathbf{TAS}} \Big|_{a_1+a_5+a_6=0} \rightarrow \mathcal{F}_{SU(3)_3}(m_0, a_1, a_5, a_6) + \mathcal{F}_{SU(3)_3}(m_0, a_2, a_3, a_4). \quad (3.33)$$

The Higgsing of the $SU(6)$ theory into two copies of $SU(3)$ theories can be understood as follows. This in turn implies that under this Higgsing, the partition function for $SU(6)_3 + 1\mathbf{TAS}$ should be expressed as a product of the partition functions of two pure $SU(3)_3$ theories

$$Z_{SU(6)_3+1\mathbf{TAS}} \Big|_{\text{Higgsing}} \rightarrow Z_{SU(3)_3}(q, A_1, A_5, A_6) Z_{SU(3)_3}(q, A_2, A_3, A_4) Z_{\text{decoupled}}(q), \quad (3.34)$$

where the parameters q and A_i are the Kähler parameters for instanton and Coulomb branch parameters, and $Z_{\text{decoupled}}(q)$ represents the overall extra terms that do not explicitly depend on the Coulomb branch moduli, which would correspond to a new decoupled mode appearing




Figure 3: A labeling of Young diagrams assigned to the horizontal edges of Figure 1.

in Figure 2. In what follows, we explicitly compute the partition function for $SU(6)_3 + 1\text{TAS}$ based on the 5-brane web and compare it with our blowup formula, and also consider this Higgsing as a consistency check.

To compute the instanton partition function based on the 5-brane web for $SU(6) + 1\text{TAS}$ given in Figure 1, we assign the Young diagrams Y_i to each horizontal edge of the web diagram as shown in Figure 3 and use the topological vertex method. For convenience, we restrict to ourselves to the unrefined case where $2\epsilon_+ = \epsilon_1 + \epsilon_2 = 0$. (See also a similar calculation done in [44].) As the web diagram in Figure 1 is left-right symmetric, it is convenient to split the web diagram as the left and right part and glue to obtain the partition function. With the Kähler parameters for Coulomb branch moduli A_i , ($i = 1, \dots, 6$) subject to $\prod_{i=1}^6 A_i = 1$, the instanton q , and the unrefined Ω -deformation parameter g , defined by

$$A_i = e^{-a_i}, \quad q = e^{-\frac{8\pi^2}{g_0^2}}, \quad g = e^{-\epsilon_-}, \quad (3.35)$$

one finds that

$$\begin{aligned} Z_{\text{Nek}} = & \sum_{(Y_1, \dots, Y_6)} q^{\sum_{i=1}^6 |Y_i|} (-A_1^6)^{|Y_1|} (-A_2^6)^{|Y_2|} (-A_2^2 A_3^4)^{|Y_3|} (-A_2^2 A_3^2 A_4^2)^{|Y_4| + |Y_5|} \\ & \times f_{Y_1}(g)^5 f_{Y_2}(g)^5 f_{Y_3}(g)^3 f_{Y_4}(g) f_{Y_5}(g)^{-1} f_{Y_6}(g)^2 Z_{\text{left}}(\vec{Y}) Z_{\text{right}}(\vec{Y}), \end{aligned} \quad (3.36)$$

where $\vec{Y} = (Y_1, Y_2, Y_3, Y_4, Y_5, Y_6)$ and the left and right parts, $Z_{\text{left}}(\vec{Y})$ and $Z_{\text{right}}(\vec{Y})$, are given by

$$\begin{aligned} Z_{\text{left}}(\vec{Y}) = & \sum_{Y_0} (-A_1^{-1} A_6^{-2})^{|Y_0|} g^{\frac{\|Y_0^t\|^2 + \|Y_0\|^2}{2}} \tilde{Z}_{Y_0}^2 f_{Y_0}^2(g) \prod_{i=1}^6 g^{\frac{\|Y_i\|^2}{2}} \tilde{Z}_{Y_i} \\ & \times R_{Y_1 Y_6}^{-1}(A_1 A_6^{-1}) R_{Y_0 Y_6^t}^{-1}(A_1^{-1} A_6^{-2}) R_{Y_1 Y_0^t}^{-1}(A_1^2 A_6) \\ & \times \prod_{2 \leq i < j \leq 5} R_{Y_i Y_j^t}^{-1}(A_i A_j^{-1}) \prod_{i=2}^5 R_{Y_0^t Y_i}(A_1 A_i A_6), \end{aligned} \quad (3.37)$$

and $Z_{\text{right}}(\vec{Y})$ is obtained from $Z_{\text{left}}(\vec{Y})$ by replacing Y_0 with Y_7 . Here, for a Young diagram $\lambda = (\lambda_1, \lambda_2, \dots)$ and its transpose λ^t , $|\lambda| = \sum_i \lambda_i$ and $\|\lambda\|^2 = \sum_i \lambda_i^2$ and

$$\tilde{Z}_\lambda = \prod_{(i,j) \in \lambda} \frac{1}{1 - g^{\lambda_i + \lambda_j^t - i - j + 1}}. \quad (3.38)$$

The framing factor $f_\lambda(g)$ is defined by

$$f_\lambda(g) = (-1)^{|\lambda|} g^{\frac{1}{2} (g^{\|\lambda^t\|^2 - \|\lambda\|^2})}, \quad (3.39)$$

and $R_{\lambda\mu}(Q) = R_{\mu\lambda}(Q)$ is defined by

$$R_{\lambda\mu}(Q) = \text{PE} \left[-\frac{g}{(1-g)^2} Q \right] \times N_{\lambda^t\mu}(Q) \quad (3.40)$$

with PE representing the Plethystic exponential and

$$N_{\lambda\mu}(Q) = \prod_{(i,j) \in \lambda} \left(1 - Q g^{\lambda_i + \mu_j^t - i - j + 1} \right) \prod_{(i,j) \in \mu} \left(1 - Q g^{-\lambda_j^t - \mu_i + i + j - 1} \right). \quad (3.41)$$

The Nekrasov partition function is then expressed as a weighted sum of the instanton partition function

$$Z_{\text{Nek}} = Z_{\text{pert}} \left(1 + \sum_{k=1}^{\infty} q^k Z_k \right), \quad (3.42)$$

where Z_{pert} is the perturbative partition function, while Z_k stands for the k -instanton partition function. The perturbative part of the partition function Z_{pert} comes from the summand of (3.36) at empty Young diagrams, i.e., $(Y_1, Y_2, Y_3, Y_4, Y_5, Y_6) = (\emptyset, \emptyset, \emptyset, \emptyset, \emptyset, \emptyset)$. It is given by

$$\begin{aligned} Z_{\text{pert}} &= Z_{\text{left}}(\emptyset, \emptyset, \emptyset, \emptyset, \emptyset, \emptyset) Z_{\text{right}}(\emptyset, \emptyset, \emptyset, \emptyset, \emptyset, \emptyset) \\ &= \text{PE} \left[\frac{2g}{(1-g)^2} \left(\frac{A_1}{A_6} + \frac{1}{A_1 A_6^2} + A_1^2 A_6 + \sum_{2 \leq i < j \leq 5} \frac{A_i}{A_j} - \sum_{i=2}^5 A_1 A_i A_6 \right) \right] \\ &\quad \times \left(\sum_{Y_0} (-A_1^{-1} A_6^{-2})^{|Y_0|} g^{\frac{\|Y_0^t\|^2 + \|Y_0\|^2}{2}} \tilde{Z}_{Y_0}(g)^2 f_{Y_0}^2(g) \right. \\ &\quad \left. N_{Y_0^t \emptyset}^{-1}(A_1^{-1} A_6^{-2}) N_{Y_0 \emptyset}^{-1}(A_1^2 A_6) \prod_{i=2}^5 N_{Y_0 \emptyset}(A_1 A_i A_6) \right)^2, \end{aligned} \quad (3.43)$$

where the last two lines can be combined into the following closed-form expression:

$$\text{PE} \left[\frac{2g}{(1-g)^2} \left(\sum_{i=2}^5 \frac{A_i}{A_i} + \sum_{i=2}^5 \frac{A_i}{A_6} - \frac{1}{A_1 A_6^2} - A_1^2 A_6 - \sum_{2 \leq i < j \leq 5} A_1 A_i A_j + \mathcal{O}(A_1^6) \right) \right]. \quad (3.44)$$

We note here that when performing the Young diagram sum over Y_0 in (3.43) to compute the Z_{pert} , we expand (3.43) in terms of A_1 and, by $\mathcal{O}(A_1^6)$, we mean that the obtained result

is valid up to $\mathcal{O}(A_1^6)$. As it is very unlikely that there will be a new term which suddenly appears in higher orders than 6 in A_1 , we believe that there are no further terms for $\mathcal{O}(A_1^6)$. It is clear then that (3.43) is manifestly consistent with the equivariant index [41] for 5d $SU(6)$ gauge theory with a hypermultiplet in the rank-3 antisymmetric representation, i.e.,

$$Z_{\text{pert}} = \text{PE} \left[\frac{2g}{(1-g)^2} \left(\sum_{1 \leq i < j \leq 6} \frac{A_i}{A_j} - \sum_{2 \leq i < j \leq 6} A_1 A_i A_j \right) \right]. \quad (3.45)$$

The 1-instanton partition function Z_1 can be obtained from the summands of (3.36) at Young diagrams satisfying $\sum_{i=1}^6 |Y_i| = 1$. There are 6 different profiles of Young diagrams. The configuration $|Y_i| = 1$ and $Y_{j \neq i} = \emptyset$ contribute to Z_1 as

$$+ \frac{g}{(1-g)^2} \frac{A_i^6}{\prod_{j \neq i} (A_i - A_j)^2} \left(-A_i \sum_{j \neq i} A_j + \sum_{j \neq i} \frac{1}{A_j} - \frac{1}{A_i} + A_i^2 \right)^2. \quad (3.46)$$

Summing over all six contributions, one finds

$$Z_1 = \sum_{i=1}^6 \frac{g}{(1-g)^2} \frac{A_i^6}{\prod_{j \neq i} (A_i - A_j)^2} \left(-A_i \sum_{j \neq i} A_j + \sum_{j \neq i} \frac{1}{A_j} - \frac{1}{A_i} + A_i^2 \right)^2. \quad (3.47)$$

which is in agreement with the blowup partition function (eqn). (overall sign: looking at the GV invariant for single W-boson + single instanton (which should be -2), the topological vertex computation seems to be correct. blowup should have an overall sign issue)

We will show the novel results for the $SO(N)$ gauge theories with spinor hypermultiplets in Section 3.2, for the gauge theories with exceptional groups in Section 3.3, and for the $SU(6)$ gauge theories with rank-3 antisymmetric hypermultiplets in Section 3.4. (See Table ???) especially for ‘exceptional’ gauge theories

4 Conclusion

Acknowledgments

We thank Saebyeok Jeong and Seok Kim for discussion. This work is supported in part by the UESTC Research Grant A03017023801317 (SSK), the National Research Foundation of Korea (NRF) Grants 2017R1D1A1B06034369 (KL, JS), and 2018R1A2B6004914 (KHL)

A One-instanton partition functions

This appendix collects the character expansion of the 1-instanton partition function Z_1 for a variety of 5d $\mathcal{N} = 1$ gauge theories. For simplicity, we display the $\hat{Z}_1 \equiv (2 \sinh \frac{\epsilon_{1,2}}{2}) \cdot Z_1$ which takes off the center-of-mass factor. They are written in terms of irreducible characters $\chi_{\mathbf{R}}^S$, whose superscript $S \in \{G, v, s, c, f, \bar{f}\}$ indicates the gauge symmetry (G) or the flavor symmetry acting on the vector (v), spinor (s), conjugate spinor (c), fundamental (f), or anti-fundamental (\bar{f}) hypermultiplets. The representation \mathbf{R} of an irreducible character $\chi_{\mathbf{R}}^S$ is specified by its Dynkin label.¹⁴ An irreducible character for the flavor symmetry is assumed to be in the orthogonal basis, such that it can be consistent with the mass parameters m_ℓ introduced in Section 2. We will often distinguish the mass parameters by the superscript $S \in \{s, c, v, f, \bar{f}\}$ according to the matter representation.

SO(8) The flavor symmetry acting on $N_s \mathbf{S} + N_c \mathbf{C} + N_v \mathbf{V}$ matter multiplets is given by $Sp(N_s)_s \times Sp(N_c)_c \times Sp(N_v)_v$. For $(N_s, N_c, N_v) = (3, 1, 0)$, the character expansion of the 1-instanton result \hat{Z}_1 is

$$\begin{aligned} \hat{Z}_1 = \sum_{n=0}^{\infty} & \left(t^{5+2n} \chi_{(0n00)}^G \chi_{(001)}^s \chi_{(1)}^v - t^{6+2n} (\chi_{(0n01)}^G \chi_{(010)}^s \chi_{(1)}^v + \chi_{(1n00)}^G \chi_{(001)}^s) \right. \\ & + t^{7+2n} (\chi_{(1n01)}^G \chi_{(010)}^s + \chi_{(0n02)}^G \chi_{(100)}^s \chi_{(1)}^v) \\ & \left. - t^{8+2n} (\chi_{(1n02)}^G \chi_{(100)}^s + \chi_{(0n03)}^G \chi_{(1)}^v) + t^{9+2n} \chi_{(1n03)}^G \right). \end{aligned} \quad (\text{A.1})$$

which was compared with the closed-form expression (2.40) up to t^{20} order. We checked that the 1-instanton partition functions Z_1 from (2.40) for $(N_s, N_c, N_v) = (3, 1, 0)$ and $(1, 3, 0)$ could be interchanged as follows:

$$Z_1^{(N_s, N_c, N_v)=(1,3,0)}(a_1, a_2, a_3, a_4) = Z_1^{(N_s, N_c, N_v)=(3,1,0)}(a_1, a_2, a_3, -a_4). \quad (\text{A.2})$$

The $SO(8)$ triality (3.13) was also confirmed as in Section 3.2. Namely, we found that

$$\begin{aligned} Z_1^{(N_s, N_c, N_v)=(3,1,0)}(\vec{a}, \epsilon_1, \epsilon_2; \vec{m}^s, \vec{m}^c, 0) &= Z_1^{(N_s, N_c, N_v)=(0,3,1)}(\vec{a}', \epsilon_1, \epsilon_2; 0, \vec{m}^s, \vec{m}^c)|_{\vec{a}' \rightarrow \vec{a}} \\ &= Z_1^{(N_s, N_c, N_v)=(1,0,3)}(\vec{a}'', \epsilon_1, \epsilon_2; \vec{m}^c, 0, \vec{m}^s)|_{\vec{a}'' \rightarrow \vec{a}} \end{aligned} \quad (\text{A.3})$$

$$\begin{aligned} Z_1^{(N_s, N_c, N_v)=(1,3,0)}(\vec{a}, \epsilon_1, \epsilon_2; \vec{m}^s, \vec{m}^c, 0) &= Z_1^{(N_s, N_c, N_v)=(0,1,3)}(\vec{a}', \epsilon_1, \epsilon_2; 0, \vec{m}^s, \vec{m}^c)|_{\vec{a}' \rightarrow \vec{a}} \\ &= Z_1^{(N_s, N_c, N_v)=(3,0,1)}(\vec{a}'', \epsilon_1, \epsilon_2; \vec{m}^c, 0, \vec{m}^s)|_{\vec{a}'' \rightarrow \vec{a}} \end{aligned} \quad (\text{A.4})$$

The character expansion for other $SO(8)$ theories with less number of hypermultiplets can be obtained from (A.1) by decoupling some mass parameters to infinity. It was checked that the general 1-instanton formula (2.40) agrees with that.

¹⁴In this paper, we follow the convention of LieART [61] to denote the Dynkin label of a representation \mathbf{R} .

SO(10) The flavor symmetry acting on $N_{\mathbf{s}}\mathbf{S} + N_{\mathbf{c}}\mathbf{C} + N_{\mathbf{v}}\mathbf{V}$ hypermultiplets is $U(N_{\mathbf{s}} + N_{\mathbf{c}})_s \times Sp(N_{\mathbf{v}})_v$, reflecting that the $SO(10)$ (conjugate) spinor representation is complex. For $N_{\mathbf{s}} + N_{\mathbf{c}} = 2$ and $N_{\mathbf{v}} = 2$, the character expansion of \hat{Z}_1 is given by

$$\begin{aligned}
\hat{Z}_1 = & t^5(\chi_{(2)_0}^s + \chi_{(01)}^v) + t^6(\chi_{(2)_{-2}}^s + \chi_{(2)_2}^s) - t^7(\chi_{(00010)}^G \chi_{(1)_{-1}}^s \chi_{(10)}^v + \chi_{(00001)}^G \chi_{(1)_1}^s \chi_{(10)}^v + \chi_{(01000)}^G \\
& + \chi_{(10000)}^G (\chi_{(2)_{-2}}^s + \chi_{(2)_2}^s)) + t^8(\chi_{(00100)}^G \chi_{(10)}^v + \chi_{(10010)}^G \chi_{(1)_{-1}}^s + \chi_{(10001)}^G \chi_{(1)_1}^s) - t^9 \chi_{(10100)}^G \\
& + \sum_{n=0}^{\infty} \left(t^{7+2n} \chi_{(0n000)}^G (\chi_{(0)_{-4}}^s + \chi_{(0)_4}^s + \chi_{(4)_0}^s) \chi_{(01)}^v - t^{8+2n} (\chi_{(0n010)}^G (\chi_{(1)_{-3}}^s + \chi_{(3)_1}^s) \chi_{(01)}^v \right. \\
& \quad + \chi_{(0n001)}^G (\chi_{(1)_3}^s + \chi_{(3)_{-1}}^s) \chi_{(01)}^v + \chi_{(1n000)}^G (\chi_{(0)_{-4}}^s + \chi_{(0)_4}^s + \chi_{(4)_0}^s) \chi_{(10)}^v) \\
& + t^{9+2n} (\chi_{(0n100)}^G (\chi_{(2)_{-2}}^s + \chi_{(2)_2}^s) \chi_{(01)}^v + \chi_{(0n020)}^G \chi_{(0)_{-2}}^s \chi_{(01)}^v + \chi_{(0n002)}^G \chi_{(0)_2}^s \chi_{(01)}^v \\
& \quad + \chi_{(0n011)}^G \chi_{(2)_0}^s \chi_{(01)}^v + \chi_{(1n010)}^G (\chi_{(1)_{-3}}^s + \chi_{(3)_1}^s) \chi_{(10)}^v + \chi_{(1n001)}^G (\chi_{(1)_3}^s + \chi_{(3)_{-1}}^s) \chi_{(10)}^v \\
& \quad + \chi_{(2n000)}^G (\chi_{(0)_{-4}}^s + \chi_{(0)_4}^s + \chi_{(4)_0}^s)) \\
& - t^{10+2n} (\chi_{(0n110)}^G \chi_{(1)_{-1}}^s \chi_{(01)}^v + \chi_{(0n101)}^G \chi_{(1)_1}^s \chi_{(01)}^v + \chi_{(1n100)}^G (\chi_{(2)_{-2}}^s + \chi_{(2)_2}^s) \chi_{(10)}^v \\
& \quad + \chi_{(1n020)}^G \chi_{(0)_{-2}}^s \chi_{(10)}^v + \chi_{(1n002)}^G \chi_{(0)_2}^s \chi_{(10)}^v + \chi_{(1n011)}^G \chi_{(2)_0}^s \chi_{(10)}^v \\
& \quad + \chi_{(2n010)}^G (\chi_{(1)_{-3}}^s + \chi_{(3)_1}^s) + \chi_{(2n001)}^G (\chi_{(1)_3}^s + \chi_{(3)_{-1}}^s) \\
& + t^{11+2n} (\chi_{(1n200)}^G \chi_{(01)}^v + \chi_{(1n110)}^G \chi_{(1)_{-1}}^s \chi_{(10)}^v + \chi_{(1n101)}^G \chi_{(1)_1}^s \chi_{(10)}^v + \chi_{(2n100)}^G (\chi_{(2)_{-2}}^s + \chi_{(2)_2}^s) \\
& \quad + \chi_{(2n020)}^G \chi_{(0)_{-2}}^s + \chi_{(2n002)}^G \chi_{(0)_2}^s + \chi_{(2n011)}^G \chi_{(2)_0}^s) \\
& \left. - t^{12+2n} (\chi_{(1n200)}^G \chi_{(10)}^v + \chi_{(2n110)}^G \chi_{(1)_{-1}}^s + \chi_{(2n101)}^G \chi_{(1)_1}^s) + t^{13+2n} \chi_{(2n200)}^G \right) \quad (\text{A.5})
\end{aligned}$$

where the $U(2)$ character $\chi_{(j)_b}^s$ is defined as (with $y_{s,i} \equiv e^{-m_i^s}$ and $y_{c,i} \equiv e^{-m_i^c}$ understood)

$$\chi_{(j)_b}^s = \begin{cases} (y_{s,1} y_{s,2})^{b/2} \cdot \sum_{a=0}^j (y_{s,1}/y_{s,2})^{-j/2+a} & \text{for } (N_{\mathbf{s}}, N_{\mathbf{c}}) = (2, 0) \\ (y_{s,1}/y_{c,1})^{b/2} \cdot \sum_{a=0}^j (y_{s,1} y_{c,1})^{-j/2+a} & \text{for } (N_{\mathbf{s}}, N_{\mathbf{c}}) = (1, 1) \\ (y_{c,1} y_{c,2})^{-b/2} \cdot \sum_{a=0}^j (y_{c,1}/y_{c,2})^{-j/2+a} & \text{for } (N_{\mathbf{s}}, N_{\mathbf{c}}) = (0, 2). \end{cases} \quad (\text{A.6})$$

Similarly, for $N_{\mathbf{s}} + N_{\mathbf{c}} = 3$ and $N_{\mathbf{v}} = 0$, the character expansion of \hat{Z}_1 is given by

$$\begin{aligned}
\hat{Z}_1 = & t^5(\chi_{(10000)}^G + \chi_{(20)_{-2}}^s + \chi_{(02)_2}^s) - t^6(\chi_{(10)_{-1}}^s + \chi_{(01)_1}^s) + t^7 \chi_{(00100)}^G \\
& + \sum_{n=0}^{\infty} \left(t^{7+2n} (\chi_{(0n000)}^G (\chi_{(00)_{-6}}^s + \chi_{(00)_6}^s + \chi_{(04)_{-2}}^s + \chi_{(40)_2}^s)) \right. \\
& \quad - t^{8+2n} (\chi_{(0n010)}^G (\chi_{(01)_{-5}}^s + \chi_{(30)_3}^s + \chi_{(13)_{-1}}^s) + \chi_{(0n001)}^G (\chi_{(10)_5}^s + \chi_{(03)_{-3}}^s + \chi_{(31)_1}^s)) \\
& \quad + t^{9+2n} (\chi_{(0n100)}^G (\chi_{(02)_{-4}}^s + \chi_{(20)_4}^s + \chi_{(22)_0}^s) + \chi_{(0n011)}^G (\chi_{(12)_{-2}}^s + \chi_{(21)_2}^s) \\
& \quad \quad + \chi_{(0n020)}^G (\chi_{(10)_{-4}}^s + \chi_{(03)_0}^s) + \chi_{(0n002)}^G (\chi_{(01)_4}^s + \chi_{(30)_0}^s)) \\
& \quad - t^{10+2n} (\chi_{(0n110)}^G (\chi_{(11)_{-3}}^s + \chi_{(12)_1}^s) + \chi_{(0n101)}^G (\chi_{(11)_3}^s + \chi_{(21)_{-1}}^s) \\
& \quad \quad + \chi_{(0n030)}^G \chi_{(00)_{-3}}^s + \chi_{(0n003)}^G \chi_{(00)_3}^s + \chi_{(0n021)}^G \chi_{(02)_{-1}}^s + \chi_{(0n012)}^G \chi_{(20)_1}^s) \\
& \quad \left. + t^{11+2n} (\chi_{(0n200)}^G (\chi_{(20)_{-2}}^s + \chi_{(02)_2}^s) + \chi_{(0n120)}^G \chi_{(01)_{-2}}^s + \chi_{(0n102)}^G \chi_{(10)_2}^s + \chi_{0n111}^G \chi_{(11)_0}^s) \right) \quad (\text{A.7})
\end{aligned}$$

$$-t^{12+2n}(\chi_{(0n210)}^G \chi_{(10)_{-1}}^s + \chi_{(0n201)}^G \chi_{(01)_1}^s) + t^{13+2n} \chi_{(0n300)}^G)$$

where the $U(3)$ character $\chi_{(mn)_c}^s$ is defined as

$$\chi_{(mn)_c}^s = (w_1 w_2 w_3)^{\frac{c-m+n}{3}} \cdot \left(\sum_{\substack{1 \leq i_1 \leq \dots \leq i_m \leq 3 \\ 1 \leq j_1 \leq \dots \leq j_n \leq 3}} \frac{w_{i_1} \cdots w_{i_m}}{w_{j_1} \cdots w_{j_n}} - \sum_{\substack{1 \leq i_1 \leq \dots \leq i_{m-1} \leq 3 \\ 1 \leq j_1 \leq \dots \leq j_{n-1} \leq 3}} \frac{w_{i_1} \cdots w_{i_{m-1}}}{w_{j_1} \cdots w_{j_{n-1}}} \right) \quad (\text{A.8})$$

with

$$(w_1, w_2, w_3) = \begin{cases} (y_{s,1}, y_{s,2}, y_{s,3}) & \text{for } (N_s, N_c) = (3, 0) \\ (y_{s,1}, y_{s,2}, y_{c,1}^{-1}) & \text{for } (N_s, N_c) = (2, 1) \\ (y_{s,1}, y_{c,1}^{-1}, y_{c,2}^{-1}) & \text{for } (N_s, N_c) = (1, 2) \\ (y_{c,1}^{-1}, y_{c,2}^{-1}, y_{c,3}^{-1}) & \text{for } (N_s, N_c) = (0, 3) \end{cases} \quad (\text{A.9})$$

Again, (A.5) and (A.7) was tested against the closed-form expression (2.40) up to t^{20} order.

SO(11) The flavor symmetry acting on $N_s \mathbf{S} + N_v \mathbf{V}$ hypermultiplets is $SO(2N_s)_s \times Sp(N_v)_v$. For $N_s = 1$ and $N_v = 3$, the character expansion of \hat{Z}_1 can be written as

$$\begin{aligned} \hat{Z}_1 = & t^5 + t^6(\chi_{(001)}^v + (y_s^2 + y_s^{-2})\chi_{(100)}^v) + t^7((y_s^2 + y_s^{-2} + 1)\chi_{(010)}^v - (y_s^2 + y_s^{-2})\chi_{(10000)}^G) \\ & - t^8((y_s + y_s^1)\chi_{(00001)}^G \chi_{(010)}^v + (y_s^2 + y_s^{-2} + 1)\chi_{(10000)}^G \chi_{(100)}^v + \chi_{(01000)}^G \chi_{(100)}^v) \\ & + t^9(\chi_{(00100)}^G \chi_{(010)}^v + \chi_{(10001)}^G (y_s + y_s^{-1})\chi_{(100)}^v + \chi_{(20000)}^G (y_s^2 + y_s^{-2} + 1) + \chi_{(11000)}^G) \\ & - t^{10}(\chi_{(10100)}^G \chi_{(100)}^v + \chi_{(20001)}^G (y_s + y_s^{-1})) + t^{11}\chi_{(20100)}^G \\ & + \sum_{n=0}^{\infty} \left(t^{8+2n}(\chi_{(0n000)}^G (y_s^4 + y_s^{-4} + 1)\chi_{(001)}^v) \right. \\ & - t^{9+2n}(\chi_{(0n001)}^G (y_s^3 + y_s^{-3})\chi_{(001)}^v + \chi_{(0n001)}^G (y_s + y_s^{-1})\chi_{(001)}^v + \chi_{(1n000)}^G (y_s^4 + y_s^{-4} + 1)\chi_{(010)}^v) \\ & + t^{10+2n}(\chi_{(0n010)}^G (y_s^2 + y_s^{-2})\chi_{(001)}^v + \chi_{(0n100)}^G (y_s^2 + y_s^{-2} + 1)\chi_{(001)}^v + \chi_{(0n002)}^G \chi_{(001)}^v \\ & \quad + \chi_{(1n001)}^G (y_s^3 + y_s + y_s^{-1} + y_s^{-3})\chi_{(010)}^v + \chi_{(2n000)}^G (y_s^4 + y_s^{-4} + 1)\chi_{(100)}^v) \\ & - t^{11+2n}(\chi_{(0n101)}^G (y_s + y_s^{-1})\chi_{(001)}^v + \chi_{(1n100)}^G (y_s^2 + y_s^{-2} + 1)\chi_{(010)}^v + \chi_{(1n010)}^G (y_s^2 + y_s^{-2})\chi_{(010)}^v \\ & \quad + \chi_{(1n002)}^G \chi_{(010)}^v + \chi_{(2n001)}^G (y_s^3 + y_s + y_s^{-1} + y_s^{-3})\chi_{(100)}^v + \chi_{(3n000)}^G (y_s^4 + y_s^{-4} + 1)) \\ & + t^{12+2n}(\chi_{(0n200)}^G \chi_{(001)}^v + \chi_{(1n101)}^G (y_s + y_s^{-1})\chi_{(010)}^v + \chi_{(2n100)}^G (y_s^2 + y_s^{-2} + 1)\chi_{(100)}^v \\ & \quad + \chi_{(2n010)}^G (y_s^2 + y_s^{-2})\chi_{(100)}^v + \chi_{(2n002)}^G \chi_{(100)}^v + \chi_{(3n001)}^G (y_s^3 + y_s + y_s^{-1} + y_s^{-3})) \\ & - t^{13+2n}(\chi_{(1n200)}^G \chi_{(010)}^v + \chi_{(2n101)}^G (y_s + y_s^{-1})\chi_{(100)}^v + \chi_{(3n100)}^G (y_s^2 + y_s^{-2} + 1) \\ & \quad + \chi_{(3n010)}^G (y_s^2 + y_s^{-2}) + \chi_{(3n002)}^G) \\ & \left. + t^{14+2n}(\chi_{(2n200)}^G \chi_{(100)}^v + \chi_{(3n101)}^G (y_s + y_s^{-1})) - t^{15+2n}\chi_{(3n200)}^G \right) \end{aligned} \quad (\text{A.10})$$

which was compared with the closed-form expression (2.40) up to t^{20} order.

SO(12) The flavor symmetry acting on $N_s \mathbf{S} + N_c \mathbf{C} + N_v \mathbf{V}$ hypermultiplets is $SO(2N_s)_s \times SO(2N_c)_c \times Sp(N_v)_v$. Here we turn off the Coulomb VEV $\vec{a} = 0$ for simplicity. The character expansion of \hat{Z}_1 at $(N_s, N_c, N_v) = (2, 0, 0)$ can be written as

$$\begin{aligned} \hat{Z}_1 = \frac{t^{18}}{(1-t^2)^{18}} & \left(-96096 (\chi_{(13)}^s + \chi_{(31)}^s) \cdot (7t^4 + 42t^2 + 72 + 42t^{-2} + 7t^{-4}) \right. \\ & + 10010 (\chi_{(24)}^s + \chi_{(42)}^s) \cdot (9t^5 + 88t^3 + 243t + 243t^{-1} + 88t^{-3} + 9t^{-5}) \\ & - 352 (\chi_{(15)}^s + \chi_{(51)}^s) \cdot (25t^6 + 474t^4 + 2169t^2 + 3504 + 2169t^{-2} + 474t^{-4} + 25t^{-6}) \\ & - 2464 (\chi_{(35)}^s + \chi_{(53)}^s) \cdot (2t^6 + 27t^4 + 108t^2 + 168 + 108t^{-2} + 27t^{-4} + 2t^{-6}) \\ & + 11 (\chi_{(06)}^s + \chi_{(60)}^s) \cdot (42t^7 + 1194t^5 + 8451t^3 + 21253t + 21253t^{-1} + \dots + 42t^{-7}) \\ & + 11 (\chi_{(26)}^s + \chi_{(62)}^s) \cdot (45t^7 + 1101t^5 + 6983t^3 + 16623t + 16623t^{-1} + \dots + 45t^{-7}) \\ & - 32 (\chi_{(17)}^s + \chi_{(71)}^s) \cdot (t^8 + 36t^6 + 336t^4 + 1176t^2 + 1764 + 1176t^{-2} + \dots + t^{-8}) \\ & + 99 \chi_{(22)}^s \cdot (5t^9 - 90t^7 + 1623t^5 + 26743t^3 + 83103t + (t \rightarrow t^{-1})) \\ & + 462 (\chi_{(02)}^s + \chi_{(20)}^s) \cdot (t^9 - 18t^7 + 153t^5 + 4059t^3 + 13485t + (t \rightarrow t^{-1})) \\ & - 32 \chi_{(33)}^s \cdot (t^{10} - 18t^8 + 450t^6 + 13340t^4 + 66977t^2 + 110772 + 66977t^{-2} + \dots + t^{-10}) \\ & + \chi_{(44)}^s \cdot (t^{11} - 18t^9 + 615t^7 + 26332t^5 + 187749t^3 + 466001t + 466001t^{-1} + \dots + t^{-11}) \\ & + (\chi_{(04)}^s + \chi_{(40)}^s) \cdot (t^{13} - 18t^{11} + 153t^9 - 816t^7 + 58115t^5 + 730170t^3 + 2129595t + (t \rightarrow t^{-1})) \\ & - 352 \chi_{(11)}^s \cdot (t^{10} - 4t^8 - 99t^6 + 2496t^4 + 18246t^2 + 32976 + 18246t^{-2} + \dots + t^{-10}) \\ & + (\chi_{(08)}^s + \chi_{(80)}^s) \cdot (t^9 + 48t^7 + 603t^5 + 2898t^3 + 6174t + (t \rightarrow t^{-1})) \\ & \left. + (t^{17} - 18t^{15} + 153t^{13} - 739t^{11} + 3753t^9 - 20195t^7 + 49881t^5 + 1203597t^3 + 4481279t + (t \rightarrow t^{-1})) \right) \end{aligned} \quad (\text{A.11})$$

It was explicitly checked that the 1-instanton partition function Z_1 at $(N_s, N_c, N_v) = (0, 2, 0)$ could be identified with the above as

$$Z_1^{(N_s, N_c, N_v)=(0,2,0)}(a_1, a_2, a_3, a_4, a_5, a_6) = Z_1^{(N_s, N_c, N_v)=(2,0,0)}(a_1, a_2, a_3, a_4, a_5, -a_6). \quad (\text{A.12})$$

Similarly, the character expansion of \hat{Z}_1 at $(N_s, N_c, N_v) = (1, 1, 0)$ can be displayed as follows:

$$\begin{aligned} \hat{Z}_1 = \frac{t^{18}}{(1-t^2)^{18}} \sum_{\pm} & \left(-2462 (y_s^{\pm 1} y_c^{\pm 4} + y_s^{\pm 4} y_c^{\pm 1}) \cdot (2t^6 + 27t^4 + 108t^2 + 168 + 108t^{-2} + 27t^{-4} + 2t^{-6}) \right. \\ & + 11 (y_s^{\pm 2} y_c^{\pm 4} + y_s^{\pm 4} y_c^{\pm 2}) \cdot (45t^7 + 1101t^5 + 6983t^3 + 16623t + (t \rightarrow t^{-1})) \\ & + 44 (y_s^{\pm 3} y_c^{\pm 3}) \cdot (23t^7 + 587t^5 + 3925t^3 + 9609t + (t \rightarrow t^{-1})) \\ & + 44 (y_s^{\pm 1} y_c^{\pm 3} + y_s^{\pm 3} y_c^{\pm 1}) \cdot (23t^7 + 2927t^5 + 26025t^3 + 70033t + (t \rightarrow t^{-1})) \\ & - 32 (y_s^{\pm 3} y_c^{\pm 4} + y_s^{\pm 4} y_c^{\pm 3}) \cdot (t^8 + 36t^6 + 336t^4 + 1176t^2 + 1764 + 1176t^{-2} + \dots + t^{-8}) \\ & - 32 (y_s^{\pm 2} y_c^{\pm 3} + y_s^{\pm 3} y_c^{\pm 2}) \cdot (t^8 + 465t^6 + 7629t^4 + 33351t^2 + 53244 + 33351t^{-2} + \dots + t^{-8}) \\ & + (y_s^{\pm 4} y_c^{\pm 4}) \cdot (t^9 + 48t^7 + 603t^5 + 2898t^3 + 6174t + (t \rightarrow t^{-1})) \\ & \left. - 32 (y_s^{\pm 1} y_c^{\pm 2} + y_s^{\pm 2} y_c^{\pm 1}) \cdot (t^{10} - 17t^8 + 1069t^6 + 44069t^4 + 234770t^2 + 393168 + 234770t^{-2} + \dots + t^{-10}) \right) \end{aligned} \quad (\text{A.13})$$

$$\begin{aligned}
& -32(y_s^{\pm 3} + y_c^{\pm 3}) \cdot (t^{10} - 17t^8 + 750t^6 + 17526t^4 + 83553t^2 + 136714 + 83358t^{-2} + \dots + t^{-10}) \\
& -32(y_s^{\pm 1} + y_c^{\pm 1}) \cdot (13t^{10} - 79t^8 + 408t^6 + 97724t^4 + 587351t^2 + 1011546 + 587351t^{-2} + \dots + 13t^{-10}) \\
& + (y_s^{\pm 4} + y_c^{\pm 4}) \cdot (t^{11} - 18t^9 + 615t^7 + 26332t^5 + 187749t^3 + 466001t + (t \rightarrow t^{-1})) \\
& + (y_s^{\pm 2} + y_c^{\pm 2}) \cdot (t^{11} + 477t^9 - 7305t^7 + 391411t^5 + 4750692t^3 + 13923764t + (t \rightarrow t^{-1})) \\
& + 4(y_s^{\pm 1} y_c^{\pm 1}) \cdot (3t^{11} + 199t^9 + 132676t^7 + 1864041t^5 + 5630341t^3 + 5630341t + (t \rightarrow t^{-1})) \\
& + (y_s^{\pm 2} y_c^{\pm 2})(t^{13} - 17t^{11} + 1136t^9 + 804t^7 + 200385t^5 + 1971471t^3 + 5450836t + (t \rightarrow t^{-1})) \\
& + (t^{15} - 17t^{13} + 214t^{11} + 1414t^9 - 33152t^7 + 704404t^5 + 11381979t^3 + 35592757t + (t \rightarrow t^{-1}))
\end{aligned}$$

in which \sum_{\pm} notation is understood as follows: $\sum_{\pm} x^{\pm 1} y^{\pm 1} = xy + xy^{-1} + x^{-1}y + x^{-1}y^{-1}$, $\sum_{\pm} x^{\pm 1} = x + x^{-1}$, and $\sum_{\pm} 1 = 1$.

SO(13) The flavor symmetry on $N_s \mathbf{S} + N_v \mathbf{V}$ matter multiplets is $SO(2N_s)_s \times Sp(N_v)_v$. The character expansion of \hat{Z}_1 at $(N_s, N_v) = (1, 1)$ is written below, after setting the Coulomb VEV $\vec{a} = 0$ to keep the expression concise.

$$\begin{aligned}
\hat{Z}_1 = & \frac{t^{20}}{(1-t^2)^{20}} \sum_{\pm} \left(y_s^{\pm 8} \chi_{(1)}^V \cdot (t^{10} + 58t^8 + 905t^6 + 5580t^4 + 15876t^2 + 22344 + 15876t^{-2} + \dots + t^{-10}) \right. \\
& - 64 y_s^{\pm 7} \chi_{(1)}^V \cdot (t^9 + 45t^7 + 540t^5 + 2520t^3 + 5292t + (t \rightarrow t^{-1})) \\
& + 26 y_s^{\pm 6} \chi_{(1)}^V \cdot (77t^8 + 2541t^6 + 22226t^4 + 74811t^2 + 110770 + 74811t^{-2} + \dots + 77t^{-8}) \\
& - 5824 y_s^{\pm 5} \chi_{(1)}^V \cdot (7t^7 + 154t^5 + 924t^3 + 2145t + (t \rightarrow t^{-1})) \\
& + y_s^{\pm 4} \chi_{(1)}^V \cdot (t^{14} - 19t^{12} + 170t^{10} + 766t^8 + 576628t^6 + 7601283t^4 + 29870761t^2 \\
& \quad \quad \quad + 46175700 + 29870761t^{-2} + 7601283t^{-4} + \dots + t^{-14}) \\
& - 64 y_s^{\pm 3} \chi_{(1)}^V \cdot (t^{11} - 20t^9 + 1256t^7 + 83074t^5 + 628311t^3 + 1580032t + (t \rightarrow t^{-1})) \\
& + 2002 y_s^{\pm 2} \chi_{(1)}^V \cdot (t^{10} - 19t^8 + 756t^6 + 15006t^4 + 66051t^2 + 105146 + 66051t^{-2} + \dots + t^{-10}) \\
& - 64 y_s^{\pm 1} \chi_{(1)}^V \cdot (13t^{11} - 51t^9 - 436t^7 + 182670t^5 + 1603925t^3 + 4218449t + (t \rightarrow t^{-1})) \\
& + \chi_{(1)}^V \cdot (t^{16} - 19t^{14} + 274t^{12} + 3185t^{10} - 73808t^8 + 1918679t^6 + 46355974t^4 + 212905247t^2 \\
& \quad \quad \quad + 342439014 + 212905247t^{-2} + 46355974t^{-4} + \dots + t^{-16}) \\
& - 13 y_s^{\pm 8} \cdot (t^9 + 35t^7 + 365t^5 + 1575t^3 + 3192t + (t \rightarrow t^{-1})) \\
& + 256 y_s^{\pm 7} \cdot (3t^8 + 80t^6 + 630t^4 + 2016t^2 + 2940 + 2016t^{-2} + \dots + 3t^{-8}) \\
& - 26 y_s^{\pm 6} \cdot (847t^7 + 15989t^5 + 89887t^3 + 203357t + (t \rightarrow t^{-1})) \\
& - 64 y_s^{\pm 5} \cdot (t^{10} - 20t^8 - 6180t^6 - 75228t^4 - 286725t^2 - 439416 - 286725t^{-2} + \dots + t^{-10}) \\
& + y_s^{\pm 4} \cdot (t^{13} - 20t^{11} + 2907t^9 - 74785t^7 - 4557934t^5 - 33690015t^3 - 83955034t + (t \rightarrow t^{-1})) \\
& - 64 y_s^{\pm 3} \cdot (t^{12} - 19t^{10} + 807t^8 - 24636t^6 - 510121t^4 - 2255129t^2 - 3592422 \\
& \quad \quad \quad - 2255129t^{-2} - 510121t^{-4} - 24636t^{-6} + \dots + t^{-12}) \\
& + 2 y_s^{\pm 2} \cdot (7t^{13} - 140t^{11} + 3189t^9 + 86972t^7 - 7685485t^5 - 71293018t^3 - 190116261t + (t \rightarrow t^{-1})) \\
& - 64 y_s^{\pm 1} \cdot (t^{12} - 84t^{10} + 2667t^8 - 36526t^6 - 1227485t^4 - 5926190t^2 - 9643046
\end{aligned}$$

$$\begin{aligned}
& -5926190t^{-2} - 1227485t^{-4} - 36526t^{-6} + \dots + t^{-12}) \\
& + (t^{15} - 20t^{13} - 602t^{11} + 5691t^9 + 495005t^7 - 22183672t^5 - 225823570t^3 - 617150913t + (t \rightarrow t^{-1})) \Big) \\
& \tag{A.14}
\end{aligned}$$

SO(14) The classical flavor symmetry on $N_{\mathbf{s}}\mathbf{S} + N_{\mathbf{c}}\mathbf{C} + N_{\mathbf{v}}\mathbf{V}$ hypermultiplets is $U(N_{\mathbf{s}})_s \times U(N_{\mathbf{c}})_c \times Sp(N_{\mathbf{v}})_v$. The character expansion of \hat{Z}_1 at $(N_{\mathbf{s}}, N_{\mathbf{c}}, N_{\mathbf{v}}) = (1, 0, 2)$ is written below, after turning off the $SO(14)$ Coulomb VEV $\vec{a} = 0$.

$$\begin{aligned}
\hat{Z}_1 = & \frac{t^{22}}{(1-t^2)^{22}} \sum_{\pm} \Big(y_s^{\pm 8} \chi_{(01)}^V \cdot (t^{11} + 69t^9 + 1309t^7 + 10065t^5 + 36828t^3 + 69300t + (t \rightarrow t^{-1})) \\
& - 64 y_s^{\pm 7} \chi_{(01)}^V \cdot (t^{10} + 55t^8 + 825t^6 + 4950t^4 + 13860t^2 + 19404 + 13860t^{-2} + \dots + t^{-10}) \\
& + 26 y_s^{\pm 6} \chi_{(01)}^V \cdot (77t^9 + 3234t^7 + 36667t^5 + 164401t^3 + 338261t + (t \rightarrow t^{-1})) \\
& - 5824 y_s^{\pm 5} \chi_{(01)}^V \cdot (7t^8 + 210t^6 + 1694t^4 + 5434t^2 + 7920 + 5434t^{-2} + \dots + 7t^{-8}) \\
& + y_s^{\pm 4} \chi_{(01)}^V \cdot (t^{15} - 22t^{13} + 231t^{11} - 1540t^9 + 614558t^7 \\
& \quad + 11510191t^5 + 62671224t^3 + 139186397t + (t \rightarrow t^{-1})) \\
& - 832 y_s^{\pm 3} \chi_{(01)}^V \cdot (33t^8 + 7744t^6 + 83776t^4 + 300104t^2 + 451192 + 300104t^{-2} + \dots + 33t^{-8}) \\
& + 2002 y_s^{\pm 2} \chi_{(01)}^V \cdot (t^{11} - 22t^9 + 621t^7 + 21262t^5 + 134245t^3 + 314181t + (t \rightarrow t^{-1})) \\
& - 832 y_s^{\pm 1} \chi_{(01)}^V \cdot (t^{12} - t^{10} - 231t^8 + 15631t^6 + 206987t^4 + 790240t^2 + 1207976 + \\
& \quad + 790240t^{-2} + 206987t^{-4} + 15631t^{-6} + \dots + t^{-12}) \\
& + \chi_{(01)}^V \cdot (t^{17} - 22t^{15} + 335t^{13} + 3179t^{11} - 84595t^9 + 1320011t^7 \\
& \quad + 63966077t^5 + 427850621t^3 + 1020096033t + (t \rightarrow t^{-1})) \\
& - 14 y_s^{\pm 8} \chi_{(10)}^V \cdot (t^{10} + 42t^8 + 539t^6 + 2948t^4 + 7854t^2 + 10824 + 7854t^{-2} + \dots + t^{-10}) \\
& + 832 y_s^{\pm 7} \chi_{(10)}^V \cdot (t^9 + 33t^7 + 330t^5 + 1386t^3 + 2772t + (t \rightarrow t^{-1})) \\
& - 2184 y_s^{\pm 6} \chi_{(10)}^V \cdot (11t^8 + 270t^6 + 2002t^4 + 6182t^2 + 8910 + 6182t^{-2} + \dots + 11t^{-8}) \\
& + 5824 y_s^{\pm 5} \chi_{(10)}^V \cdot (77t^7 + 1281t^5 + 6677t^3 + 14575t + (t \rightarrow t^{-1})) \\
& + 52 y_s^{\pm 4} \chi_{(10)}^V \cdot (33t^{10} - 726t^8 - 109153t^6 - 1133396t^4 - 3996580t^2 - 5980436 \\
& \quad - 3996580t^{-2} - 1133396t^{-4} - 109153t^{-6} + \dots + 33t^{-10}) \\
& - 64 y_s^{\pm 3} \chi_{(10)}^V \cdot (t^{13} - 22t^{11} + 868t^9 - 20559t^7 - 726341t^5 - 4583956t^3 - 10718569t + (t \rightarrow t^{-1})) \\
& + 8008 y_s^{\pm 2} \chi_{(10)}^V \cdot (49t^8 - 2102t^6 - 29678t^4 - 115094t^2 - 176638 - 115094t^{-2} + \dots + 49t^{-8}) \\
& + 4928 y_s^{\pm 1} \chi_{(10)}^V \cdot (t^{11} - 35t^9 + 217t^7 + 21505t^5 + 152866t^3 + 371316t + (t \rightarrow t^{-1})) \\
& - 8 \chi_{(10)}^V \cdot t^{10} (112t^{12} - 189t^{10} - 104258t^8 + 2855160t^6 + 46213090t^4 + 185620270t^2 \\
& \quad + 287407450 + 185620270t^{-2} + 46213090t^{-4} + \dots + 112t^{-12}) \\
& + 13 y_s^{\pm 8} \cdot (8t^9 + 229t^7 + 2101t^5 + 8393t^3 + 16401t + (t \rightarrow t^{-1})) \\
& - 5824 y_s^{\pm 7} \cdot (t^8 + 22t^6 + 154t^4 + 462t^2 + 660 + 462t^{-2} + \dots + t^{-8})
\end{aligned}$$

$$\begin{aligned}
& + 26 y_s^{\pm 6} \cdot (6075t^7 + 95425t^5 + 483483t^3 + 1042937t + (t \rightarrow t^{-1})) \\
& - 64 y_s^{\pm 5} \cdot (t^{12} - 22t^{10} + 231t^8 + 41580t^6 + 427575t^4 + 1498244t^2 + 2237312 \\
& \quad + 1498244t^{-2} + 427575t^{-4} + 41580t^{-6} + \dots + t^{-12}) \\
& + 91 y_s^{\pm 4} \cdot (11t^{11} - 473t^9 + 7623t^7 + 312675t^5 + 2010490t^3 + 4723994t + (t \rightarrow t^{-1})) \\
& + 5824 y_s^{\pm 3} \cdot (77t^8 - 2046t^6 - 32546t^4 - 129768t^2 - 200508 - 129768t^{-2} + \dots + 77t^{-8}) \\
& + 2 y_s^{\pm 2} \cdot (7t^{15} - 154t^{13} + 2475t^{11} - 93720t^9 - 257649t^7 \\
& \quad + 50128782t^5 + 390072133t^3 + 972422990t + (t \rightarrow t^{-1})) \\
& - 64 y_s^{\pm 1} \cdot (t^{14} - 22t^{12} + 231t^{10} - 24927t^8 + 317625t^6 + 7227990t^4 + 31070743t^2 + 48912688 + \\
& \quad + 31070743t^{-2} + 7227990t^{-4} + 317625t^{-6} + \dots + t^{-14}) \\
& + 154 (20t^{11} - 1740t^9 - 16109t^7 + 958563t^5 + 8046291t^3 + 20489955t + (t \rightarrow t^{-1}))
\end{aligned} \tag{A.15}$$

We also confirmed that the 1-instanton partition function Z_1 for $(N_s, N_c, N_v) = (0, 1, 2)$ could be identified with the above as follows:

$$Z_1^{(N_s, N_c, N_v)=(0,1,2)}(\vec{a}, \epsilon_1, \epsilon_2, m^c, \vec{m}^v) = Z_1^{(N_s, N_c, N_v)=(1,0,2)}(\vec{a}, \epsilon_1, \epsilon_2, m^s, \vec{m}^v)|_{m^s \rightarrow -m^c}. \tag{A.16}$$

E₆ The flavor symmetry on $N_f \mathbf{F} + N_{\bar{f}} \bar{\mathbf{F}}$ hypermultiplets is $U(N_f + N_{\bar{f}})$. The character expansion of \hat{Z}_1 at $N_f + N_{\bar{f}} = 3$ is written as follows:

$$\begin{aligned}
\hat{Z}_1 = & \frac{t^{22}}{(1-t^2)^{22}} \left((\chi_{(00)_{-9}}^f + \chi_{(00)_9}^f)(t^{11} + 56t^9 + 945t^7 + 6776t^5 + 23815t^3 + 43989t + (t \rightarrow t^{-1})) \right. \\
& - 27(\chi_{(10)_{-8}}^f + \chi_{(01)_8}^f)(t^{10} + 42t^8 + 539t^6 + 2948t^4 + 7854t^2 + 10824 + 7854t^{-2} + \dots + t^{-10}) \\
& + 351(\chi_{(20)_{-7}}^f + \chi_{(02)_7}^f)(t^9 + 28t^7 + 253t^5 + 1001t^3 + 1947t + (t \rightarrow t^{-1})) \\
& + 351(\chi_{(01)_{-7}}^f + \chi_{(10)_7}^f)(t^9 + 33t^7 + 330t^5 + 1386t^3 + 2772t + (t \rightarrow t^{-1})) \\
& + (\chi_{(30)_{-6}}^f + \chi_{(03)_6}^f)(t^{12} - 22t^{10} - 2694t^8 - 42790t^6 - 256355t^4 \\
& \quad - 712536t^2 - 994488 - 712536t^{-2} + \dots + t^{-12}) \\
& - 26(\chi_{(11)_{-6}}^f + \chi_{(11)_6}^f)(224t^8 + 4774t^6 + 32700t^4 + 96877t^2 + 137830 + 96877t^{-2} + \dots + 224t^{-8}) \\
& - 13(\chi_{(00)_{-6}}^f + \chi_{(00)_6}^f)(231t^8 + 6182t^6 + 48796t^4 + 156338t^2 + 228074 + 156338t^{-2} + \dots + 231t^{-8}) \\
& + 351(\chi_{(40)_{-5}}^f + \chi_{(04)_5}^f)(t^9 + 28t^7 + 253t^5 + 1001t^3 + 1947t + (t \rightarrow t^{-1})) \\
& - 27(\chi_{(21)_{-5}}^f + \chi_{(12)_5}^f)(t^{11} - 22t^9 - 1694t^7 - 19965t^5 - 89298t^3 - 182952t + (t \rightarrow t^{-1})) \\
& + 702(\chi_{(02)_{-5}}^f + \chi_{(20)_5}^f)(49t^7 + 707t^5 + 3399t^3 + 7150t + (t \rightarrow t^{-1})) \\
& + 702(\chi_{(10)_{-5}}^f + \chi_{(01)_5}^f)(77t^7 + 1281t^5 + 6677t^3 + 14575t + (t \rightarrow t^{-1})) \\
& - 27(\chi_{(50)_{-4}}^f + \chi_{(05)_4}^f)(t^{10} + 42t^8 + 539t^6 + 2948t^4 + 7854t^2 + 10824 + 7854t^{-2} + \dots + t^{-10}) \\
& - 351(\chi_{(31)_{-4}}^f + \chi_{(13)_4}^f)(21t^8 + 434t^6 + 2926t^4 + 8602t^2 + 12210 + 8602t^{-2} + \dots + 21t^{-8}) \\
& + 351(\chi_{(12)_{-4}}^f + \chi_{(21)_4}^f)(t^{10} - 22t^8 - 869t^6 - 6908t^4 - 21714t^2 - 31416 - 21714t^{-2} + \dots + t^{-10})
\end{aligned} \tag{A.17}$$

$$\begin{aligned}
& + 351(\chi_{(20)_{-4}}^f + \chi_{(02)_4}^f)(t^{10} - 22t^8 - 1177t^6 - 10500t^4 - 34936t^2 - 51436 - 34936t^{-2} + \dots + t^{-10}) \\
& - 1404(\chi_{(01)_{-4}}^f + \chi_{(10)_4}^f)(294t^6 + 3132t^4 + 10989t^2 + 16390 + 10989t^{-2} + 3132t^{-4} + 294t^{-6}) \\
& + (\chi_{(60)_{-3}}^f + \chi_{(06)_3}^f)(t^{11} + 56t^9 + 945t^7 + 6776t^5 + 23815t^3 + 43989t + (t \rightarrow t^{-1})) \\
& + 13(\chi_{(41)_{-3}}^f + \chi_{(14)_3}^f)(50t^9 + 1573t^7 + 15219t^5 + 62623t^3 + 124025t + (t \rightarrow t^{-1})) \\
& + (\chi_{(22)_{-3}}^f + \chi_{(22)_3}^f)(t^{13} - 22t^{11} + 231t^9 + 68530t^7 + 919589t^5 \\
& \quad + 4310670t^3 + 8985999t + (t \rightarrow t^{-1})) \\
& - 13(\chi_{(03)_{-3}}^f + \chi_{(30)_3}^f)(6t^{11} + 93t^9 - 3564t^7 - 60115t^5 - 303171t^3 - 650699t + (t \rightarrow t^{-1})) \\
& + 13(\chi_{(30)_{-3}}^f + \chi_{(03)_3}^f)(6075t^7 + 95425t^5 + 483483t^3 + 1042937t + (t \rightarrow t^{-1})) \\
& - 832(\chi_{(11)_{-3}}^f + \chi_{(11)_3}^f)(7t^9 - 154t^7 - 4095t^5 - 23683t^3 - 53471t + (t \rightarrow t^{-1})) \\
& + (\chi_{(00)_{-3}}^f + \chi_{(00)_3}^f)(t^{15} - 22t^{13} + 231t^{11} - 1540t^9 + 7315t^7 \\
& \quad + 1533042t^5 + 10536141t^3 + 24960012t + (t \rightarrow t^{-1})) \\
& - 27(\chi_{(51)_{-2}}^f + \chi_{(15)_2}^f)(t^{10} + 42t^8 + 539t^6 + 2948t^4 + 7854t^2 + 10824 + 7854t^{-2} + \dots + t^{-10}) \\
& - 351(\chi_{(32)_{-2}}^f + \chi_{(23)_2}^f)(21t^8 + 434t^6 + 2926t^4 + 8602t^2 + 12210 + 8602t^{-2} + \dots + 21t^{-8}) \\
& + 351(\chi_{(13)_{-2}}^f + \chi_{(31)_2}^f)(t^{10} - 22t^8 - 869t^6 - 6908t^4 - 21714t^2 - 31416 - 21714t^{-2} + \dots + t^{-10}) \\
& - 702(\chi_{(40)_{-2}}^f + \chi_{(04)_2}^f)(11t^8 + 270t^6 + 2002t^4 + 6182t^2 + 8910 + 6182t^{-2} + \dots + 11t^{-8}) \\
& - 27(\chi_{(21)_{-2}}^f + \chi_{(12)_2}^f)(t^{12} - 22t^{10} + 231t^8 + 34300t^6 + 334235t^4 \\
& \quad + 1139314t^2 + 1686762 + 1139314t^{-2} + \dots + t^{-12}) \\
& + 27(\chi_{(02)_{-2}}^f + \chi_{(20)_2}^f)(64t^{10} + 517t^8 - 27566t^6 - 317548t^4 - 1145354t^2 \\
& \quad - 1723106 - 1145354t^{-2} + \dots + 64t^{-10}) \\
& + 4914(\chi_{(10)_{-2}}^f + \chi_{(01)_2}^f)(7t^8 - 154t^6 - 2310t^4 - 8866t^2 - 13530 - 8866t^{-2} + \dots + 7t^{-8}) \\
& + 351(\chi_{(42)_{-1}}^f + \chi_{(24)_1}^f)(t^9 + 28t^7 + 253t^5 + 1001t^3 + 1947t + (t \rightarrow t^{-1})) \\
& + 351(\chi_{(50)_{-1}}^f + \chi_{(05)_1}^f)(t^9 + 33t^7 + 330t^5 + 1386t^3 + 2772t + (t \rightarrow t^{-1})) \\
& - 27(\chi_{(23)_{-1}}^f + \chi_{(32)_1}^f)(t^{11} - 22t^9 - 1694t^7 - 19965t^5 - 89298t^3 - 182952t + (t \rightarrow t^{-1})) \\
& + 22464(\chi_{(31)_{-1}}^f + \chi_{(13)_1}^f)(5t^7 + 77t^5 + 385t^3 + 825t + (t \rightarrow t^{-1})) \\
& + 702(\chi_{(04)_{-1}}^f + \chi_{(40)_1}^f)(49t^7 + 707t^5 + 3399t^3 + 7150t + (t \rightarrow t^{-1})) \\
& - 351(\chi_{(12)_{-1}}^f + \chi_{(21)_1}^f)(21t^9 - 462t^7 - 11605t^5 - 65983t^3 - 148071t + (t \rightarrow t^{-1})) \\
& + 351(\chi_{(20)_{-1}}^f + \chi_{(02)_1}^f)(t^{11} - 22t^9 + 231t^7 + 11516t^5 + 72799t^3 + 168707t + (t \rightarrow t^{-1})) \\
& - 702(\chi_{(01)_{-1}}^f + \chi_{(10)_1}^f)(25t^9 - 6325t^5 - 44583t^3 - 107387t + (t \rightarrow t^{-1})) \\
& + \chi_{(33)_0}^f(t^{12} - 22t^{10} - 2694t^8 - 42790t^6 - 256355t^4 - 712536t^2 - 994488 - 712536t^{-2} + \dots + t^{-12}) \\
& - 26(\chi_{(41)_0}^f + \chi_{(14)_0}^f)(224t^8 + 4774t^6 + 32700t^4 + 96877t^2 + 137830 + 96877t^{-2} + \dots + t^{-8})
\end{aligned}$$

$$\begin{aligned}
& + 26 \chi_{(22)_0}^f (25t^{10} - 550t^8 - 27030t^6 - 231990t^4 - 756657t^2 - 1107156 - 756657t^{-2} + \dots + 25t^{-10}) \\
& + (\chi_{(03)_0}^f + \chi_{(30)_0}^f)(t^{14} - 22t^{12} + 231t^{10} - 1540t^8 - 593285t^6 - 5973198t^4 \\
& \quad - 20531379t^2 - 30453456 - 20531379t^{-2} + \dots + t^{-14}) \\
& - 26 \chi_{(11)_0}^f (3t^{12} - 66t^{10} - 2002t^8 + 54670t^6 + 741975t^4 + 2786872t^2 \\
& \quad + 4232536 + 2786872t^{-2} + \dots + 3t^{-12}) \\
& + 2(1215t^{10} + 26070t^8 - 212410t^6 - 4381850t^4 - 18219943t^2 \\
& \quad - 28496524 - 18219943t^{-2} + \dots + 1215t^{-10})
\end{aligned}$$

where the $U(3)$ character $\chi_{(mn)_c}^f$ is defined as

$$\chi_{(mn)_c}^f = (w_1 w_2 w_3)^{\frac{c-m+n}{3}} \cdot \left(\sum_{\substack{1 \leq i_1 \leq \dots \leq i_m \leq 3 \\ 1 \leq j_1 \leq \dots \leq j_n \leq 3}} \frac{w_{i_1} \dots w_{i_m}}{w_{j_1} \dots w_{j_n}} - \sum_{\substack{1 \leq i_1 \leq \dots \leq i_{m-1} \leq 3 \\ 1 \leq j_1 \leq \dots \leq j_{n-1} \leq 3}} \frac{w_{i_1} \dots w_{i_{m-1}}}{w_{j_1} \dots w_{j_{n-1}}} \right) \quad (\text{A.18})$$

with

$$(w_1, w_2, w_3) = \begin{cases} (y_{f,1}, y_{f,2}, y_{f,3}) & \text{for } (N_f, N_{\bar{f}}) = (3, 0) \\ (y_{f,1}, y_{f,2}, y_{\bar{f},1}^{-1}) & \text{for } (N_f, N_{\bar{f}}) = (2, 1) \\ (y_{f,1}, y_{\bar{f},1}^{-1}, y_{\bar{f},2}^{-1}) & \text{for } (N_f, N_{\bar{f}}) = (1, 2) \\ (y_{\bar{f},1}^{-1}, y_{\bar{f},2}^{-1}, y_{\bar{f},3}^{-1}) & \text{for } (N_f, N_{\bar{f}}) = (0, 3) \end{cases} \quad (\text{A.19})$$

Again, (A.17) was tested against our general 1-instanton expression (2.40) up to t^{180} order.

E7 The flavor symmetry acting on N_f hypermultiplets is $SO(2N_f)_f$. The character expansion of \hat{Z}_1 at $N_f = 2$ is given as follows:

$$\begin{aligned}
\hat{Z}_1 = & \frac{t^{34}}{(1-t^2)^{34}} \left((\chi_{(0,12)}^f + \chi_{(12,0)}^f)(t^{17} + 99t^{15} + 3410t^{13} + 56617t^{11} + 521917t^9 \right. \\
& + 2889898t^7 + 10086066t^5 + 22867856t^3 + 34289476t + (t \rightarrow t^{-1}) \\
& - 8(\chi_{(1,11)}^f + \chi_{(11,1)}^f)(7t^{16} + 572t^{14} + 16401t^{12} + 227766t^{10} + 1759296t^8 \\
& + 8155308t^6 + 23747878t^4 + 44652608t^2 + 55026348 + 44652608t^{-2} + \dots + 7t^{-16}) \\
& + 19(\chi_{(2,10)}^f + \chi_{(10,2)}^f)(81t^{15} + 5254t^{13} + 121550t^{11} + 1376580t^9 \\
& + 8725369t^7 + 33273284t^5 + 79629972t^3 + 122510670t + (t \rightarrow t^{-1})) \\
& + 133(\chi_{(0,10)}^f + \chi_{(10,0)}^f)(11t^{15} + 760t^{13} + 18445t^{11} + 216580t^9 \\
& + 1409980t^7 + 5479474t^5 + 13273260t^3 + 20541950t + (t \rightarrow t^{-1})) \\
& - 152(\chi_{(39)}^f + \chi_{(93)}^f)(182t^{14} + 8827t^{12} + 158592t^{10} + 1426827t^8 + 7281032t^6 \\
& \left. + 22506946t^4 + 43735356t^2 + 54466776 + 43735356t^{-2} + \dots + 182t^{-14}) \right)
\end{aligned} \quad (\text{A.20})$$

$$\begin{aligned}
& - 2128(\chi_{(19)}^f + \chi_{(91)}^f)(24t^{14} + 1309t^{12} + 25454t^{10} + 241859t^8 \\
& \quad + 1281324t^6 + 4059022t^4 + 7997752t^2 + 10005112 + 7997752t^{-2} + \dots + 24t^{-14}) \\
& + (\chi_{(48)}^f + \chi_{(84)}^f)(t^{19} - 34t^{17} + 561t^{15} + 359766t^{13} + 11997546t^{11} + 161435604t^9 \\
& \quad + 1130192844t^7 + 4579505424t^5 + 11356618494t^3 + 17763983094t + (t \rightarrow t^{-1})) \\
& + 10773(\chi_{(28)}^f + \chi_{(82)}^f)(91t^{13} + 3668t^{11} + 54893t^9 \\
& \quad + 411026t^7 + 1739100t^5 + 4427038t^3 + 7011004t + (t \rightarrow t^{-1})) \\
& + 5187(\chi_{(08)}^f + \chi_{(80)}^f)(119t^{13} + 5269t^{11} + 83499t^9 \\
& \quad + 648329t^7 + 2806870t^5 + 7243122t^3 + 11543952t + (t \rightarrow t^{-1})) \\
& + 912(\chi_{(57)}^f + \chi_{(75)}^f)(t^{16} - 34t^{14} - 3597t^{12} - 78540t^{10} - 776832t^8 - 4186896t^6 \\
& \quad - 13370126t^4 - 26439556t^2 - 33110220 - 26439556t^{-2} + \dots + t^{-16}) \\
& - 56(\chi_{(37)}^f + \chi_{(73)}^f)(t^{18} - 34t^{16} + 561t^{14} + 227392t^{12} + 6213449t^{10} + 69122350t^8 + 400174169t^6 \\
& \quad + 1335305664t^4 + 2705039932t^2 + 3413732872 + 2705039932t^{-2} + \dots + t^{-18}) \\
& - 27664(\chi_{(17)}^f + \chi_{(71)}^f)(539t^{12} + 17314t^{10} + 208879t^8 + 1267860t^6 \\
& \quad + 4351490t^4 + 8949752t^2 + 11348792 + 8949752t^{-2} + \dots + 539t^{-12}) \\
& - 19\chi_{(66)}^f(7t^{17} + 217t^{15} - 24908t^{13} - 1021757t^{11} - 14769022t^9 \\
& \quad - 107322042t^7 - 444417927t^5 - 1115908152t^3 - 1755535056t + (r \rightarrow t^{-1})) \\
& - 95(\chi_{(46)}^f + \chi_{(64)}^f)(429t^{15} - 14586t^{13} - 1157156t^{11} - 20646010t^9 \\
& \quad - 168250530t^7 - 746606798t^5 - 1952106107t^3 - 3129466862t + (t \rightarrow t^{-1})) \\
& + 57(\chi_{(26)}^f + \chi_{(62)}^f)(27t^{17} - 918t^{15} + 15147t^{13} + 3520341t^{11} + 75470346t^9 \\
& \quad + 672625723t^7 + 3141068903t^5 + 8453641548t^3 + 13732731903t + (t \rightarrow t^{-1})) \\
& + (\chi_{(06)}^f + \chi_{(60)}^f)(t^{21} - 34t^{19} + 561t^{17} - 5984t^{15} + 46376t^{13} + 108842392t^{11} + 2613712872t^9 \\
& \quad + 24490191704t^7 + 117519798814t^5 + 321089011759t^3 + 525183176299t + (t \rightarrow t^{-1})) \\
& + 1296\chi_{(55)}^f(5t^{16} + 110t^{14} - 14030t^{12} - 458914t^{10} - 5440765t^8 - 32547180t^6 \\
& \quad - 110625460t^4 - 226279180t^2 - 286427492 - 226279180t^{-2} + \dots + 5t^{-16}) \\
& + 1064(\chi_{(35)}^f + \chi_{(53)}^f)(810t^{14} - 27540t^{12} - 1538126t^{10} - 21575635t^8 - 140780490t^6 \\
& \quad - 502663905t^4 - 1055162460t^2 - 1346539128 - 1055162460t^{-2} + \dots + 810t^{-14}) \\
& - 27664(\chi_{(15)}^f + \chi_{(51)}^f)(t^{16} - 34t^{14} + 561t^{12} + 62832t^{10} + 1000416t^8 \\
& \quad + 6920904t^6 + 25507174t^4 + 54425228t^2 + 69808596 + 54425228t^{-2} + \dots + t^{-16}) \\
& + \chi_{(44)}^f(t^{21} - 34t^{19} + 561t^{17} - 158136t^{15} - 1922955t^{13} + 320810876t^{11} + 7970822266t^9 \\
& \quad + 74975208858t^7 + 359889450611t^5 + 983025661861t^3 + 1607508212091t + (t \rightarrow t^{-1})) \\
& - 133(\chi_{(24)}^f + \chi_{(42)}^f)(t^{19} - 34t^{17} + 561t^{15} + 79101t^{13} - 2846514t^{11} - 102197931t^9
\end{aligned}$$

$$\begin{aligned}
& -1080814746t^7 - 5500823076t^5 - 15503708076t^3 - 25710027486t + (t \rightarrow t^{-1})) \\
& + 665(\chi_{(04)}^f + \chi_{(40)}^f)(13t^{17} + 108t^{15} - 11407t^{13} + 230758t^{11} + 11122199t^9 \\
& \quad + 125832753t^7 + 660902603t^5 + 1893530023t^3 + 3162878730t + (t \rightarrow t^{-1})) \\
& + 152\chi_{(33)}^f(6t^{18} - 204t^{16} + 18381t^{14} + 20306t^{12} - 21755987t^{10} - 387061196t^8 - 2796155121t^6 \\
& \quad - 10534894066t^4 - 22728127951t^2 - 29251476496 - 22728127951t^{-2} + \dots + 6t^{-18}) \\
& - 1064(\chi_{(13)}^f + \chi_{(31)}^f)(81t^{16} - 2754t^{14} - 49181t^{12} + 2732444t^{10} + 59237424t^8 + 458851114t^6 \\
& \quad + 1789977134t^4 + 3929114222t^2 + 5083736372 + 3929114222t^{-2} + \dots + t^{-16}) \\
& + 81\chi_{(22)}^f(91t^{17} + 41t^{15} - 356609t^{13} + 2951795t^{11} + 247685515t^9 \\
& \quad + 3029637009t^7 + 16451185429t^5 + 47931732849t^3 + 80650803640t + (t \rightarrow t^{-1})) \\
& - 1312311(\chi_{(02)}^f + \chi_{(20)}^f)(14t^{13} - 17t^{11} - 7752t^9 \\
& \quad - 103411t^7 - 581570t^5 - 1724208t^3 - 2923116t + (t \rightarrow t^{-1})) \\
& + 304\chi_{(11)}^f(11960t^{14} + 343681t^{12} - 7234554t^{10} - 208524209t^8 - 1747615980t^6 \\
& \quad - 7073563915t^4 - 15807799502t^2 - 20565064322 - 15807799502t^{-2} + \dots + 11960t^{-14}) \\
& + (t^{23} - 34t^{21} + 561t^{19} - 5984t^{17} - 192226t^{15} - 11212452t^{13} - 46556642t^{11} + 4966300623t^9 \\
& \quad + 73315010528t^7 + 427928422856t^5 + 1291626014327t^3 + 2206690491962t + (t \rightarrow t^{-1})))
\end{aligned}$$

This was tested against the closed-form expression (2.40) up to t^{280} order.

References

- [1] N. Seiberg and E. Witten, *Electric–magnetic Duality, Monopole Condensation, and Confinement in $\mathcal{N}=2$ Supersymmetric Yang-Mills Theory*, *Nucl. Phys.* **B426** (1994) 19–52, [[hep-th/9407087](#)].
- [2] N. Seiberg and E. Witten, *Monopoles, Duality and Chiral Symmetry Breaking in $\mathcal{N}=2$ Supersymmetric QCD*, *Nucl. Phys.* **B431** (1994) 484–550, [[hep-th/9408099](#)].
- [3] N. A. Nekrasov, *Seiberg-Witten Prepotential from Instanton Counting*, *Adv. Theor. Math. Phys.* **7** (2003) 831–864, [[hep-th/0206161](#)].
- [4] N. Nekrasov and A. Okounkov, *Seiberg-Witten Theory and Random Partitions*, *Prog. Math.* **244** (2006) 525–596, [[hep-th/0306238](#)].
- [5] H. Nakajima and K. Yoshioka, *Instanton counting on blowup. 1.*, *Invent. Math.* **162** (2005) 313–355, [[math/0306198](#)].
- [6] A. Braverman and P. Etingof, *Instanton Counting via Affine Lie Algebras Ii: from Whittaker Vectors to the Seiberg-Witten Prepotential*, [[math/0409441](#)].
- [7] G. W. Moore, N. Nekrasov and S. Shatashvili, *Integrating over Higgs Branches*, *Commun. Math. Phys.* **209** (2000) 97–121, [[hep-th/9712241](#)].
- [8] U. Bruzzo, F. Fucito, J. F. Morales and A. Tanzini, *Multiinstanton Calculus and Equivariant Cohomology*, *JHEP* **05** (2003) 054, [[hep-th/0211108](#)].

- [9] N. Nekrasov and S. Shadchin, *ABCD of Instantons*, *Commun. Math. Phys.* **252** (2004) 359–391, [[hep-th/0404225](#)].
- [10] M. Mariño and N. Wyllard, *A Note on Instanton Counting for $\mathcal{N}=2$ Gauge Theories with Classical Gauge Groups*, *JHEP* **05** (2004) 021, [[hep-th/0404125](#)].
- [11] F. Fucito, J. F. Morales and R. Poghossian, *Instantons on Quivers and Orientifolds*, *JHEP* **10** (2004) 037, [[hep-th/0408090](#)].
- [12] L. Hollands, C. A. Keller and J. Song, *From SO/Sp Instantons to W -Algebra Blocks*, *JHEP* **03** (2011) 053, [[1012.4468](#)].
- [13] L. Hollands, C. A. Keller and J. Song, *Towards a $4D/2D$ Correspondence for Sicilian Quivers*, *JHEP* **10** (2011) 100, [[1107.0973](#)].
- [14] S. Benvenuti, A. Hanany and N. Mekareeya, *The Hilbert Series of the One Instanton Moduli Space*, *JHEP* **06** (2010) 100, [[1005.3026](#)].
- [15] C. A. Keller, N. Mekareeya, J. Song and Y. Tachikawa, *The ABCDEFG of Instantons and W -Algebras*, *JHEP* **03** (2012) 045, [[1111.5624](#)].
- [16] C. A. Keller and J. Song, *Counting Exceptional Instantons*, *JHEP* **07** (2012) 085, [[1205.4722](#)].
- [17] A. Hanany, N. Mekareeya and S. S. Razamat, *Hilbert Series for Moduli Spaces of Two Instantons*, *JHEP* **01** (2013) 070, [[1205.4741](#)].
- [18] S. Cremonesi, G. Ferlito, A. Hanany and N. Mekareeya, *Coulomb Branch and the Moduli Space of Instantons*, *JHEP* **12** (2014) 103, [[1408.6835](#)].
- [19] C. Hwang, J. Kim, S. Kim and J. Park, *General Instanton Counting and 5D SCFT*, *JHEP* **07** (2015) 063, [[1406.6793](#)].
- [20] K. Hori, H. Kim and P. Yi, *Witten Index and Wall Crossing*, *JHEP* **01** (2015) 124, [[1407.2567](#)].
- [21] C. Cordova and S.-H. Shao, *An Index Formula for Supersymmetric Quantum Mechanics*, [1406.7853](#).
- [22] F. Benini, R. Eager, K. Hori and Y. Tachikawa, *Elliptic Genera of 2d $\mathcal{N}=2$ Gauge Theories*, *Commun. Math. Phys.* **333** (2015) 1241–1286, [[1308.4896](#)].
- [23] F. Benini, R. Eager, K. Hori and Y. Tachikawa, *Elliptic Genera of Two-Dimensional $\mathcal{N}=2$ Gauge Theories with Rank-One Gauge Groups*, *Lett. Math. Phys.* **104** (2014) 465–493, [[1305.0533](#)].
- [24] H. Nakajima and K. Yoshioka, *Lectures on instanton counting*, in *CRM Workshop on Algebraic Structures and Moduli Spaces Montreal, Canada, July 14-20, 2003*, 2003. [math/0311058](#).
- [25] H. Nakajima and K. Yoshioka, *Instanton Counting on Blowup. II. K -Theoretic Partition Function*, [math/0505553](#).
- [26] D. Gaiotto and S. S. Razamat, *Exceptional Indices*, *JHEP* **05** (2012) 145, [[1203.5517](#)].
- [27] N. A. Nekrasov, *Localizing Gauge Theories*, in *Mathematical Physics. Proceedings, 14Th International Congress, Icmp 2003, Lisbon, Portugal, July 28-August 2, 2003*, pp. 645–654, 2003.
- [28] L. Gottsche, H. Nakajima and K. Yoshioka, *K -theoretic Donaldson invariants via instanton counting*, *Pure Appl. Math. Quart.* **5** (2009) 1029–1111, [[math/0611945](#)].

- [29] L. Gottsche, H. Nakajima and K. Yoshioka, *Instanton Counting and Donaldson Invariants*, *J. Diff. Geom.* **80** (2008) 343–390, [[math/0606180](#)].
- [30] E. Gasparim and C.-C. M. Liu, *The Nekrasov Conjecture for Toric Surfaces*, *Commun. Math. Phys.* **293** (2010) 661–700, [[0808.0884](#)].
- [31] G. Bonelli, K. Maruyoshi, A. Tanzini and F. Yagi, $\mathcal{N}=2$ Gauge Theories on Toric Singularities, Blow-Up Formulae and W-Algebras, *JHEP* **01** (2013) 014, [[1208.0790](#)].
- [32] J. Gu, B. Haghighat, K. Sun and X. Wang, *Blowup Equations for 6D Scfts. I*, *JHEP* **03** (2019) 002, [[1811.02577](#)].
- [33] J. Gu, A. Klemm, K. Sun and X. Wang, *Elliptic Blowup Equations for 6D Scfts. II: Exceptional Cases*, [1905.00864](#).
- [34] T. Sasaki, *$O(-2)$ Blow-Up Formula via Instanton Calculus on Affine C^2/Z_2 and Weil Conjecture*, [hep-th/0603162](#).
- [35] Y. Ito, K. Maruyoshi and T. Okuda, *Scheme Dependence of Instanton Counting in ALE Spaces*, *JHEP* **05** (2013) 045, [[1303.5765](#)].
- [36] E. Witten, *Topological Quantum Field Theory*, *Commun. Math. Phys.* **117** (1988) 353.
- [37] M. Bershtein, G. Bonelli, M. Ronzani and A. Tanzini, *Exact results for $\mathcal{N} = 2$ supersymmetric gauge theories on compact toric manifolds and equivariant Donaldson invariants*, *JHEP* **07** (2016) 023, [[1509.00267](#)].
- [38] L. Baulieu, A. Losev and N. Nekrasov, *Chern-Simons and Twisted Supersymmetry in Various Dimensions*, *Nucl. Phys.* **B522** (1998) 82–104, [[hep-th/9707174](#)].
- [39] A. Losev, G. W. Moore, N. Nekrasov and S. Shatashvili, *Four-Dimensional Avatars of Two-Dimensional RCFT*, *Nucl. Phys. Proc. Suppl.* **46** (1996) 130–145, [[hep-th/9509151](#)].
- [40] K. A. Intriligator, D. R. Morrison and N. Seiberg, *Five-Dimensional Supersymmetric Gauge Theories and Degenerations of Calabi-Yau Spaces*, *Nucl. Phys.* **B497** (1997) 56–100, [[hep-th/9702198](#)].
- [41] S. Shadchin, *On Certain Aspects of String Theory/Gauge Theory Correspondence*. PhD thesis, Orsay, LPT, 2005. [hep-th/0502180](#).
- [42] L. Bhardwaj and Y. Tachikawa, *Classification of 4D $\mathcal{N}=2$ Gauge Theories*, *JHEP* **12** (2013) 100, [[1309.5160](#)].
- [43] P. Jefferson, H.-C. Kim, C. Vafa and G. Zafrir, *Towards Classification of 5D Scfts: Single Gauge Node*, [1705.05836](#).
- [44] H. Hayashi, S.-S. Kim, K. Lee and F. Yagi, *Rank-3 Antisymmetric Matter on 5-Brane Webs*, *JHEP* **05** (2019) 133, [[1902.04754](#)].
- [45] M. F. Atiyah, N. J. Hitchin, V. G. Drinfeld and Yu. I. Manin, *Construction of Instantons*, *Phys. Lett.* **A65** (1978) 185–187.
- [46] M. Aganagic, A. Klemm, M. Mariño and C. Vafa, *The Topological Vertex*, *Commun. Math. Phys.* **254** (2005) 425–478, [[hep-th/0305132](#)].
- [47] A. Iqbal, C. Kozçaz and C. Vafa, *The Refined Topological Vertex*, *JHEP* **10** (2009) 069, [[hep-th/0701156](#)].

- [48] D.-E. Diaconescu and R. Entin, *Calabi-Yau Spaces and Five-Dimensional Field Theories with Exceptional Gauge Symmetry*, *Nucl. Phys.* **B538** (1999) 451–484, [[hep-th/9807170](#)].
- [49] P. Jefferson, S. Katz, H.-C. Kim and C. Vafa, *On Geometric Classification of 5D SCFTs*, *JHEP* **04** (2018) 103, [[1801.04036](#)].
- [50] L. Bhardwaj and P. Jefferson, *Classifying 5D SCFTs via 6D Scfts: Rank One*, *JHEP* **07** (2019) 178, [[1809.01650](#)].
- [51] F. Apruzzi, C. Lawrie, L. Lin, S. Schfer-Nameki and Y.-N. Wang, *Fibers add Flavor, Part I: Classification of 5d SCFTs, Flavor Symmetries and BPS States*, [1907.05404](#).
- [52] H.-C. Kim, S. Kim, E. Koh, K. Lee and S. Lee, *On Instantons as Kaluza-Klein Modes of M5-Branes*, *JHEP* **12** (2011) 031, [[1110.2175](#)].
- [53] Y. Hwang, J. Kim and S. Kim, *M5-Branes, Orientifolds, and S-Duality*, *JHEP* **12** (2016) 148, [[1607.08557](#)].
- [54] S.-J. Lee and P. Yi, *D-Particles on Orientifolds and Rational Invariants*, *JHEP* **07** (2017) 046, [[1702.01749](#)].
- [55] N. Seiberg, *Five-Dimensional SUSY Field Theories, Nontrivial Fixed Points and String Dynamics*, *Phys. Lett.* **B388** (1996) 753–760, [[hep-th/9608111](#)].
- [56] J. Gu, M.-x. Huang, A.-K. Kashani-Poor and A. Klemm, *Refined BPS invariants of 6d SCFTs from anomalies and modularity*, *JHEP* **05** (2017) 130, [[1701.00764](#)].
- [57] M.-x. Huang, K. Sun and X. Wang, *Blowup Equations for Refined Topological Strings*, *JHEP* **10** (2018) 196, [[1711.09884](#)].
- [58] H.-C. Kim, S.-S. Kim and K. Lee, *5-Dim Superconformal Index with Enhanced E_n Global Symmetry*, *JHEP* **10** (2012) 142, [[1206.6781](#)].
- [59] H.-C. Kim, J. Kim, S. Kim, K.-H. Lee and J. Park, *6D Strings and Exceptional Instantons*, [1801.03579](#).
- [60] M. Del Zotto and G. Lockhart, *Universal Features of BPS Strings in Six-Dimensional SCFTs*, *JHEP* **08** (2018) 173, [[1804.09694](#)].
- [61] R. Feger and T. W. Kephart, *LieARTA Mathematica application for Lie algebras and representation theory*, *Comput. Phys. Commun.* **192** (2015) 166–195, [[1206.6379](#)].

G	matter	r_0	d_{\max}
$SU(N)_\kappa$	$N_f \times \mathbf{N}$	$d - N/2 - \kappa/2$	$0 \leq d \leq N - \kappa - 2N_f - 1(?)$
$SU(6)_3$	$1 \times \mathbf{20}$	$d - 6/2 - 3/2 + 3/2$	$1 \leq d \leq 6$
$SO(7)$	pure	$d - 5/2$	$0 \leq d \leq 5$
$SO(7)$	$1 \times \mathbf{8}$	$d - 5/2 + 1/2$	$0 \leq d \leq 4$
$SO(7)$	$1 \times \mathbf{7}$	$d - 5/2 + 1 \times 1/2$	$0 \leq d \leq 4$
$SO(7)$	$2 \times \mathbf{7}$	$d - 5/2 + 2 \times 1/2$	$0 \leq d \leq 3$
G_2	pure	$d - 4/2$	$0 \leq d \leq 4$
G_2	$1 \times \mathbf{7}$	$d - 4/2 + 1/2$	$0 \leq d \leq 3$
F_4	pure	$d - 9/2$	$0 \leq d \leq 9$
F_4	$1 \times \mathbf{26}$	$d - 9/2 + 1 \times 3/2$	$0 \leq d \leq 6$
F_4	$2 \times \mathbf{26}$	$d - 9/2 + 2 \times 3/2$	$0 \leq d \leq 3$

Table 1: list of theories

Gauge group	Hypermultiplets
$SO(14)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 1$ and $n_{\mathbf{v}} \leq 2$
$SO(13)$	$n_{\mathbf{s}} = 1$ and $n_{\mathbf{v}} \leq 2$
$SO(12)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 2$
$SO(12)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 1$ and $n_{\mathbf{v}} \leq 4$
$SO(11)$	$n_{\mathbf{s}} = 2$
$SO(11)$	$n_{\mathbf{s}} = 1$ and $n_{\mathbf{v}} \leq 4$
$SO(10)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 3$
$SO(10)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 2$ and $n_{\mathbf{v}} \leq 2$
$SO(10)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 1$ and $n_{\mathbf{v}} \leq 4$
$SO(9)$	$n_{\mathbf{s}} = 3$
$SO(9)$	$n_{\mathbf{s}} = 2$ and $n_{\mathbf{v}} \leq 2$
$SO(9)$	$n_{\mathbf{s}} = 1$ and $n_{\mathbf{v}} \leq 4$
$SO(8)$	$n_{\mathbf{s}} + n_{\mathbf{c}} + n_{\mathbf{v}} = 4$
$SO(7)$	$n_{\mathbf{s}} + n_{\mathbf{v}} = 3$

Table 2: asd

Gauge Group	Hypermultiplets	Equation No.	Attachment
$SO(8)$	$1\mathbf{S} + 1\mathbf{C} + 1\mathbf{V}$	(3.11)	
$SO(8)$	$2\mathbf{S} + 2\mathbf{C} + 2\mathbf{V}$	(3.12)	
$SO(9)$	$2\mathbf{S} + 1\mathbf{V}$	(3.17)	
$SO(10)$	$2\mathbf{S} + 4\mathbf{V}$	(A.5)	
$SO(10)$	$1\mathbf{S} + 1\mathbf{C} + 4\mathbf{V}$	(A.5)	
$SO(10)$	$2\mathbf{C} + 4\mathbf{V}$	(A.5)	
$SO(8)$	$3\mathbf{S} + 1\mathbf{C}$		
$SO(8)$	$3\mathbf{C} + 1\mathbf{V}$		
$SO(8)$	$3\mathbf{C} + 1\mathbf{V}$		

Table 3: Character expansion of $SO(N)$ theory with spinor hypermultiplets