

Instantons from Blow-up

Joonho Kim,^a Sung-Soo Kim,^b Ki-Hong Lee,^c Kimyeong Lee,^a and Jaewon Song^a

^a*School of Physics, Korea Institute for Advanced Study, Seoul 02455, Korea*

^b*School of Physics, University of Electronic Science and Technology of China,
No.4, Section 2, North Jianshe Road, Chengdu, Sichuan 610054, China*

^c*Department of Physics and Astronomy & Center for Theoretical Physics
Seoul National University, Seoul 08826, Korea*

E-mail: joonhokim@kias.re.kr, sungsoo.kim@uestc.edu.cn,
khlee11812@gmail.com, klee@kias.re.kr, jsong@kias.re.kr

ABSTRACT: The Nekrasov partition function for 4d $\mathcal{N} = 2$ or 5d $\mathcal{N} = 1$ gauge theory on the blow up of a point $\hat{\mathbb{C}}^2$ can be written in terms of the partition function on the flat space \mathbb{C}^2 . At the same time, the partition function on the blow up is identical to the partition function on a flat space for sufficiently large class of examples. This relation enables us to compute the instanton partition functions for 4d $\mathcal{N} = 2$ and 5d $\mathcal{N} = 1$ gauge theories for arbitrary gauge theory with large class of matter representations without knowing explicit construction of the instanton moduli space. Remarkably, the instanton partition function is completely determined by the perturbative part. We obtain the partition function for the previously unknown theories: exceptional gauge groups EFG with fundamental/spinor hypermultiplets and more. We also compute the case with $SU(6)$ with rank-3 antisymmetric tensor and compare with the topological vertex computation using the recently found 5-brane web configuration.

Contents

1	Introduction	1
2	Instanton Counting from Blow-up	3
2.1	Recursion formula	8
2.2	One instanton partition function	10
3	Examples	14
3.1	Known examples	15
3.2	Ki-Hong’s note	15
3.3	$SU(6)_3$ with a rank-3 antisymmetric hyper	17
4	Conclusion	19
A	$SU(N)$ theory	20

1 Introduction

The Seiberg-Witten prepotential provides a complete description for the low energy dynamics of 4d $\mathcal{N} = 2$ or 5d $\mathcal{N} = 1$ gauge theory in its Coulomb branch [1, 2]. It is a function of the vacuum expectation value (VEV) of the scalar in the vector multiplet that parameterizes the Coulomb branch moduli space. Quantum correction to the prepotential is known to be one-loop exact, while there also exist non-perturbative corrections coming from Yang-Mills instantons.

An efficient way to compute the fully quantum corrected prepotential \mathcal{F} is to study the Nekrasov partition function \mathcal{Z} on Ω -deformed \mathbb{C}^2 or $\mathbb{C}^2 \times S^1$. It can be written as the product of the classical, one-loop, and instanton contributions,

$$\mathcal{Z}(\vec{a}, \vec{m}, \epsilon_1, \epsilon_2, \mathbf{q}) = \mathcal{Z}_{\text{class}}(\vec{a}, \epsilon_1, \epsilon_2, \mathbf{q}) \mathcal{Z}_{\text{1-loop}}(\vec{a}, \vec{m}, \epsilon_1, \epsilon_2) \mathcal{Z}_{\text{inst}}(\vec{a}, \vec{m}, \epsilon_1, \epsilon_2, \mathbf{q}), \quad (1.1)$$

where the instanton piece is the fugacity sum over all multi-instanton contributions:

$$\mathcal{Z}_{\text{inst}}(\vec{a}, \vec{m}, \epsilon_1, \epsilon_2, \mathbf{q}) = 1 + \sum_{k=1}^{\infty} \mathbf{q}^k \mathcal{Z}_k(\vec{a}, \vec{m}, \epsilon_1, \epsilon_2). \quad (1.2)$$

Once the Nekrasov partition function is known, one can extract the prepotential via taking $\epsilon_i \rightarrow 0$ limit as $\mathcal{F} = \lim_{\epsilon_i \rightarrow 0} \epsilon_1 \epsilon_2 \log \mathcal{Z}$ [3–6].

The instanton part of the partition function in the Ω -background can be computed once we know appropriate instanton moduli space. For the classical gauge group, the ADHM construction of the moduli space provides a direct way compute the instanton partition function. The ADHM construction can be understood as the quantum mechanics described by the Dp - $D(p+4)$ system. The Higgs branch moduli space of the Dp system gives the desired moduli space. Matter fields can be also introduced by considering the world-volume theory on the D0-branes of the D0-D4-D8 system. By using the localization on the 1d system on the D0-branes or its dimensional reduction [7, 8], the partition function has been obtained for variety of cases: classical gauge groups [9–13], exceptional gauge groups [14–17]. 3d $N=4$ Coulomb branch realization of the instanton moduli space: [18] The precise choice of the Contour of the ADHM integral has been derived in [19–21] following the Jeffrey-Kirwan residue formula in 2d elliptic genus [22, 23].

But there is no ADHM type construction for the exceptional gauge groups or generic type of matter fields. String-theoretic picture implies that they require strong-coupling dynamics or non-Lagrangian field theories to realize instanton moduli space of exceptional group as a vacuum moduli space. There has been a few results regarding the exceptional instantons.

In this paper, we generalize the approach of Nakajima-Yoshioka (NY) [5, 24, 25] for the pure YM theory. In [16], the NY blow-up formula were used to compute the instanton partition function for exceptional gauge group and tested against the superconformal index of 4d SCFT where the Higgs branch is given by the instanton moduli space [26]. We propose a general blow-up formula for general gauge theory with arbitrary representations, under the condition that the matter representation is not too large.¹ This enables us to compute the instanton partition functions for numerous gauge theories that have not been known before, without relying on to the explicit construction of the moduli space.

The basic idea is as follows: Let us consider a one-point blow up $\hat{\mathbb{C}}^2$ of the flat space \mathbb{C}^2 . The *full* partition function on $\hat{\mathbb{C}}^2$ can be written in terms of the products of the *full* partition function of \mathbb{C}^2 . But at the same time, the partition function on the blowup is identical to that of the flat space since we can smoothly blow-down $\hat{\mathbb{C}}^2$ to \mathbb{C}^2 as long as the matter representation is not ‘too large’. This provides us a functional relation for the partition function, which turns out to be sufficient to determine the instanton partition function itself. Remarkably, this relation is completely determined by the perturbative part of the partition function. Therefore we arrive at a surprising conclusion: The *perturbative* physics determine the *non-perturbative* physics!

For example, we find the following universal expression for the 1-instanton partition function with arbitrary gauge group and matters:

$$Z_1 = \frac{e^{\epsilon_1 + \epsilon_2}}{(1 - e^{\epsilon_1})(1 - e^{\epsilon_2})} \sum_{\gamma \in \Delta_l} \frac{e^{(h^\vee - 1)a_\gamma/2} \prod_{\gamma \cdot w = 1} (1 - e^{a_w + m^{\text{phy}}})}{(e^{a_\gamma/2} - e^{-a_\gamma/2})(1 - e^{a_\gamma - \epsilon_1 - \epsilon_2}) \prod_{\gamma \cdot \alpha = 1} (e^{a_\alpha/2} - e^{-a_\alpha/2})}, \quad (1.3)$$

where ...

¹This has to do with the 1-loop beta function coefficient as we will discuss in detail.

Notation The equivariant parameters $\epsilon_1, \epsilon_2, \vec{a}$ are associated to the $U(1)^2 \times U(1)^{|G|}$ action on the k -instanton moduli space $\mathcal{M}_{k,G}$. Additional parameters \vec{m} are introduced if the gauge theory has an extra flavor symmetry. The instanton fugacity q can be written as $q \equiv \Lambda^{b_0}$ (4d) or $q \equiv \exp(2\pi i \Lambda)$ (5d) where Λ denotes the bare coupling of the 4d/5d gauge theory.

Throughout Section 2, we will characterize the partition function in terms of $m_i \equiv \mathbf{m}_i + \epsilon_+$ (where $\epsilon_+ \equiv \frac{\epsilon_1 + \epsilon_2}{2}$) as a reflection of the Donaldson-Witten twist by $SU(2)_R$ symmetry. Here m_i corresponds to the ‘physical’ mass parameter of a matter hypermultiplet. We will also stick to the ‘effective’ instanton fugacity $q \equiv q e^{b_0 \epsilon_+}$, which absorbs the $SU(2)_R$ generated mass of fermionic instanton zero modes, in describing the 5d Nekrasov partition function.

2 Instanton Counting from Blow-up

The essential idea of using the blow-up of \mathbb{C}^2 for instanton counting is that the gauge theory partition function for a 4d $\mathcal{N} = 2$ (or 5d $\mathcal{N} = 1$) theory on the blow-up of a point $\hat{\mathbb{C}}^2$ (or $S^1 \times \hat{\mathbb{C}}^2$) can be written in two different ways. This will allow us to write a recursion relation for the instanton partition function that can be solved rather easily [5, 16, 24, 25].

Localization on the blow-up $\hat{\mathbb{C}}^2$ One of the expressions for the partition function $\hat{\mathcal{Z}}$ on the blow-up $\hat{\mathbb{C}}^2$ comes from the Coulomb branch localization, which results that $\hat{\mathcal{Z}}$ can be obtained by patching together the flat-space partition function \mathcal{Z} [27].

The blow-up $\hat{\mathbb{C}}^2$ is constructed from \mathbb{C}^2 by replacing the origin with the compact 2-cycle. In particular, the geometry near the \mathbb{P}^1 is identical to a line bundle of degree (-1) on the \mathbb{P}^1 . One can parametrize $\mathcal{O}(-1) \rightarrow \mathbb{P}^1$ using three homogeneous coordinates (z_0, z_1, z_2) , satisfying the projective condition $(z_0, z_1, z_2) \sim (\lambda^{-1} z_0, \lambda^1 z_1, \lambda^1 z_2)$ for any $\lambda \in \mathbb{C}^*$, where the two-cycle $\mathbb{P}^1 \subset \hat{\mathbb{C}}^2$ corresponds to the locus $z_0 = 0$. We are interested in the $U(1)^2$ equivariant partition function, with the $U(1)^2$ action V rotating the complex coordinates (z_0, z_1, z_2) as follows:

$$(z_0, z_1, z_2) \mapsto (z_0, e^{\epsilon_1} z_1, e^{\epsilon_2} z_2). \quad (2.1)$$

Instantons are located at two fixed points of the $U(1)^2$ action, i.e., the north/south poles of the \mathbb{P}^1 , whose coordinates are $(z_0, z_1, z_2) = (0, 1, 0)$ and $(0, 0, 1)$. Around these fixed points, the local weights under the $U(1)^2$ action V are:

$$(z_0 z_1, z_2/z_1) \mapsto (e^{\epsilon_1} z_0 z_1, e^{\epsilon_2 - \epsilon_1} z_2/z_1) \quad (\text{near the north pole}) \quad (2.2)$$

$$(z_0 z_2, z_1/z_2) \mapsto (e^{\epsilon_2} z_0 z_2, e^{\epsilon_1 - \epsilon_2} z_1/z_2) \quad (\text{near the south pole}) \quad (2.3)$$

The full partition function $\hat{\mathcal{Z}}$ on $\hat{\mathbb{C}}^2$, which includes both the perturbative and instanton contributions, can be obtained by performing the localization on the Coulomb branch. On the Coulomb branch, the gauge group is generically broken to $U(1)^r$ where r is the rank of the gauge group. The $U(1)^r$ equivariant parameters \vec{a} naturally appear in the partition function. One needs to sum over all distinct field configurations with zero-sized instantons located at the north and south poles. All the inequivalent configurations are labeled by the vector \vec{k} of

the first Chern numbers, corresponding to different flux configurations on the two-cycle \mathbb{P}^1 . Here we turn off any possible external flux that can be supported on the \mathbb{P}^1 . Summing up, $\hat{\mathcal{Z}}$ can be expressed as the following formula [27–31]:

$$\hat{\mathcal{Z}}(\vec{a}, \epsilon_1, \epsilon_2) = \sum_{\vec{k} \in \Lambda} \mathcal{Z}(\vec{a} + \vec{k}\epsilon_1, \epsilon_1, \epsilon_2 - \epsilon_1) \mathcal{Z}(\vec{a} + \vec{k}\epsilon_2, \epsilon_1 - \epsilon_2, \epsilon_2) \quad (2.4)$$

where the flux sum is taken over the co-root lattice Λ of the gauge algebra. Notice that the Coulomb parameter \vec{a} gets an appropriate shift at each fixed point p , induced by the non-trivial magnetic flux \vec{k} on the blown-up \mathbb{P}^1 , with the proportionality constant $H|_p$. Values of the moment map H for the $U(1)^2$ action V , i.e., $dH = \iota_V \omega$, at the north and south poles are

$$H|_{\text{NP}} = \epsilon_1 \text{ and } H|_{\text{SP}} = \epsilon_2. \quad (2.5)$$

Partition function on $\hat{\mathbb{C}}^2$ vs \mathbb{C}^2 An alternative fact for the partition function $\hat{\mathcal{Z}}$ on the blow-up $\hat{\mathbb{C}}^2$ is that $\hat{\mathcal{Z}}$ is actually identical to the flat-space partition function \mathcal{Z} [5, 24, 25].

The blow-up $\hat{\mathbb{C}}^2$ is identical to \mathbb{C}^2 except for the origin, which is replaced by the blown-up sphere \mathbb{P}^1 . Since the Nekrasov partition function gets contributions only from the small instantons localized at the fixed points of the $U(1)^2$ equivariant action V , the size of the divisor should not affect the partition function as we smoothly shrink it. So we expect that $\hat{\mathcal{Z}} = \mathcal{Z}$. This implies the following relation: [5, 16, 24, 25, 32, 33]

$$\mathcal{Z}(\vec{a}, \epsilon_1, \epsilon_2) = \sum_{\vec{k} \in \Lambda} \mathcal{Z}(\vec{a} + \vec{k}\epsilon_1, \epsilon_1, \epsilon_2 - \epsilon_1) \mathcal{Z}(\vec{a} + \vec{k}\epsilon_2, \epsilon_1 - \epsilon_2, \epsilon_2), \quad (2.6)$$

The blow-up identity can also be generalized to orbifold partition functions [31, 34, 35]. For example, if we consider the total space of the bundle $\mathcal{O}(-2) \rightarrow \mathbb{P}^1$ and shrink the base, we land on the orbifold $\mathbb{C}^2/\mathbb{Z}_2$ with singularity at the origin. There are certain states originated from wrapped D-branes on the \mathbb{P}^1 which become massless by shrinking the \mathbb{P}^1 . They constitute to the required twisted sector of the orbifold partition function. In any case, we expect that the Nekrasov partition function still remains the same by blowing up and down.²

Background parameters The expected relation (2.6) is only a schematic expression. The partition function also depends on some background parameters, such as the gauge coupling Λ and flavor chemical potentials m . They need to be appropriately shifted at each fixed point p of the blow-up $\hat{\mathbb{C}}^2$, keeping invariant the effective mass parameters twisted by $SU(2)_R$.

One can identify the shifted parameters $\Lambda|_p$ and $m|_p$ at a fixed point p , by examining the 1-loop effective free energy of the 5d Nekrasov partition function \mathcal{Z} . For a general 5d $\mathcal{N} = 1$

²This simple picture does not necessarily hold when there are too many hypermultiplets, due to some subtle scheme dependence related to wall-crossing [28, 29, 35]. We will discuss about this issue in Section ??.

gauge theory with the Chern-Simons level κ and/or some hypermultiplets,³

$$\begin{aligned} \log \mathcal{Z} = & \frac{1}{\epsilon_1 \epsilon_2} \left[\frac{1}{2} \Lambda h_{ij} a_i a_j + \frac{\kappa}{6} d_{ijk} a^i a^j a^k + \sum_{\vec{\alpha} \in \Delta} \left(\frac{(\vec{\alpha} \cdot \vec{\alpha} + \epsilon_+)^3}{12} - \frac{\epsilon_1^2 + \epsilon_2^2 + 24}{48} (\vec{\alpha} \cdot \vec{\alpha} + \epsilon_+) + 1 \right) \right. \\ & \left. - \sum_i \sum_{\vec{\omega} \in \mathbf{R}_i} \left(\frac{(\vec{\alpha} \cdot \vec{\omega} + m_i)^3}{12} - \frac{\epsilon_1^2 + \epsilon_2^2 + 24}{48} (\vec{\alpha} \cdot \vec{\omega} + m_i) + 1 \right) \right] + \sum_{n=1}^{\infty} \frac{f(n\vec{\alpha}, n\epsilon_{1,2}, nm_i)}{n} \quad (2.7) \end{aligned}$$

in which the first 2 terms inside the square brackets constitute the classical prepotential \mathcal{F}_{cl} , whereas all the other terms come in at one-loop level. The set of all roots is denoted by Δ . The $\vec{\omega}$ runs over all weight vectors in the representation \mathbf{R}_i of the i -th hypermultiplet. First, let us consider a 5d free hypermultiplet with mass m , whose corresponding free energy is

$$\log \mathcal{Z} = -\frac{1}{\epsilon_1 \epsilon_2} \left(\frac{m^3}{12} - \frac{\epsilon_1^2 + \epsilon_2^2 + 24}{48} m + 1 \right) + \sum_{n=1}^{\infty} \frac{1}{n} \frac{e^{-nm} \cdot e^{-n\epsilon_+}}{(1 - e^{-n\epsilon_1})(1 - e^{-n\epsilon_2})}. \quad (2.8)$$

This satisfies the relation (2.6) only when it is accompanied by $m \rightarrow m \pm \frac{H}{2}|_p$. Each \pm shift preserves the combination $(m \pm \epsilon_+)$ respectively, corresponding to the $SU(2)_R$ twisted mass of the hypermultiplet [36]. Throughout this paper, we take $m_{\text{phy}} \equiv (m - \epsilon_+)$ as the invariant $SU(2)_R$ twisted mass, identifying that

$$m|_p = m - \frac{H}{2}|_p \quad (2.9)$$

at a fixed point p . Second, the twisted instanton mass can be easily read off after simplifying (2.7) based on the following identities:

$$\begin{aligned} \sum_{\vec{\omega} \in \mathbf{R}} (\vec{\alpha} \cdot \vec{\omega}) (\vec{b} \cdot \vec{\omega}) (\vec{c} \cdot \vec{\omega}) &= I_3(\mathbf{R}) d_{ijk} a^i b^j c^k, \\ \sum_{\vec{\omega} \in \mathbf{R}} (\vec{\alpha} \cdot \vec{\omega}) (\vec{b} \cdot \vec{\omega}) &= I_2(\mathbf{R}) h_{ij} a^i b^j, \\ \sum_{\vec{\omega} \in \mathbf{R}} (\vec{\alpha} \cdot \vec{\omega}) &= 0. \end{aligned} \quad (2.10)$$

Here $I_2(\mathbf{R})$ and $I_3(\mathbf{R})$ are the quadratic and cubic Dynkin indices⁴ of a given representation \mathbf{R} . In particular, $I_2(\mathbf{adj}) = h^\vee$ is the dual Coxeter number of the gauge algebra. Speaking explicitly, the rational part of the one-loop free energy (2.7) can be written as

$$\log \mathcal{Z} = \frac{1}{\epsilon_1 \epsilon_2} \left[\frac{1}{2} \Lambda_{\text{eff}} h_{ij} a^i a^j + \frac{\kappa_{\text{eff}}}{6} d_{ijk} a^i a^j a^k + (\text{independent of } \vec{a}) \right], \quad (2.11)$$

where the one-loop corrected parameters are

$$\Lambda_{\text{eff}} = \Lambda + \left(h^\vee - \sum_i \frac{I_2(\mathbf{R}_i)}{2} \right) \epsilon_+ - \sum_i \frac{I_2(\mathbf{R}_i)}{2} m_i^{\text{phy}}, \quad \kappa_{\text{eff}} = \kappa - \sum_i \frac{I_3(\mathbf{R}_i)}{2} \quad (2.12)$$

³We assume a particular Weyl chamber in the Coulomb branch, i.e., $0 < a_i < \epsilon_+ < m$ for all $i \in \{1, \dots, r\}$.

⁴The Dynkin indices are normalized such that $I_2(\mathbf{F}) = 1$ for the $SU(N)$ fundamental representation \mathbf{F} .

in which the sum is taken over all the matter multiplets. We notice that the instanton soliton carries the $SU(2)_R$ twisted effective mass, given by $m_{\text{inst}} \equiv \Lambda_{\text{eff}} - \kappa_{\text{eff}} \epsilon_+$. In addition, the effective Chern-Simons coupling κ_{eff} induces an electric charge to the instanton, contributing to its ground state energy as $E_0 = m_{\text{inst}} - \vec{a} \cdot \vec{\Pi}$, where $\vec{\Pi}$ is the $U(1)^r \subset G$ electric charge.⁵ To keep it invariant the effective instanton mass m_{inst} at a fixed point p of the blow-up $\hat{\mathbb{C}}^2$, we require the shifted gauge coupling $\Lambda|_p$ to be

$$\Lambda|_p = \Lambda + \frac{b}{2}H|_p \quad \text{with} \quad b \equiv h^\vee - \sum_i \frac{I_2(\mathbf{R}_i)}{2} - \kappa_{\text{eff}}. \quad (2.13)$$

For example, the $\mathbb{C}^2 \times S^1$ partition function of $\mathcal{N} = 1$ free Maxwell theory, given as a product between the perturbative index of a free vector multiplet,

$$\exp \left[+\frac{1}{\epsilon_1 \epsilon_2} \left(\frac{\epsilon_1 \epsilon_2 (\epsilon_1 + \epsilon_2)}{48} - \frac{\epsilon_1 + \epsilon_2}{4} + 1 \right) \right] \cdot \text{PE} \left[-\frac{e^{-\epsilon_1 - \epsilon_2}}{(1 - e^{-\epsilon_1})(1 - e^{-\epsilon_2})} \right], \quad (2.14)$$

and the non-perturbative correction from $U(1)$ multi-instantons,

$$\text{PE} \left[\frac{e^{-\Lambda} \cdot e^{-\epsilon_+}}{(1 - e^{-\epsilon_1})(1 - e^{-\epsilon_2})} \right], \quad (2.15)$$

will comply with the relation (2.6) by shifting the gauge coupling as $\Lambda \rightarrow \Lambda|_p = \Lambda + \frac{H}{2}|_p$.

In summary, the blow-up relation (2.6) should always be understood with (2.9) and (2.13).

Correlation functions In fact, (2.6) is not enough to fix the partition function completely, since there are 3 unknown functions and only one relation. More independent relations would be found from insertion of non-trivial \mathcal{Q} -closed operators [5, 25]. So we consider the correlation functions of that operator $\mathcal{O}_{\mathbb{P}^1}$ associated to the two-cycle \mathbb{P}^1 of the blow-up $\hat{\mathbb{C}}^2$.

In the 4d Donaldson-twisted theory, the \mathcal{Q} -invariant observable $\mathcal{O}_{\mathbb{P}^1}$ can be constructed by applying the descent procedure twice to the Casimir invariant $\text{Tr}(\Phi^2)$. It can be written in terms of the component fields as follows: [37]

$$\mathcal{O}_{\mathbb{P}^1} = \int \left\{ \omega \wedge \text{Tr} \left(\Phi F + \frac{1}{2} \psi \wedge \psi \right) + H \text{Tr} (F \wedge F) \right\}. \quad (2.16)$$

Here ω and H are the Kähler two-form on the \mathbb{P}^1 and the moment map $\iota_V \omega = dH$, respectively. We find it convenient to study the generating function $\langle e^{t \mathcal{O}_{\mathbb{P}^1}} \rangle$ of the correlators $\langle \mathcal{O}_{\mathbb{P}^1} \dots \mathcal{O}_{\mathbb{P}^1} \rangle$. In particular, the insertion of $\exp(t \mathcal{O}_{\mathbb{P}^1})$ causes an extra shift of the instanton mass by $2tH|_p$ at a fixed point p of the blow-up $\hat{\mathbb{C}}^2$ [5, 24, 25]. This modifies (2.13) to $\Lambda|_p = \Lambda + (\frac{b}{2} - t)H|_p$. Therefore the expectation value of the generating function is given by

$$\langle e^{t \mathcal{O}_{\mathbb{P}^1}} \rangle = \sum_{\vec{k} \in \Lambda} \frac{\mathcal{Z}^{(N),t}(\vec{k}) \cdot \mathcal{Z}^{(S),t}(\vec{k})}{\mathcal{Z}^{(0)}} , \quad (2.17)$$

⁵This agrees with the supersymmetric Casimir energy of the ADHM quantum mechanics.

where

$$\begin{aligned}
\mathcal{Z}^{(N),t}(\vec{k}) &\equiv \mathcal{Z}(\vec{a} + \vec{k}\epsilon_1, \epsilon_1, \epsilon_2 - \epsilon_1, \Lambda + (\tfrac{b}{2} - t)\epsilon_1, \vec{m} + \tfrac{1}{2}\epsilon_1) , \\
\mathcal{Z}^{(S),t}(\vec{k}) &\equiv \mathcal{Z}(\vec{a} + \vec{k}\epsilon_2, \epsilon_1 - \epsilon_2, \epsilon_2, \Lambda + (\tfrac{b}{2} - t)\epsilon_2, \vec{m} + \tfrac{1}{2}\epsilon_2) , \\
\mathcal{Z}^{(0)} &\equiv \mathcal{Z}(\vec{a}, \epsilon_1, \epsilon_2, \Lambda, \vec{m}) .
\end{aligned} \tag{2.18}$$

As we shrink the two-cycle \mathbb{P}^1 to recover the flat \mathbb{C}^2 , the effect of inserting $\mathcal{O}_{\mathbb{P}^1}^d$ turns out to give a vanishing contribution due to the selection rule. We recall that the instanton breaks the $U(1)_R$ symmetry to the discrete subgroup $\mathbb{Z}_{4h^\vee - 2\sum_i I_2(\mathbf{R}_i)}$ under which the operator $\mathcal{O}_{\mathbb{P}^1}^d$ carries a charge of $+2$. The correlation functions vanish unless the discrete charges add up to zero, modulo $4h^\vee - 2\sum_i I_2(\mathbf{R}_i)$. Therefore, expanding (2.17) in powers of t , we find

$$\langle e^{t\mathcal{O}_{\mathbb{P}^1}} \rangle = 1 + \mathcal{O}(t^{2h^\vee - \sum_i I_2(\mathbf{R}_i)}) . \tag{2.19}$$

As long as the hypermultiplet representation is not too large, i.e., when $2h^\vee - \sum_i I_2(\mathbf{R}_i) > 2$, this allows us to write 3 independent relations for the 3 unknown variables. One can expand $\langle e^{t\mu^{(C)}} \rangle$ until $\mathcal{O}(t^2)$ order and recursively solve for $\mathcal{Z}^{(0)}$ at every instanton number. So the instanton part of the partition function will be completely determined from the perturbative partition function. An explicit form of the recursion relation will be displayed in Section 2.1.

We now turn to 5d $\mathcal{N} = 1$ gauge theory wrapped on S^1 . The Casimir invariant $\text{Tr}(\Phi^2)$ and its descendants are no longer considered as well-defined observables. Instead, there are two types of \mathcal{Q} -invariant observables [38]. The first type of observables are constructed from the 5d Wilson loop on the S^1 by applying the descent procedure. The second type of observables introduce the 3d (Kähler) Chern-Simons term, which can be written as [38, 39]

$$\begin{aligned}
\mathcal{O}_{\mathbb{P}^1}^d = \exp \bigg[& d \int \left(\omega \wedge \text{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) \right. \\
& \left. + \omega \wedge \left(\phi F + \frac{1}{2} \psi \wedge \psi \right) \wedge dt + H \text{Tr} (F \wedge F) \right) \bigg] .
\end{aligned} \tag{2.20}$$

It can be viewed as the natural S^1 uplift of (2.16) via exponentiation, where the coefficient d is integer-valued as required by gauge invariance. The correlation function is given by

$$\langle \mathcal{O}_{\mathbb{P}^1}^d \rangle = \sum_{\vec{k} \in \Lambda} \frac{\mathcal{Z}^{(N),d}(\vec{k}) \cdot \mathcal{Z}^{(S),d}(\vec{k})}{\mathcal{Z}^{(0)}} \tag{2.21}$$

with the definitions (2.18) understood. For the case of 5d pure $\mathcal{N} = 1$ SYM, the correlation function turns out to be [25]

$$\langle \mathcal{O}_{\mathbb{P}^1}^d \rangle = 1 \quad \text{for} \quad 0 \leq d \leq d_{\max} \tag{2.22}$$

where $d_{\max} = h^\vee$. For $d_{\max} = h^\vee \geq 2$, there are sufficient number of algebraic relations to determine the instanton partition function \mathcal{Z} iteratively, in increasing the order of instantons. This fact was utilized in [16] to compute \mathcal{Z} in exceptional gauge theories, for which the ADHM

construction of instanton moduli space is not known. In this paper, we aim at developing the relation (2.22) for various 5d $\mathcal{N} = 1$ gauge theories with hypermultiplets, then writing down the corresponding \mathcal{Z} based on that. We will identify a certain bound on d in Section 2.2 as the *necessary* condition on (2.22) for the wide array of theories which have UV completion. See Table 1 for the list of all considered examples. The bound can be summarized as follows:

$$h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) \textcolor{red}{(add)} \quad (2.23)$$

And also, we conjecture that the bound on d is sufficient for (2.22). While we do not attempt to prove this sufficiency, we will compute an n -instanton partition function Z_n , based on the recursion formula that will be derived shortly from (2.22), and confirm the agreement with the known result obtained from an alternative method. We will call (2.22) the blowup equation.

2.1 Recursion formula

The blowup equation (2.22) can be translated to useful recursion formulae on the k -instanton contribution Z_k to the entire partition function \mathcal{Z} . To derive this, we decompose the partition function \mathcal{Z} in terms of the classical, one-loop, and instanton pieces:

$$\mathcal{Z}(\vec{a}, \epsilon_1, \epsilon_2, \Lambda, \vec{m}) = Z_{\text{class}}(\vec{a}, \epsilon_1, \epsilon_2, \Lambda, \vec{m}) \cdot Z_{1\text{-loop}}(\vec{a}, \epsilon_1, \epsilon_2, \vec{m}) \cdot Z_{\text{inst}}(\vec{a}, \epsilon_1, \epsilon_2, \Lambda, \vec{m}) \quad (2.24)$$

where Z_{inst} can be further expanded in terms of the instanton fugacity $q \equiv e^{-\Lambda}$ as

$$Z_{\text{inst}}(\vec{a}, \epsilon_1, \epsilon_2, \Lambda, \vec{m}) = 1 + \sum_{k \geq 1} q^k Z_k(\vec{a}, \epsilon_1, \epsilon_2, m) . \quad (2.25)$$

Then the blowup equation (2.22) can be written as

$$Z_{\text{inst}} = \sum_{\vec{k}} \left[\frac{Z_{\text{class}}^{(N),d}(\vec{k}) Z_{\text{class}}^{(S),d}(\vec{k})}{Z_{\text{class}}} \frac{Z_{1\text{-loop}}^{(N),d}(\vec{k}) Z_{1\text{-loop}}^{(S),d}(\vec{k})}{Z_{1\text{-loop}}} \right] Z_{\text{inst}}^{(N),d}(\vec{k}) Z_{\text{inst}}^{(S),d}(\vec{k}) , \quad (2.26)$$

where the superscript $(N/S), d$ implies the appropriate shift of parameters, specified in (2.18).

We recall the known expressions for the classical and 1-loop partition function [3, 40, 41]:

$$Z_{\text{class}} = \exp \left[\frac{1}{\epsilon_1 \epsilon_2} \left(\frac{1}{2} \Lambda h_{ij} a_i a_j + \frac{\kappa}{6} d_{ijk} a^i a^j a^k \right) \right] , \quad (2.27)$$

$$\begin{aligned} Z_{1\text{-loop}}^{\text{vec}} &= \exp \left[\frac{1}{\epsilon_1 \epsilon_2} \sum_{\vec{\alpha} \in \Delta} \left(\frac{(\vec{a} \cdot \vec{\alpha} + \epsilon_+)^3}{12} - \frac{\epsilon_1^2 + \epsilon_2^2 + 24}{48} (\vec{a} \cdot \vec{\alpha} + \epsilon_+) + 1 \right) \right] \\ &\times \text{PE} \left[- \frac{p_1 p_2}{(1-p_1)(1-p_2)} \sum_{\vec{\alpha} \in \Delta} e^{-\vec{a} \cdot \vec{\alpha}} \right] \quad \text{for the vector multiplet} \end{aligned} \quad (2.28)$$

$$\begin{aligned} Z_{1\text{-loop}}^{\text{hyp},i} &= \exp \left[- \frac{1}{\epsilon_1 \epsilon_2} \sum_{\vec{\omega} \in \mathbf{R}_\ell} \left(\frac{(\vec{a} \cdot \vec{\omega} + m_\ell)^3}{12} - \frac{\epsilon_1^2 + \epsilon_2^2 + 24}{48} (\vec{a} \cdot \vec{\omega} + m_\ell) + 1 \right) \right] \\ &\times \text{PE} \left[+ \frac{(p_1 p_2)^{\frac{1}{2}} \cdot y_\ell}{(1-p_1)(1-p_2)} \sum_{\vec{\omega} \in \mathbf{R}_\ell} e^{-\vec{a} \cdot \vec{\omega}} \right] \quad \text{for the } \ell\text{'th hypermultiplet} \end{aligned} \quad (2.29)$$

where $p_1 \equiv e^{-\epsilon_1}$, $p_2 \equiv e^{-\epsilon_2}$, $y_i \equiv e^{-m_i}$ and the PE represents the Plethystic exponential

$$\text{PE} [f(\vec{a}, \epsilon_1, \epsilon_2, \Lambda, \vec{m})] \equiv \exp \left(\sum_{n=1}^{\infty} \frac{1}{n} f(n\vec{a}, n\epsilon_1, n\epsilon_2, n\Lambda, n\vec{m}) \right). \quad (2.30)$$

Substituting them to (2.26), the combinations of three different Z 's are given by

$$\begin{aligned} \frac{Z^{(N),d}(\vec{k}) Z^{(S),d}(\vec{k})}{Z} &= q^{\frac{\vec{k} \cdot \vec{k}}{2}} (p_1 p_2)^{(\frac{b}{2}-d)(\frac{\vec{k} \cdot \vec{k}}{2}) + \frac{\kappa}{6} d_{ijk} k^i k^j k^k} \\ &\quad \times e^{-(\frac{b}{2}-d)(\vec{a} \cdot \vec{k})} e^{-\frac{\kappa}{2} d_{ijk} a^i k^j k^k} \quad \text{for } Z = Z_{\text{class}}, \end{aligned} \quad (2.31)$$

$$\frac{Z^{(N),d}(\vec{k}) Z^{(S),d}(\vec{k})}{Z} = e^{\frac{h^\vee}{2}(\vec{a} \cdot \vec{k})} \prod_{\alpha \in \Delta} \mathcal{L}_{\vec{k}, \vec{\alpha}}(\vec{a} \cdot \vec{\alpha}, \epsilon_1, \epsilon_2)^{-1} \quad \text{for } Z = Z_{1\text{-loop}}^{\text{vec}}, \quad (2.32)$$

$$\begin{aligned} \frac{Z^{(N),d}(\vec{k}) Z^{(S),d}(\vec{k})}{Z} &= e^{-\frac{I_2(\mathbf{R}_\ell)}{4}(\vec{a} \cdot \vec{k}) + \frac{I_3(\mathbf{R}_\ell)}{4} d_{ijk} a^i k^j k^k} (p_1 p_2)^{\frac{I_2(\mathbf{R}_\ell)}{8}(\vec{k} \cdot \vec{k}) - \frac{I_3(\mathbf{R}_\ell)}{12} d_{ijk} k^i k^j k^k} \\ &\quad \times y_\ell^{-\frac{I_2(\mathbf{R}_\ell)}{4}(\vec{k} \cdot \vec{k})} \prod_{\omega \in \mathbf{R}_\ell} \mathcal{L}_{\vec{k}, \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_{\text{phy}, \ell}, \epsilon_1, \epsilon_2) \quad \text{for } Z = Z_{1\text{-loop}}^{\text{hyp}, \ell}. \end{aligned} \quad (2.33)$$

Here we used $I_2(\mathbf{adj}) = 2h^\vee$, $I_3(\mathbf{adj}) = 0$, and also the fact h_{ij} and d_{ijk} are totally symmetric. The function $\mathcal{L}_k(x, \epsilon_1, \epsilon_2)$ is introduced to denote concisely the combination of the PE parts:

$$\mathcal{L}_k(x, \epsilon_1, \epsilon_2) \equiv \text{PE} \left[\left(\frac{p_1^k p_2}{(1-p_1)(1-p_2/p_1)} + \frac{p_1 p_2^k}{(1-p_1/p_2)(1-p_2)} - \frac{p_1 p_2}{(1-p_1)(1-p_2)} \right) \times e^{-x} \right] \quad (2.34)$$

One can easily check that the expression inside the PE vanishes at $k = 0, 1$. After some work, it is not difficult to find that

$$\mathcal{L}_k(x, \epsilon_1, \epsilon_2) = \begin{cases} \prod_{m+n \leq k-2} (1 - p_1^{m+1} p_2^{n+1} e^{-x}) & \text{for } k \geq +2 \\ \prod_{m+n \leq -k-1} (1 - p_1^{-m} p_2^{-n} e^{-x}) & \text{for } k \leq -1 \\ 1 & \text{for } k = 0, 1. \end{cases} \quad (2.35)$$

Combining them all, the resursion formula on the n -instanton piece Z_n can be written as

$$\begin{aligned} Z_n &= \sum_{\frac{1}{2}\vec{k} \cdot \vec{k} + \ell + m = n} \left((p_1 p_2)^{(\frac{h^\vee}{2} - \frac{\kappa_{\text{eff}}}{2} - d)(\frac{\vec{k} \cdot \vec{k}}{2}) + \frac{\kappa_{\text{eff}}}{6} d_{ijk} k^i k^j k^k} e^{(d + \frac{\kappa_{\text{eff}}}{2})(\vec{a} \cdot \vec{k})} e^{-\frac{\kappa_{\text{eff}}}{2} d_{ijk} a^i k^j k^k} \right. \\ &\quad \times \frac{\prod_l y_l^{-I_2(\mathbf{R}_l)(\vec{k} \cdot \vec{k}/4)} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k}, \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_{\text{phy}, l}, \epsilon_1, \epsilon_2)}{\prod_{\alpha \in \Delta} \mathcal{L}_{\vec{k}, \vec{\alpha}}(\vec{a} \cdot \vec{\alpha}, \epsilon_1, \epsilon_2)} \cdot p_1^{(\frac{b}{2}-d)\ell} p_2^{(\frac{b}{2}-d)m} Z_\ell^{(N)}(\vec{k}) Z_m^{(S)}(\vec{k}) \Big), \end{aligned} \quad (2.36)$$

which is generalization of the recursion formula found for the pure SYM case [25, 28]. The right-hand side of the recursion relation (2.36) only involves the instanton parts with $\ell, m < n$. Being so, one can determine the n -instanton correction Z_n recursively from $Z_0 = 1$.

2.2 One instanton partition function

At one instanton level, the formula (2.36) can be written as follows:

$$Z_1 = p_1^{(\frac{b}{2}-d)} Z_1^{(N)} + p_2^{(\frac{b}{2}-d)} Z_1^{(S)} + I_1^{(d)}, \quad (2.37)$$

where (with the fugacity variable $y_{\text{phy},l} \equiv e^{-m_{\text{phy},l}} = y_l/\sqrt{p_1 p_2}$ understood)

$$I_1^{(d)} = \sum_{\vec{k} \in \Delta_l} \left((p_1 p_2)^{(\frac{b}{2}-d)} e^{(d+\frac{\kappa_{\text{eff}}}{2})(\vec{a} \cdot \vec{k})} e^{-\frac{\kappa_{\text{eff}}}{2} d_{ijk} a^i k^j k^k} \right. \\ \left. \times \frac{\prod_l y_{\text{phy},l}^{-I_2(\mathbf{R}_l)/2} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_{\text{phy},l}, \epsilon_1, \epsilon_2)}{(1 - p_1 p_2 e^{-\vec{a} \cdot \vec{k}})(1 - p_1^{-1} e^{\vec{a} \cdot \vec{k}})(1 - p_2^{-1} e^{\vec{a} \cdot \vec{k}})(1 - e^{\vec{a} \cdot \vec{k}}) \prod_{\vec{\alpha} \cdot \vec{k} = -1} (1 - e^{-\vec{a} \cdot \vec{\alpha}})} \right). \quad (2.38)$$

We are interested in 5d $\mathcal{N} = 1$ gauge theories which are UV-complete as 5d $\mathcal{N} = 1$ SCFTs. Then the set of possible matter representation should be restricted to [42]

- fundamental representation for $SU(N)$, $SO(N)$, $Sp(N)$, G_2 , F_4 , E_6 , E_7
- antisymmetric representation for $SU(N)$, $Sp(N)$
- spinor representation for $SO(N)$ with $7 \leq N \leq 14$
- rank-3 antisymmetric representation for $Sp(3)$, $Sp(4)$, $SU(6)$, $SU(7)$
- symmetric representation for $SU(N)$.

For the $SU(N)$ symmetric representation, the inner product $\vec{k} \cdot \vec{\omega}$ can be either 0, ± 1 , or ± 2 .

$$\prod_{\omega \in \text{sym}} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_{\text{phy}}, \epsilon_1, \epsilon_2) = \prod_{\vec{k} \cdot \vec{\omega} = 2} (1 - y_{\text{phy}} p_1 p_2 e^{-\vec{a} \cdot \vec{\omega}}) \times \prod_{\vec{k} \cdot \vec{\omega} = -1} (1 - y_{\text{phy}} e^{-\vec{a} \cdot \vec{\omega}}) \\ \times \prod_{\vec{k} \cdot \vec{\omega} = -2} (1 - y_{\text{phy}} p_1^{-1} e^{-\vec{a} \cdot \vec{\omega}})(1 - y_{\text{phy}} p_2^{-1} e^{-\vec{a} \cdot \vec{\omega}}). \quad (2.39)$$

For the other cases, the inner product $\vec{k} \cdot \vec{\omega}$ can be either 0, ± 1 , such that

$$\prod_{\omega \in \mathbf{R}} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_{\text{phy}}, \epsilon_1, \epsilon_2) = \prod_{\vec{k} \cdot \vec{\omega} = -1} (1 - y_{\text{phy}} e^{-\vec{a} \cdot \vec{\omega}}). \quad (2.40)$$

Number of the blowup equations Prior to solving the equations for Z_1 , we recall that the formula (2.36) is valid only in a certain range of d , for which $\langle \mathcal{O}_{\mathbb{P}^1}^d \rangle = 1$. We want to narrow down the validity range of d by doing a simple sanity check on (2.37). Specifically, we want to examine the expansion of all single terms in (2.37) with respect to $\sqrt{p_1 p_2} \ll 1$, identifying the leading exponent as follows:

$$I_1^{(d)} \sim \begin{cases} g_0(\vec{a}, \vec{m}_{\text{phy}}) \cdot (p_1 p_2)^{\frac{b}{2}-d+1} + \dots & \text{for } N_{\text{sym}} = 0 \\ g_0(\vec{a}, \vec{m}_{\text{phy}}) \cdot (p_1 p_2)^{\frac{b}{2}-d} + \dots & \text{for } N_{\text{sym}} = 1 \end{cases} \\ Z_1 \sim g_1(\vec{a}, \vec{m}_{\text{phy}}) \cdot (p_1 p_2)^{\frac{n}{2}} + \dots \\ p_1^{(\frac{b}{2}-d)} Z_1^{(N)} \sim p_2^{(\frac{b}{2}-d)} Z_1^{(S)} \sim g_2(\vec{a}, \vec{m}_{\text{phy}}) \cdot (p_1 p_2)^{\frac{b}{4}-\frac{d}{2}+\frac{n}{4}} + \dots \quad (2.41)$$

where a numerical value of n will be discussed shortly for a variety of gauge theories (Table 1). Notice that for the equation (2.37) to be true, some terms on the right-hand side should have the leading exponent less than or equal to that of Z_1 . Therefore, the condition $d - \frac{b}{2} \geq -\frac{n}{2}$ is naturally imposed, setting a lower bound on d .

Similarly, an upper bound on d can be found from an expansion of (2.37) with respect to $1/\sqrt{p_1 p_2} \ll 1$.⁶ Each single term in (2.37) can be written as

$$\begin{aligned} I_1^{(d)} &\sim \begin{cases} h_0(\vec{a}, \vec{m}_{\text{phy}}) \cdot (1/p_1 p_2)^{d-\frac{b}{2}+1} + \dots & \text{for } N_{\text{sym}} = 0 \\ h_0(\vec{a}, \vec{m}_{\text{phy}}) \cdot (1/p_1 p_2)^{d-\frac{b}{2}} + \dots & \text{for } N_{\text{sym}} = 1 \end{cases} \\ Z_1 &\sim h_1(\vec{a}, \vec{m}_{\text{phy}}) \cdot (1/p_1 p_2)^{\frac{n'}{2}} + \dots \\ p_1^{(\frac{b}{2}-d)} Z_1^{(N)} \sim p_2^{(\frac{b}{2}-d)} Z_1^{(S)} &\sim h_2(\vec{a}, \vec{m}_{\text{phy}}) \cdot (1/p_1 p_2)^{\frac{d-b}{2}+\frac{n'}{4}} + \dots \end{aligned} \quad (2.42)$$

Again, for (2.37) to be consistent, the leading exponent of Z_1 should be greater than or equal to those of some terms on the right-hand side. Such requirement imposes the upper bound on d , namely $\frac{n'}{2} \geq d - \frac{b}{2}$. Combining the two inequalities, one can identify the following range

$$-\frac{n}{2} + \frac{b}{2} \leq d \leq \frac{n'}{2} + \frac{b}{2}, \quad (2.43)$$

as a necessary condition for (2.37). Furthermore, we explicitly checked that the k -instanton partition function Z_k actually satisfies all the $(\frac{n+n'}{2})$ recursion relations up to a certain value of k , for numerous examples (Table 1) whose Z_k is already known from alternative methods, even though the bound (2.43) itself is merely a *necessary* condition found from one-instanton analysis. Recall that the simple selection rule, governed by $\mathbb{Z}_{4h^\vee-2\sum_i I_2(\mathbf{R}_i)} \subset U(1)_R$, supports the validity of (2.19) at all instanton orders in 4d Donaldson-twisted theories. Based on the empirical observations, we claim that the 5d recursion formulae (2.36) within the above range of d are again true at all instanton orders. This implies that the non-perturbative partition function Z_{inst} is completely fixed by the perturbative partition function! Note that we have arrived at this statement not by requiring the perturbative series to be well-defined, as is often done in the resurgence analysis, but just by demanding consistency upon smooth deformation of the spacetime $\mathbb{C}^2 \times S^1$. Such consistency condition turns out to be entirely non-perturbative.

Another remarkable thing is that a numerical value of (n, n') exhibits the very simple pattern across a broad range of theories whose gauge group is not $SU(N)_\kappa$. (See Table 1.)

$$\begin{aligned} n = n' &= h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) & \text{for } G \neq SU(N)_\kappa \text{ nor } Sp(N) \\ n = n' - 2\{\frac{N_f}{2}\} &= h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) & \text{for } G = Sp(N)_{\theta=0} \\ n = n' + 2\{\frac{N_f}{2}\} &= h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + 1 & \text{for } G = Sp(N)_{\theta=\pi} \end{aligned} \quad (2.44)$$

⁶This is equivalent to assuming a different parameter regime $0 < a_i < -\epsilon_+ < m$ for all $1 \leq i \leq r$. In general, an explicit form of the 1-loop partition function (2.28)–(2.29) can change depending on a parameter regime, thus affecting (2.36). However, all the above expressions remain valid under flipping a sign of ϵ_+ , such that we can simply study the expansion of the single terms in (2.37) with respect to $1/\sqrt{p_1 p_2} \ll 1$.

As the above numerical pattern (2.44) emerges for all $G \neq SU(N)_\kappa$ examples that we studied, we speculate that (2.44) is generally true, thereby taking the recursion formulae (2.36) with

$$\begin{aligned} 0 \leq d \leq h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) & \quad \text{for } G \neq SU(N)_\kappa \text{ nor } Sp(N) \\ 0 \leq d \leq h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + \{\frac{N_f}{2}\} & \quad \text{for } G = Sp(N)_{\theta=0} \\ -\frac{1}{2} \leq d \leq h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + \frac{1}{2} - \{\frac{N_f}{2}\} & \quad \text{for } G = Sp(N)_{\theta=\pi} \end{aligned} \quad (2.45)$$

as a basic assumption to obtain the partition function \mathcal{Z} for any $G \neq SU(N)_\kappa$ gauge theory. We will derive the general expression for the 1-instanton partition function Z_1 , which applies for all $G \neq SU(N)_\kappa$ gauge theories having $h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) \geq 2$. It would be desirable to understand from the first principle the range (2.45) of d for which (2.36) holds true.

On the contrary, it turns out to be more difficult to characterize a general pattern behind (n, n') for $SU(N)_\kappa$ gauge theories, due to extra complication caused by the 5d Chern-Simons level κ . Here we consider two particular classes of $SU(N)_\kappa$ gauge theories as an illustration. In $SU(N)_\kappa + N_f \mathbf{F}$ gauge theory with $N_f + 2|\kappa| \leq 2N$, we find that

$$\begin{aligned} n &= \begin{cases} \frac{N_f}{2} & \text{if } \kappa_{\text{eff}} = N - N_f, \\ h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{F}) + |\kappa_{\text{eff}}| & \text{otherwise.} \end{cases} \\ n' &= \begin{cases} \frac{N_f}{2} & \text{if } \kappa_{\text{eff}} = -N, \\ h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{F}) + |\kappa_{\text{eff}} + \sum_l I_3(\mathbf{F})| & \text{otherwise.} \end{cases} \end{aligned} \quad (2.46)$$

Plugging these values in (2.43), the range of d becomes

$$\begin{aligned} 0 \leq d \leq N & \quad \text{if } \kappa = -N + \frac{N_f}{2} \\ 0 \leq d \leq N - \frac{N_f}{2} - \kappa & \quad \text{if } \kappa \in (-N + \frac{N_f}{2}, -\frac{N_f}{2}] \\ 0 \leq d \leq N & \quad \text{if } \kappa \in [-\frac{N_f}{2}, +\frac{N_f}{2}] \\ \frac{N_f}{2} - \kappa \leq d \leq N & \quad \text{if } \kappa \in [\frac{N_f}{2}, N - \frac{N_f}{2}) \\ 0 \leq d \leq N & \quad \text{if } \kappa = N - \frac{N_f}{2} \end{aligned} \quad (2.47)$$

including $0 \leq d \leq N$ in any case. Thus for $N \geq 2$, the recursion formula (2.36) determines the partition function \mathcal{Z} completely. In $SU(N)_\kappa + N_f \mathbf{F} + 1\mathbf{AS}$ theory with $N_f + 2|\kappa| \leq N + 4$,

$$\begin{aligned} n &= \min(h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) - (\kappa_{\text{eff}} - 2), h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + 2\{\frac{\kappa_{\text{eff}}}{2}\}) \\ n' &= \min(h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + (\kappa_{\text{eff}} + \sum_l I_3(\mathbf{R}_l) + 2), \\ & \quad h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + 2\{-\frac{\kappa_{\text{eff}}}{2} + \frac{1}{2} \sum_l I_3(\mathbf{R}_l)\}), \end{aligned} \quad (2.48)$$

from which one can identify the validity range of d by making substitution to (2.43). So long as three distinct integers are allowed for d , at a given (N, κ) , the corresponding partition function \mathcal{Z} can be solved from the recursion formula (2.36). We will also consider $SU(6)_\kappa + 1\mathbf{TAS}$ theory with $|\kappa| \leq 3$ in Section 3. This model can be Higgsed to two disjoint copies of $SU(3)_\kappa$

theory without a bifundamental hypermultiplet [43]. At the level of the partition function, the Higgsing is realized by turning off $m_{\text{phys}} = 0$ and imposing $SU(3)$ traceless conditions. As neither of them can modify n and n' , the numerical value of (n, n') must be identical to that of $SU(3)_\kappa$ gauge theory, which always contains $0 \leq d \leq 3$. Therefore, the recursion formula (2.36) sufficiently determines the full partition function \mathcal{Z} for $SU(6)_\kappa + 1\mathbf{TAS}$ theory as well.

Solving the recursion formulae We now turn to solve the recursion formulae (2.36) for the one-instanton partition function Z_1 . As long as there are at least 3 consecutive integers $\{d_0, d_0 + 1, d_0 + 2\}$ for d , the solution Z_1 to the three linear equations (2.37) is given by

$$Z_1(\vec{a}, \epsilon_1, \epsilon_2, \vec{m}) = \frac{p_1 p_2 I_1^{(d_0+2)} - (p_1 + p_2) I_1^{(d_0+1)} + I_1^{(d_0)}}{(1-p_1)(1-p_2)} \quad (2.49)$$

with $I_1^{(d)}$ defined in (2.38). It turns out to be more convenient to express (2.49) by decomposing $I_1^{(d)}$ into the flux sum, i.e., $I_1^{(d)} \equiv \sum_{\vec{k} \in \Delta_l} i_1^{(d)}(\vec{k})$, where

$$i_1^{(d)}(\vec{k}) \equiv (p_1 p_2)^{\frac{b}{2}-d} e^{(d+\frac{\kappa_{\text{eff}}}{2})(\vec{a} \cdot \vec{k})} e^{-\frac{\kappa_{\text{eff}}}{2} d_{ijk} a^i k^j k^k} \quad (2.50)$$

$$\times \frac{\prod_l (y_l^{\text{phy}})^{-I_2(\mathbf{R}_l)/2} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_{\text{phys}, l}, \epsilon_1, \epsilon_2)}{(1-p_1 p_2 e^{-\vec{a} \cdot \vec{k}})(1-e^{\vec{a} \cdot \vec{k}})(1-p_1^{-1} e^{\vec{a} \cdot \vec{k}})(1-p_2^{-1} e^{\vec{a} \cdot \vec{k}}) \prod_{\vec{\alpha} \cdot \vec{k} = -1} (1-e^{-\vec{a} \cdot \vec{\alpha}})}.$$

Using the property $i_1^{(d_0+\aleph)}(\vec{k})/i_1^{(d_0)}(\vec{k}) = (p_1 p_2)^{-\aleph} e^{\aleph(\vec{a} \cdot \vec{k})}$, the above Z_1 can be written as

$$Z_1 = \sum_{\vec{k} \in \Delta_l} \frac{(1-p_1^{-1} e^{\vec{a} \cdot \vec{k}})(1-p_2^{-1} e^{\vec{a} \cdot \vec{k}})}{(1-p_1)(1-p_2)} \cdot i_1^{(d_0)}(\vec{k}) \quad (2.51)$$

where the factor $\prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_l^{\text{phy}}, \epsilon_1, \epsilon_2)$ in (2.50) is explicitly written in (2.39) and (2.40). Notice that there are multiple options for choosing d_0 in case of $\frac{n+n'}{2} > 2$. However, it was observed that (2.51) is independent of a specific choice of d_0 . Once we choose $d_0 = 0$, for instance, which is an available choice in most cases,⁷ (2.51) becomes

$$Z_1 = \frac{(p_1 p_2)^{\frac{b}{2}} \prod_l (y_l^{\text{phy}})^{-I_2(\mathbf{R}_l)/2}}{(1-p_1)(1-p_2)} \sum_{\vec{k} \in \Delta_l} \frac{e^{\frac{\kappa_{\text{eff}}}{2}(\vec{a} \cdot \vec{k}) - \frac{\kappa_{\text{eff}}}{2} d_{ijk} a^i k^j k^k} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k} \cdot \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_l^{\text{phy}}, \epsilon_{1,2})}{(1-p_1 p_2 e^{-\vec{a} \cdot \vec{k}})(1-e^{\vec{a} \cdot \vec{k}}) \prod_{\vec{\alpha} \cdot \vec{k} = -1} (1-e^{-\vec{a} \cdot \vec{\alpha}})}. \quad (2.52)$$

We claim that (2.51) is the closed-form expression for the one-instanton partition function, which *universally* holds for any gauge theory having $\frac{n+n'}{2} \geq 2$.

More generally, the k -instanton correction Z_k can be determined by solving the following set of equations:

$$Z_k = p_1^{k(\frac{b}{2}-d)} Z_k^{(N)} + p_2^{k(\frac{b}{2}-d)} Z_k^{(S)} + I_k^{(d)} \quad \text{with an allowed range of } d, \quad (2.53)$$

⁷A numerical value of d_0 should be a half-integer for theories with $G = Sp(N)_{\theta=\pi}$.

where $I_k^{(d)}$ is defined as

$$I_k^{(d)} = \sum_{\substack{\frac{1}{2}\vec{k}\cdot\vec{k}+\ell+m=k \\ \ell, m \neq k}} \left((p_1 p_2)^{(\frac{h^\vee}{2} - \frac{\kappa_{\text{eff}}}{2} - d)(\frac{\vec{k}\cdot\vec{k}}{2}) + \frac{\kappa_{\text{eff}}}{6} d_{ijk} k^i k^j k^k} e^{(d + \frac{\kappa_{\text{eff}}}{2})(\vec{a}\cdot\vec{k})} e^{-\frac{\kappa_{\text{eff}}}{2} d_{ijk} a^i k^j k^k} \right. \quad (2.54) \\ \left. \times \frac{\prod_l y_l^{-I_2(\mathbf{R}_l)(\vec{k}\cdot\vec{k}/4)} \prod_{\omega \in \mathbf{R}_l} \mathcal{L}_{\vec{k}, \vec{\omega}}(\vec{a} \cdot \vec{\omega} + m_{\text{phy}, l}, \epsilon_1, \epsilon_2)}{\prod_{\alpha \in \Delta} \mathcal{L}_{\vec{k}, \vec{\alpha}}(\vec{a} \cdot \vec{\alpha}, \epsilon_1, \epsilon_2)} \cdot p_1^{(\frac{b}{2}-d)\ell} p_2^{(\frac{b}{2}-d)m} Z_\ell^{(N)}(\vec{k}) Z_m^{(S)}(\vec{k}) \right).$$

The k -instanton partition function Z_k is given as the solution to the three linear equations (2.53) with consecutive integers $\{d_0, d_0 + 1, d_0 + 2\}$.

$$Z_k(\vec{a}, \epsilon_1, \epsilon_2, \vec{m}) = \frac{p_1^k p_2^k I_k^{(d_0+2)} - (p_1^k + p_2^k) I_k^{(d_0+1)} + I_k^{(d_0)}}{(1 - p_1^k)(1 - p_2^k)} \quad (2.55)$$

Since $I_k^{(d)}$ only involves low-order instanton corrections, the k -instanton partition function Z_k can be constructed from $Z_{m < k}$, allowing us to obtain the full non-perturbative part Z_{inst} in a recursive manner. Again, this approach is applicable for any gauge theory having $\frac{n+n'}{2} \geq 2$.

3 Examples

As a remarkable application of the blow-up equation, we construct the instanton partition function of various 5d gauge theories on $\mathbb{C}^2 \times S^1$. The standard method to compute the multi-instanton correction to the partition function is to employ the ADHM construction of the instanton moduli space [3, 4, 44], and/or to apply the topological vertex formalism based on the geometric engineering of 5d $\mathcal{N} = 1$ gauge theory [45, 46]. An alternative approach that comes from the consistency requirement for the blow-up $\hat{\mathbb{C}}^2$ will turn out to be very efficient for bootstrapping the $\mathbb{C}^2 \times S^1$ partition function of exceptional gauge theories [16, 32, 47].

There are multiple and distinct blow-up equations (??) at different values of d , ranged over $0 \leq d \leq d_{\text{max}}$ where $d_{\text{max}} \equiv h^\vee - \frac{1}{2} \sum_i I(\mathbf{R}_i)$. We need at least three distinct equations to solve for the instanton partition function, thus the bootstrap method is applicable only for gauge theories which satisfy $d_{\text{max}} \geq 2$. Here we constrain our discussion to the theories having simple gauge groups, which are classified to have a 5d UV fixed point [42].

There are two

We will

The complete list of $\mathcal{N} = 1$ gauge theories, satisfying $d_{\text{max}} \geq 2$, is displayed in Table 2 and 3. All infinite families of theories, whose rank of the gauge group can be arbitrarily large, are summarized in Table 2. There are also finite families of theories, summarized in Table 3, which involve an exceptional gauge group or a spinor hypermultiplet.

(comparison with IMS bound, etc) (SCFT bound) Some theories in Table 3 do not belong to the

3.1 Known examples

3.2 Ki-Hong's note

Unity Blowup equations

Instanton partition functions from blowup equations From blowup equations one can compute the partition functions as follows. Rewriting the blowup equation as

$$1 = \sum_{\vec{k} \in \vec{\alpha}^\vee} f_{\vec{k}} l_{\vec{k}} \frac{Z_{\text{inst}}^{(1)} Z_{\text{inst}}^{(2)}}{Z_{\text{inst}}} \quad (3.1)$$

where $f_{\vec{k}} = Z_{\text{class}}^{(1)} Z_{\text{class}}^{(2)} / Z_{\text{class}}$ and $l_{\vec{k}} = Z_{1\text{-loop}}^{(1)} Z_{1\text{-loop}}^{(2)} / Z_{1\text{-loop}}$ with abbreviated notation

$$\begin{aligned} Z^{(1)} &= Z(\epsilon_1, \epsilon_2 - \epsilon_1, \vec{a} + \vec{k} \epsilon_1, m_i + r_i \epsilon_1, m_0 + r_0 \epsilon_1) \\ Z^{(2)} &= Z(\epsilon_1 - \epsilon_2, \epsilon_2, \vec{a} + \vec{k} \epsilon_2, m_i + r_i \epsilon_2, m_0 + r_0 \epsilon_2) \end{aligned} \quad (3.2)$$

Here note that $l_{\vec{k}}$ is independent of $Q = e^{-m_0}$, and $f_{\vec{k}}$ is some overall factor in the order of $Q^{\vec{k} \cdot \vec{k}/2}$. Expanding the equation by instanton fugacity Q , then at each Q^n level the equation is written by

$$\delta_{n,0} = p_1^{r_0} Z_n^{(1)} + p_2^{r_0} Z_n^{(2)} - Z_n + \sum_{\vec{k} \neq 0} f_{\vec{k}, r_0} l_{\vec{k}} \left(\frac{Z_{\text{inst}}^{(1)} Z_{\text{inst}}^{(2)}}{Z_{\text{inst}}} \right) \Big|_{O(Q^{n - \vec{k} \cdot \vec{k}/2})}. \quad (3.3)$$

Since each Z_k and $Z_k^{(1,2)}$ are independent of r_0 , one can solve (3.3) with three blowup equations with same r_i 's but different r_0 's.

The blowup equations for instanton partition functions of pure YM theory with generic gauge group were already studied in [16]. They are actually (??) with

$$\vec{r}_a = 0, \quad r_0 = d - h^\vee / 2 \quad (3.4)$$

where $d = 0, \dots, h^\vee$. We extend these blowup equations to the theories with matters based on pure YM blowup equations. If one restrict the cases to $\vec{r}_a = 0$, as we explained in the previous section, the r_i 's are technically required to be half intergers. Thus we look for the r_0 's that provides the correct instanton partition functions by solving (3.3) while fixing $\vec{r}_a = 0$ and $r_i = 1/2$. Here are the results.

They were tested by comparing the resulting instanton partition functions with the known results from [48] ($SO(7)$ and G_2) and [49] (F_4 with $N_{26} = 2$). They were compared numerically, putting random numbers on the fugacities. Note that matters shift the r_0 , each by one quarter of their Dynkin indices. It seems to differ from blowup formula for $SU(N)_\kappa + N_f$ instantons, where r_0 was affected only by its CS-level κ . However, one can rewrite the r_0 as

$$r_0 = d - N/2 - \left(\kappa + \frac{1}{2} N_f \right) / 2 + N_f / 4$$

$$= d - N/2 - \kappa_{\text{eff}}/2 + N_f \times I_{\text{fund}}. \quad (3.5)$$

Since fundamental matters shifts the effective CS-level, they cancel their index contributions and consequently the r_0 apparently looks independent of matters.

By above observations, we write the unity blowup equation for generic gauge groups and matter representations.

$$\begin{aligned} Z(\epsilon_1, \epsilon_2, \vec{a}, m_i, m_0) &= \sum_{\vec{k} \in \vec{a}^\vee} Z(\epsilon_1, \epsilon_2 - \epsilon_1, \vec{a} + \vec{k}\epsilon_1, m_i + \epsilon_1/2, m_0 + r_0\epsilon_1) \\ &\times Z(\epsilon_1 - \epsilon_2, \vec{a} + \vec{k}\epsilon_2, m_i + \epsilon_2/2, m_0 + r_0\epsilon_2) \end{aligned} \quad (3.6)$$

with

$$r_0 = d - h^\vee/2 - \kappa_{\text{eff}}/2 + N_{\mathbf{R}} \times I_{\mathbf{R}}. \quad (3.7)$$

Here $I_{\mathbf{R}}$ is the Dynkin index of \mathbf{R} representation.

$SU(6)_3 + 1 \times \mathbf{20}$ As a non-trivial test, we consider the instanton partition function of the $SU(6)_3 + \mathbf{20}$ whose 5-brane realization was found recently [43]. Its web-diagram is given as figure.

(Written before computing the $SU(6)_3 + \mathbf{20}$ instanton partition function.)

Rather than comparing instanton partition functions directly, we consider an interesting Higgsing procedure. We consider the $SU(3) \times SU(3) \times U(1) \subset SU(6)$ where the $SU(6)$ multiplets are decomposed by

$$\begin{aligned} A_{i\bar{j}} : \mathbf{35} &\longrightarrow (\mathbf{8}, 1)_0 \oplus (1, \mathbf{8})_0 \oplus (\mathbf{3}, \bar{\mathbf{3}})_2 \oplus (\bar{\mathbf{3}}, \mathbf{3})_{-2} \oplus (1, 1)_0, \\ \Phi_{ijk} : \mathbf{20} &\longrightarrow (\mathbf{3}, \bar{\mathbf{3}})_{-1} \oplus (\bar{\mathbf{3}}, \mathbf{3})_1 \oplus (1, 1)_3 \oplus (1, 1)_{-3}. \end{aligned} \quad (3.8)$$

Here to fit with the web-diagram, we set Φ_{156} and Φ_{234} are $(1, 1)_3$ and $(1, 1)_{-3}$. Once Φ_{156} and Φ_{234} get non-zero VEVs,

When $a_5 = -a_1 - a_6$, the web-diagram factorizes to two $SU(3)_3$ whose Coulomb VEVs are (a_1, a_5, a_6) and (a_2, a_3, a_4) . In the gauge theory, it can be seen partly from prepotential. The prepotential of $S(6)_3 + 1 \times \mathbf{20}$ is

$$\mathcal{F} = \frac{1}{2}m_0 \sum_{i=1}^6 a_i^2 + \frac{1}{2} \sum_{i=1}^6 a_i^3 + \frac{1}{6} \sum_{i < j} (a_i - a_j)^3 - \frac{1}{6} \sum_{1 < i < j} (a_1 + a_j + a_k)^3 \quad (3.9)$$

at the Weyl chamber $a_1 > \dots > a_6$. As one sets the Coulomb VEV $a_6 = -a_1 - a_5$ and $a_4 = -a_2 - a_3$, one can check

$$\mathcal{F}(m_0, a_1, a_2, a_3, a_4, a_5, a_6) = \mathcal{F}_{SU(3)_3}(m_0, a_1, a_5, a_6) + \mathcal{F}_{SU(3)_3}(m_0, a_2, a_3, a_4) \quad (3.10)$$

where

$$\mathcal{F}_{SU(3)_3}(m_0, a_1, a_2, a_3) = \frac{1}{2}m_0 \sum_{i=1}^3 a_i^2 + \frac{1}{2} \sum_{i=1}^3 a_i^3 + \frac{1}{6} \sum_{i < j} (a_i - a_j)^3. \quad (3.11)$$

It is Higgsed by

3.3 $SU(6)_3$ with a rank-3 antisymmetric hyper

(describe blow-up computation)

It was recently found in [43] that the 5d $SU(6)_3$ gauge theory with a rank-3 antisymmetric hypermultiplet can be engineered from the 5-brane web configuration, depicted in Figure 1. Given a web diagram, we utilize the topological vertex method [46, 50] to compute all genus topological string amplitudes, which is the logarithm of the 5d Nekrasov partition function on Ω -deformed $\mathbf{R}^4 \times S^1$ [45]. We will check its agreement with the blowup partition function (eqn), providing a supporting evidence to the suggested blow-up equation (eqn).

By applying the topological vertex method to Figure 1, we find that the instanton partition function can be written as the following sum over all possible 6 Young diagrams:

$$Z_{\text{Nek}} = \sum_{(Y_1, \dots, Y_6)} q^{\sum_{i=1}^6 |Y_i|} (-A_1^6)^{|Y_1|} (-A_2^6)^{|Y_2|} (-A_2^2 A_3^4)^{|Y_3|} (-A_2^2 A_3^2 A_4^2)^{|Y_4|+|Y_5|} \\ \times f_{Y_1}(g)^5 f_{Y_2}(g)^5 f_{Y_3}(g)^3 f_{Y_4}(g) f_{Y_5}(g)^{-1} f_{Y_6}(g)^2 Z_{\text{half}}(Y_1, Y_2, Y_3, Y_4, Y_5, Y_6)^2. \quad (3.12)$$

where the Kähler parameters can be identified as

$$A_i = e^{-a_i}, \quad q = e^{-\frac{8\pi^2}{g^2}}, \quad g = e^{-\epsilon_-}. \quad (3.13)$$

We briefly explain our notation: For a given Young diagram μ , $|\mu|$ denotes the total number of boxes. μ_i is the number of boxes in the i -th row of μ . μ^t is the transpose of μ . We also use

$$f_\mu(g) = (-1)^{|\mu|} g^{\frac{1}{2}(\|\mu^t\|^2 - \|\mu\|^2)}, \quad \tilde{Z}_\lambda(g) = \prod_{(i,j) \in \lambda} (1 - g^{\lambda_i + \lambda_j^t - i - j + 1})^{-1} \quad (3.14)$$

with $\|\mu\|^2 = \sum_i \mu_i^2$. The factor $Z_{\text{half}}(Y_1, Y_2, Y_3, Y_4, Y_5, Y_6)$ involves a single summation over all possible Young diagrams, i.e.,

$$Z_{\text{left}}(\vec{Y}) = \sum_{Y_0} (-A_1^{-1} A_6^{-2})^{|Y_0|} g^{\frac{\|Y_0^t\|^2 + \|Y_0\|^2}{2}} \tilde{Z}_{Y_0}(g)^2 f_{Y_0}(g)^2 \prod_{i=1}^6 g^{\frac{\|Y_i\|^2}{2}} \tilde{Z}_{Y_i}(g) \\ \times R_{Y_1 Y_6^t}^{-1}(A_1 A_6^{-1}) R_{Y_0 Y_6^t}^{-1}(A_1^{-1} A_6^{-2}) R_{Y_1 Y_0^t}^{-1}(A_1^2 A_6) \\ \times \prod_{2 \leq i < j \leq 5} R_{Y_i Y_j^t}^{-1}(A_i A_j^{-1}) \prod_{i=2}^5 R_{Y_0^t Y_i}(A_1 A_i A_6), \quad (3.15)$$

in which we introduce

$$R_{\lambda\mu}(Q) = R_{\mu\lambda}(Q) = \text{PE} \left[-\frac{g}{(1-g)^2} Q \right] \times N_{\lambda^t\mu}(Q) \quad (3.16)$$

with PE representing the Plethystic exponential and

$$N_{\lambda\mu}(Q) = \prod_{(i,j) \in \lambda} (1 - Q g^{\lambda_i + \mu_j^t - i - j + 1}) \prod_{(i,j) \in \mu} (1 - Q g^{-\lambda_j^t - \mu_i + i + j - 1}). \quad (3.17)$$

We also recall that the Nekrasov partition function is divided into the perturbative partition function Z_{pert} and the weighted sum of k -instanton partition function Z_k .

$$Z_{\text{Nek}} = Z_{\text{pert}} \left(1 + \sum_{k=1}^{\infty} q^k Z_k \right). \quad (3.18)$$

The perturbative part of the partition function Z_{pert} comes from the summand of (3.12) at empty Young diagrams, i.e., $(Y_1, Y_2, Y_3, Y_4, Y_5, Y_6) = (\emptyset, \emptyset, \emptyset, \emptyset, \emptyset, \emptyset)$. It is given by

$$\begin{aligned} Z_{\text{pert}} &= Z_{\text{half}}(\emptyset, \emptyset, \emptyset, \emptyset, \emptyset, \emptyset)^2 \\ &= \text{PE} \left[\frac{2g}{(1-g)^2} \left(\frac{A_1}{A_6} + \frac{1}{A_1 A_6^2} + A_1^2 A_6 + \sum_{2 \leq i < j \leq 5} \frac{A_i}{A_j} - \sum_{i=2}^5 A_1 A_i A_6 \right) \right] \\ &\quad \times \left(\sum_{Y_0} (-A_1^{-1} A_6^{-2})^{|Y_0|} g^{\frac{\|Y_0^t\|^2 + \|Y_0\|^2}{2}} \tilde{Z}_{Y_0}(g)^2 f_{Y_0}^2(g) \right. \\ &\quad \left. N_{Y_0^t \emptyset}^{-1} (A_1^{-1} A_6^{-2}) N_{Y_0 \emptyset}^{-1} (A_1^2 A_6) \prod_{i=2}^5 N_{Y_0 \emptyset} (A_1 A_i A_6) \right)^2, \end{aligned} \quad (3.19)$$

where the last two lines can be combined into the following closed-form expression:

$$\text{PE} \left[\frac{2g}{(1-g)^2} \left(\sum_{i=2}^5 \frac{A_1}{A_i} + \sum_{i=2}^5 \frac{A_i}{A_6} - \frac{1}{A_1 A_6^2} - A_1^2 A_6 - \sum_{2 \leq i < j \leq 5} A_1 A_i A_j + \mathcal{O}(A_1^6) \right) \right]. \quad (3.20)$$

So (3.19) is manifestly consistent with the equivariant index [41] for 5d $SU(6)$ gauge theory with a hypermultiplet in the rank-3 antisymmetric representation **20**, i.e.,

$$Z_{\text{pert}} = \text{PE} \left[\frac{2g}{(1-g)^2} \left(\sum_{1 \leq i < j \leq 6} \frac{A_i}{A_j} - \sum_{2 \leq i < j \leq 6} A_1 A_i A_j + \mathcal{O}(A_1^6) \right) \right]. \quad (3.21)$$

The 1-instanton partition function Z_1 can be obtained from the summands of (3.12) at Young diagrams satisfying $\sum_{i=1}^6 |Y_i| = 1$. There are 6 different profiles of Young diagrams. The configuration $|Y_i| = 1$ and $Y_{j \neq i} = \emptyset$ contributes to Z_1 by

$$+ \frac{g}{(1-g)^2} \frac{A_i^6}{\prod_{j \neq i} (A_i - A_j)^2} \left(-A_i \sum_{j \neq i} A_j + \sum_{j \neq i} \frac{1}{A_j} - \frac{1}{A_i} + A_i^2 \right)^2. \quad (3.22)$$

Summing over all six contributions, we find

$$Z_1 = \sum_{i=1}^6 \frac{g}{(1-g)^2} \frac{A_i^6}{\prod_{j \neq i} (A_i - A_j)^2} \left(-A_i \sum_{j \neq i} A_j + \sum_{j \neq i} \frac{1}{A_j} - \frac{1}{A_i} + A_i^2 \right)^2. \quad (3.23)$$

which is in agreement with the blowup partition function (eqn). (overall sign: looking at the GV invariant for single W-boson + single instanton (which should be -2), the topological vertex computation seems to be correct. blowup should have an overall sign issue)




Figure 1: A 5-brane web for $SU(6)_3 + 1\text{TAS}$ with massless **TAS**.




Figure 2: (a) A Higgsing from from $SU(6)_3 + 1\text{TAS}$ to two $SU(3)_3$ theories. (a) Two $SU(3)_3$ theories are painted in blue and in pink respectively. A new decoupled mode emerges.




Figure 3: A labeling of Young diagrams assigned to the horizontal lines in Figure 1.

4 Conclusion

Acknowledgments

This work is supported in part by the UESTC Research Grant A03017023801317 (SSK), the National Research Foundation of Korea (NRF) Grants 2017R1D1A1B06034369 (KL, JS), and 2018R1A2B6004914 (KHL)

A $SU(N)$ theory

We find that

$$\begin{cases} n = \frac{N_f}{2} & \text{if } \kappa_{\text{eff}} = N - N_f, \\ n = h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + |\kappa_{\text{eff}}| & \text{otherwise.} \end{cases} \quad (\text{A.1})$$

for $SU(N)_\kappa + N_f \mathbf{F}$ theory with $N_f + 2|\kappa| \leq 2N$,

$$\begin{cases} n = h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) - (\kappa_{\text{eff}} - 2) & \text{if } N: \text{ odd and } \kappa_{\text{eff}} \geq 3 - \frac{N_f}{2}, \\ n = h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) - (\kappa_{\text{eff}} - 2) & \text{if } N: \text{ even and } \kappa_{\text{eff}} > 3 - \frac{N_f}{2}, \\ n \geq h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + 2\{\frac{\kappa_{\text{eff}}}{2}\} & \text{otherwise.} \end{cases} \quad (\text{A.2})$$

for $SU(N)_\kappa + N_f \mathbf{F} + 1\mathbf{A}$ theory with $N_f + 2|\kappa| \leq N + 4$.

$$\begin{cases} n' = \frac{N_f}{2} & \text{if } \kappa_{\text{eff}} = -N, \\ n' = h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + |\kappa + \frac{1}{2} \sum_l I_2(\mathbf{R}_l)| & \text{otherwise.} \end{cases}$$

for $SU(N)_\kappa + N_f \mathbf{F}$ theory with $N_f + 2|\kappa| \leq 2N$,

$$\begin{cases} n' = h^\vee + \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + \kappa_{\text{eff}} & \text{if } N: \text{ odd and } \kappa_{\text{eff}} \leq 1 - N - \frac{N_f}{2}, \\ n' = h^\vee + \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + \kappa_{\text{eff}} & \text{if } N: \text{ even and } \kappa_{\text{eff}} < 1 - N - \frac{N_f}{2}, \\ n' \geq h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) + 2\{-\frac{\kappa_{\text{eff}} + N + N_f}{2}\} & \text{otherwise.} \end{cases} \quad (\text{A.3})$$

for $SU(N)_\kappa + N_f \mathbf{F} + 1\mathbf{A}$ theory with $N_f + 2|\kappa| \leq N + 4$.

i.e., and

In other words, This suggests that, as long as $\frac{b+b'}{2} > 2$, we can solve for the instanton partition function Z_n

In summary, there are $(b+b')/2$ blowup equations

we have found $n = n' = b = h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l)$. Thus the range of d is constrained to

$$0 \leq d \leq h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) \quad \text{for the theories with } G \neq SU(N). \quad (\text{A.4})$$

See Appendix A for the pattern on n in two classes of

Especially when $\kappa_{\text{eff}} = 0$, the value of n is same as the above.

where we observe that

$$n' = h^\vee - \frac{1}{2} \sum_l I_2(\mathbf{R}_l) \quad (\text{A.5})$$

for the theories with $G \neq SU(N)$,

We need to be a bit careful when dealing with $SU(N)_\kappa$ theories. For $SU(N)_\kappa + N_f \mathbf{F}$ theory, the validity range of d becomes

$$\begin{cases} d \in [0, N] & \text{for } \kappa = -N + \frac{N_f}{2} \\ d \in [0, N - \frac{N_f}{2} - \kappa] & \text{for } \kappa \in (-N + \frac{N_f}{2}, -\frac{N_f}{2}] \\ d \in [0, N] & \text{for } \kappa \in [-\frac{N_f}{2}, +\frac{N_f}{2}] \\ d \in [\frac{N_f}{2} - \kappa, N] & \text{for } \kappa \in [+ \frac{N_f}{2}, N - \frac{N_f}{2}) \\ d \in [0, N] & \text{for } \kappa = +N - \frac{N_f}{2} \end{cases} \quad (\text{A.6})$$

which always includes $0 \leq d \leq N$. For $SU(N)_\kappa + N_f \mathbf{F} + 1\mathbf{A}$ theory,

Substituting (n, n') to (2.43), d takes a value in the following range:

$$\begin{cases} d \in [-1 - \frac{a}{4} - \{\frac{a}{4}\}, N] & \text{for } \kappa \in [-2 - \frac{N}{2} + \frac{N_f}{2}, 1 - N - \frac{N_f}{2}] \\ d \in [-1 - \frac{a}{4} - \{\frac{a}{4}\}, N - \frac{b}{2} - \{-\frac{b}{2}\}] & \text{for } \kappa \in (1 - N - \frac{N_f}{2}, 3 - \frac{N_f}{2}) \\ d \in [-1, N - \frac{b}{2} - \{-\frac{b}{2}\}] & \text{for } \kappa \in [3 - \frac{N_f}{2}, +\frac{N_f}{2}, 2 + \frac{N}{2} - \frac{N_f}{2}] \end{cases} \quad (\text{A.7})$$

with odd N , and

$$\begin{cases} d \in [-1 - \frac{a}{4} - \{\frac{a}{4}\}, N] & \text{for } \kappa \in [-2 - \frac{N}{2} + \frac{N_f}{2}, 1 - N - \frac{N_f}{2}] \\ d \in [-1 - \frac{a}{4} - \{\frac{a}{4}\}, N - \frac{b}{2} - \{-\frac{b}{2}\}] & \text{for } \kappa \in (1 - N - \frac{N_f}{2}, 3 - \frac{N_f}{2}) \\ d \in [-1, N - \frac{b}{2} - \{-\frac{b}{2}\}] & \text{for } \kappa \in [3 - \frac{N_f}{2}, +\frac{N_f}{2}, 2 + \frac{N}{2} - \frac{N_f}{2}] \end{cases} \quad (\text{A.8})$$

with even N , where a and b are defined by $a = 2\kappa - N - N_f$ and $b = 2\kappa + N + N_f$. Although these patterns look

$$-\frac{\kappa_{\text{eff}}}{2} - \{\frac{\kappa_{\text{eff}}}{2}\} \leq d \leq N \quad \text{for } -N \leq \kappa_{\text{eff}} \leq 1 - N - \frac{N_f}{2} \quad (\text{A.9})$$




Figure 4: asd

$$-\frac{\kappa_{\text{eff}}}{2} - \left\{ \frac{\kappa_{\text{eff}}}{2} \right\} \leq d \leq N + 1 - \frac{\kappa_{\text{eff}} + N + N_f}{2} + \left\{ -\frac{\kappa_{\text{eff}} + N + N_f}{2} \right\} \quad (\text{A.10})$$

$$-1 \leq d \leq N + 1 - \frac{\kappa_{\text{eff}} + N + N_f}{2} + \left\{ -\frac{\kappa_{\text{eff}} + N + N_f}{2} \right\} \quad (\text{A.11})$$

For $SU(N)_{|\kappa| < N}$ SYM without a hypermultiplet, $n = h^\vee + |\kappa|$ so that [28]

$$-\frac{\kappa + |\kappa|}{2} \leq d \leq h^\vee + \frac{|\kappa| - \kappa}{2}. \quad (\text{A.12})$$

References

- [1] N. Seiberg and E. Witten, *Electric–magnetic Duality, Monopole Condensation, and Confinement in $\mathcal{N}=2$ Supersymmetric Yang–Mills Theory*, *Nucl. Phys.* **B426** (1994) 19–52, [[hep-th/9407087](#)].
- [2] N. Seiberg and E. Witten, *Monopoles, Duality and Chiral Symmetry Breaking in $\mathcal{N}=2$ Supersymmetric QCD*, *Nucl. Phys.* **B431** (1994) 484–550, [[hep-th/9408099](#)].
- [3] N. A. Nekrasov, *Seiberg–Witten Prepotential from Instanton Counting*, *Adv. Theor. Math. Phys.* **7** (2003) 831–864, [[hep-th/0206161](#)].
- [4] N. Nekrasov and A. Okounkov, *Seiberg–Witten Theory and Random Partitions*, *Prog. Math.* **244** (2006) 525–596, [[hep-th/0306238](#)].
- [5] H. Nakajima and K. Yoshioka, *Instanton counting on blowup. 1.*, *Invent. Math.* **162** (2005) 313–355, [[math/0306198](#)].
- [6] A. Braverman and P. Etingof, *Instanton Counting via Affine Lie Algebras Ii: from Whittaker Vectors to the Seiberg–Witten Prepotential*, [[math/0409441](#)].
- [7] G. W. Moore, N. Nekrasov and S. Shatashvili, *Integrating over Higgs Branches*, *Commun. Math. Phys.* **209** (2000) 97–121, [[hep-th/9712241](#)].
- [8] U. Bruzzo, F. Fucito, J. F. Morales and A. Tanzini, *Multiinstanton Calculus and Equivariant Cohomology*, *JHEP* **05** (2003) 054, [[hep-th/0211108](#)].
- [9] N. Nekrasov and S. Shadchin, *Abcd of Instantons*, *Commun. Math. Phys.* **252** (2004) 359–391, [[hep-th/0404225](#)].
- [10] M. Mariño and N. Wyllard, *A Note on Instanton Counting for $\mathcal{N}=2$ Gauge Theories with Classical Gauge Groups*, *JHEP* **05** (2004) 021, [[hep-th/0404125](#)].

- [11] F. Fucito, J. F. Morales and R. Poghossian, *Instantons on Quivers and Orientifolds*, *JHEP* **10** (2004) 037, [[hep-th/0408090](#)].
- [12] L. Hollands, C. A. Keller and J. Song, *From So/Sp Instantons to W-Algebra Blocks*, *JHEP* **03** (2011) 053, [[1012.4468](#)].
- [13] L. Hollands, C. A. Keller and J. Song, *Towards a 4D/2D Correspondence for Sicilian Quivers*, *JHEP* **10** (2011) 100, [[1107.0973](#)].
- [14] S. Benvenuti, A. Hanany and N. Mekareeya, *The Hilbert Series of the One Instanton Moduli Space*, *JHEP* **06** (2010) 100, [[1005.3026](#)].
- [15] C. A. Keller, N. Mekareeya, J. Song and Y. Tachikawa, *The Abcdefg of Instantons and W-Algebras*, *JHEP* **03** (2012) 045, [[1111.5624](#)].
- [16] C. A. Keller and J. Song, *Counting Exceptional Instantons*, *JHEP* **07** (2012) 085, [[1205.4722](#)].
- [17] A. Hanany, N. Mekareeya and S. S. Razamat, *Hilbert Series for Moduli Spaces of Two Instantons*, *JHEP* **01** (2013) 070, [[1205.4741](#)].
- [18] S. Cremonesi, G. Ferlito, A. Hanany and N. Mekareeya, *Coulomb Branch and the Moduli Space of Instantons*, *JHEP* **12** (2014) 103, [[1408.6835](#)].
- [19] C. Hwang, J. Kim, S. Kim and J. Park, *General Instanton Counting and 5D SCFT*, *JHEP* **07** (2015) 063, [[1406.6793](#)].
- [20] K. Hori, H. Kim and P. Yi, *Witten Index and Wall Crossing*, *JHEP* **01** (2015) 124, [[1407.2567](#)].
- [21] C. Cordova and S.-H. Shao, *An Index Formula for Supersymmetric Quantum Mechanics*, [1406.7853](#).
- [22] F. Benini, R. Eager, K. Hori and Y. Tachikawa, *Elliptic Genera of 2d $\mathcal{N} = 2$ Gauge Theories*, *Commun. Math. Phys.* **333** (2015) 1241–1286, [[1308.4896](#)].
- [23] F. Benini, R. Eager, K. Hori and Y. Tachikawa, *Elliptic Genera of Two-Dimensional $\mathcal{N} = 2$ Gauge Theories with Rank-One Gauge Groups*, *Lett. Math. Phys.* **104** (2014) 465–493, [[1305.0533](#)].
- [24] H. Nakajima and K. Yoshioka, *Lectures on instanton counting*, in *CRM Workshop on Algebraic Structures and Moduli Spaces Montreal, Canada, July 14-20, 2003*, 2003. [math/0311058](#).
- [25] H. Nakajima and K. Yoshioka, *Instanton counting on blowup. II. K-theoretic partition function*, [math/0505553](#).
- [26] D. Gaiotto and S. S. Razamat, *Exceptional Indices*, *JHEP* **05** (2012) 145, [[1203.5517](#)].
- [27] N. A. Nekrasov, *Localizing Gauge Theories*, in *Mathematical Physics. Proceedings, 14Th International Congress, Icmp 2003, Lisbon, Portugal, July 28-August 2, 2003*, pp. 645–654, 2003.
- [28] L. Gottsche, H. Nakajima and K. Yoshioka, *K-theoretic Donaldson invariants via instanton counting*, *Pure Appl. Math. Quart.* **5** (2009) 1029–1111, [[math/0611945](#)].
- [29] L. Gottsche, H. Nakajima and K. Yoshioka, *Instanton counting and Donaldson invariants*, *J. Diff. Geom.* **80** (2008) 343–390, [[math/0606180](#)].
- [30] E. Gasparim and C.-C. M. Liu, *The Nekrasov Conjecture for Toric Surfaces*, *Commun. Math. Phys.* **293** (2010) 661–700, [[0808.0884](#)].

- [31] G. Bonelli, K. Maruyoshi, A. Tanzini and F. Yagi, $\mathcal{N}=2$ Gauge Theories on Toric Singularities, Blow-Up Formulae and W-Algebras, *JHEP* **01** (2013) 014, [[1208.0790](#)].
- [32] J. Gu, B. Haghighat, K. Sun and X. Wang, Blowup Equations for 6d SCFTs. I, [1811.02577](#).
- [33] J. Gu, A. Klemm, K. Sun and X. Wang, Elliptic Blowup Equations for 6D Scfts. II: Exceptional Cases, [1905.00864](#).
- [34] T. Sasaki, $O(-2)$ Blow-Up Formula via Instanton Calculus on Affine C^2/Z_2 and Weil Conjecture, [hep-th/0603162](#).
- [35] Y. Ito, K. Maruyoshi and T. Okuda, Scheme Dependence of Instanton Counting in ALE Spaces, *JHEP* **05** (2013) 045, [[1303.5765](#)].
- [36] T. Okuda and V. Pestun, On the instantons and the hypermultiplet mass of $N=2^*$ super Yang-Mills on S^4 , *JHEP* **03** (2012) 017, [[1004.1222](#)].
- [37] E. Witten, Topological Quantum Field Theory, *Commun. Math. Phys.* **117** (1988) 353.
- [38] L. Baulieu, A. Losev and N. Nekrasov, Chern-Simons and Twisted Supersymmetry in Various Dimensions, *Nucl. Phys.* **B522** (1998) 82–104, [[hep-th/9707174](#)].
- [39] A. Losev, G. W. Moore, N. Nekrasov and S. Shatashvili, Four-Dimensional Avatars of Two-Dimensional RCFT, *Nucl. Phys. Proc. Suppl.* **46** (1996) 130–145, [[hep-th/9509151](#)].
- [40] K. A. Intriligator, D. R. Morrison and N. Seiberg, Five-Dimensional Supersymmetric Gauge Theories and Degenerations of Calabi-Yau Spaces, *Nucl. Phys.* **B497** (1997) 56–100, [[hep-th/9702198](#)].
- [41] S. Shadchin, *On Certain Aspects of String Theory/Gauge Theory Correspondence*. PhD thesis, Orsay, LPT, 2005. [hep-th/0502180](#).
- [42] P. Jefferson, H.-C. Kim, C. Vafa and G. Zafrir, Towards Classification of 5D Scfts: Single Gauge Node, [1705.05836](#).
- [43] H. Hayashi, S.-S. Kim, K. Lee and F. Yagi, Rank-3 Antisymmetric Matter on 5-Brane Webs, [1902.04754](#).
- [44] M. F. Atiyah, N. J. Hitchin, V. G. Drinfeld and Yu. I. Manin, Construction of Instantons, *Phys. Lett.* **A65** (1978) 185–187.
- [45] R. Gopakumar and C. Vafa, *M Theory and Topological Strings. 2.*, [hep-th/9812127](#).
- [46] M. Aganagic, A. Klemm, M. Mariño and C. Vafa, The Topological Vertex, *Commun. Math. Phys.* **254** (2005) 425–478, [[hep-th/0305132](#)].
- [47] M.-x. Huang, K. Sun and X. Wang, Blowup Equations for Refined Topological Strings, *JHEP* **10** (2018) 196, [[1711.09884](#)].
- [48] H.-C. Kim, J. Kim, S. Kim, K.-H. Lee and J. Park, 6D Strings and Exceptional Instantons, [1801.03579](#).
- [49] M. Del Zotto and G. Lockhart, Universal Features of BPS Strings in Six-Dimensional SCFTs, *JHEP* **08** (2018) 173, [[1804.09694](#)].
- [50] A. Iqbal, C. Kozçaz and C. Vafa, The Refined Topological Vertex, *JHEP* **10** (2009) 069, [[hep-th/0701156](#)].

G	matter	r_0	d_{\max}
$SU(N)_\kappa$	$N_f \times \mathbf{N}$	$d - N/2 - \kappa/2$	$0 \leq d \leq N - \kappa - 2N_f - 1(?)$
$SU(6)_3$	$1 \times \mathbf{20}$	$d - 6/2 - 3/2 + 3/2$	$1 \leq d \leq 6$
$SO(7)$	pure	$d - 5/2$	$0 \leq d \leq 5$
$SO(7)$	$1 \times \mathbf{8}$	$d - 5/2 + 1/2$	$0 \leq d \leq 4$
$SO(7)$	$1 \times \mathbf{7}$	$d - 5/2 + 1 \times 1/2$	$0 \leq d \leq 4$
$SO(7)$	$2 \times \mathbf{7}$	$d - 5/2 + 2 \times 1/2$	$0 \leq d \leq 3$
G_2	pure	$d - 4/2$	$0 \leq d \leq 4$
G_2	$1 \times \mathbf{7}$	$d - 4/2 + 1/2$	$0 \leq d \leq 3$
F_4	pure	$d - 9/2$	$0 \leq d \leq 9$
F_4	$1 \times \mathbf{26}$	$d - 9/2 + 1 \times 3/2$	$0 \leq d \leq 6$
F_4	$2 \times \mathbf{26}$	$d - 9/2 + 2 \times 3/2$	$0 \leq d \leq 3$

Table 1: list of theories

Gauge group	Hypermultiplets
$SU(N)_\kappa$	x
$SO(2N+1)$	$n_{\mathbf{v}} \leq 2(N-1)$
$SO(2N)$	$n_{\mathbf{v}} \leq 2(N-2)$
$USp(2N)$	$n_{\mathbf{f}} \leq 2N$
$USp(2N)$	$n_{\Lambda^2} = 1$ and $n_{\mathbf{f}} \leq 2$

Table 2: asd

Gauge group	Hypermultiplets
$SO(14)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 1$ and $n_{\mathbf{v}} \leq 2$
$SO(13)$	$n_{\mathbf{s}} = 1$ and $n_{\mathbf{v}} \leq 2$
$SO(12)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 2$
$SO(12)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 1$ and $n_{\mathbf{v}} \leq 4$
$SO(11)$	$n_{\mathbf{s}} = 2$
$SO(11)$	$n_{\mathbf{s}} = 1$ and $n_{\mathbf{v}} \leq 4$
$SO(10)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 3$
$SO(10)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 2$ and $n_{\mathbf{v}} \leq 2$
$SO(10)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 1$ and $n_{\mathbf{v}} \leq 4$
$SO(9)$	$n_{\mathbf{s}} = 3$
$SO(9)$	$n_{\mathbf{s}} = 2$ and $n_{\mathbf{v}} \leq 2$
$SO(9)$	$n_{\mathbf{s}} = 1$ and $n_{\mathbf{v}} \leq 4$
$SO(8)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 4$
$SO(8)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 3$ and $n_{\mathbf{v}} = 1$
$SO(8)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 2$ and $n_{\mathbf{v}} \leq 2$
$SO(8)$	$n_{\mathbf{s}} + n_{\mathbf{c}} = 1$ and $n_{\mathbf{v}} \leq 3$
$SO(7)$	$n_{\mathbf{s}} = 4$
$SO(7)$	$n_{\mathbf{s}} = 3$ and $n_{\mathbf{v}} = 1$
$SO(7)$	$n_{\mathbf{s}} = 2$ and $n_{\mathbf{v}} \leq 2$
$SO(7)$	$n_{\mathbf{s}} = 1$ and $n_{\mathbf{v}} \leq 3$
G_2	$n_{\mathbf{7}} \leq 2$
F_4	$n_{\mathbf{26}} \leq 2$
E_6	$n_{\mathbf{27}} + n_{\overline{\mathbf{27}}} \leq 3$
E_7	$n_{\mathbf{56}} \leq 2$
E_8	\emptyset

Table 3: asd