

RESEARCH IN ELEMENTARY PARTICLE PHYSICS

A PROPOSAL TO THE U.S. DEPARTMENT OF ENERGY

THE UNIVERSITY OF TEXAS AT ARLINGTON

Physics Department, 502 Yates Street, Arlington, Texas 76019, USA.

Lead Principal Investigator: Andrew P. White (817) 272 2812 awhite@uta.edu
Co-Principal Investigator: Kaushik De
Co-Principal Investigator: Andrew Brandt
Co-Principal Investigator: Jaehoon Yu
Co-Principal Investigator: Amir Farbin
Co-Principal Investigator: Haleh Hadavand
Co-Principal Investigator: Jonathan Asaadi
Co-Principal Investigator: David Nygren
Co-Principal Investigator: Benjamin Jones

Administrative Point of Contact: Sarah Panepinto, 817 272 0243

FOA number: **DE-FOA-0001604**

DOE/Office of Science Program Office: **High Energy Physics**

DOE/Office of Science Program Office Technical Contacts:

Energy Frontier: Abid Patwa, 301-903-0408, abid.patwa@science.doe.gov

Intensity Frontier: Glen Crawford, 301-903-9458, glen.crawford@science.doe.gov

Detector R&D: Helmut Marsiske, 301-903-6989, helmut.marsiske@science.doe.gov

PAMS Letter of Intent tracking number: LOI-0000014874

Research Subprograms:

Experimental Research at the Energy Frontier in High Energy Physics

Experimental Research at the Intensity Frontier in High Energy Physics

Detector Research and Development in High Energy Physics

Cover Page Supplement

List of Research areas:

- **Energy Frontier PI's:** Kaushik De, Andrew White, Andrew Brandt, Amir Farbin, Haleh Hadavand
- **Intensity Frontier PI's:** Jaehoon Yu, Jonathan Asaadi
- **Detector Research and Development PI's:** Andrew Brandt, Jonathan Asaadi, David Nygren, Benjamin Jones

Lead PI: Andrew White

| | Name | Research Area | Year 1 Budget | Year 2 Budget | Year 3 Budget | Total Budget |
|---------|--------------------|--------------------|---------------|---------------|---------------|--------------|
| Lead-PI | Andrew White | Energy Frontier | \$130,156 | \$132,428 | \$160,555 | \$423,139 |
| co-PI | Kaushik De | Energy Frontier | \$262,290 | \$268,294 | \$274,477 | \$805,061 |
| co-PI | Andrew Brandt | Energy Frontier | \$121,491 | \$124,144 | \$185,892 | \$431,526 |
| co-PI | Amir Farbin | Energy Frontier | \$221,356 | \$226,832 | \$232,472 | \$680,659 |
| co-PI | Haleh Hadavand | Energy Frontier | \$163,892 | \$167,227 | \$170,661 | \$501,780 |
| | Total | Energy Frontier | \$899,185 | \$918,924 | \$1,024,057 | \$2,845,165 |
| co-PI | Jaehoon Yu | Intensity Frontier | \$238,522 | \$247,962 | \$252,168 | \$738,651 |
| co-PI | Jonathan Asaadi | Intensity Frontier | \$173,432 | \$231,214 | \$234,918 | \$639,563 |
| | Total | Intensity Frontier | \$411,954 | \$479,175 | \$487,086 | \$1,378,215 |
| co-PI | Andrew Brandt | Detector R&D | \$81,177 | \$83,499 | \$45,913 | \$210,589 |
| co-PI | Jonathan Asaadi | Detector R&D | \$33,658 | \$34,258 | \$34,876 | \$102,792 |
| co-PI | David Nygren | Detector R&D | \$17,545 | \$17545 | \$17,545 | \$52,635 |
| co-PI | Benjamin Jones | Detector R&D | \$39,718 | \$40,318 | \$33,361 | \$113,397 |
| | Total | Detector R&D | \$172,098 | \$175,620 | \$131,695 | \$479,412 |
| | Grand Total | All areas | \$1,483,237 | \$1,573,718 | \$1,642,837 | \$4,699,792 |

Table 1: Name and Yearly Budget for Proposals with Multiple Research Areas.

Part I

Common Introduction

This proposal requests base funding support for the High Energy Physics (HEP) program at the University of Texas at Arlington (UTA). It covers the research plans of nine PI's in three different research subprograms. Activities of the PI's in the HEP group at UTA are highly coherent and synergistic across research areas, while maintaining the traditional independence of faculty members to pursue their own research interests. We accomplish this balance through the practices listed below.

- **Team building:** Major activities are pursued by teams comprising of multiple PI's and postdoctoral fellows. This is especially important for international scale experiments like ATLAS and LBNE, where large groups are involved from UTA.
- **Resource sharing:** Human and physical resources in the group are shared by multiple PI's. For example, postdoctoral fellows work with multiple PI's, which enhances their training while providing cost savings to the group.
- **Synergistic activities:** All activities are chosen to be closely affiliated with existing responsibilities within the group. For example, PanDA software expertise is matched with distributed computing operations experience, and the new sub-GeV dark matter searches in the intensity frontier experiments are carried out across closely coupled projects.
- **Core involvement:** All aspects of experimental work are maintained as core expertise within the HEP group, leading to successful physics results across experiments. For example, detector hardware, detector electronics, computing and statistical analysis are equally valued and expertise is maintained at UTA in each area.

The success of the HEP team effort at UTA has led the university administration to approve the formation of one of the first "Organized Research Center of Excellence (ORCE)" at UTA, the ORCE:HEP, which provides an umbrella for the research activities of the nine PI's in this proposal. The Center provides a crucial structure of organizational support for the work plan proposed here.

The shared expertise, infrastructure and organization of the HEP group at UTA provides excellent value for the investment by DOE in the base program. Every dollar in DOE funding is highly leveraged. We describe an ambitious program of work in this proposal while requesting a modest level of investment from the base program.

In subsequent sections of the proposal, the common threads described are numerous. Our expertise in detector development and computing at the Energy Frontier closely matches our new responsibilities at the Intensity Frontier. Software and computing activities directly benefit and enhance the productivity of physics results. Exploration of the Higgs leads naturally into the search for supersymmetry. Many other such synergies are described in the remainder of this proposal.

The nine PI's of this proposal are highly active and well known within the field for their expertise. Two of our PI's are DOE Outstanding Junior Investigator (OJI) or Early Career Research Program (ECRP) winners. We have world renowned expertise in hadronic calorimetry, having designed, built and operated calorimeters at many frontier experiments. Our distributed computing innovations are recognized internationally, in HEP and beyond. We have two decades of experience and leadership in SUSY. We have key responsibilities in Higgs physics. With the new emphasis on dark matter, cutting across all areas of HEP research, we are poised to pursue new discoveries vigorously using all of our combined expertise.

Part II

Research at the Energy Frontier

The PI level of effort in the Energy Frontier has increased by 1 FTE compared to our previous Comparative Review. Haleh Hadavand has joined the UTA HEP group as an Assistant Professor. She is very active in ATLAS and highly regarded in Higgs physics. She is co-leading the new HL-LHC R&D effort at UTA to construct the LVPS (Low Voltage Power Supply) system for the Scintillating Tile Calorimeter in ATLAS. We have requested additional \$164k in the EF budget to support her work. Otherwise, the budget request in the first year of this proposal is at the same level as the last year of our previous grant. For our 4 senior PI's we request flat funding even though we expect a much higher level of productivity given our new responsibilities, and since Amir Farbin is now focusing 100% on ATLAS.

In Table 2, we show the fractional contribution of each PI to experiments at the Energy Frontier (EF).

| | | 2017 | 2018 | 2019 |
|-------|----------------|------|------|------|
| ATLAS | Kaushik De | 1.0 | 1.0 | 1.0 |
| | Amir Farbin | 1.0 | 1.0 | 1.0 |
| | Haleh Hadavand | 1.0 | 1.0 | 1.0 |
| | Andrew White | 0.5 | 0.5 | 0.5 |
| | Andrew Brandt | 0.5 | 0.5 | 0.5 |
| ILC | Andrew White | 0.5 | 0.5 | 0.5 |

Table 2: Fractional level of effort for each PI by research area within the Energy Frontier.

We begin this section with individual summaries of the plans for each PI, which will be followed by details of the research plans organized by research areas and research topics. We begin with ATLAS, followed by the ILC. Within each research area, we describe the many topics pursued by our group at UTA, laying out the research objectives, proposed methods and timetables.

PI Summary: Kaushik De

De is currently supported under the DOE base program at the level of 1.05 postdoc, 1 graduate student, two months of summer salary, and a small travel and M&O budget. I request the same level of continued support to pursue an ambitious research plan for the next three years. My group proposes to deliver: support for the ATLAS Tile Calorimeter ITC originally built by De's group, urgent software development and operations support for ATLAS computing which have been identified as the highest priority by ATLAS management for the ongoing Run 2 at the LHC, completion of SUSY papers in search for stop quarks, HL-LHC Upgrade Research and Development on the TileCal Trigger and DAQ Pre-processor board, and ongoing responsibilities as Deputy US ATLAS Software and Computing Manager. We currently receive some funding from other sources for computing, operations and service activities in ATLAS: which supports a scientist from US ATLAS project funds for tile calorimetry, three computing professionals from the DOE ASCR project, five computing specialists through the US ATLAS project, and additional people through the university. These additional sources of funding dramatically enhances the primary goals of my DOE base program. As described in this proposal, the combined synergy of these multiple projects and sources of support will allow my group to make a huge contribution to the success of ATLAS, and to the physics mission of DOE.

Accomplishments: With continuous support under the DOE base program for the past 20 years, PI Kaushik De's group has played a leading role in two of the highest priority DoE supported programs: the DØ experiment at Fermilab and the ATLAS experiment at the LHC. We have contributed to all phases of these two experiments, leading to the discovery of two of the most highly sought fundamental particles: the top quark and the Higgs boson. De's leadership has been especially important during the past decade in ATLAS, where he led a team of universities to build a unique part of the ATLAS calorimeter, and pioneered a new model of distributed computing for HEP that is now used internationally by HEP, NP and other data science experiments. Over the past two decades, De has also personally contributed steadily to narrowing the path in the search for supersymmetry. As we enter the next exciting decade in particle physics, with continued DoE base support, De's group is poised to make contributions in the field of HEP at an even higher level.

De graduated a Ph.D. student in ATLAS in 2015, Dr. Smita Darmora. Her thesis explored difficult regions in stop parameter space using a deep learning (TMVA) technique. While the stop was not found, stringent limits were places and the results were included in ATLAS published papers.

Giulio Usai, a scientist in De's group, supported 55% by the base program, is the ATLAS Tile Calorimeter Upgrade co-coordinator. He is leading the TileCal Test Beam effort at CERN, while also supervising Tile Upgrade R&D for ATLAS world-wide. Usai and De are leading the HL-LHC Tile PPR TDAQi board development through the base program and US ATLAS upgrade project.

While contributions to SUSY, calorimetry and other areas are important to ATLAS, our contributions to the field of distributed computing over the past five years have had the most impact. De proposed and led the development of many major elements of the ATLAS computing model, especially the PanDA system. Thousands of physicists use PanDA daily to run millions of jobs at hundreds of computing centers worldwide. De has provided intellectual leadership for this project since the beginning. Today PanDA is supported by CERN IT, OSG, multiple international laboratories, and is used by or is under testing by AMS, ALICE, and many non-HEP communities with Big Data.

De serves on the US ATLAS Management Board as DSCM (Deputy Software and Computing Manager). In this role, he co-supervises thirty million dollars in NSF and DOE funding, and co-manages the technical work of about fifty people in three National Labs and dozens of universities.

Milestones: The three pillars of scientific discovery in energy frontier research: detector, computing and physics analysis are all covered by the program of work of De's team. The primary milestones of the program of work proposed here are:

- 2017 complete repair and re-calibration of Intermediate Tile Calorimeter, complete development and commissioning of ProdSys2, generate Monte Carlo samples for 14 TeV run, prepare and test code for quick SUSY discovery when LHC Run 2 starts, graduation of student Smita Darmora.
- 2018 maintain monthly calibration of ITC with Cs source, complete work on event server model for ATLAS computing, publish SUSY discovery or limits for third generation searches and additional new models motivated by data, graduation of student Jordan Benson.
- 2019 provide improved algorithm for high luminosity operation of TileCal, complete implementation of virtual facilities model for ATLAS computing, publish results from SUSY studies, graduation of student Mikahil Titov.

Plans: De's group is involved in extremely high priority program of work in the ATLAS experiment, in support of the detector, computing, and physics. The plan of work for each area is

coordinated in weekly meetings with various levels of ATLAS management. The overall goal is to have a successful start of Run 2 at the LHC, followed by physics measurements and new discoveries. We have sketched the plan of work in subsequent parts of this proposal. However, the scope of the entire program of work is huge. Support of De's work through the DoE base program will be much appreciated and critical to the success of ATLAS.

The training of students and postdocs is crucial to De's contributions to particle physics. All the postdoctoral scientists in his team, irrespective of their sources of support, play a leadership role in ATLAS. They are trained to contribute to the experiment in deep and meaningful ways. For example, Nurcan Ozturk is the US ATLAS Software Manager. Giulio Usai is the Calorimeter Signal Reconstruction Task Force leader. David Cote is the SUSY missing Et convener. Alden Stradling coordinates DAST. Armen Vartapetian coordinates ADCoS. Paul Nilsson is PanDA pilot lead developer. Anyone knowledgeable about ATLAS computing and physics will acknowledge the contribution of De and his team.

In summary, we present a program of work here for the next three years that is crucial to the success of ATLAS and the DoE HEP Energy Frontier mission. A modest level of base program support is requested: one student, one postdoc, two months of summer support and a small travel budget. The leveraged contributions to ATLAS will be huge and critically important. We hope that the work proposed here will lead to the next eagerly anticipated and major discovery of physics in ATLAS: supersymmetry.

PI Summary: Amir Farbin

The two drivers of my research are New Physics (NP) searches and software and computing challenges. In the past three years, these pursuits have kept me grounded in the LHC, where the frontiers of these areas reside, while my focus was on the Intensity Frontier, where I helped instantiate both the US's Long Baseline Neutrino and UTA's neutrino programs. On September 1, 2016, my term as DUNE's Deputy Software and Computing Coordinator (DS&CC) ended. I chose to not ascend to full coordinator because it required a 100% commitment that would sever me from the LHC. Free of my Intensity Frontier commitments and on sabbatical (Faculty Development Leave), I have just changed my focus back to the ATLAS experiment and the 100% Energy Frontier program I outline in this proposal.

My ATLAS group, which now consists of postdoc Louise Heelan, and recently graduated PhD student Daniel Bullock and new graduate student Leslie Rogers, has a long history of leadership (twice as subconveners) in the search for strongly produced squarks and gluinos. These searches are where the LHC derives the most stringent sparticle mass limits. We also brought the unconventional "Razor" technique to ATLAS in early Run 1 and followed through with four iterations of searches, most recently in 13 TeV data for ICHEP 2016 using a generalization of the Razor [] technique called Recursive Jigsaw [] to enhance sensitivity to compressed SUSY spectra scenarios. My interest in this technique stems from its potential to enable broader yet maximally sensitive searches, which will become more relevant as data-doubling time for Run 2 analysis get exponentially longer. Section ?? proposes a program that evolves these techniques towards more sensitive and general searches as well as targeting difficult regions, such as compressed SUSY spectra.

We also run the ATLAS software tutorials, a responsibility that I picked up while serving as ATLAS's Physics Analysis Tools (PAT) coordinator. This activity is project supported and will not be covered in detail in this proposal due to space constraints, but nonetheless is an important conduit to the latest developments in ATLAS software. A decade ago, I participated in the installation, commissioning, and operation of the Tile Hadronic Calorimeter (TileCal). In recent years, my group has continued to contribute to the study, operations, and upgrade of TileCal, through run

coordination, primary editing (i.e. writing) of the Tile performance paper, and maintenance and upgrade of front-end electronics and mobile test systems. My ATLAS roots are in that community. Heelan's roots are in Calorimetry. Minimally, my students will continue to perform service and qualify on TileCal.

My physics program in neutrinos, which will end with my student Sepideh Shahsavarani's graduation in 2018, is also focused on NP through searches for sub-GeV Dark Matter (DM) potentially produced in the neutrino beamline. Using MiniBooNE's recent dedicated off-axis run, Shahsavarani participates in the nearly complete DM-nucleon scattering search while also leading the DM-electron scattering search. She is simultaneously pursuing neutrino-Argon proton cross section measurements in LArIAT. I just concluded serving as DUNE's Deputy S&C Coordinator, where I helped design and lead the computing for the next generation of HEP experiments, bringing in experience from the LHC, and building bridges between the Frontiers. Some of these efforts will be described in section ??.

Recently, Deep Learning (DL) –an emerging branch of Machine Learning which promises better and faster algorithms that are easier to develop and scale, and enable new powerful workflows– has grabbed my interest, specifically with challenges of High Luminosity LHC (HL-LHC) computing in mind. My efforts in this area has grown from a hobby to a major thrust of research. I have been applying DL to a variety of HEP problems in several different experiments via collaborations with numerous HEP and DL colleagues. Early on, I was the first to demonstrate DL image processing potential in LArTPC neutrino experiments and Gas TPC neutrinoless double-beta decay experiments. I have jump-started many HEP collaborators by providing software support, GPUs, and producing large public data samples, while personally tackling some of the most difficult problems in collaboration with DL colleagues. Section ?? overviews my extensive program of DL projects and outlines how they naturally extend to the LHC, for example in Calorimetry. Almost all of my and fractions of other UTA PIs, will include a DL component. My considerable lead in the application of DL to HEP has yielded numerous workshop and conference presentations and organization roles in the past year [], including the only DL talk at ICHEP 2016 [].

The transition back to 100% EF provides an opportunity for me to pursue new software areas in ATLAS. For Run 3, ATLAS needs to migrate to a Multi-threaded version of their Athena framework, where I have a long history of contributions. Meanwhile ATLAS is woefully short of software expertise and manpower. I propose to dedicate my focus in ATLAS computing to aspects of this migration related specifically to the Trigger and GPUs. While no further framework migration is planned for Run 4, my now extensive experience with frameworks, Event Data Model, GPUs, and co-processors lead me to believe that a new framework is likely necessary and warranted. I will play a leading role in organizing the framework discussions in the community-wide white paper orchestrated by Pete Elmer, Mike Sokoloff (U. Cincinnati), and Mark Neubauer in preparation for a NSF SIS2 center proposal. Ideally we will organize a workshop that will pull together framework experts across experiments, many of whom I had the opportunity to closely work with as DUNE DS&CC. All of these activities will greatly benefit from my proposed deep involvement in ATLAS Core Trigger software, where I intend to assume critical responsibilities and leadership roles.

I find myself at the boundary of several distinct areas in HEP and HEP Experiments. For a long time, I've straddled the fence between Physics and Computing. My strong computing skills and expertise have lead me to assume roles generally assumed by computing oriented staff scientists and engineers who often haven't touched real data for a long time, and seldom tackled by professors, like me, with also close connections to data analysis and detectors. For the past three years, I have had one foot in the Intensity Frontier and the other in the Energy Frontier, developing a unique perspective of computing on both fronts and establishing working relationships with many core developers at Fermilab and CERN. Finally, my recent successes in the DL have pushed me to bound-

aries of cutting edge Data Science and HEP, yielding collaborations with new HEP experiments and Data Scientists. There are boundaries in HEP that are rather difficult to bridge. I believe that some of these boundaries can be resolved in software and computing, ultimately yielding better science.

[The next three years look like this...]

PI Summary: Haleh Hadavand

I started my postdoc position with Southern Methodist University on the ATLAS experiment in late 2005. I joined the online monitoring software effort at CERN and within a few months developed a histogram naming system for the online histograms and modified an existing/decommissioned software for histogram collection to work properly within a new release. I also started designing and laying down the foundations for the ATLAS data quality monitoring system. I was one of the main visionaries, designers, and developers for this system which is still used both online and offline by ATLAS. My main development contribution was the plug-in algorithms that could be loaded at any point into the system, the online display, and the online time series analysis [?, ?].

During commissioning of cosmics before LHC collisions I setup the entire online monitoring chain from histogram production to final DQ assessments of monitoring data. This required thorough understanding of the offline software, within the High Level trigger, and the online environment, software, and tools. I was the first to develop software to produce η vs ϕ plots of clusters within the LAr and Tile calorimeters online, use the gatherer to collect histograms from various monitoring nodes, display the histograms, apply DQ, and time series assessment of the data quality information. This work gives me a very thorough understanding of the monitoring system on ATLAS both online and offline.

I worked on photon physics within the exotic channels of UED diphoton + Missing Et and Randall Sundrum(RS) Graviton to diphotons. I was a main contributor to both analyses. The UED analysis was one of the first 20 published papers out of the ATLAS experiment [?] for which I served as a co-editor. For the RS Graviton analysis I provided the statistical framework for the final limits. I was a co-editor for the resulting conference note and paper [?, ?].

Since I joined UTA in July 2012 I have been working on the charged Higgs to $\tau^+ \nu$ in fully hadronic final states. I have been part of three conference results and two publications [?, ?, ?, ?, ?]. My role on the Run 1 measurement was the high mass background estimation of the $\tau \rightarrow \text{jets}$ background and the statistical treatment leading to the final limits for the analysis. I produced Run 2 projections for this analysis and determined that with only a few fb^{-1} of data evidence of signal can be observed. I also produced the first intermediate mass samples for the charged Higgs in the mass range of 160-180 GeV. This was work done with collaboration with theorists to reconcile interference terms in this region. For the Run 2 analysis I motivated using the MET trigger instead of the tau+MET trigger used in Run 1. This choice greatly reduced the trigger and background systematics. The background suffered from low statistics when using the tau+MET trigger since the trigger imposed a medium tau selection. My work on this charged Higgs final state has resulted in my appointment to the charged Higgs convener position thereby overseeing all charged Higgs activities on the ATLAS experiment.

As charged Higgs convener I have motivated new charged Higgs final state searches (five new channels) thereby expanding the groups responsibility and visibility. I have been in contact with theorists and am the main contact for MC generation of charged Higgs final states putting me in a unique position to deliver the MC samples for newly motivated final states. One of these new channels is a decay to SUSY final states which would have a signature of three leptons plus missing

ET. Two of these leptons would be same flavor and opposite sign. Several independent evidence of an excess in multi-lepton events have been reported on the ATLAS experiment [?].

In April 2016 I was appointed to the position of charge account manager (CAM or L3 manager) for the low voltage power supplies (LVPS) for the Tile calorimeter for HL-LHC. This account will manage about \$1 million of labor, travel, and materials and supplies over a 6 year period. I have been working closely with Andrew Brandt, the institutional representative for UT Arlington, on this project. More details about this project are highlighted in section ?? and the schedule is detailed in table 4.

Plans I have a very ambitious physics plan for the next three years which consists of charged Higgs physics and a new analysis endeavor into Z boson + X searches. I also have many responsibilities as charged Higgs convener and CAM for the LVPS upgrade for the Tile calorimeter for HL-LHC. My student Hussein Akafzade will graduate by mid-2019 on charged Higgs while I ramp up a new student to work on Z+X analysis starting in 2017. This new student will be funded by the grant upon Akafzade's departure. The analyses I propose will use cutting edge techniques to increase sensitivity including the use of tau polarization for distinguishing Higgs decays from W boson decays, use of Deep Learning in conjunction with Brandt and Farbin, Matrix Element Methods, and new di-tau reconstruction algorithm. The Z boson + X analysis will cover five final states within one analysis by using a parametric fit to the background in control regions and searching for new particles in the mass range of 4 GeV to several TeV. This type of combined search is unprecedented on ATLAS. I continue to motivate new charge Higgs final states as the charge Higgs convener and follow the progress of the new analyses being added to the group. There are several important milestones for the LVPS upgrade project in the next three years. These include the design and fabrication of the LVPS elevated temperature test stand (burn-in station), the version 8.2 prototype, and the testing and integration into the vertical slice test.

Milestones

- FY16 - Publish results for charged Higgs on 2016 dataset including the tau-lepton channel.
- FY16-17 - Design and fabrication of Tile LVPS burn-in station.
- FY17 - Publish charged Higgs results using τ polarization.
- FY17 - Take new graduate student for Z+X analysis.
- FY17 - Hire new postdoc to work on charged Higgs and Z+X analyses.
- FY18 - Develop analysis suite for Z+X analysis.
- FY18 - Publish Z+X results using Higgs mass constraint.
- FY18 - Publish charged Higgs results using deep learning techniques for selection and/or tau reconstruction.
- FY18-19 - LVPS V8.2 prototype.
- FY18-19 - Testing and integration of LVPS into vertical slice test.
- FY19 - Charged Higgs results with Matrix Element Methods.
- FY19 - Graduate student Akafzade graduates on charged Higgs analysis.
- FY18 - Publish Z+X results with no mass constraints.

PI Summary: Andrew P. White

Accomplishments

- Founder of UTA High Energy Physics Group 1991
- Inventor of the DZero Experiment Intercryostat Detector

- Inventor of Gas Electron Multiplier (GEM) based Digital Hadron Calorimetry
- American Physical Society Fellowship - 2011
- Recipient of UTA Distinguished Record of Research Award, May 2009
- Analysis contact for Run 1, and Run 2 Vector Boson Fusion Higgs To Invisible Decays
- Spokesperson for the SiD Detector Concept for the International Linear Collider (2010-)
- Chairman of the Physics Research Council for DESY, Hamburg/Zeuthen, Germany (2012-2015)
- Member of the U.S. Department of Energy National Reviews of Detector Research and Development at National Laboratories, 2009, 2012
- Served on national review committees and was a member of the Department of Energy, HEPAP Subpanels on the Future of Particle Physics in the United States (1998 and 2007)
- Member of the ECFA Detector Panel (2012-2016)
- Member of the ATLAS Tile Calorimeter Speakers' Committee (2014-)
- Member of the US ATLAS Analysis Support Group
- Universities Representative on the Americas Linear Collider Committee (2012-)
- N. American representative for the CALICE (Calorimetry for Linear Colliders) Collaboration
- Member of the Management Board for the Micro Pattern Gas Detector Collaboration/CERN, RD-51
- Instigator and original leader of the DZero Experiment New Phenomena physics group
- Co-designer of the ALEPH (CERN) Experiment Inner Trigger and Tracking Chamber (1982-4)
- Invited to lead numerous physics and detector working groups at large national and international high energy physics meetings

Milestones

- FY16 - Publish Run 1 results from Higgs to invisible decays analysis - done.
- FY17 - Publish Run 2 2015-16 results from Higgs to invisible decays analysis.
- FY17 - Take on new graduate student for Higgs/invisible analysis, ITC qualification tasks.
- FY17 - Complete study of ITC E-cells new segmentation, energy resolution study.
- FY17 - Complete full simulation of SiD AHCAL, barrel + endcaps.
- FY17-18 - Develop design of SiD AHCAL barrel module.
- FY17-18 - Complete plans for ATLAS ITC Tile/Fiber replacements in LS2.
- FY18-19 - Co-convene the full Run 2 Higgs to Invisible analysis.
- FY18-19 - Prepare (with ATLAS Tile Institutes) tile/fiber assemblies for installation in LS2.
- FY18-19 - Expand SiD Consortium in preparation for drafting SiD Technical Design Report.
- FY18-19 - Begin first draft of SiD Technical Design report.
- FY19 - Publish result for Run 2 Higgs to Invisible analysis, possible interpretive paper(s).
- FY19 - Contribute effort to installation and testing of the new ATLAS ITC E-cells.

Plans

With the discovery of the Higgs boson great opportunities exist for understanding this fundamentally new state of matter, and for discoveries associated with existence. In this regard, I am committed to the continuing success of the ATLAS Experiment's physics program through analysis of current and future data in the search for invisible decays of the Higgs, and through support work on the Intermediate Tile Calorimeter, and data quality validation shifts. Taking on a new graduate student, I will continue our study of the optimal segmentation of the new ITC E-cells. I will work to identify suitable replacement ITC tile and fiber materials and provide expertise in their installation and re-commissioning. I will work on the extension of the VBF-Higgs to invisible analysis, and paper publication, using existing data and, with my student, I will develop analysis strategies

for analyzing the 14 TeV data accumulated at high instantaneous luminosity and pileup. The full exploration of the Higgs sector will require precision studies of couplings at the percent level or below. The best prospect for achieving such precision measurements in the foreseeable future lies with the International Linear Collider. As Spokesperson for the SiD Detector Concept for the ILC, I will continue to lead its development and promote its realization within the global HEP community. I will work towards securing a U.S. role in the ILC project in Japan and develop the site-specific implementation of SiD at the selected Kitakami location. With the SiD Executive Committee and the Institutional Board, I will create the team that will produce the SiD Technical Design Report over a 2-3 year period, and guide the process of subsystem technology selections. I will represent SiD on the Linear Collider Physics and Detectors Executive Board, and the Americas Linear Collider Committee, and with interactions with funding agency(s). I will work to build the SiD Collaboration in all regions. Locally at UTA, I will pursue the implementation of scintillator/SiPM hadron calorimetry for SiD, together with its full simulation.

PI Summary: Andrew Brandt

1 The ATLAS Experiment

Faculty PI: Kaushik De, Andrew Brandt, Andrew White, Amir Farbin, Haleh Hadavand

The UTA HEP group joined the ATLAS collaboration in the early days of planning for the experiment, 21 years ago. Since then, we have participated in every aspect of the experiment: design, construction, commissioning, computing, management, and physics analysis. The impact of the work done by our group has benefited every aspect of ATLAS, and led to the successful physics program of the experiment. Currently, our group has N active members in ATLAS, which makes UTA one of the largest university groups in ATLAS. We have an excellent record of synergistic accomplishments among the members of our group, and among the various externally funded projects. While the DoE base program supports less than one third of our activities in ATLAS, the impact of this funding is magnified many fold in advancing the mission of the core physics goals of the DoE.

Currently, the participation of the UTA group in ATLAS is managed by 4 FTE effort among 5 PI's.

Physics at ATLAS

1.1 Higgs Physics (PI: Brandt, Hadavand, White)

On July 4, 2012 ATLAS and CMS announced the observation of a new particle consistent with a Standard Model (SM) Higgs Boson [?,?], ending the 50 year search for the elusive final SM particle. This landmark discovery fulfills the prime objective of the LHC by providing evidence for a Higgs field that gives mass to fundamental particles through their coupling to the field thus uncovering the mechanism for electroweak symmetry breaking. This Higgs-like particle weighed in with a mass of about 126 GeV, and had the highest significance and best mass resolution in the $\gamma\gamma$ and the ZZ decay channels. UTA had many contributions to the Higgs discovery, including critical roles in computing, detector building (ITC), and commissioning (ITC, MBTS, trigger).

Although the SM has been extremely successful at explaining electroweak data, so-called problems with the theory, such as the quadratic divergence of radiative corrections to the Higgs mass (a.k.a. the Hierarchy problem), and unanswered questions, such as the nature of dark matter,

drive our search for new phenomena at the TeV scale. We have strong arguments (i.e. naturalness and the “dark-matter miracle”) that solutions to these problems and questions may be found at this energy. The leading theoretical solution is Supersymmetry (SUSY) since it not only addresses the hierarchy problem, but can also provide a dark matter candidate and predicts the unification of gauge couplings at the Grand Unified Theory (GUT) scale. We also know that the uncertainty of the current Higgs measurements allow for non-SM decays of the Higgs boson of 26% [?] and that theories with SM+singlets have equivalent level of branching fraction [?].

With the discovery of the Higgs and reasons eluminated above the UTA groups focus has shifted to BSM Higgs phenomena in three fronts: A charged Higgs search in the $\tau\nu$ final state, invisible Higgs search in the VBF channel, and new proposal by Hadavand in the search for Z+X channels. The charged Higgs group is led by Brandt and Hadavand overseeing the work of postdoc Justin Griffith and Hadavand’s student Hussein Akafzade. Both Hadavand and Griffith have leadership roles in charged Higgs as the charged Higgs convener and analysis contact for the $\tau\nu$ final state. Akafzade’s thesis will be based on the Run 2 dataset of 100fb^{-1} on charged Higgs. White is the analysis co-convener for the Vector Boson Fusion Higgs to Invisible Decays for Run 2, having served in the same role for the published Run 1 analysis. White will take on a new graduate student to work on this analysis for the full Run 2 dataset. By mid-2017 Hadavand will take a new student who will work on the Z+X analysis together with a newly hired postdoc working on both charged Higgs and Z+X.

1.1.1 Charged Higgs (PI: Brandt, Hadavand)

Brandt and Hadavand will continue the search for charged Higgs in fully hadronic final state of $\tau^+\nu$. The discovery of charged Higgs would be a clear sign of BSM physics and would imply that the 125 GeV Higgs is part of a more complex Higgs sector. The Charged Higgs is predicted by several models such as ones with Higgs triplets and Two-Higgs-Doublet-Models(2HDM) [?, ?, ?]. In the MSSM, which is a type II 2 Higgs Doublet Model (2HDM), the main decay of charged Higgs at the LHC is $t \rightarrow b H^+$ for H^+ mass below m_{top} . At charged Higgs masses above m_{top} the main production at the LHC is in association with a top quark. In the MSSM the Higgs sector can be completely determined by the H^+ mass and $\tan\beta$, the ratio of vacuum expectation values of the 2HD. For masses below the top mass the $\tau\nu$ decay is dominant for $\tan\beta > 2$.

Hadavand motivated and was responsible for investigating new models for analyzing the intermediate mass region of 160-180 GeV. Only recently have theorists been able to reconcile $t\bar{t}$ interference effects supply us with NLO calculations of the cross section as a function of mass and $\tan\beta$. The addition of this mass region will close the gap mH vs $\tan\beta$ mass range and also produce competitive results at low $\tan\beta$ vs $A \rightarrow \tau\tau$ and $H^+ \rightarrow t^+b$. The measurements for this region will be performed by the end of 2016.

Hadavand and Griffith have led the charged Higgs as charged Higgs convener and analysis contact respectively. Under their leadership UTA group has produced the final results including plots and statistical interpretation, estimation of the systematic uncertainties, including theoretical, writing the analysis code, production of common ntuples, and trigger studies (with Jae Yu’s student Last Feremenga). By using simulated $t\bar{t}$ data instead of τ embedding [?] resulted in a paper with 3.2 fb^{-1} of data and a conference note at 14.7fb^{-1} of data at 13 TeV [?, ?]. This luminosity increase increased the mass reach of the analysis from 250 GeV in Run 1 to 600 GeV in Run 2. The jet to tau background statistics for the Run 2 analysis were vastly improved over Run 1 by using Hadavand’s suggestion of only using a missing ET trigger as opposed to a tau plus missing ET trigger. The use of the tau in the trigger implied that a medium selection of taus had to be used in the fake factor method, where in Run 2 a Loose selection could be used therefore greatly enhancing the statistics.




Figure 1: ICHEP 2016 hMSSM exclusions including 2015 exclusion, left; ICHEP 2016 hMSSM exclusions with 50 fb^{-1} and 100 fb^{-1} of data projections, middle; Distribution of τ polarization for Higgs and W decays, right;

Figure 1 (middle) shows that with the 2018 dataset of 100-150 GeV of data the analysis is only excluding large $\tan\beta$ values so this analysis continues to be an important measurement as more data is accumulated. The group has plans to apply several techniques to improve the analysis' sensitivity over its early Run 2 predecessor. The tasks aimed for these improvements are listed below:

Use tau polarization to reduce true tau background ; Akafzade The tau polarization can be used to separate the $W \rightarrow \tau\nu$ background from the signal. The intrinsic polarization of W^+ is -1 while that of H^+ is +1. The tau polarization is defined as $P_\tau = \frac{\sigma_R - \sigma_L}{\sigma_R + \sigma_L}$. A plot of the charge asymmetry, defined as $\Upsilon = \frac{p_T^{\text{track}}}{p_T} - 1$, is shown for the signal and background in the $\tau \rightarrow \rho\nu_\tau$ final state in figure 1, right. Other 1-prong decays will also have similar separation in the charge asymmetry. The 1-prong decays can be separated from the 3-prong decays in the final fitting machinery and added statistically together. This variable will be used as additional variable to the fit in the 1-prong case.

Use matrix element method to reduce true tau background ; Hadavand, Akafzade In addition to the tau polarization one can utilize the event kinematics by utilizing the matrix element method to distinguish the signal from the true tau background originating from $t\bar{t}$ events. This can be done by using the classifier distribution $d(x) = \frac{P_S(x)}{P_S(x) - P_B(x)}$ where P_S and P_B correspond to the weights from the signal and background hypothesis. Hadavand has performed initial tests with MadWeight for achieving this goal. This variable can then be added as one of the variables in the boosted decision trees (BDT).

Use boosted decision trees; Postdoc and Akafzade As mentioned earlier for early Run 2 results the discriminating variable used in the fit was m_T . As an alternative several variables are incorporated into a BDT which is used as the discriminating variable in the fit. Some variables that have been investigated in some preliminary studies are m_T , leading b jet p_T , τp_T , MET, $\Delta\eta(\tau\text{-leading jet})$, $\Delta\phi(\text{top-sub-leading jet})$.

Determine jet $\rightarrow \tau$ Background from Control Region ; Hadavand, Akafzade For Run 1 the systematics for the $\tau \rightarrow \text{jet}$ background were one of the largest in the analysis. Due to low statistics

at the high mass a fit above 200 GeV was necessary to get an estimate of the background in that region. This is important since this is the dominant background in the high mass region and without a background model the limit machinery would fail. As an alternative to the matrix method used for the Run 1 published paper one can use a control region with no signal and loosen the selection criteria to introduce more fake taus. This can be done by requiring a zero bjet region and only requiring a loose tau selection. The normalization of this background can then be derived within the fit to data. This should reduce both the statistical and systematic uncertainty of the analysis. The use of the tau plus MET trigger has caused some of the main issues with statistics for the QCD background in Run 1. The use of this trigger increases the statistics for the background estimation.

Apply Deep learning techniques for selection; Brandt, Farbin, Hadavand, graduate and undergraduate students The deep learning technique can be applied to this analysis in two ways. One way would be to use it to distinguish $t\bar{t}$ background from the signal by training on those samples. A comparison of the BDT method described earlier for event selection will be done with respect to this technique. Another way to apply deep learning would be in the tau reconstruction in place of the BDT technique which is used now. This work will be done utilizing Farbin's extensive experience in the field of deep learning.

Optimize Binning for Limits ; Postdoc, Akafzade The low statistics in the tail of the distribution has made the $H^+ \rightarrow \tau\nu$ channel difficult for setting limits. In many cases several systematic variations performed for pull determination would make the fits fail. This has to do with the fact that the data has low stats and is not smooth as opposed to MC samples which did have a smooth shape in the tail and could be fit successfully. By making requirements on signal over background and the number of total background in a given bin and then testing with pull estimation, Hadavand was able to optimize the best binning for both stability and sensitivity. For performing projections to higher luminosities the background is extrapolated to higher masses to determine a realistic scale of the background in this region. After performing the extrapolation the binning is re-optimized for the given luminosity projection, a technique developed by Hadavand.

Milestones

- FY16 - Publish results for charged Higgs on 2016 dataset including the tau-lepton channel.
- FY17 - Publish charged Higgs results using τ polarization.
- FY17 - Hire new postdoc to work on charged Higgs and Z+X analyses.
- FY18 - Publish charged Higgs results using deep learning techniques for selection and/or tau reconstruction.
- FY19 - Charged Higgs results with Matrix Element Methods.
- FY19 - Graduate student Akafzade graduates on charged Higgs analysis.

1.1.2 Z+X Scan (PI: Hadavand)

The increased beam energy and intensity in the LHCs Run 2 and 3 will likely provide the best opportunity to discover BSM physics for at least the next ten years. The task of discovering it, however, can be more challenging compared to the SM Higgs, whose mass was well constrained by precision electroweak data and the underlying theory, i.e. the SM, was well understood and extensively tested for over half a century [?, ?, ?, ?]. In contrast, the number of theories and models that motivate BSM Higgs searches are significantly larger and more speculative, requiring that a much wider net be cast to cover a broader range of masses and final states.

Hadavand proposes to perform Z boson + X searches using the 125 GeV Higgs to bootstrap her way to search for new exotic decays of the Higgs or other BSM states. The Higgs mass constraint will help aid the search for signal in a specific region. This method is motivated but not limited to the final states listed in table 3. The search can be further expanded in a model-independent manner by scanning for Z+X where X can be of masses not theoretically motivated. This proposal combines a search within five theoretical models each decaying to ll $\tau\tau$ and/or $\tau\tau$ ll, thereby using Hadavand's extensive experience with τ leptons. One analysis covers all of these models thereby increasing the analysis reach.

The Z+X search will primarily probe SUSY models for example Next-to-Minimal Supersymmetric Standard Model (NMSSM) that include an additional scalar singlet or SM + singlet models. It can also be used to discover other states not theoretically motivated therefore investigating new mass regions where potential new particles could exist. Additionally Hadavand proposes searching for a “dark” Z in the mass range $12 < m < 60$. A “dark” Z serves as the mediator of a new U(1) gauge symmetry that can serve as the portal to a hidden sector that constitutes dark matter [?].

| Analysis | Channels → | $\tau\tau$ ll | ll $\tau\tau$ | Theory |
|----------|-------------------------------------|---------------|---------------|------------------|
| Z Scan | $h \rightarrow ZZ^*$ | ✓ | ✓ | SM |
| | $A \rightarrow Zh$ | | ✓ | MSSM, NMSSM |
| | $h \rightarrow ZZ(\text{dark})$ | ✓ | ✓ | dark U(1) gauge |
| | $X(h) \rightarrow Za$ for a $l=\mu$ | ✓ | ✓ | NSSM, SM+singlet |

Table 3: Table showing channels and accessible final states and theoretical models for several BSM models including the standard model $h \rightarrow ZZ^*$ channel [?]. The mass of the scalar singlet a and Z(dark) are $4 < m(a) < 10\text{GeV}$ $15 < m(Z(\text{dark})) < 60\text{GeV}$ respectively.

Hadavand's strategy is to combine several final states and channels to increase statistics and look for a bump using a bump-hunting procedure Hadavand developed as a first pass. The approach removes the need for setting limits in a fine granularity in the discriminating variable, thereby greatly speeding up the statistical analysis. This speedup will allow for quicker turn around of searches in the absence of signal. In the end the full statistical treatment as recommended for ATLAS will be performed to present the final results but this first pass will allow for flexibility in the search.

For the Z scan Hadavand proposes to perform two mass scans constraining the given particle combination to the Z mass and scanning in the ‘other’ particle. For example, Z(ll) other($\tau\tau$) and Z($\tau\tau$)other(ll) would be two separate scans of the ‘other’ particle after requiring that the event has a Z boson in the final state. The mass range of the scan would go from 4 GeV to several TeV and each object would be a resonance in that mass spectrum. The ee and $\mu\mu$ channels are added together in one spectrum since they have similar resolution and hence can increase the statistics for bump-hunting. The two scans, one with taus and the other with $l=\mu,e$ can then be combined together with a statistics framework.

Of course the 4 lepton final state is one that is extensively tested for the SM ZZ* final state. This analysis would add the ll $\tau\tau$, $\tau\tau$ ll final states to this channel for the first time. The resolution will not be as favorable as the 4 lepton final state but by constraining the on shell Z to decay to $\tau\tau$ one can improve the mass resolution. These final states can add further sensitivity to this channel specially as the LHC gathers more data. With about 10 fb^{-1} of data at the center of mass energy $\sqrt{s} = 13 \text{ TeV}$ the 4 lepton analysis observed 3 events per category including a combination of electrons and muons. The efficiency for tau reconstruction is lower (30-40% as opposed to around

80%) than the electron and muon channels and in addition pose a more challenging final state due to the existence of neutrinos. However on average $35/80 \times 4 \sim 2$ ($2\tau 2e, 2\tau 2\mu, 2e 2\tau, 2\mu 2\tau$) events/ $10fb^{-1}$ can be added to this channel [?] increasing the total statistics by $2/12=17\%$.

By scanning for particles after selecting the Z boson one will get clear mass separation of the 'other' particle. The requirement of the Z boson in the event will also reduce backgrounds. Table 3 shows that many of the signals come from h, the 125 GeV Higgs. Looking for the h mass would result in an overlap between the various final states and would make it difficult to know which final state actually contributed. But by looking for the 'other' particle a clear mass separation is presented. This method is also less restrictive on the analysis since one would not put any explicit selection on the mass of the 'other' particle. In addition, another discriminating variable can be added to combine the Z and the 'other' particle and see if indeed one does reconstruct the hypothesized mass of the originating particle. For example restricting the $Z(\tau\tau)$ other(ll) mass to be within the Higgs mass window to see if it will make the peaks in the other(ll) become clearer.

The SM ZZ* is the only channel that is not a resonance in the other(ll) spectrum. It will have to be fit simultaneously as if it were a separate signal sitting on top of the Drell-Yan background. Using the bump-hunter to pick out this peak is not favorable since it is not a resonant peak and it can overlap with other signal. By studying the shapes in MC and doing a parametric fit to the data one can determine the level of SM contribution to the mass spectrum. In the end when reconstructing the 'other' mass one would see clear bumps for a, Z(dark), h in order of mass. The search can extend to high masses in case some other massive objects happens to appear in association with a Z boson.

1.1.3 Selection and Backgrounds to Z

The selection for this analysis are still to be determined but would require a di-lepton pair with $80 < m(ll) < 100$ GeV for the ee, and $\mu\mu$ channels and would require the use of the Missing Mass calculator for tau mass reconstruction to determine the selection window around the Z mass. In addition a medium/tight selection of the di-leptons will be made for the 'other' particle. When reconstructing the 'other' mass one can parameterize the fit to the background from a control region and extend it the entire range of the mass. At low mass there will be resonant standard model particles which can be modeled with a combination of simulations and the use of the control regions. The backgrounds include $J/\psi, \Upsilon, Z, t\bar{t}$ background, and Drell-Yan non-resonant background. In the intermediate mass there will be SM Z+h and ZZ* background as well.

Determine Algorithm for Boosted/Collimated di-tau Pair Identification ; Postdoc Since the proposal is a general search where an unknown particle can decay to a Z boson and another potentially unknown particle, some events will have highly boosted and/or highly collimated di-tau pairs that regular tau reconstruction does not treat. There has been work on ATLAS in this front aimed for reconstructing the channel $A \rightarrow Z(ll) h(\tau\tau)$ [?]. In this case the p_T of the di-tau system is quite high of an order of 500 GeV. In other cases the p_T of the di-tau system can be small but they can be highly collimated. This has been seen in the $h \rightarrow a(\mu\mu) a(\tau\tau)$ analysis. In fact the algorithm developed for $A \rightarrow Z(ll) h(\tau\tau)$ would not work for this channel because of the large p_T cut and the fact that it is seeded by a large R (1.0) Jet. Large R jets only have calibration available for p_T larger than 200 GeV. So Hadavand proposes to develop a high p_T di-tau algorithm and a low p_T highly collimated algorithm where standard jets of R of 0.4 can be used. A boosted decision tree (BDT) can be developed using the jet substructure information. The variables suggested to be used in the BDT or further studied within a Deep Neural Network are the following:

- core energy fractions, $f_{core}^{sub(lead)} = \frac{\sum_{cells}^{\Delta R=0.1} p_{p,cell}}{\sum_{cells}^{\Delta R=0.2} p_{T,cell}}$
- The subjet energy fractions $f(sub)lead_{subjet} = \frac{p_T^{subjet}}{p_T^{Jet}}$
- leading track momentum fractions $f(sub)lead_{subjet} = \frac{p_T^{leadTrack}}{p_T^{subjet}}$
- the maximum track distance Rmax
- the number of tracks n(sub)leading track.

Write General Analysis Suite for xAODs; Postdoc, Graduate Student This analysis has leptons including taus in the final states so the analysis suite can be generalized but always adding a constraint of having a Z boson in the final state. Since this search is inclusive where any particle can decay to Z+X, the analysis suite must be very general. However, on top of the general analysis suite one can build tools to make additional requirements on the specific states. For example if there is mass constraint that can be imposed such as the mass of the Higgs.

Determine Background Parameterization from Control Regions, $ll, l = \mu, e$ scan ; Postdoc, Graduate Student, Hadavand Two control regions (CRs) are used in this analysis to constrain the SM backgrounds in the final fit to the data. The first region, CRj, is used to constrain the low mass SM resonances and Drell-Yan dominated non-resonant background. The second region, CRb, is used to constrain the $t\bar{t}$ non-resonant background. Both regions are defined by first requiring a tagged ll candidate. CRj is further required to have at least one selected jet and not be b-tagged. CRb is required to have at least two selected jets that have been b-tagged. Any events which are in one of the SRs are excluded from the two CRs.

Determine Background Parameterization from Control Regions, $\tau\tau$ scan ; Postdoc, Graduate Student, Hadavand The control region used to determine the parameterization of the jets faking tau objects is defined as having two same sign tau leptons and inverting one of the tau identification criteria. Simulation is used to predict the shape of the contribution from events with two true leptons and hadronically-decaying tau that are either correctly reconstructed or reconstructed from leptons or leptonically-decaying taus. These two parameterizations of the two backgrounds is then used to fit to the data as a more sophisticated template method. The sidebands of the Z boson can be used to validate this approach.

Parameterize Background ; Graduate Student, Hadavand As described in section 1.1.3 the backgrounds to the multiple BSM analysis will be J/ψ , v , $t\bar{t}$, Drell-Yan, and Z tails. One can parameterize these background from the control regions described above then fit for the normalization within the signal region similar to what is done in the $\tau\tau\mu\mu$ analysis on ATLAS [?]. The SM ZZ^* is a non-peaking background which may be difficult to model. This background and the SM resonant backgrounds can be parameterized with MC and used in the fits. Hadavand has extensive experience with parameterized fits from the BABAR experiment and can guide Hadavand's graduate student to model the various SM backgrounds.

Perform two separate scans in Z final state ; Postdoc, Graduate Student In the decay of $Y \rightarrow Z + X$ the mass of X can be reconstructed as $X(ee)$, $X(\mu\mu)$, and $X(\tau\tau)$. The three Z scans look at the $X(ee)$, $X(\mu\mu)$, and $X(\tau\tau)$ mass spectrum. For the $X(l\bar{l})$, $l=e,\mu$ spectrum the Z will be reconstructed as $Z(\tau\tau)$. For the $X(\tau\tau)$ spectrum the Z will be reconstructed in $Z(l\bar{l})$, $l=e,\mu$. The $X(ee)$ and $X(\mu\mu)$ can be potentially combined if the resolution of the particles are similar enough. The $X(\tau\tau)$ will not be combined with $X(l\bar{l})$ $l=e,\mu$ since the resolution is quite different and the bump-hunter would not gain in looking for two overlapping bumps of vastly differing resolutions. The use of the bump-hunter will allow for a first statistical pass at the data to determine if there is an existence of signal in the samples.

Develop and Optimize Bump-hunter to Scan Data ; Hadavand, Graduate Student Hadavand has previously worked on a bump-hunting technique that would be independent of the signal model used, called Signal Parameter Independent Fit (SPIF). Using the parametrization described above for the background modeling, a fit is performed on the data and a window where signal is expected is removed. The background is then extrapolated within this window and the difference of the number of events in the window versus the fit extrapolation is determined. A p-value is determined for the difference between the number of events in the window and the extrapolated background within the window. A scan is done using as input the step size and a range of window sizes. Several scans are performed changing the window size and step size iteratively until the p-value no longer improves within a specified precision. Once all iterations are performed, the window(s) with the smallest p-value(s) is(are) reported. If the bump-hunter analysis reports no significant signal then a coarser binning can be used for the limit setting saving time and CPU power.

Milestones

- FY17 - Take new graduate student for $Z+X$ analysis.
- FY17 - Hire new postdoc to work on charged Higgs and $Z+X$ analyses.
- FY18 - Develop analysis suite for $Z+X$ analysis.
- FY18 - Publish $Z+X$ results using Higgs mass constraint.
- FY19 - Publish $Z+X$ results with no mass constraints.

1.1.4 Higgs Decays to Invisible Particles (PI: White)

A key question for the Higgs boson is whether it couples to the dark matter. If dark matter is indeed composed of particles, then presumably the 125 GeV Higgs, or some other Higgs boson, is at least partially responsible for their mass generation (assuming that the new particles have a weak interaction). We will continue to explore this possible connection by searching for evidence of Higgs decays to invisible particle(s). Limits on the decay of the Higgs to invisible particles from the LHC experiments can be used in turn to set limits on the process of dark matter particle-nucleon scattering, which are highly competitive with those from direct searches. The spin-independent cross-section for dark matter scattering on nucleons via Higgs exchange is directly related to the invisible width of the Higgs.

The most promising channel for this analysis is Higgs production through Vector-Boson Fusion (VBF) as shown in Figure 1.1.4 (left). The signature for this process, with an invisible Higgs decay, is two forward jets, widely separated in pseudorapidity, and large missing transverse energy. There is also a smaller signal contribution from gluon-fusion plus two jets. The two main backgrounds are $Z(\rightarrow \nu\nu) + \text{jets}$ and $W(\rightarrow \ell\nu) + \text{jets}$ (including τ decays), where the charged-lepton in the final state is not identified in the detector. There are also small backgrounds from the strong

production of jets (multijet), $t\bar{t}$, and diboson production. For the Run 1 analysis White served as contact editor together with Ketevi Assamagan(BNL), and Bill Quayle(LBNL). 20.3 fb $^{-1}$ of data at 8 TeV were analyzed and a limit of 28% was set on the branching ratio of the H(125) to invisible particles at 95% confidence level (with an expected limit of 31%) [?]. This was the best limit set by all ATLAS and CMS analyses for Run 1. The use of a $W(\rightarrow \ell\nu)$ control region, as well as the $Z(\rightarrow \ell\ell)$ control region, to estimate the $Z(\rightarrow \nu\nu)$ + jets background was a significant factor in setting our low Run 1 limit. Figure 1.1.4 (right) also shows the limit, as a function of mass, that our analysis sets on the Higgs-nucleon cross-section, and the comparison with direct search limits. White presented the ATLAS and CMS results on invisible decays of the Higgs at the LHCP2015 Conference in St. Petersburg, Russia. [?]




Figure 2: (left) Higgs production via Vector-Boson Fusion. (right) WIMP nucleon cross section as a function of the WIMP mass.

The increased Higgs production cross-section at 13 TeV, and the higher integrated luminosity from Run 2 will allow us to probe this process with significantly increased sensitivity over our Run 1 results. White (working with Alex Madsen (DESY)) is analysis contact for the Run 2 Higgs to Invisible analysis. Weekly meetings are held to coordinate the contributions to the analysis: data/Monte Carlo samples and comparisons, trigger(s), control regions specification, theory uncertainties, and limit setting. We have also been holding workshops at critical times to advance the analysis. Given the rate at which ATLAS is acquiring integrated luminosity, we expect in 2016 to reach the point at which we begin to be limited by systematic errors rather than statistically limited. In this respect investigations are ongoing into potential methods for cancellation of systematics in efficiency ratios versus the usual application of transfer factors from control to signal regions. Specific contributions from UTA to the Run 2 analysis include calculating the electroweak corrections to the the VBF Higgs signal, and providing the parton distribution function (pdf) uncertainties on the signal. For the electroweak corrections, we have set up HAWK [?] on the UTA Tier 3 in multi-threaded mode as more than 10^9 events are needed to achieve sufficiently precise corrections. HAWK provides the calculation of differential EW correction factors, which can be used as differential EW reweighting factors to improve QCD-based predictions. The results of the HAWK calculations are shown in Figure 1.1.4 - the reweighting factors have essentially a linear dependence on the Higgs Pt, and a linear fit yields the required electroweak correction.

For the calculation of the VBF signal pdf uncertainties, UTA has been running the generator Powheg v.2 [?] which allows the storage of multiple weights for each event. Our original VBF signal dataset was created using the CT10 pdf set. There are now recommended pdf sets for LHC Run 2 which combine CT14, MMHT2014, and NNPDF3.0 [?]. Currently, after exchanges with Higgs and Exotics group colleagues we have chosen to use the PDF4LHC15-nlo-30-pdfas set. This




Figure 3: Electroweak corrections to the Higgs pT distribution in the VBF selection.

is a Hessian set that allows straightforward calculation of the uncertainties, which is the variation in the number of weighted events passing the VBF selection. This work is being carried out by UTA Honors undergraduate student Darshan Chalise under White's direction. UTA will continue to provide both the electroweak and pdf uncertainties for future versions of our analysis.

The current analysis plan is to write a paper based on the 2015-16 data set, estimated to comprise about 30 fb^{-1} . The goal is to have this result ready for Moriond 2017. White will take on a new graduate student in the Fall 2016 semester. This student will work on the 2017-18 extension of the VBF Higgs to Invisible analysis. Going forward we will focus on improvements and changes to the analysis that will be necessitated by the rising instantaneous luminosity - pileup mitigation, particularly for forward jets, and reduction of the systematics for the jet energy scale and resolution. Work in these areas will form part of White's new student's ATLAS qualification tasks. Of significant physics interest, with reducing limits on the Higgs to invisible branching ratio, will be our ability to exclude models that predict an enhanced value of this ratio. There is a natural evolution of White's physics interest in this topic also with the ILC project (see the ILC section below), which will allow much stronger constraints to be set on Higgs to invisible decays.

VBF Higgs to Invisible Analysis Timetable of Activities

2017

- New UTA graduate student starts work on ATLAS
- Publish Run 2 paper based on 2015-16 dataset
- Work on "interpretation" paper for Run 2 results, DM-nucleon limits and more
- Study strategies for reduction of jet systematics
- Update Higgs to Invisible limit as more data becomes available

2018

- Graduate student pursues ATLAS qualification with work on strategies for reduction of jet systematics
- Contribute data/MC comparisons and cutflow checks
- Provide updated electroweak signal corrections, and pdf uncertainties
- Graduate student begins year at CERN
- Work on branching ratio limit from full Run 2 dataset

2019

- Publish full Run 2 result
- Work on analysis strategies beyond Run 2

1.2 SUSY Physics (PI: De, Farbin)

Despite no evidence of Supersymmetry (SUSY) in LHC Run 1 and first Run 2 data, this extension of the Standard Model’s symmetries continues to remain the most compelling mechanism to avoid the fine-tuning problem of the electroweak symmetry breaking mechanism (also referred as the hierarchy or naturalness problem), while also providing Gauge Unification and a Dark Candidate. The current highly successful LHC run, with a significant increased center of mass energy and integrated luminosity, provides a brief window where data-doubling time is short and large strides can be made in the search for new particles. As LHC nears end of Run 2, further strides will require more sensitive techniques that can target difficult regions while also searching broadly for excesses. This section overviews activities and plans of UTA SUSY group, consisting of faculty De and Farbin, postdocs Usai and Heelan, and students Bullock, Little, and Rogers.

1.2.1 Squarks and Gluino (PI: Farbin)

The most stringent limits on sparticles generally come from inclusive searches for strongly produced squark and gluinos, in particular to the all hadronic final states. In ATLAS, These searches are performed within the inclusive squark and gluino sub-group, composed of approximately 140 members and presently split into eight paper streams based on the number of jets and lepton flavors in the final state. UTA has been involved in these searches since beginning of Run 1, twice convening this subgroup (UTA postdocs David Cote and Heelan), and most recently supervising several public papers and conference notes for ICHEP 2016 [?, ?, ?, ?, ?, ?, ?, ?, ?, ?, ?, ?, ?, ?]. Heelan also directly contributed to the search for SUSY in all hadronic states, which is a flagship SUSY search.

Between 2010-15, Farbin’s ECRP grant supported introduction of novel “event view” techniques to search for SUSY, with the Razor reconstruction technique [?] in 2011 [?] and 2012 [?]. The most recent iteration of this line of searches, based on an extension called the Recursive Jigsaw Reconstruction (RJR), was fully integrated into the mainstream Run 2 search performed for ICHEP 2016 [?] and was the final topic of Bullock’s now completed PhD thesis []. The RJR groups all final state objects into hemispheres based on assumptions (or views) about how the decay is proceeding, providing an uncorrelated basis of observables consisting of energy scales (masses, transverse quantities) and scaleless variables (like angles between hemispheres and ratios of energy scale variables). RJR-based selections generally only partially overlap with standard selections (i.e. based on missing transverse momentum, H_T , etc.), for example bringing in an observed 50% additional events in the all hadronic final state and thereby extending both the phase space and the statistical reach of such searches (if properly combined).

An example of the views used in the latest analysis presented at ICHEP2016 are shown in Figure 4. Using these views signal regions were designed to search for SUSY produced by squark or gluino pair production and direct decay to final states containing jets and the lightest supersymmetric particle. While the new RJR had a standalone expected sensitivity similar to that of the standard technique, RJR extended sensitivity by also targeting difficult phase-space regions with views of Compressed Spectra (CSS) scenarios where all final state objects are assumed to recoil off an ISR jet [?]. In CSS, small mass splittings between the heavier produced sparticle and lightest SUSY particle (LSP) lead to soft final state objects and missing transverse momentum that fail traditional scale-depended background suppression selections.

As seen in Figure 5, the RJR ISR event view extended the squark and gluino exclusion limits with respect to the standard effective mass analysis for the squark-squark and gluino-pair production cases by as much as 50% and 100%, respectively, reducing mass differences sensitivity by




Figure 4: Recursive Jigsaw Reconstruction event views for three scenarios: (a) Two sparticles (P_a and P_b) are pair-produced with each decaying to one or more visible particles (V_a and V_b) which are reconstructed in the detector, and two systems of invisible particles (I_a and I_b) whose four-momenta are only partially constrained. (b) An additional level of decays can be added when requiring more than two visible objects. (c) Strong sparticle production with an ISR decay tree for use with small mass-splitting spectra, where a single sparticle system S decays to a set of visible momenta V and invisible momentum I , and recoils off of a jet radiation system ISR.




Figure 5: text

50 GeV to $\Delta m(\tilde{q} - \chi_1^0) = 100$ GeV at $m_{\tilde{q}} = 700$ GeV and $\Delta m(\tilde{g} - \chi_1^0) = 50$ GeV at $m_{\tilde{g}} = 925$ GeV. For Run 2, Farbin's student Bullock helped Rogan (Harvard) and Jackson (Adelaide) implement and commission a Razor-based high-level trigger that uses only scale-less quantities and therefore enhances CSS trigger efficiencies. This trigger is not yet used in Run 2 searches because the prescale threshold for the missing energy triggers are still below the expectations and the corresponding optimization of Razor trigger.

1.2.2 Run 2 Searches (PI: Farbin)

The unique novel tools and high level of technical proficiency that is characteristic of Farbin's SUSY group renders them extremely efficient at participating in mainstream searches while simultaneously exploring and employing leading edge techniques in complimentary efforts. Due to a long history with the Analysis Software Group (ASG), SUSY software, and ATLAS software tutorials, the

group is in many ways the best equipped and most experienced in ATLAS to easily navigate the required technical obstacles of manipulating data and performing analyses. RJR and the associated RestFrames [] toolkit are also such an asset. A new tool is Deep Learning, an area where Farbin’s breadth and depth of efforts is unmatched in HEP.

The ATLAS SUSY working group currently plans two more full rounds of searches in Run 2, which no doubt will include flagship inclusive searches for squarks and gluinos. Considering UTA’s long history with these searches, and the fact they are generally under-manned, Farbin’s group will continue to lend critical manpower to these efforts, for example by including graduate student Rogers in order to familiarize her with a large scale ATLAS analysis.

The ATLAS SUSY group is also considering a reorganization of searches from model-based (e.g. squarks and gluons, stop, electro-weak chargino, etc.) back to signature based (i.e. all hadronic, one lepton, etc.), where each signature will include a large number of signal regions targeting different models. This change will be a welcomed by Farbin’s group as it more closely mirrors their first Razor-based search [?] and Farbin’s long term plans for general searches. As this reorganization occurs, Farbin’s group will work to help define RJR-based regions for signatures beyond the all hadronic final states, by ensuring that appropriate RJR views are constructed for every search and corresponding observables are available to analyzers as they define signal regions.

As the gains in SUSY reach from Run 2’s energy and luminosity boost are exploited in the first Run 2 searches, subsequent searches will require dedicated effort to identify and target difficult phase-space regions and increasingly sophisticated techniques to extend reach. At the end of Run 1 both ATLAS and CMS scanned feasible parameters in the phenomenological MSSM (pMSSM) [?,?] to understand the strengths and weaknesses of traditional searches [?,?]. Then UTA postdoc David Cote was one of the primary drivers of this effort. In the pMSSM the 105 free parameters of the MSSM are reduced to 19 by making assumptions based on theoretical and experimental results. Thousands of SUSY signals were generated by varying those 19 free parameters uniformly (within some allowed physical range). These studies helped inform the following strategies Farbin’s group will pursue as they continue to design and carry out SUSY searches:

- **Cross-section optimized searches:** While exclusion limits on large cross-section processes, such as strongly produced squark and gluino, can be extended to higher masses (i.e. lower cross-sections) with more integrated luminosity, LHC searches are insensitive to high cross-section scenarios where traditional scale-dependent observables such as H_T (scalar sum of momenta) and missing energy are small. In addition, since searches are optimized to extend mass limits, which rely on cross-sections from Simplified Models where only relevant new particles with 100% branching fractions are added to the SM, sensitivity to smaller cross-section at lower masses are often sacrificed. Farbin’s group will rely on the Razor trigger and the RJR techniques to address the former problem, while working to define additional signal regions to address the latter.
- **Compressed spectra and initial state radiation:** The low p_T spectra (a.k.a. soft) of decay products in CSS prohibit differentiation from Standard Model processes and fail standard trigger selections. Mono-jet searches can probe these models, as they search for a single high p_T jet, presumed from initial state radiation (ISR), and missing transverse momentum. The RJR ISR technique, as applied for ICHEP2016 and detailed in [], provides the most powerful method for generically searching for any CSS scenarios without dependence on the SUSY decay. Farbin’s group will collaborate with Rogan and Jackson to develop such CSS general search.
- **High multiplicity final states:** Models that include sleptons in the decay chain are less likely

to be excluded, often coming from longer decay chains that involve charginos decaying to Standard Model bosons (V/h). These longer decay chains can contain leptons and/or a high multiplicity of jets in the final state. Farbin's group will pay special attention to ensuring these scenarios are covered with dedicated signal regions or searches.

- **Boosted Heavy SM particles:** Highly boosted W , Z , top, and Higgs are common in SUSY decays, often appearing as Jets with unique topologies. Tagging these particles is being pursued through out ATLAS, most recently using Deep Learning techniques. Beyond contributing to the development of these techniques and exploring application SUSY searches, Farbin's group will ensure these techniques are applied in the SUSY data processing and available to all SUSY collaborators.

Finally Farbin proposes to work towards the R& D of two new techniques for SUSY searches. The first attempts to classify the topology of every event. For example, consider the $b\bar{b}ll + \text{MET}$ final state which can come from a large number of SUSY and SM processes, for example stop pair production with $\tilde{t} \rightarrow t\tilde{\chi}^0$, stop pair production with $\tilde{t} \rightarrow b\tilde{\chi}^\pm$ and $\rightarrow \tilde{\chi}_o W^\pm (\rightarrow l\nu)$ or $\rightarrow l\tilde{n}u (\rightarrow \tilde{\chi}^0\nu)$, chargino pair-production with $\tilde{\chi}_2^0 \rightarrow Z (\rightarrow ll) + \tilde{\chi}_1^0$ and $\tilde{\chi}_2^0 \rightarrow h (\rightarrow bb) + \tilde{\chi}_1^0$, or fully leptonic SM $t\bar{t}$. The RJR allows constructing views and associated observables for all such topologies, enabling discrimination by directly comparison of observables from different views, a handle used in the RJR-ISR CSS technique described above. Using a Parameterized Deep Learning Classifier [], which builds a neutral network optimal for any mass in the decay tree, Farbin was able to correctly identify a few signal topologies over 95% of the time []. The vision here is a signature-based search in-line with SUSY WGs plans, where events further sorted into topologies. Farbin's group in collaboration with Rogan and Jackson will fully explore the $b\bar{b}ll + \text{MET}$ final state and lead SUSY WGs towards this approach. The Long-term vision of this thrust is to apply the technique across large number of final states, working towards a general search with the entirety of the Run 2 data during LS2.

The second technique is simply to use Deep Learning classifiers as the multivariate tool to improve background rejection. As a first step, Farbin will assist in upgrading existing multi-variate based searches, for example the stop search performed by De described in section ??, to the potentially more powerful DNNs. It is also rather straightforward to evaluate the potential of DNNs to improve other non-MVA based searches, for example those planned by Hadavand and Brandt in section ?? . While DNNs based on traditional high-level observables like masses and angles (a.k.a. features) may provide modest improvements, DNNs can also be used to identify additional handles by using 4-vectors as input. A such studies typically find that 4-vectors generally perform better than traditional features. The problem is that most searches rely on some orthogonal observable, otherwise unused in the MVA, to build sidebands for background estimation. A 4-vector based DNN's output will likely be highly correlated with any such observable, prohibiting designating unbiased sideband regions. While there are avenues to tackle this problem, its resolution is not necessary to reaping benefits of DNNs in searches. Instead, 4-vector based DNNs outperforming DNNs based on traditional variables can be studied to identify the new features (i.e. observables) than can be added to the traditional analysis. For example DNNs were able to identify additional color-flow related observable to separate light jets from W and other heavy object initiated jets [].

Service at ATLAS

1.3 Intermediate Tile Calorimeter (PI: De, White)

Intermediate Tile Calorimeter, [PI: De, White] The Intermediate Tile Calorimeter (ITC) was designed to fill the gaps between the central barrel calorimeter and the extended barrel calorime-

ters, and was built at UTA. Figure 6(a) shows a schematic view of part of the ATLAS calorimeter, showing the components of the ITC. For particles which originate at the nominal interaction point, the ITC extends over approximately $0.8 < |\eta| < 1.6$. The region $0.8 < |\eta| < 0.9$ contains 311 mm thick steel-scintillator stacks, similar in design to standard Tile Calorimeter submodules. Between $0.9 < |\eta| < 1.0$, the stacks are 96 mm in the z -direction. The combined $0.8 < |\eta| < 1.0$ region of the ITC is called the plug. In combination with the support structure for the scintillators, it is called the ITC submodule. For the forward region, the active elements of the ITC consist of scintillator only due to space constraints. The scintillators between $1.0 < |\eta| < 1.2$ are called the gap scintillators, while those between $1.2 < |\eta| < 1.6$ are called cryostat scintillators. Some additional scintillators, designated E4-prime were installed in 2015, covering $1.6 < |\eta| < 1.72$ and a small sector in ϕ , in order to explore the possibility of extending ITC coverage for improved electron and jet measurements. Together, all these ITC detectors improve the measurement of total energy in the intermediate region, thereby improving the jet and E_T resolution of ATLAS.




Figure 6: Left - Layout of the Intermediate Tile Calorimeter. Right - Average change for ITC E-modules vs. integrated luminosity for Run 1.

It is anticipated that the E cell scintillators will need to be replaced in the long shutdown in 2019, due to the expected high levels of radiation damage. White, with postdoctoral associate Park, previously carried out studies of radiation damage using Gaussian plus Landau fits to the signal distributions for all E-cell channels. This proved to be a very challenging exercise due to the high degree of variability of the signal profiles. Nevertheless, results were obtained as shown in Figure 6(b). Most of the decrease in response can be attributed to the downward response of the PMT's due to continual exposure to light during the run, as shown by the laser illumination of the photocathodes. However, there was a few percent decrease on top of this effect that was attributed to radiation damage. This approach to monitoring decreased response has now been replaced with the use of minimum bias data plus the laser measurements. The replacement of the E-cells scintillator affords the opportunity to revisit the η and ϕ segmentation of the cells. In order to determine the best layout for optimization of energy resolution in this region, UTA has been working with the Tbilisi group to examine the distributions of jet energy fractions in the E-cells. The work at UTA has been carried out by Honors undergraduate student Niyusha Davachi under White's supervision. As an example, Figure 1.3 shows the fractional energy contribution for each of the E-cells, for QCD jet events where the jet axis lies in E4, for three groupings of adjacent ϕ cells (3,5, and 7). By studying these fractional energies for jets within all the E cell η ranges, we will optimize both the use of the deposited energies for jet reconstruction, and the propose potential new layout(s) for the E cells. This work is ongoing with the goal of providing a recommended

layout of the new E-cells scintillator in late 2017/18, when the new tiles will have to be ordered.




Figure 7: Fractional contribution to jet energy by the E_i ($i = 1, 2, 3, 4$) cells for the case where the jet axis lies in the E_4 range, $1.4 < |\eta| < 1.6$.

While the scintillator replacement is not within the scope of US ATLAS work for Phase I, the UTA group has made contributions to the estimation of the expected integrated radiation dose during Run 2, to the specification and identification of radiation-hard scintillator, and will be involved in the procurement, installation and testing of the new tiles. This work will also be included in the qualification tasks for White's new graduate student.

ATLAS Tile Calorimeter Service White has been a member of the Tile Calorimeter Speakers' Committee, with responsibility for major conferences such as IEEE NSS and DIS, and for the review of Tile proceedings and notes.

ATLAS Tile Calorimeter Timetable of Activities Our work on the Tile Calorimeter/ITC will be carried out in collaboration with colleagues in CERN and Tblisi.

2017

- Complete the study of the optimal use of existing E cells for improving jet energy resolution
- Update expected radiation doses for E cells based on known and projected Run 2 luminosities.
- Study/propose alternative E cells configurations for the 2019 tile replacements
- Continue as a member of the Tile Calorimeter Speakers' Committee.
- Contribute DQ Validator shifts

2018

- Complete plans for ATLAS ITC Tile/Fiber replacements in LS2.
- Investigate the availability of radiation-hard scintillator for new ITC tiles.
- Prepare (with ATLAS Tile Institutes) tile/fiber assemblies for installation in LS2.
- Graduate student begins time at CERN, with Tile Calorimeter qualification tasks.
- Contribute DQ Validator shifts

2019

- Study effects of radiation on light yield/transmission for dismounted ITC tiles.
- Contribute effort to installation and testing of the new ATLAS ITC E-cells.
- Graduate student returns to UTA.

1.4 Trigger (PI:)

Following the recommendations of the 2014 ATLAS Future Framework Requirements Group Report [], the ATLAS Phase I Software Upgrade will entail a massive migration to athenaMT, a multi-threaded evolution of ATLAS’s current framework. This migration requires significant redesign and rewrite of existing software components and algorithms. An aggressive schedule delivers basic functionality by 2016, begins select migration in 2017, and performs bulk migration in 2018-19 in preparation for full integration in 2020 and start of data Run 3 data-taking in 2021. A cornerstone of this migration is the merger with the ATLAS trigger, where high pile-up and L1 rates necessitate use of offline algorithms. A key trigger component is a new incarnation of the Event View concept which enables partial event processing in regions of interest. Numerous technical and design issues must be addressed in both the offline and trigger for athenaMT’s first release at the end of 2017 and subsequent algorithm migration, a formidable task especially in light of a significant lack of manpower and requisite expertise.

Trigger athenaMT Migration Farbin proposes to redirect his focus, which has been very recently freed from coordinating DUNE Software & Computing, to the trigger athenaMT migration. Given Farbin’s long history with ATLAS offline software development and management, broad knowledge of HEP software framework, and extensive experience with parallel and GPU computing, the upcoming trigger coordinator Jeorg Stelzer (CERN) and core trigger software coordinators John Baines and Tomasz Bold have enthusiastically welcomed Farbin’s much needed commitment to this effort.

Farbin will commence initial work immediately during his current sabbatical, taking on migration of the e/gamma trigger chain with tracking, calorimetry, and hypothesis starting with TopoCluster calorimeter clustering algorithm and then continuing on through the Trigger Software Upgrade’s extensive workplan []. The in-depth experience gained here will allow Farbin to gradually shift from algorithm migration to design and implementation of core components relating to Event Views, Menu, Decision, Prescales, Monitoring, and so on. We propose that a new UTA postdoc begin assuming Farbin’s projects in mid-2017, dedicating 50% of his/her effort and freeing Farbin for bigger contributions to this area, possibly through management and leadership roles similar to his convenership of the Physics Analysis Tools group.

GPU Trigger Demonstrator As R&D for HL-LHC, Farbin and Hilliard will also initiate an effort to revisit the ATLAS GPU Trigger Demonstrator [], which implemented an example of a full tracking and calorimetry algorithm chain for CPU and GPU. In 2015 this study found that while the GPU algorithms ran up to two orders of magnitude faster, data conversions necessary for the GPU eliminated any cost benefits. Meanwhile the Deep Learning renaissance and rapid co-processor evolution have significantly shifted the future software and hardware landscape away from the trajectory set by the Future Framework Requirements Group Report. It is imperative that HEP investigates the potential of the industry’s Deep Learning infrastructure to address HL-LHC computing challenges. And considering that Deep Learning can improve and accelerate reconstruction algorithms, close integration of DL with HEP software will no doubt be a requirement of HL-LHC HEP software.

After deploying the GPU Demonstrator on Farbin’s GPU systems, Farbin and Hilliard will migrate it to TensorFlow (TF), Google’s open source distributed computation framework developed for Deep Learning applications. In contrast to the schedulers in multi-threaded HEP frameworks, TF utilizes data flow programming, representing computation as directed acyclic graphs that it optimizes and dispatches for processing across cores, co-processors (including Google’s ultra fast

and efficient Tensor Processing Units), and HPC nodes. The system will be studied on Farbin’s 48 core, 7 GPU, 10 Gb linked systems, with data pre-converted to tensors that flow through TF ops wrapping kernels of the existing GPU-Demonstrator algorithms. The results will be documented in a report intended to help shape HL-LHC software and computing planning. It is noteworthy that Hilliard had obtained support from Fermilab (under supervision of Jim Kowalkowski) to perform a similar R&D exercise with LArTPC event processing during summer 2016, but instead took a Silicon Valley internship.

Computing at ATLAS

Computing costs for HL-LHC are estimated to be at least a factor of 5 more than current LHC [], and likely much more due to the stalling of Moore’s Law. While greater use of HPCs in HEP will ideally alleviate some of the problem, HEP software has yet to harness the parallelism in the processors, like GPUs, where these systems get most of their horsepower. Farbin’s success in Deep Learning has its roots in his projects with undergrads on GPUs and the GPU systems they build and share with XX colleagues and collaborators. [The primary machine (2x 12-core Xeon, 128 GB RAM, 50 TB disk) hosts one previous generation and three older NVidia GPUs.] Farbin’s initial work with these GPUs was aimed at understanding how to efficiently use them in the ATLAS software framework resulting in 4 unfunded NSF collaborative proposals. Section ?? describes how Farbin will direct this thrust of work towards migration of ATLAS trigger to multi-thread version of Athena and integration of GPUs.

Meanwhile, the world is in the midst of a renaissance in Machine Learning and Artificial Intelligence, known as Deep Learning, driven by the emergence of large data sets, powerful Graphical Processing Units (GPUs) processors, and new techniques to train billion-neuron multi-layer artificial neural networks. Systems trained on raw and sometimes unlabeled data, can now recognize objects, detect human emotion and intent, play video games, translate between languages, generate mathematical proofs, sometimes better than humans and most importantly with minimal engineering. Developed in University Machine Learning labs, DL has led Google, Facebook, and other industry-leading companies to rethink everything, building Artificial Intelligence teams, software, processors, and cloud services, and demonstrating impressive feats. DL is the first multi-domain application that requires and easily harnesses the power of specialized processors, like GPUs, and supercomputers (i.e. HPCs). Section ?? describes Farbin’s efforts to apply DL to HEP.

1.5 Core Trigger Software (PI: Farbin)

Following the recommendations of the 2014 ATLAS Future Framework Requirements Group Report [], the ATLAS Phase I Software Upgrade will entail a massive migration to athenaMT, a multi-threaded evolution of ATLAS’s current framework. This migration requires significant redesign and rewrite of existing software components and algorithms. An aggressive schedule delivers basic functionality by 2016, begins select migration in 2017, and performs bulk migration in 2018-19 in preparation for full integration in 2020 and start of data Run 3 data-taking in 2021. A cornerstone of this migration is the merger with the ATLAS trigger, where high pile-up and L1 rates necessitate use of offline algorithms. A key trigger component is a new incarnation of the Event View concept which enables partial event processing in regions of interest. Numerous technical and design issues must be addressed in both the offline and trigger for athenaMT’s first release at the end of 2017 and subsequent algorithm migration, a formidable task especially in light of a significant lack of manpower and requisite expertise.

Trigger athenaMT Migration Farbin proposes to redirect his focus, which has been very recently freed from coordinating DUNE Software & Computing, to the trigger athenaMT migration. Given Farbin’s long history with ATLAS offline software development and management, broad knowledge of HEP software framework, and extensive experience with parallel and GPU computing, the upcoming trigger coordinator Jeorg Stelzer (CERN) and core trigger software coordinators John Baines and Tomasz Bold have enthusiastically welcomed Farbin’s much needed commitment to this effort.

Farbin will commence initial work immediately during his current sabbatical, taking on migration of the e/gamma trigger chain with tracking, calorimetry, and hypothesis starting with TopoCluster calorimeter clustering algorithm and then continuing on through the Trigger Software Upgrade’s extensive workplan [1]. The in-depth experience gained here will allow Farbin to gradually shift from algorithm migration to design and implementation of core components relating to Event Views, Menu, Decision, Prescales, Monitoring, and so on. We propose that a new UTA postdoc begin assuming Farbin’s projects in mid-2017, dedicating 50% of his/her effort and freeing Farbin for bigger contributions to this area, possibly through management and leadership roles similar to his convenership of the Physics Analysis Tools group.

GPU Trigger Demonstrator As R&D for HL-LHC, Farbin and Hilliard will also initiate an effort to revisit the ATLAS GPU Trigger Demonstrator [2], which implemented an example of a full tracking and calorimetry algorithm chain for CPU and GPU. In 2015 this study found that while the GPU algorithms ran up to two orders of magnitude faster, data conversions necessary for the GPU eliminated any cost benefits. Meanwhile the Deep Learning renaissance and rapid co-processor evolution have significantly shifted the future software and hardware landscape away from the trajectory of set by the Future Framework Requirements Group Report. It is imperative that HEP investigates the potential of the industry’s Deep Learning infrastructure to address HL-LHC computing challenges. And considering that Deep Learning can improve and accelerate reconstruction algorithms, close integration of DL with HEP software will no doubt be a requirement of HL-LHC HEP software.

After deploying the GPU Demonstrator on Farbin’s GPU systems, Farbin and Hilliard will migrate it to TensorFlow (TF), Google’s open source distributed computation framework developed for Deep Learning applications. In contrast to the schedulers in multi-threaded HEP frameworks, TF utilizes data flow programming, representing computation as directed acyclic graphs that it optimizes and dispatches for processing across cores, co-processors (including Google’s ultra fast and efficient Tensor Processing Units), and HPC nodes. The system will be studied on Farbin’s 48 core, 7 GPU, 10 Gb linked systems, with data pre-converted to tensors that flow through TF ops wrapping kernels of the existing GPU-Demonstrator algorithms. The results will be documented in a report intended to help shape HL-LHC software and computing planning. It is noteworthy that Hilliard had obtained support from Fermilab (under supervision of Jim Kowalkowski) to perform a similar R&D exercise with LArTPC event processing during summer 2016, but instead took a Silicon Valley internship.

1.6 Deep Learning (PI: Farbin)

Space constraints prohibit pedagogical discussion of Deep Learning and details of much of the activity described and planned out in this section. Surveys of DL activity in HEP and much of these details can be found in Farbin’s recent talks [3], for example at HEP Software Foundation [4] HSF, ATLAS Machine Learning workshop [5], or the Harvard Big Data Conference [6]. This section is

organized into DL thrusts that sometimes span multiple experiments: Simple Classification, Matrix Element Acceleration, Imaging Detector Reconstruction, and Tracking.

Simple Classification with Deep Learning The first application of DL in HEP demonstrated that (fully-connected) DNNs out-perform shallow networks and derive new “features” from 4-vectors beyond traditional observable. Farbin recently demonstrated this simple application of DNNs with help of Chris Rogan (Harvard) and Paul Jackson (Adelaide). They showed that 4-vector based DNNs out-performed a Jigsaw-based DNNs in correctly distinguishing between different SUSY-like decay topologies. This work is in a step in an effort to build a DNN that generically classifies events and enables unsupervised general searches for new phenomena. They are also now applying DNNs to compressed-SUSY scenarios, building on experience of the ICHEP 2016 Jig-saw result and following Rogan and Jackson’s recent paper [1]. Section ?? provides additional details on these topics in the context of SUSY. [Hadavand and Brandt will apply similar techniques to their work...]

Signal and background classification tasks are the first and simplest and can be easily enhanced by replacing traditional classifiers with DNNs. Groups in ATLAS are applying this idea to tasks like b-jet or boost-object tagging. Farbin has developed a Deep Learning Tutorial covering a variety of DNN tasks, such as event or object classification with any dataset. The tutorial introduces a thin framework (DLKit) developed by Farbin that simplifies running large-scale DL studies using Keras, a powerful python-based DL framework which supports both Theano and TensorFlow backends. Postdoc Heelan, who manages the ATLAS software tutorials and workbook, will introduce the DL tutorial at the October 3rd session, with subsequent iterations adding all of the additional tasks Farbin is currently performing in DLKit, such as Matrix Element Regression and 2D and 3D Imaging detector classification and Energy regression. Farbin, Brandt, Hadavand, Asaadi, and Jones have committed students to applying DNNs to various problems, and this DLKit and tutorials are integral to these projects.

While the tutorial helps address one obstacle to broad adoption of DL, namely familiarity and full end-to-end working examples, access to GPUs, which give two orders of magnitude acceleration of DL training with respect to CPUs, remain as the next biggest obstacle. For the past few years, Farbin has granted access to the GPU system he has built with his undergrads to XX HEP scientists from YY experiments (LArIAT, DUNE, MicroBooNE, NEXT, CMS, ATLAS). Farbin performed an early study of turning regression problems into classification problems on the Oakridge’s Titan HPC via PANDA WMS with the help of Sergey Panitkin (BNL), the first use of Titan’s GPUs in HEP. A subsequent iteration of the DLKit tutorial will include submission of Hyperparameter scan DL jobs to Oakridge’s Titan computer and ideally any other HPC accessible via PANDA and with suitable software.

Matrix Element Acceleration with Deep Learning The Universality Theorem [2] states that a sufficiently large Neural Network can represent an arbitrary function of any input dimensions. The theorem implies that fast DNNs can potentially encapsulate prohibitively expensive computations, for example Matrix Elements in context of NLO and NNLO computations, which are becoming increasingly important for the LHC, or the application of Matrix Element Method (MEM), which is in principle the most powerful search technique but is sparsely applied due to technical complexity and computational demands. The idea is to perform the computations once using significant computing resources, for example an HPC, and train a DNN to efficiently (in both memory and speed) accurately reproduce the same computation for subsequent event generation, integration for cross-sections, or application of MEM.

Farbin has been pursuing both of these problems in collaboration with Tancredi Carli (CERN), ATLAS’s next Deputy Physics Coordinator and Tobias Golling’s group (U. Geneva). Carli has produced a one billion event sample of NLO $t\bar{t}$ in Sherpa which he has been studying on Farbin’s GPU system due to memory requirements. He has been attempting to encapsulate event weights in a high dimensional (i.e. incoming and outgoing 4-vectors) adaptive binning histogram using the foam [] method. Simultaneously Farbin has developed a Matrix Element DNN (MEDNN) to achieve the same task. Since Golling’s group, who initially relied on Farbin’s GPU system, is pursuing a similar task for Matrix Element Method based analysis, Farbin joins Golling’s group meetings and shares ideas and code. DNN training for such a regression task is essentially a high dimensional fit, which due to the non-linearity of NNs, is often biased unless a proper cost function is applied. This problem is discussed in the Imaging Detector Energy Regression description below. For MEDNN, the state-of-the-art turns the regression (i.e. fitting) into a classification problem by binning the target with bin learned bin edges.

Imaging Detector Reconstruction with Deep Learning Starting in 2014 Deep Convolution Neural Networks (CNNs) have exponentially improved performance on ImageNet, a one-million image classification challenge [], to now super-human performance []. An obvious application of CNNs is the classification tasks, such as particle identification, in “imaging” detectors such as Time-Projection Chambers (TPCs), Cherenkov Imaging detectors, and high granularity calorimeters. The application in HEP of this idea was in the Nova collaboration, where they were able to obtain 40% better electron efficiency for same background rate as their best techniques [].

LArTPC In fall of 2015, Farbin CNN-based demonstrated particle identification in LArTPC with simulation of the LArIAT Experiment []. Despite a great deal of multi-experiment (e.g. ArgoNeuT, LArIAT, MicroBooNE, DUNE, ...) effort, event reconstruction in LArTPC has proven be challenging, with performance still far from expectation and poor neutrino reconstruction. Farbin’s Inception [] based CNN treated raw data from the TPC as images, and was able to achieve significantly better performance than traditional reconstruction in both particle and neutrino reconstruction [] (give example numbers??). This effort has now evolved to a collaboration with computer scientists Pierre Baldi and Peter Sadowski at UCI. While the ultimate goal is demonstrate end-to-end DL-based Neutrino reconstruction, the first goal is to design optimized networks for particle and neutrino classification and energy regression. The first step was to produce large samples of simulated events with flat energy spectrum. Using Farbin’s local machines Shahsavari and undergrad Hilliard have so far produced 15 million events of every particle species, which is a largest LArTPC sample ever produced (?). This dataset is public, is intended to facilitate collaborations, and is already being used by independently by HEP and DL researchers.

The collaboration with UCI has yielded several particle and neutrino classification and regression networks, including Siamese [] Inception and ResNet [] based networks that seem to require less depth for LArTPC tasks than image classification. Obtaining better classification performance than automatic reconstruction has proven to be rather easy. Recently MicroBooNE presented some promising first studies [], based on work that was initiated on Farbin’s GPU system. Farbin et al we able to obtain better performance, training for a week with $O(100k)$ down-sampled raw events, for example achieving 2% fake rate at 80% for electron versus neutral pion discrimination, a critical benchmark for separation of charged current from neutral current neutrino interactions. However obtaining the design performance (1% fake rate) will likely require full resolution and deeper networks train for long time on the significantly larger training sample Farbin et al have produced.

Obtaining good energy reconstruction has proven to be difficult. In order to avoid bias, Farbin

et al employ likelihood based cost functions that also provide a per-event resolution estimate. First studies with small networks on down-sampled data obtain poor electron resolution with 11% sampling term, significantly worse than the expected 3%. Meanwhile Muon energy reconstruction relies on measuring multiple scattering angle, and thereby requires full detector resolution. Again, these studies are at the stage of much more challenging training requirements and are underway, with the goal of paper submission by end of 2016.

Farbin and Asaadi propose to implement the network architectures and techniques developed in these studies as DNN-based algorithms in the software framework common to nearly all LArTPC experiments, LArSoft []. This work will be carried out primarily by Shahsavarani and undergrad [Asaadi insert name here] in collaboration with Robert Sulej (CERN, Fermilab, ...), who has been pursuing DNN-based electromagnetic versus hadronic hit identification, also on Farbin’s GPU system. As these algorithms are implemented, they can be tested or trained on LArIAT data. The goal will be to provide full end-to-end DNN-based neutrino reconstruction in LArSoft by the end of 2017.

Gas TPC The neutrinoless double beta ($0\nu\beta\beta$) experiment NEXT, relies on high pressure xenon (HPXe) Gas TPC, read out by SiPMs to produce 3-D images of beta decays. NEXT relies on topological signatures to separate signal events (two electrons) from background events (mainly due to single electrons with kinetic energy comparable to the end-point of the $0\nu\beta\beta$ decay). Late 2015, Farbin, with the assistance of Josh Renner (?), ... (?), demonstrated that CNNs can nearly perfectly separate this signal and background in an idealized toy simulation. Since then, they have shown CNNs out perform the traditional technique in full simulation. They also used 2D DNNs to easily compare different detector granularities and understand the relative contribution of physics processes to the fake-rate. This activity is detailed in a paper (with Farbin as co-author along with the NEXT collaboration) that is nearly ready for submission. Jones is committing undergrad XXX and graduate student XXX, to upgrade the technique to 3D and attempt to identify the decay point and angle, in an effort to test Lorentz invariance [Ben, do I put this in?].

Calorimeters The ATLAS Electromagnetic and Hadronic calorimeters produce 3D images in of variable granularity in η , ϕ versus depth (i.e. the LAr presampler to the Tile D layers) of energy deposits. Particles, such as photons, are identified by their characteristic shower profiles, with their energies determined via a weighted fit of layer-wise deposits calibrated to test beam and Z decays. Any improvement, for example in photon identification or energy resolution, can dramatically improve searches and measurements, for example producing narrower Higgs peaks with less background. This calorimeter (as opposed to CMS’s crystal EM calorimeter) is ideally suited for application of CNN for classification and energy regression. Indeed several factors give hope that significant improvements can be achieved. For example, energy reconstruction in the LAr EM calorimeter currently does not use shower shape information and is not correcting for variations in the LAr calorimeter’s characteristic accordion structure. Similarly the hadronic calibration does not use sampling information.

The variable granularity of the ATLAS calorimeter can likely be built into the CNN architectures. Since simulation does not faithfully reproduce the shower shapes, simulation trained CNNs can be initially calibrated on $Z \rightarrow ee$ or testbeam data. While training on data is also possible, for example training on data Z decays using likelihood-based cost functions that account for Z line-shape and calorimeter resolution, an interesting direction is hybrid training with adversarial networks. An example of this technique simultaneously trains the classification or regression network with and an adversarial network learning to distinguish data and simulation. The full network

is then trained until data and simulation cannot be distinguished.

Another potentially high impact of DNNs to calorimetry is fast showering. Full Geant4 shower simulation in the ATLAS calorimeter takes of order of an hour. Fast shower techniques such as shower libraries or high dimensional binning of shower observable generally suffer from intractable memory requirements. Zach Marshall et al (Berkeley) have more efficiently stored shower observables in shallow neural networks. DNNs may provide a much more powerful technique. Starting with examples only, Generative DNNs have been demonstrated to generate new images of faces or hotel rooms, or text in style of the example author (e.g. Shakespeare). Two techniques are likely relevant to calorimetry. The first is Generative Adversarial Networks, which starting from random input trains a network that simultaneously produces the desired output and attempts to distinguish generated from real examples. The network is trained when it no longer can make the distinction. Variational Autoencoders train a 2 part network: one that “encodes” real examples into a latent lower dimensional representation of Gaussian distributed variables, and another that “decodes” to an image trained to reproduce the original. Once trained, the decoder can be primed to then generate new examples.

A significant obstacle to applying CNNs to the ATLAS calorimeter is accessibility to data. The energy deposits into the 200k cells require significant storage and are only retained in the difficult to access Event Summary Data (ESD). With a baseline of ≈ 1 in 10^4 jet rejection, CNN training will require large samples of specially filtered EM-like jet backgrounds, finely binned to compensate for drop in jet-cross section. Finally, due to ATLAS’s strict data-sharing policies, collaboration with DL experts is difficult. Farbin and graduate student Leslie Rogers are working to address these issues and assemble appropriate ATLAS training sets,

In meantime, Farbin has been collaborating with CMS colleagues Maurizio Pierini (CERN) and Jean-Roch Vlimant (CalTech) to generate simpler public datasets where application of DNNs to calorimetry can be explored in collaboration with DL experts and without unnecessary complications. Starting with the high granularity LCD CLIC calorimeter concept, they have so far simulated 2 million photon and neutral pions and presented first classification studies were presented in July 2016 [1]. With the goal a paper by the end of 2016, Farbin is currently applying the energy regression techniques developed in LArTPC to this dataset.

Ideally, the very clean environment of the LArTPC and Gas TPC and this simple LCD dataset, can serve as a stepping stone for developing the classification, regression, and generation techniques described in this section to the ATLAS calorimeter and the pile-up ridden LHC environment. Exploring and implementing these techniques in ATLAS will be a part of Rogers’ PhD thesis, building on Farbin’s ATLAS software and DL experience and Heelan’s extensive background with the calorimeter (e.g. she is the editor of the Tile performance paper) and test beam. The goal is to first tackle photon identification and then photon/electron calibration. Of critical importance will be handling of pile-up. If successful, the technique will be implemented in ATLAS’s framework for further study and use by other collaborators. Extension of the technique to EM/hadronic cluster identification and calibration should be straightforward next step. Postdoc Griffiths, who contributes significantly to τ identification, will investigate CNN applications to τ s. Simultaneously these efforts will explore applying the generative DNN fast showering technique developed in the context of LCD, to ATLAS.

HEP is confronted with two fundamental problems in the current generation of HEP experimental software: the inability to take advantage and adapt to the rapidly evolving processor landscape, and the difficulty in developing and maintaining increasingly complex software and computing systems by physicists. DL and DL software systems provide a paradigm-shifting solution to these problems. It may be possible to replace the millions lines of code, meticulously crafted by thousands of physicists, with large neural networks that are simply trained on raw simulated

data. Not only are these networks much simpler and faster to compute than traditional HEP algorithms, they inherently make efficient use new processors. These systems can provide scalable implementations of existing HEP algorithms.

Tracking Pattern recognition rate in particle tracking scales quadratically with hits in the tracking detector. As a result, tracking in 200 pile-up HL-LHC events is one of biggest challenges for the HL-LHC, where some tracking and vertex finding at 40 MHz bunch-crossing might be required for the trigger. While some are investigating dedicated hardware, such as GPUs, FPGAs, or associated memory, a group of ATLAS and CMS physicists, including Farbin, are hoping that by presenting the HL-LHC tracking problem as a Machine Learning challenge (TrackingML) with a prize, solutions arise that scales better with number of hits [1]. One source of inspiration is DeepMind’s AlphaGo [2] artificial intelligence agent, which was able to assess moves by looking at the whole board with a DNN instead of performing a look-ahead tree search. This DNN was trained through human expert and self-play games.

Preparing this challenge requires generating HL-LHC like events in an appropriate detector, providing baseline traditional tracking software for comparisons, and developing mechanisms to benchmark and assess the performance of submitted algorithms. The majority of the work so far has been carried out by Andreas Sulzberger (CERN), who is currently in charge of ATLAS reconstruction and formerly a tracking expert. He has developed a standalone simulation and tracking framework, known as ATLAS Common Tracking Software (ACTS). Farbin’s contribution to the project so far has been data conversion to more ML friendly HDF5 format. Farbin and undergrad Hilliard are assuming the responsibility of developing the automatic benchmarking mechanisms, likely using Docker containers on hardware Farbin will dedicate to the project.

Upgrade at ATLAS

1.7 Tile HL-LHC Upgrade (PI: De, White)

1.8 Tile Low Voltage Power Supply(LVPS) Upgrade (PI: Brandt, Hadavand)

In preparation for High Luminosity LHC (HL-LHC), various upgrades of the ATLAS detector to cope with the new conditions are being planned. One of these projects, involving the replacement of the 2048 low voltage power supplies (LVPS) required for operation of the ATLAS Tile hadronic calorimeter, became available last year. Building on UTA’s connections in the TileCal since its inception, Hadavand and Brandt assembled a group and proposed that UTA take over this project. Our proposal was well-received, consequently, Hadavand is now the L3, or deliverable manager, of this project for the ATLAS detector while Brandt is the institute contact. The construction phase of this 6-year 1 million dollar project is expected to be covered by an NSF MREFC proposal in progress.

Starting in 2023 the LHC will be shutdown for 2.5 years in preparation for HL-LHC upgrade to achieve $5.0 \times 10^{34} / cm^2 s$. Hadavand and Brandt have assumed the responsibility for the testing and production of half of the entire 2048 low voltage power supplies (LVPS) needed for HL-LHC running, a project that will cost about 1 million dollars over the next few years, the bulk of which is anticipated to be provided by an NSF MREFC grant. Hadavand is the deliverables manager or CAM for the low voltage power supplies overseeing the work of UT Arlington as well as that of Northern Illinois University which is responsible for integrating the UTA power supplies into boxes that will go into the detector. The UT Arlington group also has responsibilities on the Tile Pre-processor for HL-LHC. We currently share an electrical associate (EA), Seyedali Moeyadi, on these two projects. The EA has worked closely with our research faculty Giulio Usai at CERN and has been involved

in the June TileCal test beam. We have also procured funding from the university for an electrical engineering (EE) PhD student, Michael Hibbard, whose thesis will be based on this project.

We are currently in a transition phase assuming responsibility for the project as Argonne National Labs, who designed and built previous versions of power supplies is assuming responsibilities in the upgraded tracking detector. They have produced a working prototype, but the final design of the power supplies including interface changes to account for other system upgrades is scheduled for 2018, and will be our responsibility. A timeline of the project is as follows (more details are shown in table 4): During 2017 the research and development component of the project will be in the design of the elevated temperature testing apparatus (Burn-in Station) for the power supplies. Such a system is necessary to age the components and identify power supplies with early mortality rates. A temperature analysis of the system has also been performed to determine whether the cooling capabilities of the system are sufficient for running the power supplies in redundant mode, where the current is twice the nominal. This is a fallback mode in case of a power supply failure, the other supply in the drawer can provide power to the full drawer. After determining changes needed from components interfacing to the LVPS such as the ELMB++ a small number of prototype LVPS will be made in 2019 with the new design. These LVPS will be assembled into boxes by NIU and sent to CERN for testing in the vertical slice test. From 2019-2022 we will be producing LVPS in batches testing and burning them in for quality control. This is a huge undertaking and will require a detailed and well-documented plan that a set of undergraduate students will follow for testing the LVPS.

| Task | Personnel | Start Date | End Date |
|---|-----------------|------------|----------|
| New Burn-in Station Design and Fabrication | Hibbard, EA, EE | 01/30/17 | 12/15/17 |
| Prototype | - | 01/29/18 | 08/30/19 |
| Procurement of brick components for prototype V8.2 bricks | Hibbard, EA | 01/29/18 | 05/11/18 |
| Basic Check-out and burn-in (Prototype) | Hibbard, EA | 04/22/19 | 05/24/19 |
| Prototype Review | EE, EA | 05/27/19 | 08/30/19 |
| Integration and testing of vertical slice | Hibbard, EA | 06/03/19 | 07/26/19 |
| Finalize V8.3 pre-production design | EA, EE | 07/29/19 | 08/23/19 |
| LVPS pre-production design review | EA, EE | 08/26/19 | 08/30/19 |
| Pre-production | - | 09/2/19 | 12/30/20 |
| Check-out and burn-in tests on pre-production bricks | EA, undergrads | 07/06/20 | 09/25/20 |
| Ship bricks to CERN | EA | 09/28/20 | 10/02/20 |
| Integration tests at CERN and final review | EA | 10/05/20 | 12/31/20 |
| Production | - | 01/04/21 | 04/08/22 |
| Production PCB Assembly | EE, EA | 04/12/21 | 06/04/21 |
| Check-out and burn-in (16 to start and find any issues) | undergrads | 04/19/21 | 04/30/21 |
| Check-out and burn-in (64 to confirm production) | undergrads | 05/03/21 | 05/28/21 |
| Check-out and burn-in (1000) | undergrads | 05/31/21 | 04/08/22 |

Table 4: The HL-LHC Tile LVPS upgrade schedule.

1.9 International Linear Collider Project (PI: White)

The study of the Higgs sector continues to be a major part of the HEP physics program, and will continue to be for the next 10 - 20 years. The full exploration of this sector will require the combination of the upgraded, high luminosity LHC plus precision results from a lepton collider. The lepton collider that is closest to being realized is the International Linear Collider in Japan. UTA has been engaged, since 2002, in studies for the SiD Detector Concept for the ILC. This work has been an investment in the future of HEP for our group, and is consistent with our approach of having an involvement in each generation of colliders. The SiD Detector Concept was one of two designs validated by the International Detector Advisory Group. Since 2012, White has served as co-Spokesperson for SiD, with co-Spokesperson Marcel Stanitzki(DESY). The SiD design study achieved an important milestone with the production of the Detailed Baseline Design (DBD) [?] together with the TDR for the ILC Accelerator. The DBD is a comprehensive statement of an advanced detector design (see Figure 1.9 (left)) [?] that will enable the study of a full range of TeV scale physics. This includes precision Higgs measurements, a detailed study of top physics, which is closely connected to the phenomenon of electroweak symmetry breaking, and a sensitive probe of new physics, such as that from additional gauge bosons and extra dimensions at high mass scales, in a program that is complimentary to that at the LHC. The reach of the ILC in precision measurements of Higgs couplings is shown in Figure 1.9 (right). The branching ratio for Higgs to invisible decays can be probed at the 1/2% level at the ILC. This is the level at which theoretical studies (Ref. xx) show that significant effects from new physics can be expected. The ILC is thus a discovery machine both through direct searches and precision measurements.




Figure 8: (left) The SiD Detector (right) Higgs couplings projection for ILC and LHC.

Since the writing of the DBD, many further physics and detector optimization studies have been carried out, including the following areas:

- **Pixel tracker option.** Studies of the benefits of a coherent pixel technology implementation for the vertex detector and silicon tracker combination.
- **Hadron calorimeter** New baseline technology choice for the hadron calorimeter: scintillator/steel instead of RPC/steel. UTA's role in the SiD hadron calorimeter is discussed in the next section.
- **Flux return/muon steel** Redesign of the solenoid flux return to reduce the external field and facilitate easier handling and transportation of steel components.
- **Backgrounds** Studies of the effects of pair, $\gamma\gamma$, and beam muon backgrounds on reconstruction of tracks, jets.
- **Multiplicity studies** Evaluation of the required pipeline depth for the KPix chip - used in all systems except the vertex detector.

- **Forward region** Redesign of the LumiCal, BeamCal, and other components to reflect the new common L* for SiD and ILD.
- **SiD Simulation** A complete reworking of the SiD simulation in the DD4HEP framework - with UTA providing the details of the hadron calorimeter.

Recently, SiD decided to form the SiD Consortium to formalize the membership of this pre-collaboration. SiD now has 30 institutions from the U.S.(55%), Europe (40%) and Asia (5%). We have a set of rules for membership, application procedures for potential new members, and an Institutional Board that meets at each major ILC event. As SiD Spokesperson White has the following roles:

- **Overall Leadership.** Responsible for the physics and technical leadership of the SiD Detector Concept, including appointment of sub-group leaders and promotion of SiD within the Linear Collider and High Energy Physics community within the U.S. and worldwide.
- **Head of the SiD Executive Committee.** Responsible for the overall strategy and guidance of the SiD Consortium, and calling and chairing weekly meetings.
- **SiD Representation** External committees and at ILC conferences. SiD Spokespersons are members of, or report to, a number of external committees such as the Executive Committee of the Physics and Detectors section of the Linear Collider Organization.
- **Americas Linear Collider Committee** White also serves as the at-large universities' representative on the Americas Linear Collider Committee.

In their report, P5 described the ILC physics case as "extremely strong", and support for the ILC Project was included in all scenarios. Earlier a HEPAP Facilities Panel (on which White served) concluded that "The ILC accelerator and detectors enable a research program that will address questions of very great scientific importance, and both the accelerator and the detectors are *absolutely central*.". The Project is currently under evaluation by the Japanese MEXT organization. Recently a U.S. DoE - MEXT Working Group was established, and a list of priority areas identified for focus while the project is under evaluation. There is very strong political, industrial, and community support for the ILC in Japan and a decision is expected in the next 2 years. There has been a series of high-level political and industrial visits by Japanese delegations to the U.S. and White has participated as a representative of the U.S. HEP community. ICFA has restructured the Linear Collider Collaboration with a format designed to carry the project through the next period prior to the final construction decision. If the ILC project is to be realized, the involvement of the U.S. is essential - there simply is not the required cryomodule production capacity in the rest of the world. In order to prepare for potential participation, it is therefore critical that the U.S. maintain its long term intellectual leadership in the project by continued investment in Physics and Detector studies.

Hadron Calorimetry for SiD, [PI: White] UTA has for 14 years worked on the development of hadron calorimetry for a future lepton collider. Our graduate and undergraduate students have benefitted significantly from participation in this work, and have made valuable contributions. Future experiments make severe demands on achieving excellent jet energy resolution. Much of the new physics program for the ILC requires high-precision measurements of jet energies and jet-jet invariant masses. Studies have shown that there is a basic requirement of at least $30\%/\sqrt{E}$, at 100 GeV or 3-4% in general for jet energy resolution. This need arises from the requirement for precision Higgs studies, and precision measurements of the masses and other properties of new particles that may be discovered in the next few years at the LHC. A prime physics example of the need for this precision is the reconstruction and separation of hadronic decay modes of W and Z bosons, on an event-by-event basis. Hence there has existed a need for the development of a new type of hadron calorimetry.

The hadron calorimeter is an essential component of the Particle Flow Algorithm (PFA) approach to achieving the required jet energy resolution. The PFA approach has been developed and

extensively used in physics studies by both of the main ILC detector groups, SiD and ILD. It has also been used in LHC jet reconstruction. Most of the calorimeter development work to provide technical solutions to implementing PFA-oriented systems has been carried out by the CALICE Collaboration [?]. White and Felix Sefkow (DESY) were asked to be the authors/main editors of a Reviews of Modern Physics paper on "Experimental tests of particle flow calorimetry". This paper was published [?] in early 2016 and represents a statement and synthesis of developments of PFA calorimetry. As reflected by this paper, there has emerged a choice of technologies in which to implement the hadron calorimeter for SiD. The technology must support individual charged particle tracking through the calorimeter, allow detailed imaging of energy depositions for track-shower association and separation of closeby showers, while providing good energy resolution for the direct measurement of the energies of neutral particles. Highly granular calorimetry, as required for Particle Flow, has recently been adopted by CMS for an upgrade of its forward calorimetry. White proposed [?] a digital hadron calorimeter implementation using Gas Electron Multiplier (GEM) technology [?]. GEM offers a robust solution with sufficient amplification in a double-foil configuration, and a thin active layer. This last requirement is important since a typical lepton collider hadron calorimeter would have about forty layers, with the entire system contained inside superconducting coil – the cost of which must be limited.

Figure 9(a) shows the basic GEM-DHCAL idea introduced by White. The concept of embedding the front-end electronics as part of the active layer has been realized through the development of the KPiX 1024-channel chip by SLAC (with a series of earlier versions also having been tested at UTA). KPiX is specifically designed for the time structure of the ILC beam, and requires a synchronous environment to operate at full efficiency. To implement a full digital hadron calorimeter system would clearly require large area chambers. UTA has developed a series of prototype GEM-DHCAL chambers (see Figure 9(b)) with fine ($1\text{cm} \times 1\text{cm}$) cells. A key characteristic of an active layer suitable for use in a digital hadron calorimeter is a low hit multiplicity per track per layer while maintaining a high hit efficiency. Figure 9(c) shows that our GEM chamber(s) indeed have this desired performance characteristic. The GEM-DHCAL development work has given us very valuable experience with respect to the implementation of the hadron calorimeter for SiD.




Figure 9: (a) GEM Digital Hadron Calorimeter Schematic. (b) large chamber under construction. (c) Multiplicity vs. hit efficiency.

The recently adopted, and current, baseline technology for the SiD hadron calorimeter, for detector simulation and physics studies, is small scintillator tiles with SiPM readout, and steel absorber plates. This approach has been the subject of considerable work by CALICE.

UTA plans to take a leading role within SiD in the development, design, prototyping, and construction of the SiD hadron calorimeter system based on the scintillator/SiPM/steel technology. As a first step, UTA has undertaken the implementation of this technology in the DD4HEP [?]

framework for the SiD simulation. White has been working with two undergraduate students to specify the details of the hadron calorimeter simulation. Figure 10(a) shows the details of the implementation of the active layer of the hadron calorimeter in the SiD simulation. CALICE studies have shown that $3\text{cm} \times 3\text{cm}$ cells yield the required jet energy resolution, and we have used this cell size for SiD. Figure 10(b) shows the first result for the barrel simulation with a single incident charged particle. Once the full hadron calorimeter simulation is complete, we will make extensive comparisons between our simulations and the results of the CALICE beam tests for a variety of incident particles. SiD has chosen a 12-fold design for the hadron calorimeter (see Figure 10(c)). Taking the CALICE active layer design as input, we will develop a 40-layer module design. This will require the construction and testing of a representative size prototype of an active layer, specification of layer assembly techniques (possibly automated), and mechanical module prototypes.




Figure 10: (a) SiD AHCAL Barrel layer schematic. (b) SiD AHCAL simulation - single pion. (c) SiD AHCAL Barrel 12-sided structure.

All this activity will be a precursor to the SiD Technical Design Report, the timescale for which depends on the overall progress on the ILC project. We anticipate 2-3 years to complete the TDR, once resources are available.

SiD Timetable of Activities A decision on the ILC Project is expected in 2018. This timetable reflects this anticipated schedule. The activities below reflect White's role as SiD Spokesperson and the planned work at UTA on the SiD hadron calorimeter.

2017

- Guide and pursue the optimization studies for all SiD subsystems
- Initiate and guide further physics studies in response to new LHC results
- Expand the SiD Consortium, particularly in Europe and Asia
- Work with the U.S.- Japan Caucus and the ALCC to prepare the case for U.S. support of the ILC
- Represent SiD at national and international Linear Collider conferences
- Complete the full simulation of the SiD baseline hadron calorimeter, and verify performance vs. CALICE results
- Develop initial hadron calorimeter barrel and endcap module concepts

2018

- Continue the first five tasks from 2017
- Work with Japanese colleagues to provide input from SiD to inform the anticipated ILC Project decision by MEXT

- Begin the framework for the SiD Technical Design Report
- Develop production and assembly procedures for SiD hadron calorimeter active layers, and module integration

2019 Subject to the ILC Project proceeding:

- Continue the general tasks from 2017/18
- Prepare the SiD Experiment initial submission in response to anticipated call for expressions of interest/proposals
- Establish initial areas of responsibilities for SiD member institutes in the transition from the SiD Consortium to the SiD Collaboration
- Begin full engineering design and prototyping for SiD hadron calorimeter modules

Part III

Research at the Intensity Frontier

Executive Summary

The Intensity Frontier (I.F.) group of the University of Texas at Arlington started in 2014 with 0.5 FTE each of PI Jaehoon Yu and of PI Amir Farbin aiming for a balanced program between US-based and non-US based experiments. In order to continue to build up a strong I.F. program, the group recently hired a full time I.F. junior faculty, Dr. Jonathan Asaadi. The overall strength of the group has grown double to two full PI's with the successful transition of Yu to full time I.F. despite Farbin transitioning back to the Energy Frontier (E.F.).

Beyond the addition of a new faculty member, the UTA I.F. group has made significant contributions to the Liquid Argon In A Test beams (LArIAT), Long Baseline Neutrino Experiment (LBNE)/Deep Underground Neutrino Experiment (DUNE) and MiniBooNE experiments. Farbin played a key role of deputy computing coordinator for the DUNE project and Yu has served as a co-convener for the LBNE R&D Coordination group. Yu's role has evolved to a co-convenership of the DUNE Beyond the Standard Model (BSM) physics group in September 2015. UTA also hosted the first off-Fermilab-site DUNE collaboration meeting in January, 2016, in which over 150 collaborators participated.

Farbin and Yu have been supervising a student (Sepideh Shahsavarani) who has been contributing to the MiniBooNE beam-dump dark matter search with first results expected in fall 2016. Asaadi has continued his roles on MicroBooNE, serving as the convener of the Astro-Particle and Exotics group through August 2016 and a lead TPC-Expert for the experiment. Asaadi has also played a leadership role on LArIAT serving as the analysis coordinator in 2015 and 2016 leading to the first measurement of the π^- -Argon cross-section and now will transition to serve as a co-spokesperson (starting fall 2016). The UTA group has also joined the SBND and the ICARUS experiment (with Yu as institutional board member) and has been contributing to the refurbishment of the light detection system with the hire of a new post-doctoral researcher (an existing ICARUS collaborator), Dr. Andrea Falcone.

In this document, we propose to complete the ongoing efforts on MiniBooNE (Yu) and LArIAT (Asaadi, Yu) and continue/add significant contributions to SBND (Asaadi, Yu), MicroBooNE (Asaadi), ICARUS (Asaadi, Yu) and DUNE (Asaadi, Yu) with vital contributions to the two proto-DUNE projects. These experiments are carefully selected to leverage our group's growing technical and analysis expertise on liquid argon (LAr) time projection chambers (TPC). Moreover, in order to accomplish the work laid out in this proposal, the UTA group will leverage Asaadi's start-up to provide an additional (beyond what is requested here) post-doctoral researcher (Dr. Andrea Falcone) in year one and two of this proposal. Dr. Falcone will play a leadership role on the ICARUS and SBND experiments during construction, installation and commissioning, moving to Fermilab in time with the transfer of the refurbished ICARUS detector.

The work on MiniBooNE is limited to data analysis using the beam dump data taken in 2014 for a low mass dark matter search. This is anticipated to complete in the early stage of this renewal proposal period with the graduation of Sepideh Shahsavarani. This student will stay on the I.F. program for another two years till she completes her Ph.D. under the joint advisement of Farbin, Yu, and Asaadi and is expected to contribute to the other I.F. efforts of the group. Asaadi will continue to play a leadership role on LArIAT together with Yu on data analysis and operations with the expected winding down of the project in the next 1-2 years.

UTA Strategic Plan

The UTA group will have contributions and responsibilities across the Fermilab short-baseline neutrino (SBN) and the long-baseline neutrino (LBN) programs. The primary strategic goal of the group throughout the period of this proposal is to contribute to the success of the construction and execution of these experiments, focusing on LArTPC technology. We aim to position ourselves to make major contributions to the DUNE project by playing leadership roles in the design and construction of the single phase (SP) and dual phase (DP) prototype detectors, protoDUNE's at CERN neutrino platform. This work is intended to allow UTA to be a world leading institution for the construction and data analysis of the DUNE detector. Synergistic with this goal, our active participation in the construction, commissioning and operation of the SBN experiments - MicroBooNE, SBND, and ICARUS - allows us to enhance our expertise in LArTPC technology, produce valuable physics results, and gain experience in near/far oscillation analyses utilizing the data from these detectors.

Among the various tasks described below, we consider protoDUNE to be under the most restrictive timeline and deserve immediate attention early in the period of this renewal proposal. Since the availability of charged particle beams at CERN's neutrino platform is dictated by the CERN accelerator complex upgrade schedule, having the protoDUNE detectors ready before the start of LHC heavy ion run in October 2018 is critical. Moreover, the protoDUNE projects are on the critical path for the success of the U.S. flagship experiment, DUNE. The SBN experiments are also seen by the UTA group as a high priority, but given that Fermilab will continue providing beams to these experiments through at least 2021, they are not on as critical of a schedule driven path as protoDUNE's.




Figure 11: Timeline of the proposed projects

Figure 11 summarizes the timeline for the projects to be executed during the funding period of this proposal based on the latest information at the time of writing this proposal. These projects

are described in greater detail in subsequent sections. The overarching strategic goal is to have Asaadi focus on the short and mid-term experiments such as MicroBooNE and SBND with a smaller portion of his time on ICARUS and DUNE initially and then ramping it up over time. Meanwhile, Yu will focus on the long and mid-term experiments such as DUNE and ICARUS with a gradually increasing portion over time on the whole SBN program. This enable our group to maintain a constant level of contribution from both the PI's and the post-doctoral and graduate researchers on all four experiments.

Tables 5 summarizes the proposed projects and associated lead P.I. who will take a lead role on this proposal and where the effort will be located. The division of projects and P.I. effort is designed to find synergy between the timeline and when each effort is to take place. This will allow Asaadi and Yu to maximize their impact across the neutrino program.

IF Summary of Proposed Work

| Experiment | Project | Location | Lead PI |
|------------|--|----------|---------|
| DUNE | protoDUNE SP-APA QA/QC and installation | UTA/CERN | Asaadi |
| | BSM Physics | UTA | Yu |
| | protoDUNE DP-FC Construction | UTA/CERN | Yu |
| SBND | Cold Electronics TPC Test-stand | UTA/FNAL | Asaadi |
| | Detector Construction, Installation, and Commissioning | FNAL | Yu |
| | Cross-Section Data Analysis | UTA/FNAL | Asaadi |
| MicroBooNE | TPC Detector Expert | UTA/FNAL | Asaadi |
| | Detector Operations | UTA/FNAL | Asaadi |
| | Cross-Section Data Analysis | UTA/FNAL | Asaadi |
| ICARUS | Detector Installation and Commissioning | FNAL | Asaadi |
| | NuMI Off-Axis Cross-Sections | UTA/FNAL | Yu |
| MiniBooNE | Beam Dump Dark Matter Search | UTA/FNAL | Yu |

Table 5: Overview of the UTA projects across the Intensity Frontier

Table 6 summarizes the projects and associated personnel resources that will execute the research described in greater detail in the subsequent sections. The requested post-doctoral efforts are divided evenly between DUNE and the SBN programs with an additional post-doctoral researcher coming from Asaadi's start-up funds. The graduate students will contribute to hardware on both DUNE and the SBN programs, but will focus on the SBN for their thesis analyses.

MicroBooNE (Asaadi): Asaadi has been an essential member of the construction, commissioning, operation and data analysis on MicroBooNE. With Asaadi being a young-tenure track faculty member, it is essential for him to be able to produce physics results in the first years of the proposal. Given this, Asaadi will focus on operations and data analysis, presented in greater detail in Section 2.5.

- **Detector Operations and TPC Detector Expert:** Section 2.6
- **Cross-section Data Analysis:** Section 2.7

SBND Tasks (Asaadi, Yu): UTA's effort on SBND is presented in greater detail later. Asaadi will play the lead role as institutional representative with Yu in discussions with the SBND management for joining the collaboration prior to the start of this funding period.

- **Cold Electronics TPC Test-stand (Asaadi):** Section 2.2
- **Detector Construction, Installation, and Commissioning (Yu):** Section 2.3

Summary of PI, Postdoc, and Graduate Personal

| Personnel | Associated Task | Years Supported | Source of Support |
|---|--|----------------------------|--------------------------------------|
| Postdoc 1 (Animesh Chatterjee) | protoDUNE SP/DP SBND Operations and Data Analysis ICARUS Operations and Data Analysis | 2017 - 2020 | UTA Base Grant |
| Posdoc 2 (TBN) | protoDUNE SP/DP MicroBooNE Operations and Data Analysis SBND Cold Electronics Test Stand | 2017 - 2020 | UTA Base Grant |
| Postdoc* (Andrea Falcone) | ICARUS/SBND Installation and Commissioning MicroBooNE Operations and Data Analysis | 2017 - 2019 | UTA Start-up funds |
| Graduate Student 1 (Garrett Brown) | protoDUNE SP/DP SBND/ICARUS Operations and Data Analysis | 2017 - 2020 | UTA Base Grant |
| Graduate Student 2a (Sepideh Shahsavarani) | MiniBooNE Data Analysis protoDUNE DP | 2017 - 2019 | UTA Base Grant |
| Graduate Student 2b * (Zack Williams) | SBND Cold Electronics Test Stand, SBND Construction, Installation, and Commissioning MicroBooNE Operations and Data Analysis | 2017 - 2019 2019 - 2020 | UTA Start-up funds UTA Base Grant |
| Prof. Jonathan Asaadi | SBN/DUNE | 2017 - 2020 | UTA Base Grant |
| Prof. Jae Yu | DUNE/SBN | 2017 - 2020 | UTA Base Grant |

Table 6: Table summarizing the personnel working on the project described in this proposal. Note: Personnel marked with “*” denote that their effort is supported for some phase of the project utilizing Asaadi’s start-up funds. This is done to maximally leverage the UTA group across both DUNE and the SBN program.

- **Cross-Section Data Analysis (Asaadi):** Section 2.4

ICARUS (Asaadi, Yu): UTA group has joined the experiment as a member with Yu as the institutional board representative as one of the handful of U.S. groups to participate in it. Our efforts have been augmented by utilizing Asaadi’s start-up funds to support a post-doctoral researcher to help lead the refurbishment and integration of ICARUS, described in more detail in Section 2.8.

- **Installation, Commissioning, and Detector Operations (Asaadi):** Section 2.9
- **NuMI Off-Axis Cross-Sections & Dark Matter Searches (Yu):** Section 2.10

DUNE (Asaadi, Yu): The UTA group will be contributing to both the protoDUNE Dual Phase (DP) and Single Phase (SP) detectors. Asaadi and Yu expect to play major roles in various detector component construction, installation, and commissioning as well as data taking during the first two years of this proposal. This work will lay the strong foundation to contributing to the DUNE experiment and is described in greater detail in Section 3.1. Along with this, UTA hopes to play a critical role in the physics profile for the DUNE experiment through contributions to the physics working groups.

- **ProtoDUNE Single Phase-Anode Plane Assembly QA/QC (Asaadi):** Section 3.2
- **ProtoDUNE Dual Phase-Field Cage Construction (Yu):** Section 3.3
- **DUNE Beyond the Standard Model Physics (Yu):** Section 3.4
- **LBNF Beam Simulation (Yu):** Section 3.5

With this strategic plan in place, the activities proposed aim to ensure synergy between the SBN and LBN efforts and to optimize our use of resources. What follows is a broad introduction to the compelling physics which motivates this research program as well as a more detailed sketch of the program of work which is intended to be executed in the intensity frontier.

Physics Introduction

The discovery that neutrinos undergo oscillation in their flavor, and thus are massive particles, serves as one of the first pieces of evidence for physics beyond the Standard Model (SM) of particle physics. The prevailing description of neutrino oscillations provided by the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix characterizes the flavor change as a result that the neutrino flavor eigenstates (ν_e, ν_μ, ν_τ) are a linear combination of the neutrino mass eigenstates (ν_1, ν_2, ν_3). The rotation from the mass eigenstates to the flavor eigenstates is governed by three angles $\theta_{i,j}$, where i and j correspond to the mass eigenstates with $i < j$, and a phase δ which determines magnitude of charge-parity (CP) violation within the neutrino sector. The exact ordering of the neutrino mass states, known as the mass hierarchy, as well as the size of the CP-violating phase δ are, as yet, unknown. These quantities remain one of the last major pieces of the Standard Model of particle physics and offer the opportunity to answer such fundamental questions as:

- 1) **What is the origin of the matter/antimatter asymmetry in the universe?**
- 2) **Do we understand the fundamental symmetries of the universe?**
- 3) **Is the three-flavor paradigm of the Standard Model for neutrino oscillation the accurate description for neutrino interactions?**

Into this experimental landscape, there exists a set of series of experimental measurements which suggest that the three-flavor paradigm of neutrino oscillations is incomplete. Two general classes of anomalous observations may point to additional physics beyond SM in the neutrino sector.

- **The disappearance signal in low energy electron anti-neutrinos from reactor neutrino experiments [?] (“Reactor Neutrino Anomaly”) and Mega-Curie radioactive electron neutrino sources in Gallium [?,?] (“Gallium Anomaly”)**
- **The electron-like excess from muon neutrino (and anti-neutrino) particle accelerators (“LSND/MiniBooNE Anomaly”) [?,?]**

Neither of these anomalies can be explained by the standard three-flavor oscillations of SM and may hint at the existence of additional neutrino states with larger mass difference ($\Delta m_{new}^2 \geq 0.1\text{eV}^2$) which participate in the mixing of the flavour states (referred to as “sterile neutrinos”). Definitive evidence of the existence of new neutrino states would be a revolutionary discovery with broad implications for both particle physics and cosmology. Moreover, in order for future accelerator based neutrino experiments to disentangle the mass hierarchy and search for CP-violation, the oscillation framework must be concretely known and precisely measured.

Additionally, recent theoretical interest has been paid to the possibility of studying the models for Low-mass Dark Matter (LDM) production at low-energy, fixed-target experiments (see Refs. [?, ?, ?, ?]). High flux neutrino beam experiments, such as those used to address these questions in neutrino physics, have potential to provide coverage of LDM + mediator parameter space which currently are not covered by either direct detection or collider experiments. Upon striking the target, the proton beam can produce the dark photons either directly through $pp(pn) \rightarrow \mathbf{V}$ or

indirectly through the production of a π^0 or a η meson which then promptly decays into two SM photons of which one couples to the heavy dark photon. For the case where $m_V > 2m_{DM}$, the dark photons will quickly decay into a pair of DM particles. These relativistic DM particles from the beam will travel along the beamline (similar to the neutrinos) and could be detected in a sufficiently sensitive neutrino detector and provide evidence of the existence of LDM. These particles can then be detected through neutral-current like interactions either with electrons or nucleons in the detector.

Liquid Argon Time Projection Chambers (LArTPCs) offer fine-grain tracking as well as powerful calorimetry and particle identification capabilities making them ideal detectors for studying neutrino-nuclei interactions. The combination of millimeter scale tracking capabilities, outstanding calorimetry through a fully active/sampling detector, and powerful particle identification made by combining the ionization along the particle trajectory (dE/dX) and the topological information, have made LArTPC's the premier neutrino detector technology choice for the future and provide an excellent detector to search for non-standard interactions in the existing beamlines. For these reasons, this detector technology has been chosen for both the study of neutrino oscillations over relatively short baselines (< 1 km) and long baselines (> 1000 km).

PI Summary: Jaehoon Yu

Accomplishments

Since I have been in transition in the past three years, my accomplishments Energy Frontier (EF) are also summarized here along with those in I.F. **Accomplishments in Energy Frontier Program**

- Graduated a Ph.D. student, Dr. Heeyeun Kim in 2015
- Expect to graduate another Ph.D. student, Mr. Last Feremenga by the end of the current funding cycle
- Served on various Higgs editorial boards as a member
- Contributed to publication of a couple of $H \rightarrow WW$ papers as an author
- Contributed to publication of a couple $H^+ \rightarrow \tau + \nu$ papers as an author
- Contributed to publication of exclusive production of Higgs paper as an author
- Served as an ATLAS leader shifter in 2016 for 13 TeV data taking
- Serving as a member of the search committee for the deputy US ATLAS Physics Support Manager

Accomplishments in Intensity Frontier Program

- Co-lead Detector R&D Coordination group of LBNE
- Contributed to the transition from LBNE to DUNE working on collaboration governance
- Organized and hosted the first off-Fermilab-site DUNE collaboration meeting at UTA
- Co-leading the Beyond the Standard Model physics group of DUNE
- Contributed to LBNE Physics Book
- Contributed to various beam systematics and optimization studies
- Contributed to LBNE/DUNE hadron monitor development
- Contributed to WA105 $3 \times 1 \times 1$ m³ prototype construction during the sabbatical stay in Sept. 2015 - May 2016
- Lead UTA group to join WA104, ICARUS experiment
- Supervised numerous undergraduate students in their contribution to the LBNF beam line systematics and optimization studies

- Supervised postdoctoral fellow Dr. Animesh Chatterjee for his contributions to DUNE, LArIAT, and MiniBooNE

Milestones

DUNE BSM Physics

- Complete Initial DUNE BSM Physics list document: Late 2016
- Complete Systematic Sensitivity Studies: Late 2017/Early 2018
- Refine Sensitivity Studies: Mid 2018
- Contribute to writing the DUNE Technical Design Report (TDR): Mid 2018

protoDUNE Experiment

- Complete DP Field Cage (FC) Design and establish QA/QC Plan: Late 2016/Early 2017
- Complete FC submodule assembly, functional prototype testing and shipping: Mid 2017
- Detector Installation, Commissioning, and Operations: Late 2017 - Mid 2018
- Data Taking and Analysis: Mid 2018 - 2019
- Contribute to DUNE TDR: Mid 2018

SBND Experiment

- Join the experiment: Early 2017
- Detector Construction and Installation: Early 2018 - Mid/late 2018
- Commissioning and Operation: Starting Mid/late 2018
- Data Analysis: 2019

ICARUS Experiment

- Installation and Commissioning: Early 2017
- Data Taking and Expert Training: Early 2018
- NuMI Off-Axis Cross-Sections & Dark Matter Searches: Early 2019
 - Sensitivity Study: Late 2017
 - Preliminary Data Analysis for Conferences: Late 2018
 - Publication Result: Late 2019

Plans

During this funding cycle, I will be leading the DUNE program, protoDUNE DP and the ICARUS detector during the construction, installation and commissioning of them at various stages. In particular, I am responsible for assembly, functional testing and installation of the protoDUNE DP field cage (FC). The submodules of the protoDUNE DP FC will be built and tested at UTA and be shipped to CERN for installation by mid 2017. I plan on being stationed at CERN in fall 2017 together with a postdoctoral fellow and a graduate student for installation of them. My schedule will be coordinated with P.I. Asaadi to maintain a constant level of P.I. effort on SBN and protoDUNE and to maximize the impact of our personnel resources. Once I return to the U.S. after the stay at CERN in 2018, I will be focusing on SBN program to ensure the success of these programs and our group's role in SBND and ICARUS experiments on their construction, installation, commissioning and operation. This plan will enable our group to maintain respectful number of personnel on both DUNE and SBN programs and enhance necessary expertise to play crucial roles in construction and eventual physics data analysis of DUNE.

The physics I will focus on in this proposal is the Low-mass Dark Matter searches in DUNE and SBN experiments, along with the bread and butter cross-section analyses across the SBN jointly

with P.I. Asaadi. In particular, based on the production angle of LDM in high intensity proton beams, it is also feasible to perform a parasitic experiment in ICARUS off of the NuMI neutrino beams. A systematic sensitivity study for DUNE LDM searches will be completed for an inclusion into the DUNE TDR. I will lead the DUNE BSM subgroup to complete systematic and realistic sensitivity studies of various topics within the group to contribute to the DUNE TDR.

1.9.1 Search for Low-mass Dark matter

As described in Section ??, high power proton beams necessary for high flux neutrino beams enable experiments to search for BSM phenomena. The particular interest of our group is the potential for discovering LDM particles in high precision detectors. The LDM particles produced in the target through the decay of dark photons can be detected through neutrino neutral-current like interactions either with electrons or nucleons in the detector. Since the signature of DM events looks just like those of the neutrinos, the neutrino beam provides the major source of background for the LDM signal.

Several ways have been proposed to suppress neutrino backgrounds by using the unique characteristics of the DM in the beam. Since DM will travel much slower than the neutrinos due to its much higher masses, the arrival time of the LDM signal in the near detector can be used to distinguish LDM from neutrinos. In addition, since the electrons struck by LDM will be much more forward direction than those from neutrino interactions, the angle of these electrons may be used to reduce backgrounds, taking advantage of fine angular resolution DUNE near detector can provide. Finally, a special run can be devised to turn off the focusing horn to significantly reduce the charged particle flux that will produce neutrinos. If DUNE near detector were LArTPC, since the entire detector volume will be active, the effective number of DM events detected will be much higher with the detector of the same mass. A major goal of this study is to include the sensitivity study into the DUNE Technical Design Report to be released in mid 2019.

In addition, given the 600t LAr fiducial mass of ICARUS detector and its location with respect to NuMI target, it is also feasible to search for LDM in it off of NuMI neutrino beams. Since ICARUS is single phase LArTPC, such a study in ICARUS will enable a more realistic and systematic study in preparation for searches in DUNE, if not outright discovering LDM. This idea requires more thorough study in its feasibilities.

PI Summary: Jonathan Asaadi

Accomplishments

ArgoNeuT Experiment

- Lead Analyzer on the “Measurement of ν_μ and $\bar{\nu}_\mu$ Neutral Current $\pi^0 \rightarrow \gamma\gamma$ Production in the ArgoNeuT Detector” : First measurement of NC- π^0 production on argon
- Contributer to “First Observation of Low Energy Electron Neutrinos in a Liquid Argon Time Projection Chamber”: Paper in preparation

MicroBooNE Experiment

- TPC Commissioning Coordinator: Developed and executed the commissioning plan for the MicroBooNE TPC
- Lead TPC Expert: Develop the expert documentation for the TPC and coordinate the expert shifts

- **Astro-Particle and Exotics Working Group Convener:** Lead one of the MicroBooNE working groups and contribute to a preliminary proton decay background analysis [].

LArIAT Experiment

- **LArIAT Co-Spokesperson:** Beginning September 2016
- **LArIAT Analysis Coordinator:** Lead the first measurement of the inclusive π^- -Argon cross-section shown in Figure 12 (paper in preparation).




Figure 12: (Left) First inclusive π^- -Argon cross-section.

University of Texas Arlington

- **Establishment of Cryogenic Laboratory:** Building a facility capable of purifying and recirculating liquid argon for use in detector R&D and testing.

Milestones

protoDUNE Experiment

- **APA QA/QC Documentation and Testing:** Early 2017
- **Detector Installation, Commissioning, and Operations:** Late 2017-2018
- **Data Taking and Analysis:** Mid 2018 - 2019

SBND Experiment

- **Cold Electronics Test Stand:** Delivery to FNAL Summer 2017
- **Detector Construction:** Mid 2017 - Mid 2018
- **Commissioning and Data Taking:** Starting Mid 2018
- **Cross-Section Analysis:** Preliminary studies beginning late 2019

MicroBooNE Experiment

- **Charged Current Coherent π Analysis:** 2017 - 2019
 - Sensitivity Study: Summer 2017
 - Preliminary Data Analysis for Conferences: Mid 2018
 - Publication Result: Early 2019
- **Data Taking and Expert Training:** 2017 through 2020

ICARUS Experiment

- **Installation and Commissioning:** Early 2017
- **Data Taking and Expert Training:** Mid 2017 - 2020
- **NuMI Off-Axis Cross-Sections**
 - Sensitivity Study: Late 2017

- Preliminary Data Analysis for Conferences: Late 2018
- Publication Result: Late 2019

Plans

During this funding cycle, I plan on leading UTA on the SBN program during the construction, installation and commissioning of the various SBN detectors. In parallel, I intend to contribute to the single phase protoDUNE program by being stationed at CERN during the crucial months in the leadup to first beam. By coordinating with P.I. Yu, the UTA I.F. group can maintain a constant P.I. effort on SBN and protoDUNE and strategically place our graduate students and postdoctoral researchers where the need is greatest during each of the three years. Moreover, by leveraging my start-up I intend to have one post-doc beyond what is requested in this funding to be stationed full time at FNAL during the ramp-up phase of the SBN. This will also help in the training of new detector experts and smooth the transition from commissioning to operations by having a single person present throughout. In tandem, a new remote operation station at UTA will allow researchers to contribute to data taking and expert shifts while still stationed at UTA.

The physics I will focus on during the years of this proposal will be the cross-section analyses across the SBN, as outlined in the subsequent sections. This selection both plays to my experience from ArgoNeuT and LArIAT, where I have lead two cross-section analyses which have been the first of their kind, as well as allows me to produce publication quality results during my early years as an assistant professor. Understanding of the cross-sections is both crucial for the LArTPC detector technology by providing a “standard candle” for reconstruction and identification of events, but also plays a major role in understanding the systematics associated with the flagship oscillation analyses to be performed.

2 The Fermilab Short-Baseline Neutrino Program

The conclusive redress of the experimental hints of sterile neutrinos becomes high priority for the field of neutrino physics. The Fermilab Short-Baseline Neutrino (SBN) program offers the unique opportunity to definitely address the “LSND/MiniBooNE” anomaly through the utilization of three LArTPCs detectors and well characterized Booster Neutrino Beam (BNB). The SBN program offers a rich physics program with the ability to perform the most sensitive search to date for the existence of sterile neutrinos at the eV mass-scale. The Short-Baseline Near Detector (SBND) will be a new 112 ton LArTPC and serve as the near detector to the SBN program located 110 meters downstream of the BNB target. SBND will measure the un-oscillated neutrino flux from the BNB and enable searches in both the neutrino appearance and disappearance channels. The MicroBooNE detector is an 89 ton active mass LArTPC located 470 meters downstream of the BNB target (just in front of the MiniBooNE experiment). MicroBooNE serves as the pioneer LArTPC experiment on the BNB and will lay the groundwork for the oscillation analysis. The far detector will utilize the upgraded ICARUS-T600 experiment, previously installed and operated at the Gran Sasso Laboratory, and will be located in a new building 600 meters from the BNB target. ICARUS’s large detector mass provides the SBN program with the experimental sensitivity to definitively search for the existence of eV mass-scale sterile neutrinos.

2.1 Short-Baseline Near Detector (SBND) (PI: Asaadi, Yu)

The SBND experiment is designed to build upon the many years of LArTPC R&D and serve as a test-bed for the future long baseline neutrino experiment. SBND's design is to construct a membrane cryostat in a new experiment hall located 110 meters from the BNB target. The cryostat will house the full TPC consisting of one central cathode plane assembly (CPA) and four anode plane assemblies (APAs) which will have three wire planes with three millimetre spacing (similar to the ICARUS design) and the first two induction planes oriented at $\pm 30^\circ$ to the beam axis and the final plane oriented vertically. SBND will be a $5.0\text{ m} \times 4.0\text{ m} \times 4.0\text{ m}$ ($l \times w \times h$) TPC with 112 tons of active volume. SBND will also have a light detection system based on a hybrid of the ICARUS cryogenic PMT's and the proposed DUNE light-guide with silicon photomultiplier (SiPMs) on the end. This light detection system will be embedded behind the APA structure on both sides of the TPC.

One new unique aspect of the SBND detector will be the inclusion of the entire front end readout chain being moved into the liquid argon. The front end electronics are composed of 16-channel analogue front end ASIC which provides amplification and shaping, a 16 channel analogue to digital converter ASIC which provides digitization, buffering, and multiplexing as well as a cold FPGA which provides second multiplexing and voltage regulation. This technical improvement in readout electronics will provide improved signal-to-noise as well as allow for the development of an efficient zero-suppression scheme implemented in the FPGA to greatly reduce the total data volume. Many bench tests of the readout electronics have been performed and show excellent performance. The full integration test with an operating TPC, however, has not been successfully performed (given the many problems seen by the DUNE 35ton prototype) and is an absolutely necessary service task the UTA group is planning to spearhead.

2.2 Cold Electronics Teststand

Figure 13 shows a schematic of what this test-stand would look like utilizing the "Blanche" cryostat currently installed at the Proton Assembly Building (PAB) at Fermilab. This cryostat is engineered to have delivered purified liquid argon into the cryostat as well as circulate, re-condense, and purify boil-off argon. A duplicate of this cryostat is currently being built and will be delivered in late 2016 to UTA. This cryostat will work in conjunction with the liquid argon purification system currently in operation at UTA. This system is built using start-up funds from PI Asaadi and will allow UTA to play an important role in the ability to do detector R&D in the coming years.

Inside this cryostat, a small scale TPC equipped with prototype cold readout electronics from SBND installed along side a pair of light guide bars can be deployed and readout through a 14" inch cold signal feedthrough as designed for the SBND detector. External to the cryostat, scintillator paddles can be positioned to act as an external trigger. Since the cryostat is a copy of an existing one at FNAL, upon its completion the top flange and TPC detector will be shipped to FNAL for longer term use for SBND (and potentially other future LArTPCs). Additional material costs, such as the electronics, power supplies, and cabling are expected to be provided by SBND project funds and Prof. Asaadi's start-up funds.

This test-stand can allow for a robust set of tests for the integration of many new readout components prior to their deployment in the experiment. This critical step was not taken during the MicroBooNE assembly and as a result a number of problems with the readout electronics were not detected in advance of attempting to commission the detector. These problems included the bias voltage line for the TPC wires behaving in an unexpected way and causing pick-up noise to be seen on the electronics, cross-talk between the light detection system and the TPC readout, and the




Figure 13: Conceptual design of the SBND cold electronics vertical slice test-stand. Integration of both cold or warm electronics, light collection system, cosmic ray paddles, as well as warm interface electronics allows for complete testing of the entire readout system prior to deployment in the experiment and a platform for DAQ debugging outside the actual the experiment. The current design has 768 channels (256 collection, 512 induction) utilizing six SBND designed motherboards.

incorrect configuration of electronics settings because of software bugs. While many of these issues were able to be solved during the commissioning phase, they slowed the progress of transitioning to data taking and caused unnecessary harm to the experiment. Moreover, this test-stand will provide a platform for testing and debugging of the DAQ software and readout electronics configuration without interrupting the operation of the SBND experiment.

The postdoctoral researcher and graduate student supported by this work will be developing the readout software into a common DAQ software package used by other LArTPC based experiments known as the artDAQ framework. This common platform ensures that the work done by those supported in this proposal can have a greater impact on future planned LArTPCs as well as allowing them to benefit from the work that has already been done by others.

Finally, such a test-stand can provide an R&D platform for long term testing of future readout components as well as software development for online triggers and zero-suppression schemes without risking downtime on operating neutrino detectors. The trigger schemes envisioned include utilizing multi-core graphical processing units to do online TPC based triggering for rare search events such as proton decay and supernova neutrino triggering. Moreover, by providing a platform for the development of LArTPC's DAQ systems into a common platform such as artDAQ, a greater push to the integration of the data, simulation, and analysis into one common software platform can be accomplished. The events processed utilizing the artDAQ software are immediately readable by the common liquid argon software framework known as LArSoft. Thus, working on this system and the associated neutrino detectors DAQ help promote the use of a common software framework.

2.3 Construction, Installation, and Commissioning

The UT Arlington group is positioned to play a major role in the construction and commissioning of the SBND detector and data acquisition system. Given the groups experience from MicroBooNE and LArIAT (where Asaadi played a lead role of TPC expert during construction, commissioning, and operations for both experiments) as well as the experience of the post-doctoral researchers already with the group (Falcone from ICARUS and Chatterjee from LArIAT) UTA hopes to offer hands on

leadership during the construction phase.

During the construction phase for the TPC (foreseen in mid 2017 through mid 2018), Falcone is expected to be in residence at FNAL and will be spending 50% of his time on SBND. During that same time, a to-be-named post-doc will join him at FNAL with a focus on SBND TPC construction. Having these two present will allow for a rapid ramp-up on the project. At the same time, the Cold-Electronics test-stand is expected to be moved from UTA to FNAL for operations. The graduate student Zach Williams is expected to go with this project in the summer of 2017 and be in residence at FNAL. With his time already spent on this project, continuing to contribute to the DAQ development and aiding in the construction and installation of the cold electronics on the TPC is a natural fit. PI Asaadi is expecting to spend a significant portion of his time at FNAL during the second half of 2017 to help oversee these activities and contribute to the construction and phase.

Following the construction and installation phase, one post-doc and one graduate student are expected to stay with the SBND experiment a significant portion of their time to play a role as detector experts during the commissioning and initial data taking. The aim here is to share their expertise across the SBN program, but to have a reliable source of experts in residence at FNAL for SBND. While stationed at UTA, graduate and undergraduate students will have the opportunity to take remote shifts on this experiment utilizing UTA's remote shift station currently residing in P.I. Asaadi's cryogenic lab. This remote station has already been used to take shifts on the LArIAT experiment and is currently being commissioned to be able to take remote shifts for the MicroBooNE experiment. This station should be online well in advance of SBND data taking and offer the ability for researchers to contribute both on-site and off-site.

2.4 SBND Data Analysis

SBND will provide important physics measurements during its early operations in addition to providing an overall flux normalization to the key SBN oscillation analysis. Critically, SBND will collect very quickly statistics to confirm the nature of the MiniBooNE excess as measured by MicroBooNE. If MicroBooNE were to confirm the MiniBooNE excess as originating from electron-like sources, SBND could quickly measure if there is an oscillation component to the electron-like signal by measuring the rate as seen in the near detector. Conversely, if MicroBooNE were to determine the MiniBooNE excess as originating from photon-like sources, SBND can cross-check if the source is an unaccounted for beam like background or coming from cosmogenic like backgrounds. Regardless of the outcome, SBND will play a critical role in quickly collecting high statistics data as the near detector to the SBN program.

SBND will also provide critical neutrino cross-section measurements at a statistical precision unprecedented by any other LArTPC. SBND will collect approximately two million neutrino interactions per 2.2×10^{20} protons on target (roughly one year of running). With 1.5 million ν_μ charged current interactions and 12,000 ν_e charged current interactions in one year. With such statistics, many precision cross-section measurements (e.g. double differential) become possible and improvements on first and second generation analyses from MicroBooNE can be explored in the first year of data taking.

Furthermore, by collecting approximately 100,000 $\text{NC}\pi^0$ events per year a full characterization of the leading background cross-section to the long baseline CP-violation analysis can be performed. Figure 14 shows the only neutral current π^0 result on an argon nucleus from the ArgoNeuT collaboration [1]. This result was lead by P.I. Asaadi and thus positions the UTA group to uniquely lead this next generation analysis with a much larger statistics sample. The elimination of this systematic uncertainty in the cross-section will improve the experimental reach of the future planned DUNE experiment as well as benefit the short-baseline ν_e oscillation analysis.




Figure 14: The only existing NC- π^0 measurement on argon from the ArgoNeuT collaboration. P.I. Asaadi was the leading analyzer on this result and intends to leverage the large statistics sample to be collected by SBND to provide a low energy measurement of this cross-section, which is one of the leading systematics in the both the short and long-baseline oscillation results.

2.5 Micro-Booster Neutrino Experiment (MicroBooNE) (PI: Asaadi)

The MicroBooNE experiment serves as the first detector deployed at the SBN facility and represents the next step in LArTPC technology. MicroBooNE is a $10.3\text{ m} \times 2.5\text{ m} \times 2.3\text{ m}$ TPC with 89 tons of active volume. The TPC has three instrumented wire planes with the first two induction planes oriented at $\pm 30^\circ$ to the beam axis and the final plane oriented vertically. Both the pitch and wire spacing is chosen to be 3 mm, which provides superb resolution for imaging interactions inside the detector. Additionally there are 32, 8" cryogenic photomultiplier tubes (PMTs) which provide the t_0 for an interaction by recording the scintillation light produced when the charged particles interact in the argon. In the summer of 2015 MicroBooNE was filled with liquid argon and began commissioning. With the rapid success of the system, MicroBooNE has now transitioned from commissioning to neutrino data taking starting in October of 2015. Figure 15 shows an example of an automatically identified neutrino candidate event collected by utilizing both light and charge information.




Figure 15: (Left) One of the first automatically identified neutrino events utilizing the light and charge readout in MicroBooNE.

One of the most compelling measurements MicroBooNE will make is to confirm or refute the nature of the MiniBooNE low-energy electron neutrino excess. Utilizing the particle identification powers of the LArTPC (specifically the dE/dX discrimination), MicroBooNE will be able to differentiate the electron-like electromagnetic showers from photon-like electromagnetic showers. Moreover, the dominant background in the MiniBooNE analysis, neutral current π^0 production, can be extremely reduced using the powerful imaging techniques of a LArTPC. The analysis techniques developed for the low energy excess search will be developed in the common software framework known as LArSoft. This software framework is common amongst many of the LArTPC experiments, helping ensure that the reconstruction techniques and analysis strategies developed on MicroBooNE will have applicability to future experiments.

MicroBooNE will also be able to measure many high-statistic cross-sections at $E_\nu < 1\text{GeV}$. At this energy range, the impact of various nuclear effects such as final state interactions and short-range nucleon correlation are poorly understood. These nuclear effects can change the classification of neutrino nucleus interaction, and thus change the measured cross-section. The fine grain tracking offered by LArTPCs allows for the classification of neutrino-nucleon interaction in terms of final state particles instead of using simplifications such as the quasi-elastic scattering assumption. Moreover, with a proton threshold measured as low as 21 MeV of kinetic energy [?], these nuclear effects can even be measured with high statistics using neutrinos as a probe. The broader neutrino cross-section community is anticipating how the results measured by MicroBooNE compare to previous measurements.

2.6 MicroBooNE Operations

UT Arlington group will continue to play a major role in the data taking and operations of the MicroBooNE detector. PI Asaadi has served as the TPC commissioning leader and now as the TPC operations expert. Asaadi has only recently stepped down as Astro-Particle and Exotics working group convener, but remains active in this group for the foreseeable future where a natural synergy exists within the UTA group given PI Yu's role as BSM convener on DUNE. MicroBooNE will explore the physics capabilities of LArTPC including classification of low energy events as a background for supernova neutrinos and searching for cosmogenic backgrounds related to proton decay analysis.

One post-doctoral researcher supported by this proposal will part of their time working on the MicroBooNE operations and is expected to be trained to serve as the TPC operations expert. This is in addition to the effort expected to be present from the postdoc supported by startup funds in the first two years of the proposal. In addition to data taking shift requirements, the he/she is also expected to play a role in the MicroBooNE online DAQ/data quality management as training for the future planned work on the SBND DAQ. With MicroBooNE just finishing the commissioning of their continuous readout data stream ("supernova data stream"), UTA hopes to play a role in supporting the analysis and improvement of this system. The graduate student supported by this work is also expected to take shifts on MicroBooNE and play a supporting role on the expert training. While stationed at UTA, graduate and undergraduate students will have the opportunity to take remote shifts on this experiment utilizing UTA's remote shift station currently residing in P.I. Asaadi's cryogenic lab. This remote station has already been used to take shifts on the LArIAT experiment and is currently being commissioned to be able to take remote shifts for the MicroBooNE experiment.

2.7 MicroBooNE Data Analysis

Being a driving force on early neutrino cross-section analysis is a good way to have impact on the physics program at MicroBooNE. The postdoctoral researcher and graduate student are expected to work on neutrino cross-section analysis using the data taken in the first years of running in addition to measurements done once the full SBN program is operational. This data set will provide many first glimpses into the short-baseline analysis. Following up on previous low statistics cross-sections measured by ArgoNeuT is one way which the UTA can have immediate impact and leverage our previous experience where Asaadi played a major role.

An example of one such cross-section measurement that is of immediate interest, shown in Figure 16, is the charged current coherent charged pion production (CC Coh- π). This result is of particular interest because it is an example of a relatively simple topology, but one where theory and experiment do not agree. Previous attempts to measure this cross-section at low energy by the SciBooNE and K2K collaborations have proven unsuccessful. However, the analogous neutral current process (NC Coh- π^0) has been observed at low energy. To further complicate the picture, two higher energy measurements from ArgoNeuT and Minerva both show observation of CC Coh- π although somewhat at odds with various modern neutrino generator predictions. The initial MicroBooNE data set will be a valuable tool in disentangling this and many other cross-section oddities and allow for a better construction of ν -Ar scattering models. Researchers from UTA are expected to take a leading role in the analysis of CC Coh- π data sample in MicroBooNE and begin exploration of the NC Coh- π production. This analysis is expected to build on the work began in ArgoNeuT [] and build on the groups expertise for charge pion identification developed through the work on LArIAT.




Figure 16: Recent results from T2K [] showing the tension which exists from the low-energy and high energy measurements of the charged-current coherent π production. With the SciBooNE and K2K experiments showing no evidence of this process at $E_\nu < 1$ GeV but ArgoNeuT and Minerva both measuring this process at higher energy. Moreover, the recent results from T2K disagree with the current cross-section models.

The tools developed for data analysis and reconstruction in MicroBooNE will have transferability to the other SBN LArTPC experiments through the use of the common software package, LArSoft. The UTA group has developed expertise with this software package (as evidenced by Asaadi contributing to the development and planning of this software package) and will continue to contribute to its development as a tool to perform a synthesized analysis across the SBN.

2.8 ICARUS Experiment (PI: Asaadi, Yu)

The ICARUS-T600 detector is the largest LArTPC experiment ever actualized containing 760 tons of purified liquid argon (476 tons of active mass). Comprised of two 300 ton modules, the T600 detector initially tested in Pavia, Italy in 2001 where one of the two modules was exposed to surface running for a three month period. Extensive system testing was performed before the complete system was transported to the underground Gran Sasso National Laboratories (LNGS). In 2010, the entire T600 detector was brought online at Gran Sasso where it completed a three year neutrino run in the Cern to Gran Sasso (CNGS) neutrino beam corresponding to 8.6×10^{19} protons-on-target. The successful operation of a large LArTPC experiment in an underground facility with > 90% data taking efficiency (collecting ~ 3000 neutrino events) and achieving high argon purity and long argon lifetime represents a major technological milestone for LArTPC's.

In 2014 the ICARUS-T600 detector was decommissioned and transported to CERN to undergo a refurbishment and upgrade in anticipation of its future non-underground operation at Fermilab's SBN program. Each module in the ICARUS detector is comprised of a common cathode and a TPC with dimensions $18.0 \text{ m} \times 1.5 \text{ m} \times 3.2 \text{ m}$ ($l \times w \times h$). The TPC has three instrumented wire planes with the first two induction planes oriented at $\pm 60^\circ$ to the beam axis and the final plane oriented horizontally. Both the pitch and wire spacing is chosen to be 3 mm which provides superb resolution for imaging interactions inside the detector.

The importance of the ICARUS-T600 success to the experimental reach of the SBN program is shown in Figure 17. Plotted is the significance with which an experimental configuration covers the 99% confidence level (C.L.) for the allowed sterile neutrino mixing from the LSND experiment as a function of Δm^2 (the mass difference between the active and sterile neutrinos) for the simplest 3+1 model. The gray bands represent ranges of Δm^2 where LSND reports no allowed regions at 99% C.L. The presence of the ICARUS-T600, by providing a large sensitive mass at the far detector location, is absolutely imperative for the SBN program to achieve a definitive (5σ) coverage of the LSND allowed region.




Figure 17: The experimental sensitivity for $\nu_\mu \rightarrow \nu_e$ oscillations including backgrounds and systematics assuming a nominal three year exposure in the BNB for the SBND and ICARUS experiments and a six year exposure for the MicroBooNE experiment.

For these reasons, the UTA group has already begun contributing to the ICARUS experiment with the continued stationing of the post-doctoral researcher Andrea Falcone at CERN to continue

his contributions to the upgrade of the ICARUS light detection system. The upgraded light detection system is currently being installed with 90-PMTs per TPC providing an estimated 5% photo-cathode coverage. Testing of the PMT's and readout electronics at CERN is shown in Figure 18. The increased coverage will allow for excellent trigger efficiency for neutrino induced events as well as providing cosmogenic background rejection. Falcone is also leading the work to develop the readout electronics for the light detection system and integrating them into the common data acquisition system (artDAQ) used by the other two SBN experiments.



Figure 18: PMT Testing at CERN where UTA postdoctoral researcher Falcone has played a lead role in the design, installation and (in the immediate future) commissioning of this system.

2.9 Installation, and Commissioning

By participating in both the installation, commissioning, and data taking of the SBND and MicroBooNE as well as leveraging on the experience of Dr. Falcone's work on ICARUS, the researchers supported by this proposal will be well positioned to contribute to the installation, commissioning, and first data analysis of the ICARUS LArTPC upon its arrival at Fermilab. Having a robust team of researchers based at Fermilab to provide expertise and support for the ICARUS system will ensure the successful execution of the SBN program.

Falcone is expected to travel with the ICARUS detector when it moves to Fermilab in early 2017 and play a key role in the commissioning and data acquisition of the light system. At the same time he will be training the other (TBN) post-doctoral and graduate researchers on the ICARUS system to ensure a smooth first data taking in mid to late 2017. These detector experts are expected to remain in residence at FNAL during the initial operations of the detector and play a key role in early data analysis. The graduate and undergraduate students will have the opportunity to take remote shifts on this experiment utilizing UTA's remote shift station currently residing in P.I. Asaadi's cryogenic lab. This remote station has already been used to take shifts on the LArIAT experiment and is currently being commissioned to be able to take remote shifts for the MicroBooNE experiment. This remote shift station will allow UTA to have a presence on the experiment both at FNAL and at UTA.

2.10 ICARUS Data Analysis

In addition to providing the necessary sensitivity in the $\nu_\mu \rightarrow \nu_e$ oscillation channel, the large mass and long length of the detector allow for more complete containment of high energy muons and electromagnetic showers due to $\pi^0 \rightarrow \gamma\gamma$ decays. Using this, and the deployment of a near

detector in the BNB beamline, a complimentary sterile neutrino search looking for muon neutrino disappearance as well as neutral current disappearance becomes possible. Contributing to both of these oscillation analyses is a natural fit for the UTA group given their efforts on MicroBooNE and SBND cross-sections. The extended length of the ICARUS-T600 detector provides better π/μ separation (since pions have a higher cross-section to interact) as well as more accurate muon energy reconstruction (since more muons will be fully contained) thus extending the sensitivity in the muon disappearance channel.

Similarly, by targeting a clearly identifiable neutral current process (such as $\text{NC}\pi^0$ production) the disappearance rate can be measured at both the near and far detector to search for the sterile neutrino signature in a complimentary way to the ν_e appearance. ICARUS's large volume ensures near complete photon shower containment and thus increases the statistics available for a $\text{NC}\pi^0$ disappearance search.

On top of the three detector SBN program, the stand-alone T600 detector can offer physics insight through the study of neutrino cross-sections at energies pertinent to the future planned Deep Underground Neutrino Detector (DUNE). The ICARUS experiment can due this because it will see a significant off-axis component of the Neutrinos from the Main Injector (NuMI) beam. The NuMI beam uses 120 GeV protons to produce a higher energy neutrino beam than the BNB. ICARUS is expected to collect one neutrino event every 150 seconds from the NuMI beam in the energy range of 0-3 GeV. The neutrino energy distribution for ν_e and ν_μ charged current interactions from the NuMI beam in ICARUS is shown in Figure 19. Such high energy neutrino cross-section data on an argon target will provide valuable input to the DUNE experiment and offer experimental measurements of detector efficiencies and event reconstruction techniques at these higher energies.




Figure 19: Neutrino rates taken from Reference [] for the NuMI beamline for one year of exposure ($\sim 3.0 \times 10^{20}$ POT. Left: CC- ν_e energy distribution originating from kaon decay. Right: CC- ν_μ energy distribution with one component coming from pion decay and another broader distribution coming from kaon decay.

3 The Fermilab Long-Baseline Neutrino Program

The Deep Underground Neutrino Experiment (DUNE) [?] aims to utilize massive LArTPC's to measure the CP phase in the neutrino sector as well as determine the neutrino mass hierarchy. DUNE will use a high power wide band beam capable of producing neutrinos and antineutrinos directed from Fermi National Accelerator Laboratory (FNAL) outside of Chicago, Illinois towards massive underground LArTPC detectors located in the Sanford Underground Research Facility (SURF) 1300

km away in Lead, South Dakota. By measuring the asymmetry between appearance of electron neutrinos from a beam of muon neutrinos ($P(\nu_\mu \rightarrow \nu_e)$) compared to the appearance of electron antineutrinos from a beam of muon antineutrinos and $P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e)$) as well as the precise measurement of the ν_e energy spectrum measured at the far detector, a measurement both the CP violating phase and the mass hierarchy can be done in the same experiment. In order to achieve this measurement, DUNE will require three essential components:

1) High power neutrino beam:

A neutrino beamline designed to provide sufficient intensity and appropriate energy range to exaggerate the sensitivity to the first and second oscillation maximum. The beam comes from a conventional, horn-focused neutrino beamline generated from 60 GeV - 120 GeV protons from the Fermilab Main Injector designed for initial operation at proton-beam power of 1.2 MW, with the capability to support an upgrade to 2.4 MW. The beam shall be sign-selected to provide separate neutrino and antineutrino beams with high purity to enable measurement of CP violation, the neutrino mass hierarchy, and precision oscillation measurements.

2) Large mass underground far detector:

The far detector is designed to be a 40 kiloton LArTPC consisting of four 10 kT detectors. These detectors are to be deployed 4850 feet below the surface in caverns located at SURF in order to reduce the number of cosmic rays in time with the neutrino beam to $\sim 1\%$ of the expected background. These detectors are capable of precision $\nu_\mu/\bar{\nu}_\mu$ and $\nu_e/\bar{\nu}_e$ identification and energy measurements to provide definitive measurement of the CP-phase and mass hierarchy.

2) Precision near detector:

The near detector, which is exposed to an intense flux of neutrinos, also enables a wealth of fundamental neutrino interaction measurements. The current reference design for the DUNE near detector includes a NOMAD-inspired [?] fine-grained tracker consisting of a 3.5 m \times 3.5 m \times 6.4 m; central straw-tube tracker, a lead-scintillator sampling electromagnetic calorimeter, a 4.5 m \times 4.5 m \times 8.0 m large-bore warm dipole magnet surrounding the straw tube tracker and calorimeter providing a magnetic field of 0.4 T, and RPC-based muon detectors located in the steel of the magnet as well as upstream and downstream of the tracker.

3.1 Deep Underground Neutrino Experiment (DUNE) (PI: Asaadi, Yu)

We describe in detail the project listed in the strategic plans. The primary goal of the projects listed in this section are to ensure the group to play leadership role in construction of first two 10 kt modules of DUNE as well as in physics topics of the group's interests.

3.2 APA QA/QC

Given that the highest priority of DUNE is to establish and demonstrate the functionality of its baseline technology, namely the single phase LArTPC, Asaadi will be focusing on the quality assurance, quality control, and commissioning of the anode plane assemblies (APA's) for protoDUNE SP. This work will help building up infrastructure, expertise, and capabilities necessary to contribute and lead in the DUNE SP far detector construction.

3.3 Field Cage for protoDUNE Dual Phase

Field cages provide uniform electric fields for ionization electrons to drift to anodes for the detection in the Time Projection Chamber. The baseline design for DUNE LArTPC is that of the single phase in which the ionization electrons created by the secondary particles resulting in neutrino-nucleon interactions drift in LAr and get detected on the anode plane that resides inside the liquid phase of argon. An alternative technology is the LArTPC that the ionization electrons drift through LAr but then extracted through the strong extraction field at the top of the liquid and detected in the anode in gaseous phase of argon after a signal amplification via a large area gas electron multiplier (LEM), hence the dual phase.

During the period of his sabbatical stay at CERN, Yu has begun to work on WA105, a dual phase $6m \times 6m \times 6m$ prototype testing project at CERN through the participation in the smaller prototype cosmic ray detector in $1m \times 1m \times 3m$. This work provided an opportunity for UTA IF group to join WA105 as the first U.S. group and to position itself well to play a leading role in dual phase. While dual phase technology is currently at a lower priority to the single phase LAr TPC, it is clear that U.S. groups' participation in alternate technology is beneficial in many perspectives, including that of strengthening the international nature of DUNE collaboration an essential ingredient in its success.

As the schedule for DUNE experiment and for the two protoDUNE experiments get clearer, it became apparent that UTA will be able to play an importnat role in design and construction of these experiments. The overarching strategy in identifying the construction was to ensure UTA's contribution to any of the two protoDUNE experiment would aim to direct participation in DUNE from the first 10kt module in early 2020s. One such component easily identifiable is the field cage for which the collaboration is targeting to utilize as much common components as possible for single phase and dual phase protoDUNE detectors. With this premise, Yu has discussed with the DUNE management and has agreed to take the responsibility in design and construciton of dual phase field cage together with the University of Zurich group.

Figure 20 shows the current conceptual design of the field cage for the dual phase protoDUNE experiment. The primay concept of this field cage is to use compartmentalized structure of submodules that consist of several straight profiles to provide voltage differentials for the generation of uniform fields. The use of straight profiles made of either aluminium or stainless steel makes the preassembly and shipment of submodules convenient. Each of these submodules will be prepared to the quality to hold high voltages at 180kV (for single phase) or 360kV (for dual phase) over the drift length.

At present, it has been agreed that either CERN or Fermilab will be responsible for purchasing and production of the necessary mechanical parts, including the I-beams that act as the spine of the submodule and the each profile. The project funds will pay for the purchase electrical parts - voltage divider resistors and varistors for surge arresting - and the production of electronic boards the field cage. UTA will work with the single phase field cage group, including BNL team, and the single phase field cage electronics group at Louisiana State University as well as the University of Zurich group on design of both the mechanical and electrical part of the dual phase field cage. We will be responsible for pre-assembly of the field cage submodules, mounting of the electric boards, testing of each submodules and performing functional prototype testing with as large a field cage as possible before shipping the dissembled submodules out to CERN for installation.

For the completion of the design of the dual phase field cage, our postdoctoral fellow, Animesh Chatterjee will be stationed at CERN starting from October 2016 through mid January 2017. During this period, Chatterjee will work with the ETH and CERN groups to finalize both mechanical and electrical design of the field cage for dual phase protoDUNE and ensure as much a commonality




Figure 20: Conceptual schematic drawing of the field cage assembly for dual phase protoDUNE.

as possible with that of the single phase. Once these design parameters are agreed with the single phase field cage groups, a production could proceed. The parts for the dual phase field cage will be shipped to UTA for the clean up and preparation of each of the parts, assembly of submodules, mounting of electrical components, and electrical and mechanical testing for a submodule qualification. A sizeable functional prototype field cage of size $6m \times 6m$, an electrically independent unit, will be put together and be subject under high voltage, though it may not be as high as 360kV given that the testing will be performed in air. Once the functional testing completes, the functional prototype will be disassembled and shipped to CERN for the installation.

Since protoDUNE time scale has a hard limit of completing the beam data taking before the LHC heavy ion run in October 2018, it is essential for our group to operate under this time restriction in mind as presented in the strategic plan in section III. To meet these goals, Yu plans on staying at CERN in fall 2017, thanks to the remaining funds from the agreements with LAPP and ETH which covers costs for local stay at CERN and the teaching buyout, respectively. A UTA postdoctoral fellow will also be stationed at CERN together with a graduate student to help with both single and dual phase protoDUNE installation and commissioning. Yu will then return to the U.S. at the end of year 2017 at which time Asaadi will be stationed at CERN for the first half of 2018 to continue fulfilling our responsibilities in both the protoDUNE experiments.

At the time of writing this renewal proposal, the dual phase protoDUNE field cage group is working closely with the SP-FC groups (BNL, Stony Brook and Louisiana State University) to agree on common design parameters for mechanical and electrical components of the two field cages, such as the dimension of each profile bars of the field cage, the material, the electrical board design, quality and tolerance of resistors and surge arresters, quality control and assurance criteria, etc.

3.4 Beyond the Standard Model physics group leadership - Yu

In addition to the standard neutrino physics topics, neutrino mass hierarchy and CP violation measurements in the neutrino sector, the expected high intensity proton beams provides ample opportunities for DUNE to look for physical phenomena beyond the Standard Model. UTA has been a leading participant in searching for low mass dark matter in high intensity proton beams from the start of the I.F. group in 2014. For this work, Yu has been leading the BSM physics working group of DUNE since September 2015 and has grown the group to play a significant role within the collaboration. Yu plans on ensuring various BSM topics be included in the DUNE Technical Design Report to be released in summer 2019.

Yu quickly organized the group into five subgroups based on primary physics interests and to provide necessary simulation and analysis tools specialized to support the BSM group physics analysis. The five subgroups are the simulation and software group led by our postdoctoral fellow, Chatterjee and four physics topics what cover Low Mass Dark Matter search (LDM), the Sterile Neutrino Search, the Non-standard Neutrino Interactions and Heavy Neutrino searches. Additional physics topics would continued to be added to the group's interest but these four physics topics are the primary topics to be studied in the coming 1.5 to 2 year time scale with the goal to provide the results for DUNE Technical Design Report (TDR) scheduled to compete by summary 2019. In preparation for TRD, the group plans on producing the write up for the initial list of topics and tasks to complete to provide sensitivity studies for TDR, along with the milestones for the group to follow. We anticipate that the list of tasks will be essential for the new members to participate in BSM group's planned studies.

3.5 Beam Simulation Tasks for LBNF

UTA group has contributed to beam simulations for optimization and systematic studies of the Long Baseline Neutrino Facility (LBNF). All new students joining the group are required to learn root and beam simulations as part of their training process. Since most of these tasks are well defined, each student can be assigned to the given task until they complete the task and write up the report. Many undergraduate students were able to make helpful contributions in these tasks and made presentations at beam simulation group meetings for tangible contributions. We plan on continue contributing to the beam simulation group's tasks for various studies, including an improved decay pipe radius dependence of the CPV sensitivity, target dimension and material dependence of neutrino flux and magnetic field map computations and display. New students will be assigned of additional tasks that are helpful for beam optimization group based on the discussion with the leadership of the group. This will allow students to continue improving their analysis skills while working on hardware projects.

Part IV

Research in Detector R& D

PI Summary: David Nygren

PI Summary: Benjamin Jones

SiPMWheel: a large-area, position-sensitive, energy-resolving light collector (Asaadi, Jones and Nygren.)

4 Introduction

The designs of scintillation light collection systems for noble element time projection chambers (TPCs) are driven by two main requirements:

- Photons with very short wavelengths (128 nm in Ar, 175 nm in Xe) must be collected.
- Large surface areas must be instrumented to collect as much light as possible, with a channel count kept low in order not to drive up the system cost.

Although some VUV-sensitive light detectors are available [?, ?], their quantum efficiency at these wavelengths is typically not high, and their surface area per channel is not large. To sensitize visible light detectors to VUV photons a wavelength shifter is often employed, absorbing in the UV and emitting in the visible. A common choice is tetra-phenyl butadiene (TPB) [?, ?, ?, ?, ?, ?, ?]. A fraction of the visible light thus emitted can then be detected by a standard photon detector like a silicon photomultiplier (SiPM) or a photomultiplier tube (PMT). Several geometries have been considered, including through-plate systems [?, ?], high-reflectivity foils [?, ?], and light-guides [?, ?, ?]. Light guide systems, have the advantage that large areas can be sensitized with only a moderate channel count. However, it has the disadvantage that light losses through non totally-internally-reflected rays, and surface scattering and re-absorption effects [?] are significant, and that the collection efficiency depends on the geometrical position of light arrival, making calorimetric reconstruction of localized events difficult.

We propose to develop a new large-area wavelength-shifting detector based on the light-guiding concept, with significantly improved collection efficiency and calorimetric performance. This concept is motivated by the needs of the NEXT neutrinoless double beta decay experiment [?] and by low energy physics analyses such as those of supernova neutrinos, proton decay and solar neutrinos in large liquid argon TPC detectors like DUNE [?, ?]. The pervasiveness of noble-element TPCs in particle physics is so widespread that light collection solutions with strong position and / or strong calorimetric resolution potential are likely to be widely applicable. Use cases as a primary scintillation detector may include noble element dark matter searches and other surface and underground liquid argon TPC detectors, and as an electroluminescent energy plane may include the DUNE two-phase far detector and possible argon gas near detector.




Figure 21: Left: Existing high pressure xenon gas purification and recirculation system in the lab of Nygren and Jones. Right: Existing liquid argon purification system in lab of Asaadi.

5 Research team, existing facilities, and details of request

This project will be led by Profs. Jonathan Asaadi, Ben Jones, and David Nygren at UTA. It will make use of existing liquid argon and high pressure xenon gas purification systems which are already operational at UTA and shown in Figure 21.

This project also leverages the experience of these researchers. Nygren is the inventor of the time projection chamber [?] and a pioneer of electroluminescent xenon detectors for neutrinoless double beta decay [?, ?, ?, ?, ?, ?]. He is spokesperson of the NEXT collaboration and developed a test stand that demonstrated energy resolution near the intrinsic limit of xenon gas [?] (1% FWHM at 662 keV), the worlds most precise energy resolution from a xenon detector. Asaadi is a prominent member of the MicroBooNE [?], SBND [?], DUNE [?] and LArIAT [?] collaborations, with expertise in liquid argon TPC detector design, development and construction [?]. Jones has extensive experience in noble element light collection, including the developing “Wunderbar” light-guide detectors for large LArTPCs [?, ?, ?]; assembling and operating the Bo liquid argon optical test stand at Fermilab [?, ?, ?]; exploring wavelength shifter properties and photochemistry [?]; and simulating light in liquid argon [?, ?].

Most supporting equipment for this project is already available or will be purchased from University funds, including the test stands, data acquisition systems and SiPMs, all of which were already acquired for other projects. To pursue this research we request support for:

1. Personnel: one FTE graduate student (in the form of two students at 50% effort level each), and one undergraduate.
2. M&S costs: to include argon and xenon supply, as well as fluors and plate materials.




Figure 22: Top: Example of operation of bar detectors, like the “Wunderbar” - image from [?]. Bottom: Drawing of plate detector we propose to develop.

6 Detection concept and comparison to existing devices

The detector we plan to develop will use an array of silicon photomultipliers (SiPMs) coupled around the perimeter of a TPB coated plate. As with bar-type light-guide detectors (hereafter referred to as “bars”), shown in Figure 22, top, VUV photons absorbed at the coating surface are re-emitted in the blue, some of them into the totally internally reflected modes of the polymer plate. In contrast to bar detectors, the SiPMWheel is instrumented at many points around the perimeter. This provides significant advantages which we hope to demonstrate: 1) the sensitive surface area is maximized relative to the allowed path-length between emission and detection, which optimizes light collection efficiency against losses during propagation; 2) the fraction of solid angle outside the totally internally reflected range is much reduced, leading to a higher trapped light yield 3) by reading out all SiPMs, geometrical information about the event can be extracted - as well as being intrinsically useful, this position information allows for a correction to be applied to improve calorimetric response.

In this section we derive some quantitative comparisons between our proposed SiPMWheel detector and the more typical bar-type geometry. We assume the same coating properties can be achieved over a 2D surface for both plates and bars (fabrication of the “Wunderbar” is easily generalizable [?]) and that the bar length / plate radius are free parameters to be optimized for each device.

When comparing different light collection technologies it is important to define a useful Figure of merit (FOM). The following FOMs, though by no means exclusive, appear to represent reasonable ways to assess the light collector performances for our intended use cases:

1. For illumination by a distant light source, how many photons can be captured per SiPM?
2. For illumination by a distant light source, how many photons can be captured per detection unit (one plate or one bar with many coupled SiPMs)?




Figure 23: Comparison of plate-type to bar-type detectors with various figures of merit described in the text. The plate significantly outperforms the bar-type detector in all cases. Top left: photons per SiPM. Top right: Photons per detector unit. Bottom left: Photons per unit area. Bottom right: total yield and spread from localized mono-energetic sources.

3. For illumination by a distant light source, how many photons can be captured per unit surface area?
4. For localized light deposits at different positions, what is the collection efficiency and how uniform is it?

The parameters of the detector geometry may be optimized differently to satisfy each criterion for both bar and SiPMWheel detectors. As simplifying assumptions we assume that the thickness of the plastic sheet used to make both the bar and plate is equal to the SiPM width, which we take to

be 5mm. We assume both can be prepared with the same coating quantum efficiency ϵ , are cut from the same material (acrylic with refractive index $n=1.5$), and that attenuation in the light guide is exponential in light-ray length parallel to the coated surface (this is known to be invalid at very short distances but is a reasonable approximation for longer path lengths [?]). We consider two values of the parallel attenuation length λ that appear reasonable based on past studies, $\lambda = 50$ cm and $\lambda = 100$ cm [?, ?]. Finally we assume that the 5 mm SiPMs are placed with 5 mm spacing between each, which gives three SiPMs per bar, or as many as can fit around the radius of the plate detector. The FOMs above are compared using the output of a simple ray tracing simulation.

FOM (1) is compared in Figure 23, top left. For both detector types, the collection efficiency increases as the device becomes larger, saturating at a distance comparable to the attenuation length, as expected. The plate-type detector has consistently higher collection efficiency and a higher saturated value. This is primarily due to the loss of supercritical rays in the bar detector, which the plate detector does not suffer from.

Whether the most useful Figure of merit is the light yield per channel or the light yield per detector unit depends on which factor is limiting in the experimental design or budget. A moderate improvement in FOM (1) corresponds to an enormous improvement in FOM (2) because each plate detector has a large number of SiPM channels, whereas each light guide detector has only 3. This comparison is shown in Figure 23, top right (note log scale). The improvement in FOM (3), the light collected per unit area, is intermediate between these two cases, and is shown in Figure 23, lower left. In all cases, our proposed detector represents a major improvement.

FOM (4), the stability of the light yield for light at different locations, is quantified in Figure 23, lower right, which shows example total light yield distributions for localized deposits in random positions across a device with 50 cm length / diameter and 100 cm attenuation length. Note that no photon counting fluctuations are included in these distributions - they show only the changing light yield due to differences in the detector response in different locations. Though much improved over the bar-type detector, the energy resolution obtained by simply integrating photons is still not sufficient for sub-% precision calorimetry. However, the light yield is correlated with the light source position, which in the case of the SiPMWheel, can be extracted from the distribution of light between SiPMs. The position resolution and hence the quality of the correction depend strongly on the number of photons detected, and will vary between applications, improving into the sub-percent regime as the detected photon count becomes increasingly large. The quality of this correction and the optimal method for applying it is something we plan to explore in both simulation- and hardware-based studies if this proposal is funded.

7 Electroluminescent TPC use case: The NEXT Experiment

The NEXT collaboration is a primarily US-European collaboration with the goal of developing a ton-scale, ultra-low-background neutrinoless double beta decay detector using high pressure ^{136}Xe gas (GXe) as the active medium. This technology has energy resolution far surpassing other xenon-based detectors, and a reconstructable topological signature for neutrinoless double beta decay events which is absent in liquid xenon (LXe) or xenon-doped liquid scintillator (LSXe). The projected background indices, which will ultimately limit experimental sensitivity at the ton scale, are 9 counts per ton per year per ROI (ctyR) for GXe, 130 ctyR for LXe and 210 ctyR for LSXe, as assessed by an independent review [?].

The NEXT detector is based on an electroluminescent TPC concept. Ionization charge is drifted towards a high-field region where it is amplified through nearly fluctuation-less electroluminescent gain. Each electron is accelerated in the field of the amplification region, creating excited




Figure 24: Left: Existing asymmetric TPC design, where energy must be recorded using the PMT-based “energy plane”. Right: Symmetric design that could be realized using a high-resolution “plate detector”. Plate detectors may be deployed at the anode region, the cathode region, or both.

xenon atoms which decay radiatively, emitting 175 nm light. This light is collected by two subsystems. Directly behind the electroluminescent region is a tracking array of SiPMs on a 1 cm grid. These record an image of the amplified event and allow for event topological reconstruction. Their placement is sufficiently sparse that the integrated light yield per MeV depends on the precise geometry of the event too strongly to provide a calorimetric measurement with the required precision of $\sim 1\%$ FWHM - this is shown schematically in Figure 7, top left. Addition of more SiPMs to give a complete tiling is possible but costly. Even if this were implemented, the dark rate of the many SiPMs would likely produce fluctuations in the measured energy that prevent intrinsic resolution from being achieved.

To circumvent this limitation, in the present generation of the NEXT detector, the calorimetric reconstruction of the event is handled by a different subsystem consisting of low-radioactivity PMTs at the cathode end. Light emitted in the electroluminescent region is reflected around the detector by PTFE foils and shifted to the blue by TPB coatings, and detected by the PMTs of the “energy plane”. With this arrangement, energy resolutions corresponding to 0.63% FWHM at $Q_{\beta\beta}$ have been demonstrated [?]. A sketch is shown in Figure 24, top.

This two-plane solution is not without drawbacks. Even the low radioactive photomultiplier tubes represent a significant fraction of NEXT-100’s radioactivity budget, contributing approximately 0.4 counts per ton per keV per year in each of bismuth and thallium backgrounds at $Q_{\beta\beta}$, representing the largest absolutely measured background contribution. The PMTs must be operated outside of the high-pressure region which introduces an engineering challenge, requiring an evacuated volume to be optically coupled to the active region at 15 bar. Finally, the PMTs must be




Figure 25: Left: Track energy measurement with tracking plane only (top), and tracking plane + plate (bottom). Right: Focussing effect of plate on transmitted electroluminescence light, which may improve tracking resolution in NEXT.

operated in a low-field region which leads to the HV being graded down in a short “buffer region”, wasting valuable xenon mass and introducing a region of larger HV stress.

A highly efficient calorimetric plate detector as we have described would allow a significant improvement to the NEXT design. Instead of measuring energy at the cathode, a wavelength-shifting plate between the electroluminescent mesh and the tracking plane could be used to integrate light emission from the mesh. Since the plate has a uniform collection surface, dependencies of the light yield on event geometry does not spoil its resolution, as shown in Figure 7, bottom left. The light which is not guided into the plate escapes through the back surface to be used for tracking. The plate also provides a focussing effect shown in Figure , right, removing high-angle rays from electroluminescence and potentially improving the tracking resolution of the detector. It is also plausible to add another energy plane detector behind the cathode transparent cathode. This adds the electrical complications of SiPMs being operated at high voltage, but increases the calorimetric area by a factor of two. Finally, because the vacuum region and buffer region are no longer necessary, this design allows a symmetric TPC to be realized, using the GXe volume in a highly efficient way and simplifying the delivery of HV. This concept is shown in Figure 24, bottom.

8 Single-phase LArTPC use case: DUNE

Light collection in surface-based TPCs plays a critical role of identification of cosmogenic backgrounds which would swamp true neutrino events in the absence of an optical trigger [?]. In deep underground detectors like DUNE, where cosmogenic backgrounds are much reduced, the main goal of light collection systems is fundamentally different. Rather than being primarily a tool to reject energetic off-beam cosmic ray events, the light collection system allows extension of the physics

program to low energy, non-beam physics.

The importance of light collection for non-beam physics is primarily related to establishing the position of the event in the drift direction. This is vital in order to apply a lifetime correction and thus obtain a well calibrated energy for the event from the TPC. Most of the off-beam neutrino physics goals of DUNE rely on energy reconstruction, either to identify the signal events or to learn about the physics of their sources.

The following are cases where a sensitive light collection system is vital for achieving the physics goals of DUNE:

- *Detection of supernova neutrinos* [?]. A high efficiency for detecting 5 MeV electrons has been cited as the detector goal for adequately performing this physics. This is to be contrasted to the design goal of the MicroBooNE optical system, the largest running LArTPC optical system in the USA, which was to efficiently trigger on 40 MeV protons across the (much smaller) fiducial volume. Clearly, to meet DUNE’s ambitious off-beam physics goals, high light-yield technologies surpassing existing systems are required.
- *Studies of solar neutrinos with DUNE* [?] have been discussed. This also requires sensitivity to few-MeV energy deposits across the fiducial volume, with the physics capability extending as the achievable trigger threshold is reduced. This physics will be greatly enhanced by any improvement in light collection efficiency.
- *Proton decay* [?]. Golden channels for proton decay in DUNE include $p \rightarrow K^+ \nu$, $p \rightarrow K^0 \mu^+$ and $p \rightarrow K^+ \mu^- \pi^+$. Detecting these modes requires not only to trigger on the off-beam events (likely not too challenging due to the large Q-value in the decays), but also identification of the kaon and muon daughters. Reliable identification is difficult with the TPC alone, since in many cases the “kink” in the outgoing track where the daughter particle decays is not strongly pronounced. It is thus of benefit to access the detailed time-structure of the event, and reconstruct the muon, and potentially even kaon events in time. A high collection efficiency with the optical system may allow this temporal reconstruction.

The present baseline design for the DUNE optical system is a system based on bar detectors. We have shown in Section 6 that the SiPMWheel is expected to improve upon the collection efficiency of similarly prepared bars when measured either per-SiPM, per-unit-area, or per-detector. The SiPMWheel also provides positional information - this will be valuable in cases where multiple events arrive within one drift window, as, for example, during the initial peak of flux from a nearby supernova. As with bars, the installation of SiPMWheels between mostly-transparent anode plane assemblies is possible as a deployment strategy. Two-side-coated as well as one-side-coated devices are also possible for this application.

9 Proposed program of work

The request in this proposal is primarily for personnel to develop this technology using already existing resources. We hope to acquire funding for two graduate students who will spend 50% of their research time for 3 years. The other 50% of each will be dedicated to analysis work and funded from other sources. One undergraduate will also support the team.

In the first year, development will focus on bench-top work, not involving noble element test stands. This includes learning to produce high quality optical coatings and testing them for efficiency and attenuation length in air, closely following and improving upon previous work with bar

coatings (student 1, working primarily with Jones); and commissioning of a DAQ system capable of reading out large SiPM arrays and efficiently processing the data from these (student 2, working primarily with Asaadi). Possible improvements beyond the present state-of-the-art include the addition of coating stabilizing additives to improve fluorescence yield and the exploration of high refractive index polymers. Simulation topics relating to detector optimization and expected performance will be instigated as an undergraduate project in the first year. In the second year, bench-top experience will transition into noble element environments; with Nygren and Jones, one graduate student will build a subsystem as part of an existing high pressure xenon gas test stand whereby localized electroluminescent emission can be produced near the SiPMWheel surface at various positions to study its energy and position resolution. The other student will work with Asaadi to integrate the SiPMWheel detector with his planned liquid argon calibration test stand, where an independent program of work to deploy radioactive calibration sources in large liquid argon TPCs will already be underway. With these sources, the plate performance in liquid argon will be studied. The undergraduate will assist with one or both activities. The final year will involve a program of optimization of the detector, potentially along separate trajectories for use in LAr and GXe. At the end of the three year program we hope to have demonstrated strong energy- and position-reconstruction performance and suitability of the SiPMWheel as both an electroluminescence and primary scintillation light detector.

Part V

Appendix: Biographical Sketch

Curriculum Vitae

Dr. Andrew P. White

Department of Physics, University of Texas at Arlington, Box 19059, Arlington, TX 76019

E-mail:awhite@uta.edu

Phone: (817) 272-2812 (office), (817) 272-2266 (Physics Department)

FAX: (817) 272-3637

9.1 EDUCATION AND TRAINING

- Ph.D. in Experimental High Energy Physics, Westfield College, University of London (1969-1972)
- B.Sc (Honours) Physics, University of Southampton, U.K., (1965-1969)

9.2 RESEARCH AND PROFESSIONAL EXPERIENCE

- Professor of Physics, University of Texas at Arlington (1991-present)
- Research Scientist, High Energy Physics Group, University of Florida (1985-1991)
- Staff Physicist, High Energy Physics Group, Imperial College, University of London (1973-1985) (this included a period of work based at SLAC 1974-1976)
- Research Associate, HEP, Westfield College, University of London (1972-1973)
- Research Assistant, HEP, Westfield College, University of London (1969-1972)

9.3 PUBLICATIONS:

1. ATLAS Collaboration, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”, Phys. Lett. B 716 (2012) 1-29.
2. ATLAS and CMS Collaborations, “Search for invisible decays of a Higgs boson using vector-boson fusion in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector” JHEP01 (2016) 172.
3. ATLAS and CMS Collaborations, “Search for Invisible Decays of the Higgs boson at the LHC”, A. White on behalf of the ATLAS and CMS Collaborations, Proceedings of LHCP 2015 Conference, St. Petersburg, Russia, August 2015
4. International Linear Collider, Technical Design Report, Volume 4, Detectors; <http://www.linearcollider.org/ILC/Publications/Technical-Design-Report>
5. “Experimental tests of particle flow calorimetry”, Felix Sefkow, Andy White, et al, Rev. Mod. Phys. 88, 015003 (2016)
6. ”Development of a Gas Electron Multiplier-based Digital Hadron Calorimeter”, A White et al 2012 J. Phys.: Conf. Ser. 404 01203

9.4 SYNERGISTIC ACTIVITIES

1. **D0 Intercryostat Detector:** Invented the Intercryostat Detector for D0 used to correct the energies of particle/jets due to substantial losses in dead material.
2. **GEM-based Digital Hadron Calorimetry:** Invented and developed the concept of Gas Electron Multiplier based digital calorimetry for high resolution jet energy measurements at colliders
3. **SiD Spokesperson:** Spokesperson for the SiD Detector Concept for the International Linear Collider; leading and guiding all aspects of the concept towards its realization
4. **CALICE Collaboration:** North American Representative for the CALICE Collaboration developing all aspects of calorimetry for future linear colliders.
5. **CERN RD51 Collaboration:** Deputy Chair and Member of the Management Board for RD51 - Micro-pattern Gas Detector Collaboration.

9.5 COLLABORATORS AND CO-EDITORS

- **ATLAS Experiment:** Ketevi Assamagan BNL, Joey Huston (MSU), Bill Quayle - U.Wits, Young Kee Kim - U.Chicago, Tae Hong - U.Pittsburgh, Elliot Lipeles - U. Penn., Alexander Madsen - DESY
- **SiD Detector Concept:** M.Breidenbach, J.Jaros, T. Barklow SLAC, M.Demarteau - ANL, H.Weerts ANL, J. Brau - U.Oregon, J.Strube - PNNL, M.Stanitzki (DESY)
- **CALICE Collaboration:** F.Sefkow - DESY, J.Repond - ANL, K.Kawagoe - Kyushu U.
- **RD51 Collaboration:** M. Hohlman - FIT

9.6 GRADUATE AND POSTDOCTORAL ADVISORS AND ADVISEES

1. **Mark Sosebee**, Ph.D., *University of Texas at Arlington*,
 2. **Richard Bonde**, Ph.D., *University of Texas at Arlington, 2015*
 3. **Carlos Medina**, M.S., *Colorado School of Mines, 2010*
 4. **Fajer Jafaari**, M.S., *University of Texas at Arlington and Tarrant County College, 2010*
-
1. **Dr. Seongtae Park**, Postdoctoral Fellow
University of Texas at Arlington, 2010-2014.
 2. **Dr. Mark Sosebee**, Postdoctoral Fellow
University of Texas at Arlington, 1996-present.

Curriculum Vitae

Kaushik De

Department of Physics, University of Texas at Arlington, Box 19059, Arlington, TX 76019

E-mail: kaushik@uta.edu, URL: <http://heppc1.uta.edu/kaushik/index.htm>

Phone: (817) 272-2813 (office), -2266 (physics dept.), -2824 (FAX), (682)521-5323 (cell)

Education and Training

B.A./M.A., Physics/Honors Curriculum, Summa Cum Laude,

Hunter College of CUNY, Class Rank 1/734 **1978-81**

Sc.M., Physics, Brown University **1982**

Ph.D., Physics, Brown University **1988**

Research and Professional Experience

Director, Center of Excellence in HEP, UT Arlington **2011-**

Professor, University of Texas at Arlington **2003-**

Associate Dean, Honors College, UT Arlington **1999-2003**

Associate Professor, University of Texas at Arlington **1997-2003**

Assistant Professor, University of Texas at Arlington **1993-1997**

Research Fellow, University of Michigan **1989-1992**

Research Associate, Indiana University **1988-1989**

Publications – closely related to proposed project

1. The ATLAS Experiment at the CERN Large Hadron Collider, The ATLAS Collaboration, G. Aad et al., JINST 3 (2008) S08003.
2. The ATLAS Simulation Infrastructure, The ATLAS Collaboration, G. Aad et al., Eur. Phys. J. C (2010) 70: 823–874.
3. Contributions to CHEP15 (eight papers on Computing in HEP):
<http://indico.cern.ch/event/304944/session/10/contribution/100/author/2>.
4. SUSY1
5. SUSY2

Synergistic Activities

- a) **Leadership in Physics at the New Frontier:** leading a large group of researchers and students at UTA in the cutting edge research projects in the ATLAS experiment at the Large Hadron Collider at CERN, Geneva, Switzerland, since 1995. Many masters and Ph.D. students and postdocs in Physics supervised.
- b) **Big Data Innovation:** led the development of a new paradigm in computing over the past decade: the PanDA software, which provides physicists automatic access to hundreds of supercomputing centers internationally. Thousands of physicists analyze data and publish results in multiple High Energy Physics (HEP) experiments using PanDA. Supervised/co-supervised many masters and Ph.D. theses in Computer Science on PanDA.

- c) **New Discoveries:** played key roles in many aspects of the HEP experiments that discovered two fundamental particles in physics over the past two decades: the top quark at the Tevatron, and the Higgs boson at the LHC.
- d) **New Physics searches:** early proponent of the search for the supersymmetric partner of the top quark in both the D0 and the ATLAS experiments at the LHC. Supervised multiple Ph.D. students who completed theses in D0 and ATLAS on this topic.
- e) **Supercomputing technology:** founding director of the SouthWest Tier 2 supercomputing center, located at UTA and Oklahoma University. Funded by multiple grants from National Science Foundation, and the Department of Energy.

Collaborators

The D0 collaboration (see <http://www-d0.fnal.gov/~madaras/authorlist.html>)

The ATLAS collaboration (see

<http://graybook.cern.ch/programmes/experiments/lhc/ATLAS.html>)

Graduate and Postdoctoral Advisors

Prof. Mildred Widgoff (Brown University), Prof. Andrej Zieminski (Indiana University), Prof. Homer Neal (University of Michigan).

Graduate Student Advisees

Yan Song (IBM), Barry Spurlock (UTA), Rishiraj Pravahan (AT&T), Smita Darmora (UTA), Jared Little (UTA), Ted Eltzroth (unknown), Nevzat Guler (unknown), Richard Kaiser (NRC), Yu Xia (unknown).

Postdoctoral Associates

Elizabeth Gallas (Oxford), Jia Li (deceased), Mark Sosebee (UTA), Armen Vartapetian (UTA), Nurcan Ozturk (UTA), Paul Nilsson (BNL), Alden Stradling (UTA), Giulio Usai (UTA), David Cote (Ciena).

ANDREW BRANDT, PH.D.
UNIVERSITY OF TEXAS AT ARLINGTON

A. Education and Training:

| | | |
|---------------------------------------|---------------------|-------------|
| College of William and Mary | Physics, Economics | B.S., 1985 |
| University of California, Los Angeles | Physics | M.S., 1988 |
| University of California, Los Angeles | Physics | Ph.D., 1992 |
| Fermi National Accelerator Laboratory | Postdoctoral Fellow | 1992-1995 |
| Fermi National Accelerator Laboratory | Wilson Fellow | 1996-1999 |

B. Research and Professional Experience:

- 2010 – Professor, The University of Texas at Arlington, Texas
2004-2010 Associate Professor, The University of Texas at Arlington, Texas
1999-2004 Assistant Professor, The University of Texas at Arlington, Texas

C. Publications related to the proposed project:

1. ``Search for charged Higgs bosons decaying via $H^+ \rightarrow \tau\nu$ in fully hadronic final states using pp collision data at $\sqrt{s} = 8$ TeV, with the ATLAS detector," (ATLAS Collaboration), JHEP03 88 (2015); arxiv:1412.6663."
2. "Search for charged Higgs bosons produced in association with a top quark and decaying via $H^+ \rightarrow \tau\nu$ using pp collision data recorded at $\sqrt{s} = 13$ TeV by the ATLAS detector, Phys. Lett. B 759 (2016) 555-574, arxiv:1603.09203.
3. "Expected Performance of the ATLAS Experiment: Detector Trigger and Physics," ATLAS Collaboration, arXiv:0901.0512 CERN-OPEN-2008-020. ISBN: 978-92-9083-321-5.
4. "Expected pileup values at the HL-LHC," ATL-UPGRADE-PUB-2013-014.
5. "The FP420 R&D Project: Higgs and New Physics with Forward Protons at the LHC," FP42Collaboration, arXiv:0806.0302v2 [hep-ex]; J. Inst.: 2009_JINST_4_T10001. "A Forward Proton Detector at DZero," A. Brandt *et al.*, Fermilab Proposal P-900, FERMILAB-PUB-97-377.
6. ``Letter of Intent for the Phase-I Upgrade of the ATLAS Experiment," CERN-LHCC-2011-012 LHCC-I-020; <https://cds.cern.ch/record/1402470>
7. "A Particle Consistent with the Higgs Boson Observed with the ATLAS Detector at the Large Hadron Collider", Science 338 (2012) 1576-1582"
8. ``Design of Cherenkov bars for optical part of Time-of-flight detector in Geant 4," A.Brandt, L. Nozka; Optics Express. <http://www.opticsinfobase.org/oe/upcomingissue.cfm>
9. ``Charged-particle multiplicities in pp interactions measured with the ATLAS detector at the LHC," New J. Phys. 13 053033 (2011); CERN-PH-EP-2010-079; arXiv:1012.5104.
10. "Search for the Standard Model Higgs Boson in Tau Lepton Pair Final States" V. M. Abazov et al. (D0 Collaboration), Phys. Rev. D 88, 052005;arXiv:1203.4443.

D. Synergistic Activities:

1. Established and led ATLAS Trigger Rates Group (2008-2010)
2. U.S ATLAS L2 Manager for AFP in Phase 1 Upgrade CD-0 (2012-2013)
3. Supervised grad students on Diffractive Higgs, SM Higgs, and post-doc on charged Higgs (2010-2016)
4. Spokesman of T958, Fermilab Test Beam Experiment on picosecond timing (2006-2015)
5. Proposed, initiated, and led DZero Forward Proton Detector group (1996-2006)

E. Collaborators and Other Affiliations

Collaborators: **ATLAS Collaboration**, Hal Evans (Indiana), Elliot Lipeles (Penn), Mark Oreglia (Chicago), Michael Reeijssenbeek (Stony Brook)

Graduate and Postdoctoral Advisors:

Graduate – H. Montgomery (Jefferson Lab)
Postdoctoral – P. Schlein (UCLA, deceased)

Thesis Advisor and Postgraduate-Scholar Sponsor:

Past graduate students:

Michael Strang (Ph.D. Physics, 2005) Ohio State; Pedro Duarte (M.Sc. Physics, 2007) Data Engineer Altx, Berkeley, CA; Arnab Pal (Ph.D. Physics, 2013) Ninah Consulting, Gurgaon, India; Ian Howley (Ph.D. Physics, 2013) NASA Marshal Space Flight Center.

Post-doctoral Fellows:

Christophe Royon (2000-2002) Kansas, Pierrick Hanlet (2001-2002) Illinois Institute of Technology, Duncan Brown (2004-2007) Fermilab, Edward Sarkisyan-Grinbaum (2008-2012) CERN, Justin Griffiths (2013-) UTA

Amir Farbin

University of Texas Arlington
Department of Physics
P.O. Box 19059
Arlington, TX, 76019

e-mail: afarbin@uta.edu

Education and Training

Massachusetts Institute of Technology Cambridge, MA
Sep 93 - Jun 1997
S.B. in Physics. Thesis advisor: Louis Osborne.
University of Maryland College Park, MD
Aug 97 - June 03
Ph.D. in Physics. Thesis advisor: Hassan Jawahery.

Postdoctoral Research Assistant BaBar Group- UMD
June 03 - April 04

Research Associate ATLAS- University of Chicago
May 04 - Feb 05

Research Fellow ATLAS- CERN
March 05 - August 07

Research and Professional Experience

Associate Professor (Tenured) University of Texas Arlington
September 2012-Present
Collaborator on ATLAS, DUNE, MiniBooNE, and LArIAT experiments. DUNE Deputy Computing Coordinator. ATLAS contributions include Supersymmetry searches, physics analysis tool leadership and development, Tile Hadronic Calorimeter operations and leadership, and ATLAS software tutorials and documentation.

Assistant Professor University of Texas Arlington
August 07 - September 2012
2009 DOE Early Career Research Program (ECRP) Award for “Model-Independent Dark-Matter Searches at the ATLAS Experiment and Applications of Many-core Computing to High Energy Physics”.

Recent Relevant Publications and Presentations

- ATLAS Collaboration, *Further searches for squarks and gluinos in final states with jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector*, Tech. Rep. ATLAS-CONF-2016-078, CERN, Geneva, August, 2016. <http://cds.cern.ch/record/2206252>
- ATLAS Collaboration, *Search for squarks and gluinos in final states with jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector*, Eur.Phys.J. C76 (2016) no.7, 392, arXiv:1605.03814.
- ATLAS Collaboration, *Summary of the searches for squarks and gluinos using $\sqrt{s} = 8$ TeV pp collisions with the ATLAS experiment at the LHC*, JHEP 10 (2015) 054, arXiv:1507.05525.
- ATLAS Collaboration, *Multi-channel search for squarks and gluinos in $\sqrt{s} = 7$ TeV proton-proton collisions with the ATLAS Detector*, Eur.Phys.J. C73 (2013) 2362, arXiv:1212.6149.

- R.Acciarri *et al.* [DUNE Collaboration], “Long-Baseline Neutrino Facility (LBNF) and Deep Underground Neutrino Experiment (DUNE) Conceptual Design Report Volume 2: The Physics Program for DUNE at LBNF”.

Collaborations

- Member of the ATLAS Collaboration (2004-present).
- Member of the LBNE and DUNE Collaborations (2013-present).
- Member of the LArIAT Collaboration (2012-present).
- Member of the miniBooNE Collaboration (2012-present).
- Member of the NEXT Collaboration (2016-present).
- Member of the BaBar Collaboration (1998-2005).

Advisors and Mentorship

- Undergraduate Advisors: T. Arias (MIT), L. Osborne (MIT).
- Ph.D. Advisor: H. Jawahery (University of Maryland).
- Postdoctoral Advisors: H. Jawahery (University of Maryland), F. Merritt (University of Chicago).
- Postdoctoral Mentorship: (all at University of Texas at Arlington): A. R. Stradling, R. Pravahan (AT&T Foundry, Palo Alto), Louise Heelan (UTA), David Cote (Ciena Telecommunications, Ottawa, Canada).
- Graduate Mentorshop (all at University of Texas at Arlington): P. C. Vajhula (current affiliation unknown), Heather Brown (Hewlett-Packard), Daniel Bullock (UTA), Sepideh Shahsavari (UTA).

Haleh Hadavand

Education and Training

Undergraduate: University of Maryland College Park

Sep 1995 - Jun 1999

Degree: BS in Physics, June 1999.

College Park, MD

Graduate: University of California San Diego

Aug 1999 - Sept 2005

Degree: Ph.D. in High Energy Physics Dissertation: "The Measurement of CP Asymmetries in the Three-body Charmless B_d Meson Decay to $K_S^0 K_S^0 K_S^0$ at BABAR "

Thesis adviser: David MacFarlane.

La Jolla, CA

Research and Professional Experience

Assistant Professor

Nov 2014 - Present

University of Texas Arlington

Research Faculty

July 2012 - Oct 2014

University of Texas Arlington

Postdoctoral Fellow

Sept 2005-July 2012

Southern Methodist University

Graduate Research Assistant

Jun 1999 - Sept 2005

UCSD on BABAR experiment

Relevant Publications

- [1] The ATLAS Collaboration, Search for charged Higgs bosons produced in association with a top quark and decaying via $H^\pm \rightarrow \tau\nu$ using pp collision data recorded at $\sqrt{s} = 13$ TeV by the ATLAS detector, Phys. Lett. B **759** (2016) 555-574, arxiv:1603.09203.
- [2] The ATLAS Collaboration, "Search for charged Higgs bosons decaying via $H^\pm \rightarrow \tau^\pm\nu$ in fully hadronic final states using pp collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector", JHEP03 (2015) 088.
- [3] The ATLAS Collaboration, "A Search for Charged Higgs Bosons in the $\tau +\text{jets}$ Final State with pp Collision Data Recorded at $\sqrt{s} = 8$ TeV with the ATLAS Experiment", ATLAS-CONF-2013-090 (2013).
- [4] The ATLAS Collaboration, "Search for extra dimensions using diphoton events in 7 TeV proton-proton collisions with the ATLAS detector", Phys. Lett. B **710**, 538-556 (2012).
- [5] The ATLAS Collaboration, "Search for Diphoton Events with Large Missing Transverse Energy in 7 TeV Proton-Proton Collisions with the ATLAS Detector", Phys. Rev. Lett. **106**, 121803 (2011).
- [6] Ilchenko, Y., Cuenca-Almenar, C. , Corso-Radu, A., Hadavand, H. Kolos, S, Slagle, K., Taffard, A., "Data Quality Monitoring Display for ATLAS experiment", J. Phys. Conf. Ser. **219**, 022035 (2010).
- [7] H. Hadavand [ATLAS Collaboration], "Commissioning of the ATLAS offline software with cosmic rays", J. Phys. Conf. Ser. **119**, 032021 (2008).
- [8] S. Kolos, A. Corso-Radu, H. Hadavand, M. Hauschild, R. Kehoe, "A software framework for Data Quality Monitoring in ATLAS", J. Phys. Conf. Ser. **119**, 022033 (2008).
- [9] B. Aubert *et al.* [BABAR Collaboration], "Branching Fraction and CP Asymmetries in $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ ", Phys. Rev. Lett. **95**, 011801 (2005).

- [10] B. Aubert *et al.* [BABAR Collaboration], “Measurement of the B^+/B^0 production ratio from the $\Upsilon(4S)$ meson using $B^+ \rightarrow J/\psi K^+$ and $B^0 \rightarrow J/\psi K_S^0$ decays”, Phys. Rev. D **69**, 071101 (2004). “New Physics results from the BABAR Collaboration”, July 2005.

Synergistic Activities

Invited Talks

- [11] Lake Louise Winter Institute, Lake Louise, Alberta, Canada, “Beyond-the-Standard Model Higgs and Invisible Higgs Decays Using the ATLAS Experiment”, February 2014.
- [12] Photon 2011 Conference at Spa, Belgium, “New Physics Searches with Photons at the ATLAS and CMS Experiments”, May 2011.
- [13] Beyond Standard Model Physics Conference, Boston, MA., “Beyond the Standard Model Photon Physics at the ATLAS and CMS Experiments at the Large Hadron Collider”, June 2009.

Leadership Experience

- Charged Higgs convenor on the ATLAS experiment.
- Co-editor of paper and conference note on extra dimension diphotons resonances [3, 4].
- co-editor and analysis contact for paper *Search for Diphoton Events with Large Missing Transverse Energy in 7 TeV Proton-Proton Collisions with the ATLAS Detector* [5].

Collaborators and Co-editors

Maria Pilar Casado, Universitat Autònoma de Barcelona, Arnaud Ferrari, Uppsala University, Thomas Junk, Fermilab, Anna Kopf, Freiburg University, Allison McCarn, University of Michigan, Henrik Ohman, Uppsala University, Michael Pitt, Weizmann Institute of Science, John Parsons, Columbia University, Nikolaos Rompotis, University of Wisconsin, Jana Schaarschmidt, Weizmann Institute of Science, Stephen Sekula, Southern Methodist University, Camila Rangel Smith, Uppsala University, Michelle Stancari, Fermilab, Micheal Pitt, Weizmann Institute of Science, Dhiman Chakraborty, Northern Illinois University.

Graduate and Postdoctoral Advisors and Advisees

Yuriy Ilchenko, UT Austin, David MacFarlane, Stanford University, Bob Kehoe, Southern Methodist University, Ryszard Stroynowski, Southern Methodist University.

Jaehoon Yu

Department of Physics, University of Texas at Arlington, Box 19059, Arlington, TX 76019

E-mail:jaehoonyu@uta.edu, URL:<http://www-hep.uta.edu/~yu/>

Phone: (817) 272-2814 (office), -2266 (physics dept.), (817)808-9605 (cell)

FAX: (817) 272-3637 (dept)

EDUCATION AND TRAINING

- **Research Associate**, Fermi National Accelerator Laboratory, **1996-1998**
- **Research Fellow**, University of Rochester, **1993-1996**
- **Ph.D.**, Physics, State University of New York, Stony Brook, **1993**
- **M.S.**, Physics, State University of New York, Stony Brook, **1992**
- **M.A.**, Physics, Korea University, Seoul, South Korea, **1985**
- **D.S.**, Physics, Korea University, Seoul, South Korea, **1983**

RESEARCH AND PROFESSIONAL EXPERIENCE

- **Professor**, University of Texas at Arlington, **2012–present**
- **Associate Professor**, University of Texas at Arlington, **2006–2012**
- **Assistant Professor**, University of Texas at Arlington, **2001–2006**
- **Associate Scientist**, Fermi National Accelerator Laboratory, **1998-2001**

SELECTED PUBLICATIONS

1. ATLAS Collaboration, Measurement of exclusive $\gamma\gamma \rightarrow W^+W^-$ Production and search for exclusive Higgs boson production in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector, *Phys. Rev. D* **94**, 032011 (2016)
2. ATLAS Collaboration, Measurements of the Higgs boson production and decay rates and coupling strengths using pp collision data at $s=7$ and 8 TeV in the ATLAS experiment, arXiv:1507.04548 [hep-ex]
3. Georges Aad et al., ATLAS Collaboration, Study of (W/Z)H production and Higgs boson couplings using HWW decays with the ATLAS detector, *JHEP* **1508** (2015) 137.
4. Keisuke Fujii et al., Linear Collider Physics Panel, Physics Case for the International Linear Collider, arXiv:1506.05992 [hep-ex].
5. CMS and ATLAS Collaborations, Combined Measurement of the Higgs Boson Mass in pp Collisions at $s=7$ and 8 TeV with the ATLAS and CMS Experiments, *Phys.Rev.Lett.* **114** (2015) 191803.
6. Georges Aad et al., ATLAS Collaboration, Search for a Charged Higgs Boson Produced in the Vector-Boson Fusion Mode with Decay HWZ using pp Collisions at $s=8$ TeV with the ATLAS Experiment, *Phys.Rev.Lett.* **114** (2015) 23, 231801.
7. C. Adams et al., LBNE Collaboration, The Long-Baseline Neutrino Experiment: Exploring Fundamental Symmetries of the Universe, arXiv:1307.7335 (2014)
8. H. Frisch, C. Hast, E. Ramberg, M. Artuso, A. Seiden, M. Wetstein, M.C. Sanchez & J. Fast, et al, Compendium of Instrumentation Frontier Whitepapers on Technologies for Snowmass 2013
9. Georges Aad et al., ATLAS Collaboration, Search for dark matter in events with heavy quarks and missing transverse momentum in pp collisions at $s = 7$ TeV with the ATLAS detector arXiv:1410.4031 (2014)
10. Georges Aad et al., ATLAS Collaboration, Search for neutral Higgs boson of the minimal supersymmetric standard model in pp collisions at $s = 8$ TeV with the ATLAS detector *JHEP* **1411** (2014) 056

SYNERGISTIC ACTIVITIES

1. **Aug. 2016 present:** Executive Board member, WA105 experiment at CERN
2. **Nov. 2015 present:** Institutional Board member, ICARUS experiment at CERN/Fermilab
3. **Sept. 2015 present:** DUNE Beyond the Standard Model physics group co-convener
4. **Mar. 2015 present:** Institutional Board representative, Deep Underground Neutrino Experiment at Fermilab
5. **Aug. 2013 Jan. 2015:** LBNE R&D Coordinating committee co-convener

COLLABORATORS:ATLAS, CALICE, SiD, LBNE, LArIAT and ORKA Collaborations
GRADUATE AND POSTDOCTORAL ADVISORS

- **Dr. Robert Bernstein**, Fermilab, PostDoctoral Advisor: **1996–1998**
- **Prof. Frederick Lobkowitz (deceased)**, Univ. of Rochester, PostDoctoral Advisor: **1993–1996**
- **Prof. Robert L. McCarthy**, SUNY Stony Brook, Thesis Advisor: **1988–1993**

POSTDOCTORAL ADVISEES

1. **Dr. Animesh Chatterjee**, Postdoctoral Fellow, University of Texas at Arlington, **2014-present**.
2. **Dr. Justin Griffiths**, Postdoctoral Fellow, University of Texas at Arlington, **2012-present**.
3. **Dr. Seongtae Park**, Senior Postdoctoral Fellow, University of Texas at Arlington, **2010-2014**
4. **Dr. Hyunwoo Kim**, Postdoctoral Fellow, University of Texas at Arlington, **2004-2007**, currently an associate scientist at Fermilab
5. **Mr. Sudhamshi Reddy**, Software Engineer, University of Texas at Arlington, **2007-2009**, currently on UTA Computer Science and Engineering Ph.D. candidate

GRADUATE STUDENT ADVISEES

1. **Garrett Brown**, Ph.D., Univ. of Texas at Arlington, **2016 - present**
2. **Last Feremenga**, Ph.D., Univ. of Texas at Arlington, expected to graduate in **2016**
3. **Heeyeun Kim**, Ph.D., Univ. of Texas at Arlington, **2015 - Researcher at Harvard Medical School**
4. **Dr. Jacob Smith**, Ph.D., Univ. of Texas at Arlington, **2013–PostDoc at U. of Maryland**
5. **Dr. Hyeonjin Kim**, Ph.D., Univ. of Texas at Arlington, **2010–PostDoc at U. of Stockholm, Sweden**
6. **Dr. Venkatesh Kaushik**, Ph.D., Univ. of Texas at Arlington, **2007– EMC²**
7. **Jacob Smith**, M.S., Univ. of Texas at Arlington, **2010–Continued into the Ph.D. program at UTA**

Biographical Sketch Jonathan Asaadi

Education and Training

| Institution | Location | Major | Degree & Year |
|---------------------------|----------------------|-----------|---------------|
| Undergraduate Institution | University of Iowa | Physics | B.S. 2004 |
| Graduate Institution | Texas A&M University | Physics | M.S. 2007 |
| Graduate Institution | Texas A&M University | Physics | PhD. 2012 |
| Postdoctoral Institution | Syracuse University | Neutrinos | 2012-2015 |

Research and Professional Experience

| | | |
|-------------------------|-------------------------------|----------------|
| Assistant Professor | University of Texas Arlington | 2015 – Present |
| Postdoctoral Researcher | Syracuse University | 2012 – 2015 |

Publications

- “*Measurement of ν_μ and $\bar{\nu}_\mu$ Neutral Current $\pi^0 \rightarrow \gamma\gamma$ Production in the ArgoNeuT Detector*”, **Submitted to PRD (2014), arXiv:1511.00941 (Primary author and primary analyzer)**
- “*Testing of High Voltage Surge Protection Devices for Use in Liquid Argon TPC Detectors*”, **JINST 9 P09002 (2014), arXiv:1406.5216 (Primary author and primary analyzer)**
- “*The detection of back-to-back proton pairs in Charged-Current neutrino interactions with the ArgoNeuT detector in the NuMI low energy beam*” **Phys. Rev. D 90, 012008 (2014), arXiv:1405.4261 (Reviewer and collaborator)**
- “*Measurements of Inclusive Muon Neutrino and Antineutrino Charged Current Differential Cross Sections on Argon in the NuMI Antineutrino Beam*” **Phys. Rev. D 89, 112003 (2014), arXiv:1404.3698 (Collaborator)**
- “*A Proposal for a Three Detector Short-Baseline Neutrino Oscillation Program in the Fermilab Booster Neutrino Beam*” **arXiv:1503.01520 (Collaborator)**
- “*ArgonCube: a novel, fully-modular approach for the realization of large-mass liquid argon TPC neutrino detectors*” **CERN-SPSC-2015-009 ; SPSC-I-243 (Contributing author and analyzer)**
- “*LAr1-ND: Testing Neutrino Anomalies with Multiple LArTPC Detectors at Fermilab*” **Snowmass White Paper SNOW13-00176, arXiv:1309.7987 (Collaborator)**
- “*Signature-based search for delayed photons in the exclusive photon plus missing transverse energy events from proton anti-proton collisions with center of mass energy = 1.96 TeV*” **Phys. Rev. D 88, 031103 (2013), arXiv:1307.0474 (Primary author and primary analyzer)**
- “*LArIAT: Liquid Argon In A Testbeam*” **arXiv:1406.5560 (Collaborator)**

Synergistic Activities

- **Neutrino Detector R&D Facilities Workshop**

Organizing Committee Member, January 2016

- **The Liquid Argon TPC Reconstruction Assessment and Requirement Workshop**

Organizing Committee Member, November 2015

- **Albert Einstein Center Visiting Fellow 2014**, Laboratory for High Energy Physics (LHEP),
University of Bern Switzerland

- **Coordinating Panel for Advanced Detectors (CPAD) Instrumentation Frontier Meeting**

Invited Talk “New Technologies for Neutrino Oscillations”, October 2015

- **25th Workshop on Weak Interactions and Neutrinos (WIN2015)**

Invited Talk “The Fermilab Short-Baseline Neutrino Program”

Collaborators

Collaborators and Co-Editors:

| | | |
|-------------------|---------------------------------------|--------------------------|
| Adam Aurisano | University of Cincinnati | Collaborator |
| Bruce Baller | Fermilab | Collaborator |
| Tim Bolton | Kansas State University | Collaborator |
| Carl Bromberg | Michigan State University | Collaborator |
| Flavio Cavanna | Fermilab | Collaborator |
| Eric Church | Pacific Northwest National Laboratory | Collaborator |
| Janet Conrad | Massachusetts Institute of Technology | Collaborator |
| Bhaskar Dutta | Texas A&M | Graduate Advisor |
| Antonio Ereditato | Bern University | Collaborator |
| Bonnie Fleming | Yale University | Collaborator |
| Teruki Kamon | Texas A&M University | Graduate Advisor |
| Igro Kreslo | Bern University | Collaborator |
| Ornella Palamara | Fermilab | Collaborator |
| Jennifer Raaf | Fermilab | Collaborator |
| Brian Rebel | Fermilab | Collaborator |
| Mitch Soderberg | Syracuse University | Post-doctoral Advisor |
| Josh Spitz | University of Michigan | Collaborator |
| Andrzej Szlec | Manchester University | Collaborator |
| David Toback | Texas A&M University | Graduate Advisor (Chair) |
| Michele Weber | Bern University | Collaborator |
| Tingjun Yang | Fermilab | Collaborator |
| Geralyn Zeller | Fermilab | Collaborator |

Graduate Advisors and Postdoctoral Sponsors

Prof. David Toback (Texas A&M)

Prof. Mitch Soderberg (Syracuse University)

David Nygren

Biographical Sketch

Education and Training

| | | |
|---|-------------------------|-------------|
| Whitman College, Washington | Mathematics and Physics | B.A., 1960 |
| University of Washington, Seattle, Washington | Physics | Ph.D., 1967 |
| Nevis Laboratories, Columbia University, New York | Postdoctoral researcher | 1967-1969 |

Research and Professional Experience

- 2014 – Presidential Distinguished Professor, The University of Texas at Arlington
2014-2015 Division Director, Acting, LBNL Physics Division
2004-2014 Distinguished Scientist, LBNL
1999-2009 LBNL Senior Staff
1997-2004 Physics Division Fellow, LBNL
1992-1997 Senior Research Associate, Nevis Laboratories
1990-1992 Assistant Professor, Columbia University

Selected Publications (ordered by relevance)

1. "Operation and first results of the NEXT-DEMO prototype using a silicon photomultiplier tracking array", NEXT Collaboration, *Journal of Instrumentation* 8 (2013) P09011
2. "Ionization and scintillation response of high-pressure xenon gas to alpha particles", NEXT Collaboration, *Journal of Instrumentation* 8 (2013) P05025,
3. "Near-Intrinsic Energy Resolution for 30 to 662 keV Gamma Rays in a High Pressure Xenon Electroluminescent TPC", NEXT Collaboration, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* 708 (2013) 101-114,
4. "High-pressure Xenon Gas Electroluminescent TPC for 0-v $\beta\beta$ Decay Search", D. Nygren, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **603** (2009) p337-348
5. "Optimal detectors for WIMP and 0-v $\beta\beta$ searches: Identical high-pressure xenon gas TPCs" D. Nygren, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*. **581** (2007) 632-642
6. "First Observation of PeV-energy Neutrinos with IceCube", IceCube Collaboration, *Physical Review Letters* 111 (2013) 021103, arXiv:1305.5356
7. "A negative-ion TPC with ultra-high energy resolution for 0-v double beta decay search in ^{136}Xe ", D. R. Nygren. *Third Symposium on Large TPCs for Low-energy Rare Events* (Paris, France, 11-12 December 2006) *Journal of Physics: Conference Series* **65** (2007) 012003 doi:10.1088/1742-6596/65/1/012003
8. "High Resolution X-ray Imaging Using a Silicon Strip Detector", E. Beuville, R. Cahn, B. Cederstrom, M. Danielsson, A. Hall, B. Hasegawa, L. Luo, M. Lundqvist, D. Nygren, E. Oltman, J. Walton, *IEEE Transactions on Nuclear Science* 45: 3059-3063 (1998).
9. "Measurement of the Kaon Content of Three Prong τ Decays", by TPC/Two Gamma Collaboration, *Physical Review D* 50:13 (1994).
10. "Performance of a Time Projection Chamber", D. Fancher, H.J. Hilke, S. Loken, P. Martin, J.N. Marx, D.R. Nygren, P. Robrish, G. Shapiro, M. Urban and W. Wenzel, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* 161:383 (1979).

Synergistic and Public Service Activities

1. DOE Review of LZ -CD2/3 April 2016
2. DOE Review of National Laboratory R&D
3. DOE Institutional Review of FNAL - February 2015
4. DOE Particle Detector R&D Review Panel- November 2013
5. DPF Snowmass 2013: Advisor for Gas Detection Techniques
6. DOE Early Career Award: Proposal Review Committee 2013
7. DOE Review of Samurai TPC Project MSU, 2013

Collaborators

IceCube Collaboration
NEXT Collaboration

Collaborators and Co-editors:

Azriel Goldschmidt, LBNL, Nuclear Science Division
Juan Jose Gomez-Cadenas, IFIC, Valencia, Spain
Francis Halzen, University of Wisconsin, Madison
John Hauptman, Iowa State University
James Siegrist, Director, Office of High Energy Physics, DOE
Bob Stokstad, LBNL, Nuclear Science Division, retired

Graduate and Postdoctoral Advisors and Advisees:

Graduate – Robert W. Williams, University of Washington, Seattle
Postdoctoral – Jack Steinberger, Columbia University, New York

Thesis Advisor and Postgraduate-Scholar Sponsor:

Graduate Students:

Jordan Benson (current), Marjorie Shapiro (now at UC Berkeley), Nick Hadley (now at University of Maryland), John Huth (now at Harvard University)

Postgraduate-scholars:

Ben Jones (now at UTA), Carlos A. B. de Oliveira

Benjamin Jones

Biographical Sketch

Education and Training

| | | |
|-------------------------------|----------------------------------|-----------|
| Mass. Institute of Technology | Ph.D. in Physics | 2010-2015 |
| University of Bristol | Undergraduate research assistant | 2009-2010 |
| Cambridge University | B.A. M.Sci (Nat Sci) | 2004-2009 |

Research and Professional Experience

| | | |
|-------------------------|----------------------------------|--------------|
| Assistant Professor | University of Texas at Arlington | 2016-present |
| Postdoctoral researcher | University of Texas at Arlington | 2015-2016 |

Selected Publications (ordered by relevance)

1. *Improved TPB-coated Light Guides for Liquid Argon TPC Light Detection Systems*, Z. Moss, L. Bugel, G. Collin, J.M. Conrad, B.J.P. Jones, J. Moon, M. Toups and T. Wongjirad JINST 10 (2015) no.08, P08017
2. *Demonstration of a Lightguide Detector for Liquid Argon TPCs*
L. Bugel, J.M. Conrad, C. Ignarra, B.J.P. Jones, T. Katori, T. Smidt and H.-K. Tanaka, Nucl.Instrum.Meth. A640 (2011) 69-75
3. *A Simulation of the Optical Attenuation of TPB Coated Light-guide Detectors*
B.J.P. Jones JINST 8 (2013) C10015
4. *Photodegradation Mechanisms of Tetraphenyl Butadiene Coatings for Liquid Argon Detectors*, B.J.P. Jones, J.K. VanGemert, J.M. Conrad, A. Pla-Dalmau, JINST 8 (2013) P01013
5. *The Effects of Dissolved Methane upon Liquid Argon Scintillation Light*
B.J.P. Jones, T. Alexander, H.O. Back, G. Collin, J.M. Conrad, A. Greene, T. Katori, S. Pordes and M. Toups, JINST 8 (2013) P12015
6. *A Measurement of the Absorption of Liquid Argon Scintillation Light by Dissolved Nitrogen at the Part-Per-Million Level*, B.J.P. Jones, C.S. Chiu, J.M. Conrad, C.M. Ignarra, T. Katori and M. Toups JINST 8 (2013) P07011
7. *Single Molecule Fluorescence Imaging as a Technique for Barium Tagging in Neutrinoless Double Beta Decay*, B.J.P. Jones, A.D. McDonald, D.R. Nygren, arXiv1609:04019, (2016) submitted to JINST
8. *Searches for Sterile Neutrinos with the IceCube Detector*, M.G. Aartsen *et al.* (IceCube Collaboration), Phys.Rev.Lett. 117 (2016) no.7, 071801
9. *Dynamical pion collapse and the coherence of conventional neutrino beams*
B.J.P. Jones, Phys.Rev. D91 (2015) no.5, 053002
10. *Testing of High Voltage Surge Protection Devices for Use in Liquid Argon TPC Detectors* J. Asaadi, J.M. Conrad, S. Gollapinni, B.J.P. Jones, H. Jostlein, J. M. St. John, T. Strauss, S. Wolbers and J. Zennamo, JINST 9 (2014) P09002

Synergistic Activities

1. Served as reviewer for Department of Energy SBIR (Small Business Innovation and Research) proposals.
2. Field cage project leader for the NEXT experiment
3. Served as analysis reviewer for two IceCube oscillations working group physics analyses.
4. Organizer of UTA high energy physics summer school 2016
5. Recent outreach talks:
“*A Taste of Research at UTA*”, Talk for high school students visiting UTA physics department
“*Neutrino Telescopes and New Physics*” Lunch Talk for the UTA Society of Physics Students
“*Detecting Neutrinos with Liquid Argon*”, public talk and museum event for the Neutrino2014 conference
“*From Symmetries to Neutrinos*”, after-school talk on physics for 16-18 year olds at King Edwards School, Stratford-Upon-Avon, UK
“*Discovering the Ingredients of the Universe*”, Guest lecture for “From Big Bangs to Black Holes” physics course at Roosevelt University, Chicago, IL

Collaborators

| | |
|--------------------------|----------------|
| IceCube Collaboration | 2014 – present |
| NEXT Collaboration | 2015 – present |
| DUNE Collaboration | 2013 – present |
| MicroBooNE Collaboration | 2010 – 2015 |

Collaborators and Co-editors:

Arguelles Delgado, Carlos (MIT), Back, Henning (Princeton), Grant, Darren (University of Alberta), Gollapini, Sowjanya (KSU), Halzen, Francis (UW Madison), Jostlein, Hans (Fermilab), Karle, Albrecht (UW Madison), Katori, Teppei (Queen Mary University of London), Koskinen, Jason (NBI Copenhagen), Lockwitz, Sarah (Fermilab), Mufson, Stuart (Indiana University), Pordes, Stephen (Fermilab), Salvado Serra, Jordi (IFIC Valencia), St John, Jason (Fermilab), Strauss, Thomas (Fermilab), Toups, Matthew (Fermilab), Wolbers, Steven (Oregon State University)

Graduate and Postdoctoral Advisors and Advisees:

Conrad, Janet (MIT) – Graduate Advisor
Nygren, David (UT Arlington) – Postdoctoral Advisor

Current and Pending Support: Andrew Brandt

| | | | | |
|---|---|----------------------------------|--|--|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending | | |
| Sponsor: DOE | Award/Identifying Number: 209151 | | | |
| Proposal Title: Research in Experimental Elementary Particle Physics (co-PI) | | | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$2,520,000 | | | | |
| Award Period: 5/01/2014- 3/31/2017 | | | | |
| Number of Person-months per year to be devoted to the project: 1.0 | | | | |
| Abstract: Base funding for the UTA HEP group to support their summer salaries, post-docs, students, and travel. This umbrella proposal encompasses various activities in the energy frontier primarily for ATLAS: data analysis in Higgs and SUSY, trigger development, major leadership roles in computing, etc., with a modest effort in ILC development and leadership; and the intensity frontier ranging from Lariat to Dune. | | | | |

| | | | | |
|--|---|----------------------------------|--|--|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending | | |
| Sponsor: DOE | Award/Identifying Number: 215078 | | | |
| Proposal Title: Development of a Long Life Photomultiplier Tube for High Flux Applications (PI) | | | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$125,000 | | | | |
| Award Period: 6/01/2015- 3/31/2017 | | | | |
| Number of Person-months per year to be devoted to the project: 1.0 | | | | |
| Abstract: This project seeks is concerned with the development of long-life microchannel plate (MCP) photomultiplier tubes (PMTs), capable of high rate operation. Its goals are the optimization of lifetime testing methods including the efficacy of multiple lifetime measurements per device, expedited lifetime measurements, and after-pulsing studies that seek to correlate lifetime with the amount of specific heavy ions. | | | | |

| | | | | |
|---|---|----------------------------------|--|--|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending | | |
| Sponsor: Department of Education | Award/Identifying Number: | | | |
| Proposal Title: Reaching Goals in Physics with GAANN Fellowships (Co-Pi) | | | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$536,688 | | | | |
| Award Period: 9/01/2016- 8/31/2019 | | | | |
| Number of Person-months per year to be devoted to the project: 0.5 | | | | |
| Abstract: This proposal provides funding for physics graduate students with demonstrated need for financial aid, and includes a supervised teaching requirement. | | | | |

| | | | | |
|---|----------------------------------|---|--|--|
| Support: | <input type="checkbox"/> Awarded | <input checked="" type="checkbox"/> Pending | | |
| Sponsor: DOE | Award/Identifying Number: | | | |
| Proposal Title: Research in Experimental Elementary Particle Physics (co-PI) | | | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$ | | | | |
| Award Period: 4/01/2017- 3/31/2020 | | | | |
| Number of Person-months per year to be devoted to the project: 2.0 | | | | |
| Abstract: Renewal of base funding for the UTA HEP group to support their summer salaries, post-docs, students, and travel. This umbrella proposal encompasses various activities in the energy frontier primarily for ATLAS: data analysis in Higgs and SUSY, trigger development, major leadership roles in computing, and TileCAl, with a modest effort in ILC development and leadership; and the intensity frontier ranging from Lariat to Dune. | | | | |

Current and Pending Support: Kaushik De

| | | |
|---|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: | NSF | |
| Award Number: NSF PHY-1119200 | | |
| Title of the Funded Research Project: The U.S. ATLAS Research Program: Empowering U.S. Universities for Discoveries at the Energy Frontier | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$1,611,368 | | |
| Award Period: 10/01/15 - 9/30/16 | | |
| Number of Person-months per year to be devoted to the project by the PI: 2.0 | | |
| <p>Abstract: UTA is a sub-contractor of the NSF US ATLAS Operations program cooperative agreement managed by Columbia University. This cooperative agreement supports M&O, S&C and R&D activities on the ATLAS experiment at the LHC. Activities at UTA include the operation of the SouthWest Tier 2, PanDA software development, US Computing Operations, Analysis support and documentation, and TileCal detector operation and upgrade R&D. These support activities are critical to the success of the ATLAS physics program.</p> | | |

| | | |
|---|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: | BNL Contract #229206 | |
| Title of the Funded Research Project: The U.S. ATLAS Research Program | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$282,000 | | |
| Award Period: 10/01/14 - 9/30/16 (NCE) | | |
| Number of Person-months per year to be devoted to the project by the PI: 0 | | |
| <p>Abstract: UTA receives DOE funding for M&O and S&C activities in support of the US ATLAS Research Program through Brookhaven National Laboratory. Supported activities at UTA include the operation of the SouthWest Tier 2, and TileCal detector operation and upgrade R&D. These support activities are critical to the success of the ATLAS physics program.</p> | | |

| | | |
|--|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: | DOE | |
| Award Number: DOE DE-SC0011686 | | |
| Title of the Funded Research Project: High Energy Physics Base Funding | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$890,000 | | |
| Award Period: 05/01/16 - 04/31/17 | | |
| Number of Person-months per year to be devoted to the project by the PI: 2.0 | | |
| <p>Abstract: This proposal requests support for a program of research in elementary particle physics at The University of Texas at Arlington. We propose studies of the recently discovered Higgs boson, and searches for new particles in nature which may be responsible for dark matter, at the ATLAS Experiment at the European Center for Nuclear Research (CERN) in Geneva, Switzerland, and the Silicon Detector (SiD) at the proposed International Linear Collider. The SiD is a novel concept for a future experiment in particle physics. Our program of work involves detector research and development, and distributed computing innovations. Together, the ATLAS Experiment and SiD can provide a deep understanding of two fundamental forces of nature: electromagnetism and the weak nuclear force, in addition to allowing for the discovery of associated new particles suggested by theory. In a new direction for the group, support is also requested for participation in the future Long Baseline Neutrino Experiment (LBNE), which will explore the masses of the neutrinos that are involved in the weak nuclear interactions and search for low-mass dark matter in the beam, and the ORKA Experiment, that will search for signs of new physics in the rare decays of the K-meson, a particle only produced in high energy collisions. Finally, we propose to carry out theoretical studies of the dark matter that exists in large quantities around and between galaxies, in terms of its interactions with astrophysical objects, and its possible creation in low-energy, high beam intensity experiments.</p> | | |

| | | |
|---|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: DOE | Award Number: DOE DE-SC008635 | |
| Title of the Funded Research Project: Next Generation Workload Management and Analysis System for Big Data | | |
| Award Period: 9/01/12 - 08/31/16 (NCE) | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$746,908 | | |
| Number of Person-months per year to be devoted to the project by the PI: 1.0 | | |
| <p>Abstract: One of the largest scientific collaborations ever assembled, the ATLAS experiment at the Large Hadron Collider (LHC), is designed to explore the fundamental properties of matter for the next decade. An important foundation underlying the impressive success of ATLAS data processing and analysis is the Production and Distributed Analysis (PanDA) workload management system. We propose here a program to develop a generic version of PanDA which can be easily used by many data intensive sciences. With a modest investment of effort, we can enable easy adoption of PanDA by others. We propose generalizing PanDA as a meta-application, providing location transparency of processing and data access, for High Energy Physics, other data-intensive sciences, and a wider exascale community.</p> | | |

| | | |
|--|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: DOE | Award Number: DOE DE-SC0016280 | |
| Title of the Funded Research Project: Big PanDA Workflow Management on Titan for High Energy and Nuclear Physics and for Future Extreme Scale Scientific Applications | | |
| Award Period: 7/01/16 - 06/30/18 | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$1,063,000 | | |
| Number of Person-months per year to be devoted to the project by the PI: 1.0 | | |
| <p>Abstract: Scientific priorities in High Energy and Nuclear Physics continue to serve as drivers of integrated computer and data infrastructure. The lack of scalable and extensible workload management capabilities across heterogeneous computing infrastructure, however presents a barrier to the scientific progress. BigPanDA represents important conceptual advances and novel capabilities to workload management. We propose to deploy and bring into production BigPanDA workflow management techniques on the Oak Ridge Leadership Computing Facility (OLCF) Titan supercomputer. This will significantly and positively impact scientific communities in High Energy and Nuclear Physics, and beyond, for current and future leadership computing facilities.</p> | | |

Current and Pending Support: Andrew P. White

| | | |
|---|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: DOE | Award Number: DOE DE-SC0011686 | |
| Title of the Funded Research Project: | Research in Elementary Particle Physics | |
| Total Award Amount for the Entire Award Period (including indirect costs): | \$890,000 | |
| Award Period: | 04/01/16 - 03/31/17 | |
| Number of Person-months per year to be devoted to the project by the PI: | 2.0 | |
| Abstract: | This project supports the work of the UTA HEP group for the Energy and Intensity Frontiers. For the Energy Frontier, activities for the ATLAS experiment include leadership in computing and software, support for the operation and calibration of the Tile Calorimeter, physics studies in the SUSY and Higgs sectors, and upgrade work on TDAQi and low voltage power supplies. Also for the Energy Frontier, we have a Spokesperson role in the SiD Consortium for the International Linear Collider. Activities for SiD include the development of the design of the SiD Detector, establishing the SiD Consortium as a precursor to a full detector collaboration, promotion and coordination all aspects of detector R&D and physics and performance studies, and representation of SiD within the HEP community nationally and internationally. For the Intensity Frontier efforts include optimization of LBNF beam line for DUNE, Design and construction of proton beam alignment monitor (PBAM) for DUNE, aka hadron monitor, optimization of Optical Coupling for DUNE photo detectors, DUNE 35t Data Analysis and Operations, studies for subGeV Dark Matter, Phase I LArIAT experiment data analyses, MiniBooNE beam dump data analysis, and contributions to Fermilab onsite long baseline experiments. | |

| | | |
|---|---|---|
| Support: | <input type="checkbox"/> Awarded | <input checked="" type="checkbox"/> Pending |
| Sponsor: DOE | Award Number: | |
| Title of the Funded Research Project: | Research in Elementary Particle Physics | |
| Total Award Amount for the Entire Award Period (including indirect costs): | \$????? | |
| Award Period: | 04/01/17 - 03/31/20 | |
| Number of Person-months per year to be devoted to the project by the PI: | 2.0 | |
| Abstract: | | |

Current and Pending Support: Haleh Hadavand

| Current and Pending Support | | |
|--|---|--|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: | NSF ATLAS Project Funds | Award/Identifying Number: NSF PHY-1119200 |
| Title of the Proposal: The U.S. ATLAS Research Program: Empowering U.S. Universities for Discoveries at the Energy Frontier | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$58,000 | | |
| Award Period: 1/30/16-9/30/16 | | |
| Number of Person-months per year to be devoted to the project: 0.0 | | |
| Abstract: UTA is a sub-contractor of the NSF US ATLAS Operations program cooperative agreement managed by Columbia University. This cooperative agreement supports M&O, S&C and R&D activities on the ATLAS experiment at the LHC. Activities at UTA include the operation of the SouthWest Tier 2, PanDA software development, US Computing Operations, Analysis support and documentation, and TileCal detector operation and upgrade R&D. These support activities are critical to the success of the ATLAS physics program. | | |

| Current and Pending Support | | |
|---|----------------------------------|---|
| Support: | <input type="checkbox"/> Awarded | <input checked="" type="checkbox"/> Pending |
| Sponsor: | NSF | Award/Identifying Number: 1654772 |
| Title of the Proposal: Search for Beyond Standard Model Phenomena with Tau Leptons and Higgs | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$810,011 | | |
| Award Period: 2017 – 2021 | | |
| Number of Person-months per year to be devoted to the project: 2 months/year | | |
| Abstract: This proposal investigates search for Beyond Standard Model Phenomena with Tau Leptons and Higgs on the ATLAS experiment. This includes charged Higgs searches in the $t\bar{t}$ final state and a Z boson + X search including leptons and taus in the final states. The Z+X combines five analyses into one scan by using a parametric fit to the background over a range of 4 GeV to a few TeV. This proposal also includes the R&D and production efforts for High Luminosity LHC for the Tile Scintillating Calorimeter low voltage power supplies. | | |

| | | |
|---|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: | Department of Education | Award/Identifying Number: |
| Proposal Title: Reaching Goals in Physics with GAANN Fellowships (Co-Pi) | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$536,688 | | |
| Award Period: 9/01/2016- 8/31/2019 | | |
| Number of Person-months per year to be devoted to the project: | | |
| Abstract: This proposal provides funding for physics graduate students with demonstrated need for financial aid, and includes a supervised teaching requirement. | | |

Current and Pending Support: Jaehoon Yu

| | | |
|--|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: DOE | Award Number: DOE DE-SC0011686 | |
| Title of the Funded Research Project: Research on Elementary Particle Physics | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$890,000 | | |
| Award Period: 04/01/16 - 03/31/17 | | |
| Number of Person-months per year to be devoted to the project by the PI: 2.0 | | |
| <p>Abstract: This proposal requests support for a program of research in elementary particle physics at The University of Texas at Arlington. We propose studies of the recently discovered Higgs boson, and searches for new particles in nature which may be responsible for dark matter, at the ATLAS Experiment at the European Center for Nuclear Research (CERN) in Geneva, Switzerland, and the Silicon Detector (SiD) at the proposed International Linear Collider. The SiD is a novel concept for a future experiment in particle physics. Our program of work involves detector research and development, and distributed computing innovations. Together, the ATLAS Experiment and SiD can provide a deep understanding of two fundamental forces of nature: electromagnetism and the weak nuclear force, in addition to allowing for the discovery of associated new particles suggested by theory. In a new direction for the group, support is also requested for participation in the future Long Baseline Neutrino Experiment (LBNE), which will explore the masses of the neutrinos that are involved in the weak nuclear interactions and search for low-mass dark matter in the beam, and the ORKA Experiment, that will search for signs of new physics in the rare decays of the K-meson, a particle only produced in high energy collisions. Finally, we propose to carry out theoretical studies of the dark matter that exists in large quantities around and between galaxies, in terms of its interactions with astrophysical objects, and its possible creation in low-energy, high beam intensity experiments.</p> | | |

| | | |
|---|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: National Cancer Institute, National Health Institute | Award Number: 1R15CA199020-01A1 | |
| Title of the Funded Research Project: Boosting photo-induced cancer therapies through real-time image guidance | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$415,336 | | |
| Award Period: 04/01/16 - 03/31/19 | | |
| Number of Person-months per year to be devoted to the project by the PI: 0.2 | | |
| <p>Abstract: We propose to use the position-sensitive gas electron multiplier (GEM) detector and advanced spatiotemporal image processing to enable real-time image guided PITs. The GEM technology is a recent advance of the revolutionary digital imaging of gas detectors using multiwire proportional chambers (MWPC), which won Georges Charpak a Nobel Prize for Physics in 1992. The advantages of GEM-based devices include: intrinsic spatial resolution of 50 μm or better; rate capability larger than 1MHz/mm²; easy achievable gains above 10^5; allowing detection of single electrons; efficiency for minimum ionizing particles close to 100%. In addition to its excellent detection performance, the flexibility of GEM can be used for a miniature device with the easy integration of an NIR fiber for therapeutic purpose. In this project, for the first time, we propose to develop a multifunctional device using GEM technology for PITs, called "Beta Image Guided Light-Induced Therapeutic device (BIGLITE)", which can achieve simultaneous imaging and photo-induced therapy.</p> | | |

| | | |
|---|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: NSF | Award Number: 1639157 | |
| Title of the Funded Research Project: Support for Biennial African School of Fundamental Physics 2016 | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$28,215 | | |
| Award Period: 07/01/16 - 06/30/17 | | |
| Number of Person-months per year to be devoted to the project by the PI: 0.01 | | |
| <p>Abstract: This proposal is in support of the forth school in the biennial series. The aim of the school is to build the capacity to harvest, interpret, and exploit the results of current and future physics experiments with particle accelerators, and to increase proficiency in related applications such as medicine, and technologies, such as grid computing. The schools are based on a close interplay between theoretical, experimental and applied physics. The organizing committee consists of a number of people key in the above areas, from both inside and outside Africa. Sub-Saharan Africa is under-represented in sub-atomic physics and this school will serve to provide more opportunities for students to become aware of and to participate in this field.</p> | | |

| | | |
|--|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: Fermi National Accelerator Laboratory | Award Number: N/A | |
| Title of the Funded Research Project: Application for Neutrino Physics Center Fellowship | | |
| Award Period: 07/01/16 - 06/30/17 | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$10,000 | | |
| Number of Person-months per year to be devoted to the project by the PI: 1.5 | | |
| <p>Abstract: The major goals of this project are to understand the behavior of the membrane cryostat, develop and design the beam hadron monitor and optimize the beam line components for DUNE experiment. These funds enable the PI to contribute directly to DUNE experiment through an extended stay at FNAL.</p> | | |

| | | |
|---|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: CNRS, France | Award Number: 1263101510 | |
| Title of the Funded Research Project: MOU for Research on WA105 Dual Phase Detector and DUNE | | |
| Award Period: 10/01/15 - 12/31/17 | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$36,728 | | |
| Number of Person-months per year to be devoted to the project by the PI: 1.0 | | |
| <p>Abstract: The major goal of this project is to contribute to the setup of WA105 and understanding DUNE cryostats.</p> | | |

| | | |
|--|---|----------------------------------|
| Support: | <input checked="" type="checkbox"/> Awarded | <input type="checkbox"/> Pending |
| Sponsor: | Brookhaven National Lab (DOE) | Award Number: |
| Title of the Funded Research Project: Development of SiPM Coupling with Scintillation Counters for Range Stack Detector | | |
| Award Period: 10/01/13 - 09/30/16 | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$46,000 | | |
| Number of Person-months per year to be devoted to the project by the PI: 0.1 | | |
| <p>Abstract: The Range Stack (RS) in ORKA detector plays an essential role in particle identification, especially the pions from Kaon decays from muons. It must be able to measure the energy, range and decay sequence of charged particles emerging from the target with a good position resolution. In addition, it must be able to assist photon veto (PV) detector by identifying them with good efficiency for the photons converting before getting into the PV system. These funds have been repurposed to support LBNE/DUNE photo detector R&D of the same topic.</p> | | |

Current and Pending Support: Jonathan Asaadi

| Current and Pending Support | | | | |
|---|--|---|--|--|
| Support: | <input type="checkbox"/> Awarded | <input checked="" type="checkbox"/> Pending | | |
| Sponsor: NSF | Award/Identifying Number: 1654507 | | | |
| Title of the Proposal: CAREER: A novel fully modular liquid argon neutrino detector for the Deep Underground Neutrino Experiment | | | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$1,114,875 | | | | |
| Award Period: 2017 - 2021 | | | | |
| Number of Person-months per year to be devoted to the project: 2 months/year | | | | |
| Abstract: This proposal puts forward the development of a new modular liquid argon time projection chamber (LArTPC) neutrino detector to be used as a near detector for the Deep Underground Neutrino Experiment (DUNE). The ultimate goal of this project is to demonstrate the feasibility of constructing and operating identical but separate LArTPC modules in a common bath of liquid argon. Each module features a relatively short drift length and at a fully independent TPC with its own readout, light detection system, cryogenics, and services. | | | | |

| | | | | |
|---|----------------------|-----------|--|--|
| Support: | Awarded | X Pending | | |
| Sponsor: DOE | Award Number: | | | |
| Title of the Funded Research Project: Research in Elementary Particle Physics | | | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$????? | | | | |
| Award Period: 04/01/17 - 03/31/20 | | | | |
| Number of Person-months per year to be devoted to the project by the PI: 2.0 | | | | |
| Abstract: | | | | |

Current and Pending Support: Benjamin Jones

| | | |
|---|---------|----------------------|
| Support: | Awarded | X Pending |
| Sponsor: | DOE | Award Number: |
| Title of the Funded Research Project: Research in Elementary Particle Physics | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$????? | | |
| Award Period: 04/01/17 - 03/31/20 | | |
| Number of Person-months per year to be devoted to the project by the PI: 0.0 | | |
| Abstract: | | |

Current and Pending Support: David Nygren

| | | |
|---|---------|----------------------|
| Support: | Awarded | X Pending |
| Sponsor: | DOE | Award Number: |
| Title of the Funded Research Project: Research in Elementary Particle Physics | | |
| Total Award Amount for the Entire Award Period (including indirect costs): \$????? | | |
| Award Period: 04/01/17 - 03/31/20 | | |
| Number of Person-months per year to be devoted to the project by the PI: 0.0 | | |
| Abstract: | | |

FACILITIES AND OTHER RESOURCES

The University of Texas (UTA) is the second largest university in the UT system with around 35,000 students. It is a comprehensive doctoral university located in the Dallas-Ft. Worth metroplex. HEP was selected as one of the first "Organized Research Center of Excellence" at UTA in 2011. PI De is the Director of the ORCE:HEP Center, which also includes faculty from cosmology, astrophysics, space sciences, and computational sciences. The combined synergy of these activities, along with substantial commitment of university resources, provides strong support to the core DOE HEP mission at UTA. Overall, the university has invested over two million dollars to support HEP research activities.

A prime example of UTA's investment in science was the provision of the 120,000 sq.ft. Physics and Chemistry Research Building in 2006. This building houses a high bay area for HEP, our ATLAS Tier 2 center, three detector development laboratories, an HEP conference room, faculty offices, and postdoc and graduate student offices. The building houses an excellent Physics mechanical workshop with the capabilities to manage large scale detector construction..

One finished lab space at UT Arlington's Physics and Chemistry Research Building is a 700 sq. feet lab space with the necessary ventilation for cryogenic experiments to take place. This lab space has recently been completed with a purification and pressure based gas recirculation system for liquid argon detector R&D. This finished lab space also houses desk space, computers, soldering stations, and work space for the undergraduate detector sensor lab as well as a intensity frontier remote operations station. This remote operation station has already been used to take shifts on the LArIAT experiment and is being expanded for remote shift taking on MicroBooNE, SBND, and ICARUS.

A 700 sq foot unfinished lab space adjacent to the purified liquid argon lab and located off the high-bay area has a 3 ton crane for detector construction and assembly. This lab space is located directly adjacent to the UTA physics department machine shop which can be used during detector testing and construction.

In addition to this lab space, the UTA HEP group have retained our previous office suite in Science Hall, and this area has been renovated as the ATLAS Tier 2 operations and visitors area. The lab space in the basement of Science Hall now houses the purified gaseous xenon system as well as a laboratory space for testing Single Molecule Fluorescence Imaging.

UTA also hosts the SouthWest Tier 2 center (SWT2) for ATLAS, which is one of the largest computing centers for ATLAS, providing over 3000 cores and 3 petabytes of storage. The UTA HEP Tier 3 center is co-located with the Tier 2, providing easy access to ATLAS data.

Equipment

9.7 Intensity Frontier Program Budget Justification

Preamble: This section provides the budget justifications for PIs Yu and Asaadi. Detailed description for each of the items is given in Year 1. A canonical cost of living adjustment rate of 4% is applied to all salaries and STEM tuition in the subsequent years. Thus, Year 2 and Year 3 budget justifications will not contain as much details as year 1 but the base rate, any significant changes for each item and total amount are listed.

The tasks carried out on-campus versus off-campus incur substantially different indirect rates 51.5% on-campus versus 26% off-campus the requests are made with the specifications based on the personnel allocation plan specified in the table below.

| PI Name | Category | ON/OFF Campus | Year 1 | Year 2 | Year 3 |
|-----------------|----------|---------------|--------|--------|--------|
| Jaehoon Yu | PI | ON | 100% | 100% | 100% |
| | Postdoc | ON | 50% | 0% | 0% |
| | Postdoc | OFF | 100% | 100% | 100% |
| | GRA | ON | 50% | 50% | 0% |
| | GRA | OFF | 50% | 50% | 100% |
| | UG | ON | 75% | 55% | 75% |
| | UG | OFF | 25% | 25% | 25% |
| Jonathan Asaadi | PI | ON | 100% | 100% | 100% |
| | Postdoc | ON | 50% | 0% | 0% |
| | Postdoc | OFF | 100% | 100% | 100% |
| | GRA | ON | 50% | 50% | 0% |
| | GRA | OFF | 50% | 50% | 100% |
| | UG | ON | 75% | 55% | 75% |
| | UG | OFF | 25% | 25% | 25% |

Table 7: UTA Intensity Frontier Personnel Resource Allocation Plan

9.7.1 PI: Jaehoon Yu

1. Year 1

- **Senior Personnel:** Two months summer salary for Yu is requested at the current rate of \$121,607 per 9 month academic year, equivalent to \$13,512 per month. The total request for this item is \$27,024. The fringe benefit rate is 30% of the request. The indirect rate for this item is the agreed on-campus rate of 51.5.
- **Postdoctoral Researcher:** A request for one postdoctoral fellow is made using the base salary of \$54,000 per annum. The fringe benefit rate is 30% of the request. Since it is anticipated to have the postdoctoral fellow 50% on campus and 50% off campus for tasks at CERN, the indirect rate for this cost is 51.5% on-campus and 26% off-campus for the relevant portion of the cost. Currently Dr. Animesh Chatterjee who has been with the group for 21 months is supported through this request.
- **Graduate Students:** Support for one graduate student is requested at the base rate of \$24,000 per annum. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 50% on campus and 50% off campus for tasks at Fermilab, the indirect rate for this cost is 51.5% on-campus and 26% off-campus for the relevant portion of the cost. Yus student Garrett Brown is currently supported via teaching assistant support in the department. Yu is in the process of recruiting a couple of additional students strategically spaced in time.
- **Undergraduate Students:** Undergraduate students contribute to well defined tasks for the project, such as systematic studies of beam line components for LBNF. A request for two un-

dergraduate support is requested at the base rate \$5,000, equivalent to \$12.5 per hour, 10 hours a week for 40 weeks per year. The total cost for this request is \$10,000. The fringe benefit rate for an undergraduate student is 8.5%. Since it is anticipated to have undergraduate students to spend 75% on campus and 25% off campus for tasks at Fermilab and at CERN during the summer, the indirect rate for this cost is 51.5% on-campus and 26% off-campus for the relevant portion of the cost. The undergraduate students currently in the group are R. Musser, E. Amador, M. Avilla, N. Smith and V. Cervantes.

- **Travel and Cost of Living Adjustment:** We request a total of \$17,950 for travel and COLA support for postdoctoral fellow and graduate student. Of this amount, the travel support \$5,000 is requested to be allocated to the funds at UTA and be subject to on-campus rate of 51.5%. Of the remaining \$12,950, we request \$5,000 to be placed in our groups LSA at Fermilab to minimize the indirect cost at UTA. The remaining \$7,950 is for COLA and is subject to off-campus rate of 26%. COLA request to defray the cost differentials between Arlington, TX and Fermilab or CERN is computed based on \$300 per month for graduate students and \$450 for postdoc for Fermilab. The rate for CERN is higher at \$1,000 per month for students and \$1,600 per month for postdoctoral fellow. These COLA rates are consistent with that of Energy Frontier program to ensure the fair treatment of the group personnel.
- **STEM Tuition:** Graduate student tuition support for one student is request at the rate of \$9,140 per annum. This cost does not incur indirect cost.
- **M&S:** A modest maintenance and services cost of \$2,500 per annum is requested to support various costs. This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$27,557.
- **Total Indirect:** The total indirect cost computed using the proportion of the on-campus (51.5%) and off-campus (26%) described above is \$66,351.
- **Grand Total for Year 1:** The grand total request for year 1 for Yu is \$238,522.

2. Year 2

- **Senior Personnel:** Two months summer salary for Yu is requested after applying a 3% canonical cost of living adjustment, at the rate of \$125,255 per 9 month academic year. The total request for this item is \$27,835. The fringe benefit rate is 30% of the request. The indirect rate for this item is the agreed on-campus rate of 51.5%.
- **Postdoctoral Researcher:** A request for one postdoctoral fellow is made using the base salary of \$55,620 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 30% of the request. Since it is anticipated to the postdoctoral fellow is 100% off campus for tasks 50% at Fermilab and 50% at CERN, the indirect rate for this cost is 26%.
- **Graduate Students:** Support for one graduate student is requested at the base rate of \$24,720 per annum after a 3% canonical cost of living adjustment. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 50% on campus and 50% off campus for tasks at CERN, the indirect rate for this cost is 51.5% on-campus and 26% off-campus for the relevant portion of the cost.
- **Undergraduate Students:** This request for two undergraduate support is at the base rate \$5,000. The total cost for this request is \$10,000. The fringe benefit rate for an undergraduate student is 8.5%. The proportion of undergraduates on and off campus tasks is the same as year 1, and the indirect cost is applied the same as that of year 1.
- **Travel and Cost of Living Adjustment:** We request a total of \$28,300 for travel and COLA support for postdoctoral fellow and graduate student. Of this amount, the travel support \$5,000 is requested to be allocated to the funds at UTA as in year 1. Of the remaining \$23,300, we request \$5,000 to be placed in our groups LSA at Fermilab to minimize the indirect cost at UTA. The remaining \$18,300 is for COLA and is subject to off-campus rate of 26%. COLA request to defray the cost differentials between Arlington, TX and Fermilab or CERN is computed based on \$300 per month for graduate students and \$450 per month for postdoctoral fellow at Fermilab. The rate for CERN is higher at \$1,000 per month for students and \$1,600 per month for postdoctoral fellow. These COLA rates are consistent with that of Energy Frontier program to ensure the fair treatment of the group personnel.

It is worthwhile to note that this increase in travel cost request compared to year 1 is offset by the reduction in the indirect cost, thanks to taking advantage of the 26% off-campus rate.

- **STEM Tuition:** Graduate student tuition support for one student is request at the rate of \$9,140 per annum. This cost does not incur indirect cost.
- **M&S:** A modest maintenance and services cost of \$2,500 per annum is requested to support various costs. This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$28,358.
- **Total Indirect:** The total indirect cost computed using the proportion of the on-campus (51.5%) and off-campus (26%) described above is \$61,488, reduced compared to year 1 due to the allocation of personnel to off-campus.
- **Grand Total for Year 2:** The grand total request for year 2 for Yu is \$247,962.

3. Year 3

- **Senior Personnel:** Two months summer salary for Yu is requested after applying a 3% canonical cost of living adjustment, at the rate of \$129,013 per 9 month academic year. The total request for this item is \$28,670. The fringe benefit rate is 30% of the request. The indirect rate for this item is the agreed on-campus rate of 51.5%.
 - **Postdoctoral Researcher:** A request for one postdoctoral fellow is made using the base salary of \$57,289 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 30% of the request. Since it is anticipated to the postdoctoral fellow is 100% off campus for tasks 50% at Fermilab and 50% at CERN, the indirect rate for this cost is 26%.
 - **Graduate Students:** Support for one graduate student is requested at the base rate of \$25,462 per annum after a 3% canonical cost of living adjustment. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 100% off campus for tasks at Fermilab (50%) and at CERN (50%), the applied indirect is 26% off-campus rate.
 - **Undergraduate Students:** This request for two undergraduate support is at the base rate \$5,305, after a 3% canonical cost of living adjustment. The total cost for this request is \$10,610. The fringe benefit rate for an undergraduate student is 8.5%. The proportion of undergraduates on and off campus tasks is the same as year 1, and the indirect cost is also applied the same as that of year 1.
 - **Travel and Cost of Living Adjustment:** We request a total of \$30,100 for travel and COLA support for postdoctoral fellow and graduate student. Of this amount, the travel support \$5,000 is requested to be allocated to the funds at UTA as in year 1. Of the remaining \$25,100, we request \$5,000 to be placed in our groups LSA at Fermilab to minimize the indirect cost at UTA. The remaining \$20,100 is for COLA and is subject to off-campus rate of 26%. COLA request to defray the cost differentials between Arlington, TX and Fermilab or CERN is computed based on \$300 per month for graduate students and \$450 per month for postdoctoral fellow at Fermilab. The rate for CERN is higher at \$1,000 per month for students and \$1,600 per month for postdoctoral fellow. These COLA rates are consistent with that of Energy Frontier program to ensure the fair treatment of the group personnel.
- It is worthwhile to note that this increase in travel cost request compared to year 1 is offset by the reduction in the indirect cost, thanks to taking advantage of the 26% off-campus rate.
- **STEM Tuition:** Graduate student tuition support for one student is request at the rate of \$9,140. This cost does not incur indirect cost.
 - **M&S:** A modest maintenance and services cost of \$2,500 per annum is requested to support various costs. This request is subject to on-campus indirect rate of 51.5%.
 - **Total Fringe Benefit:** The total cost for the fringe benefit is \$29,184.
 - **Total Indirect:** The total indirect cost computed using the proportion of the on-campus (51.5%) and off-campus (26%) described above is \$59,825, reduced compared to year 1 or 2 due to the greater allocation of personnel on off-campus.
 - **Grand Total for Year 3:** The grand total request for year 3 is \$252,168.

9.7.2 PI: Jonathan Asaadi

1. Year 1

- **Senior Personnel:** Two months summer salary for Asaadi is requested at the current rate of \$84,460 per 9 month academic year, equivalent to \$7,038 per month. The total request for this item is \$18,768. The fringe benefit rate is 30% of the request. The indirect rate for this item is the agreed on-campus rate of 51.5%.
- **Postdoctoral Researcher:** A request of \$27,000 for half postdoctoral fellow is made using the base salary of \$54,000 per annum. This is because we anticipate that it will take about half a year to fill the position. The fringe benefit rate is 30% of the request. Since it is anticipated to have the postdoctoral fellow is going to be located at Fermilab (25%), the indirect rate for this cost is 26% off-campus for the relevant portion of the cost.
- **Graduate Students:** Support for one graduate student is requested at the base rate of \$24,000 per annum. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 50% on campus and 50% off campus for tasks at Fermilab, the indirect rate for this cost is 51.5% on-campus and 26% off-campus for the relevant portion of the cost. Asaadi's student, Zach Williams, is currently supported via teaching assistant support in the department. Asaadi is in the process of recruiting an additional students strategically spaced in time.
- **Undergraduate Students:** Undergraduate students contribute to well defined tasks for the project, LArTPC slow control component work utilizing arduino based micro-controllers. A request for two undergraduate support is requested at the base rate \$5,000, equivalent to \$12.5 per hour, 10 hours a week for 40 weeks per year. The total cost for this request is \$10,000. The fringe benefit rate for an undergraduate student is 8.5%. Since it is anticipated to have undergraduate students to spend 75% on campus and 25% off campus for tasks at Fermilab and at CERN during the summer, the indirect rate for this cost is 51.5% on-campus and 26% off-campus for the relevant portion of the cost. The undergraduate students currently in the group are Ilker Parmaksiz and Nhan Pham.
- **Travel and Cost of Living Adjustment:** We request a total of \$21,400 for travel and COLA support for postdoctoral fellow and graduate student. Of this amount, the travel support \$5,000 is requested to be allocated to the funds at UTA and be subject to on-campus rate of 51.5%. Of the remaining \$16,400, we request \$5,000 to be placed in our group's LSA at Fermilab to minimize the indirect cost at UTA. The remaining \$11,400 is for COLA and is subject to off-campus rate of 26%. COLA request to defray the cost differentials between Arlington, TX and Fermilab or CERN is computed based on \$300 per month for graduate students and \$450 per month for postdoctoral fellow at Fermilab. The rate for CERN is higher at \$1,000 per month for students and \$1,600 per month for postdoctoral fellow. These COLA rates are consistent with that of Energy Frontier program to ensure the fair treatment of the group personnel.
- **STEM Tuition:** Graduate student tuition support for one student is requested at the rate of \$9,140 per annum. This cost does not incur indirect cost.
- **M&S:** A modest maintenance and services cost of \$2,500 per annum is requested to support various costs. This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$16,980.
- **Total Indirect:** The total indirect cost computed using the proportion of the on-campus (51.5%) and off-campus (26%) described above is \$43,644.
- **Grand Total for Year 1:** The grand total request for year 1 is \$173,432.

2. Year 2

- **Senior Personnel:** Two months summer salary for Asaadi is requested after applying a 3% canonical cost of living adjustment, at the rate of \$86,994 per 9 month academic year. The total request for this item is \$19,331. The fringe benefit rate is 30% of the request. The indirect rate for this item is the agreed on-campus rate of 51.5%.
- **Postdoctoral Researcher:** A request for one postdoctoral fellow is made using the base salary of \$55,620 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 30% of the request. Since it is anticipated to the postdoctoral fellow is 100% off campus for tasks 50% at Fermilab and 50% at CERN, the indirect rate for this cost is 26%.

- **Graduate Students:** Support for one graduate student is requested at the base rate of \$24,720 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 50% on campus and 50% off campus for tasks at CERN, the indirect rate for this cost is 51.5% on-campus and 26% off-campus for the relevant portion of the cost.
- **Undergraduate Students:** This request for two undergraduate support is at the base rate \$5,000. The total cost for this request is \$10,000. The fringe benefit rate for an undergraduate student is 8.5%. The proportion of undergraduates on and off campus tasks is the same as year 1, and the indirect cost is also applied the same as that of year 1.
- **Travel and Cost of Living Adjustment:** We request a total of \$28,300 for travel and COLA support for postdoctoral fellow and graduate student. Of this amount, the travel support \$5,000 is requested to be allocated to the funds at UTA as in year 1. Of the remaining \$23,300, we request \$5,000 to be placed in our groups LSA at Fermilab to minimize the indirect cost at UTA. The remaining \$18,300 is for COLA and is subject to off-campus rate of 26%. COLA request to defray the cost differentials between Arlington, TX and Fermilab or CERN is computed based on \$300 per month for graduate students and \$450 per month for postdoctoral fellow at Fermilab. The rate for CERN is higher at \$1,000 per month for students and \$1,600 per month for postdoctoral fellow. These COLA rates are consistent with that of Energy Frontier program to ensure the fair treatment of the group personnel.
It is worthwhile to note that this increase in travel cost request compared to year 1 is offset by the reduction in the indirect cost, thanks to taking advantage of the 26% off-campus rate.
- **STEM Tuition:** Graduate student tuition support for one student is request at the rate of \$9,140. This cost does not incur indirect cost.
- **M&S:** A modest maintenance and services cost of \$2,500 per annum is requested to support various costs. This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$25,807.
- **Total Indirect:** The total indirect cost computed using the proportion of the on-campus (51.5%) and off-campus (26%) described above is \$55,795, reduced compared to year 1 due to the allocation of personnel to off-campus.
- **Grand Total for Year 2:** The grand total request for year 2 for Asaadi is \$231,214.

3. Year 3

- **Senior Personnel:** Two months summer salary for Asaadi is requested after applying a 3% canonical cost of living adjustment, at the rate of \$89,604 per 9 month academic year. The total request for this item is \$19,911. The fringe benefit rate is 30% of the request. The indirect rate for this item is the agreed on-campus rate of 51.5%.
- **Postdoctoral Researcher:** A request for one postdoctoral fellow is made using the base salary of \$57,289 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 30% of the request. Since it is anticipated to the postdoctoral fellow is 100% off campus for tasks 50% at Fermilab and 50% at CERN, the indirect rate for this cost is 26%.
- **Graduate Students:** Support for one graduate student is requested at the base rate of \$25,462 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 100% off campus for tasks at Fermilab (50%) and at CERN (50%), the applied indirect is 26% off-campus rate.
- **Undergraduate Students:** This request for two undergraduate support is at the base rate \$5,000. The total cost for this request is \$10,000. The fringe benefit rate for an undergraduate student is 8.5%. The proportion of undergraduates on and off campus tasks is the same as year 1, and the indirect cost is also applied the same as that of year 1.
- **Travel and Cost of Living Adjustment:** We request a total of \$30,100 for travel and COLA support for postdoctoral fellow and graduate student. Of this amount, the travel support \$5,000 is requested to be allocated to the funds at UTA as in year 1. Of the remaining \$25,100, we request \$5,000 to be placed in our groups LSA at Fermilab to minimize the indirect cost at UTA. The remaining \$20,100 is for COLA and is subject to off-campus rate of 26%. COLA request to defray the cost differentials between Arlington, TX and Fermilab or CERN is computed based on \$300

per month for graduate students and \$450 per month for postdoctoral fellow at Fermilab. The rate for CERN is higher at \$1,000 per month for students and \$1,600 per month for postdoctoral fellow. These COLA rates are consistent with that of Energy Frontier program to ensure the fair treatment of the group personnel.

It is worthwhile to note that this increase in travel cost request compared to year 1 is offset by the reduction in the indirect cost, thanks to taking advantage of the 26% off-campus rate.

- **STEM Tuition:** Graduate student tuition support for one student is request at the rate of \$9,140. This cost does not incur indirect cost.
- **M&S:** A modest maintenance and services cost of \$2,500 per annum is requested to support various costs. This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$26,556.
- **Total Indirect:** The total indirect cost computed using the proportion of the on-campus (51.5%) and off-campus (26%) described above is \$53,961.
- **Grand Total for Year 3:** The grand total request for year 3 is \$234,918.

9.8 Detector R&D Program Budget Justification

Preamble: This section provides the budget justifications for PIs Brandt, Nygren, Asaadi, and Jones related to the Detector R&D proposal. Detailed description for each of the items is given in Year 1. A canonical cost of living adjustment rate of 3% is applied to all salaries in the subsequent years. Therefore, Year 2 and Year 3 of the budget justification does not contain as much details as year 1 and instead the base rates and significant changes for each item and total amount are listed. Only PI Brandt is seeking any salary support from the Detector R&D activities, Nygren, Asaadi, and Jones are seeking support for personnel and M&S.

The tasks carried out on-campus incur an indirect rate of 51.5%.

| PI Name | Category | ON/OFF Campus | Year 1 | Year 2 | Year 3 |
|------------------------|----------|---------------|--------|--------|--------|
| Andrew Brandt | PI | ON | 100% | 100% | 100% |
| | Postdoc | ON | 100% | 100% | 0% |
| | Postdoc | OFF | 0% | 0% | 0% |
| | UG | ON | 0% | 0% | 100% |
| | UG | OFF | 0% | 0% | 0% |
| David Nygren | PI | ON | 100% | 100% | 100% |
| | UG | ON | 100% | 100% | 100% |
| | UG | OFF | 0% | 0% | 0% |
| Jonathan Asaadi | PI | ON | 100% | 100% | 100% |
| | GRA | ON | 100% | 100% | 100% |
| | GRA | OFF | 0% | 0% | 0% |
| Benjamin Jones | PI | ON | 100% | 100% | 100% |
| | GRA | ON | 100% | 100% | 100% |
| | GRA | OFF | 0% | 0% | 0% |

Table 8: UTA Detector R&D Personnel Resource Allocation Plan

9.8.1 PI: Andrew Brandt

1. Year 1

- **Senior Personnel:** One month summer salary for Brandt is requested at the current rate of \$110,646 per 9 month academic year, equivalent to \$12,294 per month. The total request for this item is \$27,024. The fringe benefit rate is 30% of the request. The indirect rate for this item is the agreed on-campus rate of 51.5.
- **Postdoctoral Researcher:** A request for one-half of a postdoctoral fellow is made using the base salary of \$54,000 per annum. The fringe benefit rate is 30% of the request. Since this postdoctoral fellow will be based on campus the indirect rate for this cost is 51.5% for the relevant portion of the cost.
- **Graduate Students:** No support for a graduate student is sought in this year.
- **Undergraduate Students:** No support for undergraduates is sought in this year.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** No STEM tuition is sought in this year.
- **M&S:** A modest maintenance and services cost of \$2,500 per annum is requested to support various costs. This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$11,788.
- **Total Indirect:** The total indirect cost computed using the on-campus (51.5%) rate is \$27,595.
- **Grand Total for Year 1:** The grand total request for year 1 for Brandt is \$81,177.

2. Year 2

- **Senior Personnel:** One month summer salary for Brandt is requested after applying a 3% canonical cost of living adjustment, at the rate of \$113,967 per 9 month academic year. The total request for this item is \$12,663. The fringe benefit rate is 30% of the request. The indirect rate for this item is the agreed on-campus rate of 51.5%.
- **Postdoctoral Researcher:** A request for one-half of a postdoctoral fellow is made using the base salary of \$55,620 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 30% of the request. Since this postdoctoral fellow will be based on campus the indirect rate for this cost is 51.5% for the relevant portion of the cost.
- **Graduate Students:** No support for a graduate student is sought in this year.
- **Undergraduate Students:** No support for undergraduates is sought in this year.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** No STEM tuition is sought in this year.
- **M&S:** A modest maintenance and services cost of \$2,500 per annum is requested to support various costs. This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$12,142.
- **Total Indirect:** The total indirect cost computed using the on-campus (51.5%) rate is \$28,384.
- **Grand Total for Year 2:** The grand total request for year 2 for Brandt is \$83,499.

3. Year 3

- **Senior Personnel:** One month summer salary for Brandt is requested after applying a 3% canonical cost of living adjustment, at the rate of \$117,387 per 9 month academic year. The total request for this item is \$13,043. The fringe benefit rate is 30% of the request. The indirect rate for this item is the agreed on-campus rate of 51.5%.
- **Postdoctoral Researcher:** No support for a postdoctoral researcher is request for this year.
- **Graduate Students:** No support for a graduate student is sought in this year.
- **Undergraduate Students:** This request for support of two undergraduates at the base rate \$5,000 equivalent to \$12.5 per hour, 10 hours a week for 40 weeks per year. The total cost for this request is \$10,000. The fringe benefit rate for an undergraduate student is 8.5% with the indirect rate for this cost is 51.5% on-campus.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** No STEM tuition is sought in this year.
- **M&S:** A modest maintenance and services cost of \$2,500 per annum is requested to support various costs. This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$4,763.
- **Total Indirect:** The total indirect cost computed using the on-campus (51.5%) rate is \$15,607.
- **Grand Total for Year 3:** The grand total request for year 3 for Brandt is \$45,913.

9.8.2 PI: David Nygren

1. Year 1

- **Senior Personnel:** No summer salary for Nygren is requested during this year.
- **Postdoctoral Researcher:** No support for a postdoctoral researcher is request for this year.
- **Graduate Students:** No support for a graduate student is sought in this year.
- **Undergraduate Students:** Undergraduate students contribute to well defined tasks for the project, running various tests, assembly and construction, and data collection. A request for two undergraduate support is requested at the base rate \$5,000, equivalent to \$12.5 per hour, 10 hours a week for 40 weeks per year. The total cost for this request is \$10,000. The fringe benefit rate for an undergraduate student is 8.5%. Since it is anticipated to have undergraduate students on campus, the indirect rate for this cost is 51.5% . The undergraduate students currently in the group are I. Safa, and N. Davachi.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** No STEM tuition is sought in this year.
- **M&S:** No maintenance and services cost is requested in this year.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$850.
- **Total Indirect:** The total indirect cost using the on-campus rate (51.5%) is \$6,695.
- **Grand Total for Year 1:** The grand total request for year 1 for Nygren is \$17,545.

2. Year 2

- **Senior Personnel:** No summer salary for Nygren is requested during this year.
- **Postdoctoral Researcher:** No support for a postdoctoral researcher is request for this year.
- **Graduate Students:** No support for a graduate student is sought in this year.
- **Undergraduate Students:** Undergraduate students contribute to well defined tasks for the project, running various tests, assembly and construction, and data collection. A request for two undergraduate support is requested at the base rate \$5,000, equivalent to \$12.5 per hour, 10 hours a week for 40 weeks per year. The total cost for this request is \$10,000. The fringe benefit rate for an undergraduate student is 8.5%. Since it is anticipated to have undergraduate students on campus, the indirect rate for this cost is 51.5% . The undergraduate students currently in the group are I. Safa, and N. Davachi.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** No STEM tuition is sought in this year.
- **M&S:** No maintenance and services cost is requested in this year.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$850.
- **Total Indirect:** The total indirect cost using the on-campus rate (51.5%) is \$6,695.
- **Grand Total for Year 1:** The grand total request for year 2 for Nygren is \$17,545.

3. Year 3

- **Senior Personnel:** No summer salary for Nygren is requested during this year.
- **Postdoctoral Researcher:** No support for a postdoctoral researcher is request for this year.
- **Graduate Students:** No support for a graduate student is sought in this year.
- **Undergraduate Students:** Undergraduate students contribute to well defined tasks for the project, running various tests, assembly and construction, and data collection. A request for two undergraduate support is requested at the base rate \$5,000, equivalent to \$12.5 per hour, 10 hours a week for 40 weeks per year. The total cost for this request is \$10,000. The fringe benefit rate for an undergraduate student is 8.5%. Since it is anticipated to have undergraduate students on campus, the indirect rate for this cost is 51.5% . The undergraduate students currently in the group are I. Safa, and N. Davachi.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** No STEM tuition is sought in this year.
- **M&S:** No maintenance and services cost is requested in this year.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$850.
- **Total Indirect:** The total indirect cost using the on-campus rate (51.5%) is \$6,695.
- **Grand Total for Year 1:** The grand total request for year 3 for Nygren is \$17,545.

9.8.3 PI: Jonathan Asaadi

1. Year 1

- **Senior Personnel:** No summer salary for Asaadi is requested during this year.
- **Postdoctoral Researcher:** No support for a postdoctoral researcher is request for this year.
- **Graduate Students:** Support for one-half a graduate student is requested at the base rate of \$24,000 per annum. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 100% on campus, the indirect rate for this cost is 51.5%. Asaadis student, Hunter Sullivan, is currently supported via teaching assistant support in the department and will be seeking outside support for the other one-half.
- **Undergraduate Students:** No support for a undergraduate student is sought in this year.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** Graduate student tuition support for one-half student is request at the rate of \$9,140 per annum, coming to \$4,570 for year 1. This cost does not incur indirect cost.
- **M&S:** A maintenance and services cost of \$6,000 per annum is requested to support the running of the cryogenic argon facility. This request will pay for approximately 30 operating days of argon as well as filter material and supplies. This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$1,200.
- **Total Indirect:** The total indirect cost computed using the on-campus (51.5%) rate is \$9,888.
- **Grand Total for Year 1:** The grand total request for year 1 for Asaadi is \$33,658.

2. Year 2

- **Senior Personnel:** No summer salary for Asaadi is requested during this year.
- **Postdoctoral Researcher:** No support for a postdoctoral researcher is request for this year.
- **Graduate Students:** Support for one-half a graduate student is requested at the base rate of \$24,720 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 100% on campus, the indirect rate for this cost is 51.5%.
- **Undergraduate Students:** No support for a undergraduate student is sought in this year.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** Graduate student tuition support for one-half student is request at the rate of \$9,140 per annum, coming to \$4,570 for year 2. This cost does not incur indirect cost.
- **M&S:** A maintenance and services cost of \$6,000 per annum is requested to support the running of the cryogenic argon facility. This request will pay for approximately 30 operating days of argon as well as filter material and supplies. This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$1,236.
- **Total Indirect:** The total indirect cost computed using the on-campus (51.5%) rate is \$10,092.
- **Grand Total for Year 1:** The grand total request for year 2 for Asaadi is \$34,258.

3. Year 3

- **Senior Personnel:** No summer salary for Asaadi is requested during this year.
- **Postdoctoral Researcher:** No support for a postdoctoral researcher is request for this year.
- **Graduate Students:** Support for one-half graduate student is requested at the base rate of \$25,462 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 100% on campus, the indirect rate for this cost is 51.5%.
- **Undergraduate Students:** No support for a undergraduate student is sought in this year.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** Graduate student tuition support for one-half student is request at the rate of \$9,140 per annum, coming to \$4,570 for year 3. This cost does not incur indirect cost.
- **M&S:** A maintenance and services cost of \$6,000 per annum is requested to support the running of the cryogenic argon facility. This request will pay for approximately 30 operating days of

argon as well as filter material and supplies. This request is subject to on-campus indirect rate of 51.5%.

- **Total Fringe Benefit:** The total cost for the fringe benefit is \$1,273.
- **Total Indirect:** The total indirect cost computed using the on-campus (51.5%) rate is \$10,302.
- **Grand Total for Year 3:** The grand total request for year 3 for Asaadi is \$34,867.

9.8.4 PI: Benjamin Jones

1. Year 1

- **Senior Personnel:** No summer salary for Jones is requested during this year.
- **Postdoctoral Researcher:** No support for a postdoctoral researcher is request for this year.
- **Graduate Students:** Support for one-half a graduate student is requested at the base rate of \$24,000 per annum. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 100% on campus, the indirect rate for this cost is 51.5%. Jones student, still to be named, will be supported half via teaching assistant support in the department.
- **Undergraduate Students:** No support for a undergraduate student is sought in this year.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** Graduate student tuition support for one-half student is request at the rate of \$9,140 per annum, coming to \$4,570 for year 1. This cost does not incur indirect cost.
- **M&S:** A maintenance and services cost of \$10,000 in year one is requested. This resource will support the plate materials and associated machining (\$2,000), the necessary flours, solvents, and chemistry supplies (\$2,000), SiPM board development along with associated cables and connectors (\$2,500), the mechanical mounting and frame machining (\$2,500) as well as a supply of gas argon for preliminary tests (\$1,000). This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$1,200.
- **Total Indirect:** The total indirect cost computed using the on-campus (51.5%) rate is \$11,948.
- **Grand Total for Year 1:** The grand total request for year 1 for Jones is \$39,718.

2. Year 2

- **Senior Personnel:** No summer salary for Jones is requested during this year.
- **Postdoctoral Researcher:** No support for a postdoctoral researcher is request for this year.
- **Graduate Students:** Support for one-half a graduate student is requested at the base rate of \$24,720 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 100% on campus, the indirect rate for this cost is 51.5%.
- **Undergraduate Students:** No support for a undergraduate student is sought in this year.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** Graduate student tuition support for one-half student is request at the rate of \$9,140 per annum, coming to \$4,570 for year 2. This cost does not incur indirect cost.
- **M&S:** A maintenance and services cost of \$10,000 in year two is requested. This resource will support further plate materials and associated machining (\$2,000), a supply of purified xenon gas (\$5,000), as well as a supply of gas argon for preliminary tests (\$1,000) the necessary flours, solvents, and chemistry supplies for continued testing (\$2,000). This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$1,236.
- **Total Indirect:** The total indirect cost computed using the on-campus (51.5%) rate is \$12,152.
- **Grand Total for Year 1:** The grand total request for year 1 for Jones is \$40,318.

3. Year 3

- **Senior Personnel:** No summer salary for Jones is requested during this year.
- **Postdoctoral Researcher:** No support for a postdoctoral researcher is request for this year.
- **Graduate Students:** Support for one-half graduate student is requested at the base rate of \$25,462 per annum after applying a 3% canonical cost of living adjustment. The fringe benefit rate is 10% of the request. Since it is anticipated to have the graduate student 100% on campus, the indirect rate for this cost is 51.5%.
- **Undergraduate Students:** No support for a undergraduate student is sought in this year.
- **Travel and Cost of Living Adjustment:** No COLA or travel is sought in this year.
- **STEM Tuition:** Graduate student tuition support for one-half student is request at the rate of \$9,140 per annum, coming to \$4,570 for year 3. This cost does not incur indirect cost.

- **M&S:** A maintenance and services cost of \$5,000 in year three is requested to support the final tests associated with the detector development. This request will pay for a supply of xenon gas (\$5,000). This request is subject to on-campus indirect rate of 51.5%.
- **Total Fringe Benefit:** The total cost for the fringe benefit is \$1,273.
- **Total Indirect:** The total indirect cost computed using the on-campus (51.5%) rate is \$9,787.
- **Grand Total for Year 3:** The grand total request for year 3 for Jones is \$33,361.

Intensity Frontier Data Management Plan

The projects described in this proposal do not produce any data on their own; they merely make use of data generated by the activities related to the MicroBooNE, SBND, ICARUS, and DUNE experiments. All experiments have a data management policy consistent with the Department of Energy's data management plan (found here <http://science.energy.gov/funding-opportunities/digital-data-management/>). This policy conforms with that of the Data Preservation in High Energy Physics (DPHEP) study group, which has described a hierarchy for the types of data that particle-physics experiments produce, and given recommendations for how such data should be preserved for future use.

The types of data these experiments produce include the raw data produced by the detectors, the reconstructed version of the raw data, and simulated events. MicroBooNE, SBND, ICARUS, and DUNE have their own mechanisms for archiving data, including producing both digital and tape storage of raw data, and any data that resides at our institution will merely be a copy of data that is stored permanently by the experiment. Thus, there is no need for this project to separately manage or archive any experimental data. No personally identifiable information is expected to be generated during the execution of the project, and thus no explicit plans to protect confidentiality or personal privacy are used.

The analysis of the experimental data is described in published, peer-reviewed journal articles; summaries of data analyses that are released to the public (often as contributions to conferences); and notes that are circulated internally within I MicroBooNE, SBND, ICARUS, and DUNE as well as public technical notes made available using Fermilab's Technical Publication. The journal articles are archived by the journals themselves, and are also typically available through the arxiv.org e-print archive. The public analysis summaries and internal notes are archived by I MicroBooNE, SBND, ICARUS, and DUNE experiments and are available through Web interfaces. All collaborations encourage frequent and timely publication of results related to the research described in this proposal and thus data generated by the activities described here are expected to have near annual publication releases (through peer reviewed papers and contribution to conferences).