AP CALCULUS AB

Unit 1 • Self Check 1.4 • Lessons 1-7 to 1-9

Self Check 1.4 • Lessons 1-7 to 1-9

Name: _____ Date: _____

Directions

Answer every question. Multiple-choice questions have one best answer. On free-response questions, show your work and justify your answers. Questions labeled **NO CALCULATOR** must be done without a calculator; questions labeled **CALCULATOR** require a graphing calculator. For questions with a graph, read the values you need from the grid.

Question 1

NO CALCULATOR

$\lim_{h\to 0} \frac{\frac{1}{2+h} - \frac{1}{2}}{h}$ equals $k'(2)$ for $k(x) = \frac{1}{x}$. Its value is

(A) $\frac{1}{2}$

(B) $-\frac{1}{4}$

(C) $-\frac{1}{2}$

(D) $\frac{1}{4}$

Question 2

NO CALCULATOR

Where is $f(x) = \sqrt[3]{x}$ continuous but not differentiable?

(A) $x = 1$

(B) $x = 0$

(C) $x = -1$

(D) nowhere

Question 3

NO CALCULATOR

For $k(x) = \frac{1}{x}$ on $[1, 4]$, the MVT gives $c =$

(A) 2.5

(B) $\sqrt{2}$

(C) 2

(D) 3

Question 4 (Free response)

NO CALCULATOR

A differentiable function has $f(1) = 2$, $f(3) = 8$, $f(6) = -1$. Which theorem guarantees each, and why? (a) $f(c) = 5$ for some c in $(1, 3)$. (b) $f'(c) = 3$ for some c in $(1, 3)$. (c) $f'(c) = -3$ for some c in $(3, 6)$.