The Selling closure

Petrus H. Zwart *

Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA.

E-mail: phzwart@lbl.gov

Selling reduction; obtuse superbase; Delone sorts; lattice symmetry; reindexing; pseudo-symmetry

Abstract

A lattice with a right angle has up to sixteen Selling-reduced cells. With the integer matrices between them they form the Selling closure, which holds every lattice symmetry and every operator between alternative reduced cells. Symmetries relate only cells of the same shape, and in primitive lattices with two or three right angles most of the point group relabels no reduced cell, although a search over lattice directions finds every symmetry. Zero conorms fix the members and operators, equal conorms the symmetries, and Delone's 24 sorts are reproduced as the closed patterns of equalities. For a measured cell, near-congruence within a band τ is decided by the widened closure, Selling reduction with its boundary at $-\tau$; when $\lambda_1 > 32\tau$ it is computed completely by zeroing every conorm up to 4τ, enumerating and filtering, and walks from the measured cell miss members; for anisotropic cells the band is a strain, the same in every basis, and projecting onto every face within the band, enumerating and filtering recovers needles and plates. One reduced cell of a new measurement then gives its reindexing operators onto a reference, with a frame-specific guarantee.

1. Introduction

Two questions recur in indexing and data processing. What are the symmetries of a lattice? And are two cells the same lattice, and if so by which integer matrix? Both are usually answered by searching: for symmetry, a scan for near-twofolds within an angular tolerance (Le Page, 1982); for reindexing, a list of small integer matrices, or the operators of the point group just found. The symmetry search is complete within its index bound; the reindexing search is complete only if the list happens to contain the operator. Both are complicated by the same fact: a lattice with a right angle has more than one reduced cell, and the alternatives are related by integer matrices that are not symmetries.

Selling reduction makes the multiplicity explicit. A lattice on a right angle has several obtuse superbases, as Delaunay (1933) noted and Patterson & Love (1957) made practical; Kurlin (2022) proved that there are never more than sixteen and listed them by zero pattern. We take the whole set, with the integer matrices between its members, as the object of study and call it the *Selling closure*; the term is introduced here. What is added to the classical material is: the accounting of which symmetries relabel a reduced cell and which are operators to other cells, with the observation that no symmetry relates cells of different shape; the sorts and their closures obtained from the operators alone; the widened closure of a measured cell with its characterisation superbase by superbase; the idealisation at 4τ with a completeness theorem, and the demonstration that walks from the measured cell are incomplete; and the enumeration of reindexing operators from one reduced cell of a new measurement, with a guarantee in the reference frame. The ingredients are classical. Selling (1874) gave the reduction and the identity of Appendix A; Voronoi (1908) the relevant vectors; Delaunay (1933) the 24 sorts and the observation, made practical by Patterson & Love (1957) and tabulated in *International Tables* (Burzlaff & Zimmermann, 2006), that the reduced

superbase is not unique when a conorm vanishes. Conway & Sloane (1992) supplied the conorm language and the theorem that a superbase is obtuse exactly when its partial sums are the Voronoi vectors, with the five zero patterns as the five types of Voronoi polyhedron. Kurlin (2022) enumerated the obtuse superbases of each pattern. Andrews, Bernstein & Sauter (2019a,b; Andrews & Bernstein, 2023a) cast the same objects as S^6 with its boundary transformations. On the Niggli side the multiplicity of Buerger cells (Buerger, 1960; Gruber, 1973) and the instability of the reduced cell under error (Andrews *et al.*, 1980; Andrews & Bernstein, 1988) are the phenomenon treated here, and Gruber's (1997, 2006) genera refine both the characters and the sorts. Symmetry from the metric tensor with a tolerance is Le Page (1982), Himes & Mighell (1987) and Sauter *et al.* (2004); the operators relating two settings form a coset of the symmetries (Flack, 1987). Closest in aim are the S^6 distance and SELLA of Andrews, Bernstein & Sauter, which carry the boundary images of a cell to measure its distance to each sort, and Oishi-Tomiyasu's error-stable Bravais-lattice determination (2012, 2024), alongside the lattice-character quality indices of Kabsch (1993). She gives a bound on the error under which a finite list of basis transforms is provably enough, which is the role $\lambda_1 > 32\tau$ plays here, and she names the failure that the difference-wall members of Section 4.2 resolve: a noisy cell that is nearly reduced while the true cell is not reduced in the same basis. On the seven measured cells she distributes, the prices of the closure's operators, taken in increasing order, reproduce her ranking of Bravais types (Appendix C).

Section 2 defines the closure and works through one monoclinic cell; Section 3 states the computations; Section 4 gives what the closure contains, how it is computed for a measured cell, and what that gives for reindexing. Proofs are in Appendix A, the conversion of a measurement error into a band in Appendix B, catalogues that the main text only refers to in Appendix C, a description of the operators with no lattice

in Appendix D, and every computed number in the text carries a mark [z*nn*] naming the check that reproduces it (Appendix E).

2. The Selling closure

2.1. Superbases, conorms and the metric tensor

Selling reduction works with four vectors rather than three: the cell vectors a, b, c and $d = -a - b - c$. The four form a *superbase*; any three of them are a basis of the lattice. The six negated pair products

$$p_{da} = -d \cdot a, \quad p_{db} = -d \cdot b, \quad p_{dc} = -d \cdot c, \quad p_{ab} = -a \cdot b, \quad p_{ac} = -a \cdot c, \quad p_{bc} = -b \cdot c$$

are the *conorms* (Selling, 1874; Conway & Sloane, 1992); the Selling parameters of the S^6 formalism (Andrews *et al.*, 2019a,b) are the same numbers with the opposite sign. Conorm vectors are written in this slot order, da, db, dc, ab, ac, bc, throughout, except that the sort patterns of Table 3 use the S^6 order bc, ac, ab, ad, bd, cd so that they can be compared with Andrews & Bernstein (2023b) and Andrews *et al.* (2023). A superbase is *obtuse*, which is to say Selling-reduced, when all six conorms are non-negative. The metric tensor in the basis (a, b, c) is linear in the conorms,

$$\mathbf{G} = \begin{pmatrix} p_{ab} + p_{ac} + p_{ad} & -p_{ab} & -p_{ac} \\ -p_{ab} & p_{ab} + p_{bc} + p_{bd} & -p_{bc} \\ -p_{ac} & -p_{bc} & p_{ac} + p_{bc} + p_{cd} \end{pmatrix}, \tag{1}$$

so the conorms are an alternative set of six cell parameters, and a change to the conorms is a change to the cell with nothing to reconstruct. A conorm that is exactly zero is a right angle between two of the four vectors; this paper calls a zero conorm a *wall*, after the boundary it forms in the space of cells, and the set of slots that vanish the *zero pattern* of the cell. Only six zero patterns are possible if the four vectors are to remain connected by non-zero conorms: no zero; one; two sharing a vector; two disjoint; three forming a triangle; three forming a path. The last two occur in the same lattices — a lattice with three right angles has reduced cells of both kinds — so there

are five kinds of closure, which correspond to the five types of Voronoi polyhedron (Voronoi, 1908; Conway & Sloane, 1992).

2.2. The Selling operators

Two kinds of integer operator act on a superbase, and every operator in this paper is a product of them.

Flips. A flip at the conorm p_{ab} replaces a by $-a$ and adds the old a to the two other vectors: $(a, b, c, d) \mapsto (-a,\ b,\ c+a,\ d+a)$. The new conorms are $p'_{ab} = -p_{ab}$, $p'_{ac} = p_{ad} + p_{ab}$, $p'_{da} = p_{ac} + p_{ab}$, $p'_{bc} = p_{bc} + p_{ab}$, $p'_{db} = p_{db} + p_{ab}$ and $p'_{dc} = p_{dc} - p_{ab}$ (the boundary transformation of Andrews *et al.*, 2019a), so the sum of the six conorms drops by $|p_{ab}|$ when $p_{ab} < 0$; when p_{ab} is exactly zero the flip merely exchanges the slots ac and da, so all reduced cells of one lattice carry the same six conorm values. The vector negated is the first of the pair in the order d, a, b, c: a flip at da, db or dc negates d. Selling reduction applies flips at negative conorms until none remain; the sum of the conorms decreases at each step, so it stops, at an obtuse superbase. A flip at a conorm that is exactly zero carries one obtuse superbase to another obtuse superbase of the same lattice, which is why the reduced cell of a lattice with a right angle is not unique. With the basis (a, b, c) each of the six flips is a fixed 3×3 integer matrix of determinant -1 that is its own inverse (Table 1); a word of flips such as *ab.bc.db* is applied left to right, each flip read in the current labelling, and its matrix is the product $\mathbf{F}_{ab}\mathbf{F}_{bc}\mathbf{F}_{db}$ of the matrices of Table 1, columns being the new basis vectors in the old basis. The flips are the boundary transformations of Andrews *et al.* (2019a), written on the basis rather than on the six Selling parameters.

Table 1. *The six Selling flips as* 3×3 *integer matrices in the basis* (a, b, c), $d = -a - b - c$; *columns are the new basis vectors in the old basis. Each has determinant* -1 *and squares to the identity. The second column gives the flip as a setting change in the notation of crystallographic software; for the flips at da, db, dc the vector d is negated and the new* a, b, c *are listed.*

flip at	new basis	matrix
ab	$(-a,\ b,\ a+c)$	$\begin{pmatrix} -1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$
ac	$(-a,\ a+b,\ c)$	$\begin{pmatrix} -1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$
bc	$(a+b,\ -b,\ c)$	$\begin{pmatrix} 1 & 0 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$
da	$(a,\ -a-c,\ -a-b)$	$\begin{pmatrix} 1 & -1 & -1 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}$
db	$(-b-c,\ b,\ -a-b)$	$\begin{pmatrix} 0 & 0 & -1 \\ -1 & 1 & -1 \\ -1 & 0 & 0 \end{pmatrix}$
dc	$(-b-c,\ -a-c,\ c)$	$\begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ -1 & -1 & 1 \end{pmatrix}$

Swaps. A superbase is an unordered set of four vectors up to a common sign, so one reduced cell can be written in $4! \times 2 = 48$ ways. The 48 reorderings and sign changes are the swaps. They change which three vectors are called a, b, c and in which order, and nothing else. A swap is written by listing which vectors of the superbase become the new d, a, b, c, with a leading sign for the common sign change: $-(a, c, b, d)$ makes the old a the new d, the old c the new a, keeps b, makes the old d the new c, and negates all four. In matrices a swap is a signed permutation of the basis, with the fourth vector d allowed to enter it; the 48 swap matrices are written $\mathbf{Q}$ below.

Products, and symmetries among them. A sequence of flips at zero conorms followed by a swap carries a reduced cell to a reduced cell of the same lattice, possibly one of different shape, and every integer matrix that carries a reduced cell to a reduced cell is such a product (Kurlin, 2022; Lemma A7). A lattice symmetry is an integer matrix g with $g^{\mathsf{T}}\mathbf{G}g = \mathbf{G}$ (Himes & Mighell, 1987); it carries reduced cells to reduced cells,

so it is one of these products. Which products are symmetries depends on the conorm values, not on the operators: a product is a symmetry for exactly those lattices whose conorms it leaves unchanged. The twofold rotations are the symmetries that matter in practice: together with the inversion they generate the point group of any lattice, which is the basis of the Le Page (1982) search, of the tolerance-ranked subgroups of Sauter *et al.* (2004), and of the twinning operators of Lebedev *et al.* (2006).

2.3. The closure, and one cell used throughout

The *Selling closure* $\mathcal{C}(\mathbf{G})$ of a lattice is the set of all its obtuse superbases, each counted once up to swaps, together with the products of flips and swaps between them. That the obtuse superbase is unique except when a conorm vanishes is Delaunay's (1933) observation, set out by Patterson & Love (1957) and by Conway & Sloane (1992); Kurlin (2022) proved that the set has 1, 2, 3, 6 or 16 members according to the pattern of zero conorms and listed them as integer combinations of the reduced cell vectors, the same for every lattice with the same zero pattern (Lemmas 4.1–4.5 of the arXiv version); that flips at zero conorms reach all of them is read off those lists and verified here.[z52] Lemma A0 gives the counts independently. A reduced cell in this paper is always an obtuse superbase, not a Niggli cell: the two coincide for a Niggli cell of type II and differ by one fixed operator for type I (Appendix D), and an orthorhombic lattice has obtuse superbases that are monoclinic-looking cells while its Niggli cell is the orthogonal one. Two members have the same *shape* when their conorm vectors agree up to a swap; in its own basis each member is a cell, and members of different shape are differently shaped cells. The operators can also be characterised with no lattice at all, as the integer matrices that map a face of the orthant of non-negative conorms — the region where all six conorms are ≥ 0, a face being where a chosen set of them is 0 — into that orthant (Appendix D).

Take $a = 30$, $b = 40$, $c = 50\,\text{Å}$, $\beta = 100°$. Its reduced superbase is $M_0 = (d, a, b, c)$ with conorms

$$p_{da} = 639.5, \quad p_{db} = 1600.0, \quad p_{dc} = 2239.5, \quad p_{ab} = 0, \quad p_{ac} = 260.5, \quad p_{bc} = 0\ \text{Å}^2.$$

The two walls are the right angles α and γ at the twofold axis b, present in every primitive monoclinic lattice. No other conorm is small, so the lattice is not near a higher symmetry.[z46] The closure has six members (Table 2); each has two walls, each flip leads to a neighbour, and the six form a ring M_0–M_1–M_5–M_4–M_3–M_2–M_0. All six carry the same six conorm values in different positions, and they fall into three shapes: $\{M_0, M_4\}$ is 30, 40, 50 Å, $\beta = 100°$; $\{M_1, M_3\}$ is 30, 40, 53.7 Å, $\beta = 113.4°$, the alternative choice $a + c$; $\{M_2, M_5\}$ is the third cell choice, on c and $a + c$.[z46]

Table 2. *The Selling closure of the monoclinic cell $a = 30$, $b = 40$, $c = 50$ Å, $\beta = 100°$: six obtuse superbases, each up to the order and common sign of its vectors, the flips at zero conorms that connect them, and where the four symmetries of the point group $2/m$ carry M_0.*[z46]

member	four vectors	zero-conorm flips lead to	symmetries carry M_0 here
M_0	$-a-b-c,\ a,\ b,\ c$	M_1, M_2	1, $\bar{1}$ (swaps of M_0)
M_1	$-b-c,\ -a,\ b,\ a+c$	M_0, M_5	
M_2	$-a-c,\ a+b,\ -b,\ c$	M_0, M_3	
M_3	$a+c,\ b-c,\ -b,\ -a$	M_2, M_4	
M_4	$a-b+c,\ -c,\ b,\ -a$	M_3, M_5	2 along b, m normal to b
M_5	$a-b,\ c,\ b,\ -a-c$	M_1, M_4	

The point group is $2/m$ and each of its four operations carries M_0 to a member (last column of Table 2).[z46] The identity and the inversion are swaps of M_0, visible in it. The twofold sends $(a, b, c) \mapsto (-a, b, -c)$ and so carries $(-a - b - c, a, b, c)$ to $(a - b + c, -a, b, -c)$: not a reordering of M_0 but the member M_4, three flips away, and the mirror does the same. Read in the twofold's order M_4 has the metric tensor of (a, b, c); read in the order of Table 2 it has a different one, with 639.5 and 2239.5 exchanged. As a product, the twofold is the flips ab, bc, db — the path $M_0 \to M_1 \to M_5 \to M_4$, which ends at M_4 in the labelling $(c,\ -a+b-c,\ -b,\ a)$ — followed by the swap $-(a, c, b, d)$, which gives $(a - b + c, -a, b, -c)$; the mirror is the

same flips with the swap (a, c, b, d).[z46] Half of the point group is therefore not visible in the reduced cell: it consists of operators between reduced cells (Figure 1). "Visible" means only that the operation reorders the reduced superbase; a search over lattice vectors, as Le Page's, finds the twofold either way, since it tests directions rather than relabellings.




Fig. 1. The closure of the 100° cell: six members on a ring, joined by flips at the two walls (edge labels are the flip in the labelling of Table 2), coloured by cell shape; the twofold and the mirror carry M_0 to M_4 along three flips and a swap, so they are operators between members rather than relabellings. Placeholder drawing.

2.4. Delone sorts

Delaunay (1933; the same author as Delone) classified lattices into 24 sorts by the pattern of zeros and equalities among the reduced Selling parameters, each sort lying in one Bravais type and some Bravais types comprising several sorts (Delone *et al.*, 1975; Burzlaff & Zimmermann, 1985, 2006); the pattern table and the symbols C1 to

A3 used below are those of Andrews & Bernstein (2023b) and SELLA (Andrews *et al.*, 2023), which also treats the sorts as subspaces of S^6 with their images under the reflections. The sorts are the Selling counterpart of the 44 Niggli lattice characters (Niggli, 1928; de Wolff & Gruber, 1991; de Wolff, 2006): they split a Bravais type by the shape of the cell, not by its symmetry. Since the members and operators of a closure depend only on the zero pattern and its symmetries on the equalities, the sort is the unit for describing closures; Section 3.1 obtains the sorts and their closures from the operators alone.

2.5. Measured cells

A measured cell is never exactly on a wall: a right angle is measured as 90.25° and the conorm is a small number of either sign. The band τ, in Å^2, is the tolerance within which a conorm counts as zero; Appendix B says how a declared error η on the metric tensor, its Frobenius norm in the reduced frame, is converted into τ; the examples use $\tau = 3\eta$. The *widened closure* of a metric at band τ is

$$\mathcal{S}_\tau(\mathbf{G}) = \left\{ F \ : \ \text{all six conorms of } F \text{ under } \mathbf{G} \text{ are } \geq -\tau \right\}, \tag{2}$$

the superbases that Selling reduction would accept with its boundary moved from 0 to $-\tau$. The *depth* of a superbase is the size of its most negative conorm, so it belongs to the widened closure when its depth is at most τ, and its depth is the band at which it enters. Two cells are τ-*congruent* when some reduced superbase of one, carried by an integer operator $\mathbf{P}$ onto the other, has each of its six conorms agreeing to within τ under the two metrics; exact congruence, $\mathbf{P}^\mathsf{T}\mathbf{G}\mathbf{P} = \mathbf{G}'$, is the case $\tau = 0$; the relation is a proximity, not transitive, and is used only between a reference and a new cell. Finally, λ_1 denotes the squared length of the shortest lattice vector, and the *price* of an operator $\mathbf{P}$ between two metrics is the residual $\|\mathbf{P}^\mathsf{T}\mathbf{G}\mathbf{P} - \mathbf{G}'\|_F$ with $\mathbf{G}' \approx \mathbf{P}^\mathsf{T}\mathbf{G}\mathbf{P}$; with $\mathbf{G}' = \mathbf{G}$ it is the amount by which $\mathbf{P}$ fails to be a symmetry. Depth and price

are both in Å^2 but belong to different things: the depth of a cell says whether the measurement supports that cell as a reduced cell of its lattice; the price of an operator says how far the operator is from a symmetry, or from a match between two cells. The measured examples below use the 100° cell of Section 2.3 with β moved to 92°, so that it sits on two walls and near a third; this 92° *cell*, perturbed by a metric error of 14.4 Å^2, is the measured worked cell throughout.

3. Methods

3.1. Closures, sorts and symmetries from the operators

For each zero pattern Z the operator set O_Z is the set of integer matrices whose action on conorm vectors maps the face $\{p \geq 0,\ p_Z = 0\}$ into the orthant, a sign condition on integer matrices that is enumerated exactly (Appendix D); its members are O_Z modulo the 48 swaps. On the face every operator permutes the six conorm slots, so each operator fixes a subspace of conorm vectors, and the sorts are the closed sets of equalities obtained by intersecting those fixed subspaces and keeping the intersections that meet the interior of the face (Proposition A9); the same sorts are obtained in the other direction by fixing a subgroup of operators and taking its fixed subspace. For a sort V the point group is the set of operators that fix V pointwise; the symmetries visible in a reduced cell are those that are swaps of it; two members have the same shape when their six conorms, as linear forms on V, agree up to a slot permutation; and the orbits of the point group on the members are computed by composition. Every entry of Table 3 is the result of this linear algebra on integer matrices, with "visible" read for the member whose zeros and equalities are placed as the pattern is written.[z54] The same counts were also obtained on lattices with arbitrary parameter values as a check.[z46]

3.2. Measured cells: idealise, enumerate, filter

The widened closure of equation (2) is computed as follows.

1. Selling-reduce the measured cell to a superbase F_0.

2. Set every conorm of F_0 that is at most 4τ to zero, all at once, and form the metric tensor $\mathbf{G}^*$ of the result by equation (1). This is the *idealised lattice* Λ^*; it has an exact zero pattern. If $\mathbf{G}^*$ is singular, or λ_1 of the measured cell — the smallest squared length among a, b, c, d and $a+b, a+c, b+c$ of F_0 — is not above 32τ, the hypothesis of Theorem A6 (Appendix A) fails and the procedure stops here: the shortest cell edge must be long compared with the band, which for an isotropic cell means an angular error below about 1.8° (Section 4.2).

3. Enumerate the exact closure of Λ^* by flips at its zeros, keeping each member once up to swaps as integer combinations of F_0 — equivalently, keeping the operators $\mathbf{W}_1, \ldots, \mathbf{W}_m$ from F_0 to each member.

4. Evaluate each member with the *measured* metric tensor and keep those whose six conorms are all at least $-\tau$. Record each member's depth.

The operators of step 3 are integer matrices independent of the metric, so they are the same for Λ^* and for the measured cell; the measured metric enters only in step 4. Section 4.2 states the theorem that this returns exactly $\mathcal{S}_\tau(\mathbf{G})$. The enumeration and the selection are separate: steps 2 and 3 may be run once at the widest band $\tau_{\max}$ for which the idealised lattice exists, and the widened closure at any $\tau \le \tau_{\max}$ is then the members of depth at most τ, with no new enumeration, since the closure of the wider idealisation contains that of every narrower one (Lemma A8) and the completeness theorem needs $\lambda_1 > 32\tau$ only for the band selected at. Each stored member carries its depth and, for a given operator, its price, and either can be the selection criterion at lookup time.

The strain norm. One band on the conorms is the wrong measure for a cell whose edges differ in length: for a needle the error on a conorm between the two long vectors is hundreds of times that between the short ones. The measure used for measured cells is the relative change of the metric tensor,

$$\|\Delta\mathbf{G}\|_{\mathbf{G}} = \|\mathbf{G}^{-1/2}\,\Delta\mathbf{G}\,\mathbf{G}^{-1/2}\|_F, \tag{3}$$

a strain, the affine-invariant norm on positive definite matrices (Pennec *et al.*, 2006). It is unchanged by a change of basis, since $\mathbf{G} \mapsto \mathbf{P}^{\mathsf{T}}\mathbf{G}\mathbf{P}$ and $\Delta\mathbf{G} \mapsto \mathbf{P}^{\mathsf{T}}\Delta\mathbf{G}\mathbf{P}$ leave it as it is, so it is the same in every frame and every operator is an isometry for it; and a perturbation of norm ϵ changes the dot product of any two lattice vectors by at most $\epsilon|u||v|$ (Proposition A11), so an angle by at most about ϵ radians and a squared edge by at most a fraction ϵ, whatever the shape of the cell. The widened set in this norm, $\mathcal{S}_\epsilon(\mathbf{G})$, is the set of superbases F for which the nearest metric that makes F obtuse is within ϵ of $\mathbf{G}$; the distance is a non-negative least-squares problem in the six conorms of F.

Project, enumerate, filter. For a measured cell and a band ϵ:

1. Selling-reduce the measured cell to a superbase F_0.

2. For each of the 127 kinds of operator of the universal set (Appendix D), read the measured cell in that frame, and for each of the 38 faces of the Selling cone (sets of zero conorms whose other slots connect the four vectors) project it onto the face: the nearest metric in the norm (3) with those conorms zero and the others non-negative.

3. For every frame and face whose projection is within 4ϵ, enumerate the exact closure of the projected lattice and carry its members to the coordinates of F_0. Stop if a projected lattice is singular.

4. Keep the members within ϵ of being obtuse for the measured metric. The answer is the union over the projections.

This is Proposition A1 made constructive: the widened set is a union, superbase by superbase, over the lattices near the measurement, and the projections are those lattices. A single projection, onto the most special face within 4ϵ, is not enough: on the needle of Section 4.2 at a small band the members built on $a + c$ belong to a lattice in which $(a + c) \cdot b$, not $a \cdot b$, is zero, which a small change of the angle between b and the long axis produces.[z58] The cost is 127×38 small least-squares problems and a few closures, seconds per cell.

Zeroing as a special case. With one conorm band in the frame of F_0, the projection onto a face sets its conorms to zero and steps 2–4 become the four steps above, whose completeness is Theorem A6. Bands per slot, $\tau_{ij} = \sigma|u_i||u_j|$, applied by zeroing approximate the strain norm and are cheaper, but zeroing in the frame of F_0 alone is not exact (Section 4.2); zeroing on the closure — whenever a member's own frame shows a conorm within four times its band, the conorms of F_0 that enter it are zeroed too — was never wrong in the cases tested, and ordered by the threshold it is a filtration up to a geometric limit at which the idealised lattice becomes singular, but it refuses plates and long needles that the projection returns.[z57]

Ties take no band. Only zeros are idealised. Every sort lies on one zero pattern and every symmetry of every lattice on that pattern is one of its operators (Proposition A9), so a near-equality — the $a \approx b$ of a pseudo-tetragonal cell, or the equal edges behind a pseudo-hexagonal C-centred cell — never adds a candidate: the operator that would be its symmetry is already enumerated, and whether it is accepted is decided at selection, by its price. A flip does subtract, $p'_{dc} = p_{dc} - p_{ab}$ for the flip at p_{ab}, so a tie between opposite slots is a zero in the flipped superbase, the perpendicular diagonals of a rhombus with equal edges; but that superbase has $p'_{ab} = -p_{ab}$, inside

the band only when p_{ab} is itself near zero, so within the widened closure a flip only ever subtracts a near-zero and a tie is a zero only in a conventional cell outside it.

Choosing the band. In the norm (3) the classical tolerances are one kind of number. Changing an angle between two vectors by δ is a strain of $\sqrt{2}\sin\delta$, and changing one edge by a fraction f a strain of about $2f$; Le Page's 3° (Le Page, 1982) and a 5% edge tolerance are strains of 0.074 and 0.10. Enumeration may be generous and selection strict: steps 2–3 at a wide band, selection at the measurement's precision. Ties enter only at selection, through the price of the operator that exchanges the near-equal conorms, which can be measured in the same norm, $\|\mathbf{P}^{\mathsf{T}}\mathbf{G}_R\mathbf{P} - \mathbf{G}_N\|_{\mathbf{G}_N}$, frame-free.

3.3. Reindexing from one reduced cell

A reference R is stored as its widened closure in integer coordinates, the operators $\mathbf{W}_k$ of step 3 that survive step 4. A centred reference, and a centred new cell, are first taken to primitive cells, and the centring matrices are composed with the operator found, which is then rational in the centred bases; the result is carried to the conventional setting of the target space group as a last step, outside the closure. A new cell N is reduced once, to a basis B_N in its own coordinates. Let B_R be the reduction matrix of the reference (its reduced basis in the reference's input basis, columns) and B_N that of the new cell. For each stored $\mathbf{W}_k$ and each of the 48 swaps $\mathbf{Q}$ the candidate operator, from the reference's input basis to the new cell's, is $\mathbf{P} = B_R\mathbf{W}_k\mathbf{Q}\,B_N^{-1}$, so that $(a_N\, b_N\, c_N) = (a_R\, b_R\, c_R)\mathbf{P}$, $\mathbf{G}_N \approx \mathbf{P}^{\mathsf{T}}\mathbf{G}_R\mathbf{P}$ and Miller indices transform as $h_N = \mathbf{P}^{\mathsf{T}}h_R$; candidates are ranked by their price in R's frame. Since a scale error common to all three axes (a detector distance or a wavelength) shifts every conorm in proportion and is larger than any wall for a large cell, the price is computed after the scale that minimises $\|s\,\mathbf{P}^{\mathsf{T}}\mathbf{G}_R\mathbf{P} - \mathbf{G}_N\|_F$ has been applied, and prices are reported as multiples of η. Candidates come in pairs $\pm\mathbf{P}$, and those of determinant -1 are

improper; for a chiral crystal they are discarded. The operators relating two settings of one lattice are one operator composed with each lattice symmetry, a coset of the point group, and near a wall the pseudo-symmetries enlarge it. Membership of the true operator in the candidate set is what is guaranteed; the price does not single it out within that coset, which is a matter for the intensities, as in a twinning test. The band is set from the declared error η: the worked examples use $\tau = 3\eta$, which covers every stored member in the frames that occur there but is not a worst-case guarantee, and the 180-cell test of Section 4.3 uses the coverage bound τ_{cov} of Appendix B; with a covariance for the metric tensor the band can be set slot by slot, which is the route for anisotropic cells (Section 4.2). Programs report standard deviations on $a, b, c, \alpha, \beta, \gamma$ rather than on $\mathbf{G}$; the metric is a smooth function of those six, so its covariance follows from the Jacobian of equation (1) composed with that of the cell parameters, with two cautions: a symmetry-constrained refinement reports zero uncertainty on the constrained parameters and none on the scale, and the error between two data sets is dominated by the detector distance and wavelength, which per-sweep standard deviations omit.

Two tiers. For reindexing, the band of the reference is set Le Page-style: a new cell is accepted as the same lattice if, carried by the operator, it differs from the reference by no more than one cell angle changed by 3°, a strain of $\epsilon_1 = \sqrt{2}\sin 3^\circ = 0.074$ in the norm (3). Membership is then one-sided, as before: if the new cell and the reference differ by at most ϵ_1 under the true operator, the new cell's reduced superbase, carried to the reference, is within ϵ_1 of obtuse for the reference (Corollary A12a), so the reference alone is widened and the new cell is reduced once. The widened closure of the reference is computed exactly from the finite set of Lemma A12, and it lies in an adjacency layer of Theorem A12. In *tier 1* it lies in the first layer, the 127 superbases related to the reduced cell by the universal operator set; this holds

whenever ϵ_1 is below the reference's adjacency radius δ_1, which for cells of ordinary shape exceeds a quarter strain, about 10°, and then the candidates are those superbases filtered, a few dozen at most. In *tier 2*, the needles and plates whose δ_1 is below the band, it lies in the second layer of 2755 superbases, or, beyond δ_2, in the ball of Lemma A12 itself; the same filter applies, and the distances to the 24 sorts (Section 4.2) are reported with the candidates, since such a cell's pseudo-symmetries are part of the question. Either tier returns every operator within the band, ranked by its strain price $\|\mathbf{P}^{\mathsf{T}}\mathbf{G}_R\mathbf{P}-\mathbf{G}_N\|_{\mathbf{G}_N}$ after the best common scale; the tier records how far the closure had to reach, not how far it can be trusted. The procedure is implemented as `agentsg.cell.strain_closure`.

3.4. Verification

Closure memberships were checked against brute force: every basis with coordinates in $[-2,2]$ relative to F_0 (135 408 bases of determinant ±1), completed to a superbase by its fourth vector, tested against equation (2). Every operator between reduced cells has a labelling whose three basis vectors lie in this window (Appendix D), so the window contains every member of every exact closure, and, under the hypothesis of Theorem A6, every member of the widened closure, since those are members of the idealised closure.[z52] Beyond the hypothesis the window is not proved complete for the widened set: on twelve needles and plates at a 3° band whose idealisation is refused, coordinates up to 3 were needed once, and $[-4,4]$ added nothing to $[-3,3]$ in the six cells checked;[z57] the checks with per-slot bands beyond the hypothesis therefore use $[-3,3]$. Test lattices were specified by their conorms. The measured worked cell is 30.02, 39.97, 50.03 Å, 90.25°, 92.05°, 89.80°. It is the 92° cell of Section 2.5, 30–40–50 Å, $\beta=92°$, perturbed by 14.4 Å^2 in the Frobenius norm of its metric tensor. Its six members correspond to M_0 to M_5 of the 100° cell. It is treated at $\tau=43.2$ Å^2,

a band chosen large enough to show the filter at work, and at $\tau = 20\,\text{Å}^2$, within the hypothesis of Theorem A6. The checks are the test files of Appendix E.

4. Results

4.1. What the closure contains

Table 3 gives the closure of each sort. Out of each member there are exactly as many symmetry operators as the point group has elements, and in every sort that number is the order of the holohedry: 48, 24, 16, 12, 8, 4 or 2.[z54] The twofolds, with the inversion, generate the group in every sort, so finding the twofolds suffices.[z46]

Table 3. *The Selling closure of each Delone sort, computed from the operators alone (Section 3.1). Patterns are Selling parameters in the* S^6 *order* $b\cdot c, a\cdot c, a\cdot b, a\cdot d, b\cdot d, c\cdot d$*; equal letters denote equal values and* 0 *a zero. Members are counted up to swaps. "Visible" is the number of symmetries that carry the reduced superbase to itself; the rest carry it to another member. "Shapes" is the number of distinct cell shapes among the members, equal in every case to the number of orbits of members under the symmetries. "Visible" depends on which member one reduces to; values are for the pattern as written.*[z54]

sort	Bravais	pattern	zeros	members	|group|	visible	shapes
C1	cI	rrrrrr	0	1	48	48	1
C3	cF	rr0rr0	2	3	48	16	1
C5	cP	000rrr	3	16	48	12	2
H4	hP	00rrrs	2	6	24	4	1
T1	tI	rrsrrs	0	1	16	16	1
T2	tI	rr0rrs	1	2	16	8	1
T5	tP	000rrs	3	16	16	4	3
R1	hR	rrrsss	0	1	12	12	1
R3	hR	rr0sr0	2	3	12	4	1
O1A	oF	rrsrrt	0	1	8	8	1
O1B	oI	rstrst	0	1	8	8	1
O2	oI	rs0srt	1	2	8	4	1
O3	oI	rs0rs0	2	3	8	8	2
O4	oS	00rsst	2	6	8	4	2
O5	oP	000rst	3	16	8	2	4
M1A	mC	rrsttu	0	1	4	4	1
M1B	mC	rstrsu	0	1	4	4	1
M2A	mC	rs0stu	1	2	4	2	1
M2B	mC	rs0rst	1	2	4	4	2
M3	mC	rs0ts0	2	3	4	4	2
M4	mP	00rstu	2	6	4	2	3
A1	aP	rstuvw	0	1	2	2	1
A2	aP	rs0tuv	1	2	2	2	2
A3	aP	rs0tu0	2	3	2	2	3

Zero pattern and sort fix different parts. Which members exist and which operators

join them is fixed by the zero pattern; which of the operators are symmetries is fixed by the sort, and the abstract group by the Bravais type. Primitive monoclinic, C-centred orthorhombic and primitive hexagonal lattices all have two zeros sharing a vector and six members and differ only in how many operators are symmetries (4, 8, 24). Within one sort the shape of the cell plays no part: the members, the operators and the symmetries are the same, and the lattice supplies only the numbers attached to them.

Symmetry visible in one cell. In the sorts with walls the symmetries that relabel the reduced cell are a minority: 2 of 8 for primitive orthorhombic, 4 of 16 for primitive tetragonal, 12 of 48 for primitive cubic, 4 of 24 for hexagonal, 4 of 12 for rhombohedral R3.[z46] For the twofolds alone: none of the three twofolds of a primitive orthorhombic lattice relabels its reduced cell, nor the one twofold of primitive monoclinic; of the five, nine and seven of primitive tetragonal, cubic and hexagonal, one, three and one do.[z46] The rest are short flip words to other members.

Cell shapes. Two members are related by a symmetry exactly when they have the same shape: a symmetry preserves the metric tensor and hence the conorm multiset, and conversely the change of basis that carries one member's conorm vector onto another's preserves the metric tensor by equation (1). So the number of shapes equals the number of symmetry orbits (last column of Table 3), and no symmetry relates cells of different shape.[z54] Two measurements of the monoclinic lattice that reduce to M_0 and to M_1 report different cell parameters and are related by no symmetry, only by the flip at $a \cdot b$. Ten of the 24 sorts have more than one shape; primitive orthorhombic has four, the orthogonal cell and three monoclinic-looking cells with two right angles, one for each axis traded for a face diagonal. Reduction of a measured orthorhombic cell stops at whichever of the sixteen members its errors favour; twelve of them are monoclinic-looking cells on a face diagonal, and the orthogonal cell is

up to three flips away, reached by no twofold of the lattice. Every extra shape looks like a conventional cell of some class, and the three shapes of a primitive monoclinic lattice are the three cell choices of *International Tables* (Hahn, 2006, Section 2.2.16); the catalogue is Table C1 of Appendix C. The same algebra gives the conditions under which a lattice of one sort becomes a lattice of another. Every such condition of codimension one is a conorm vanishing or two conorms becoming equal; the classical values $c/a = \sqrt{2}$, $b = \sqrt{3}\,a$ and $a = c$ in a monoclinic cell are among them, and the monoclinic conditions come in threes, one per cell choice (Table C2, Appendix C).[z50] The two kinds behave differently under error. A near-equality, such as $c/a \approx \sqrt{2}$ or a C-centred monoclinic cell one equality from body-centred tetragonal, changes no zero pattern: the member set is already complete and the pseudo-symmetry appears as a priced operator. A near-wall, such as a C-centred monoclinic cell one zero from primitive tetragonal, adds members, and those are what the idealisation of Section 3.2 recovers.

4.2. Near-congruence within a band

The measured problem. For the measured worked cell of Section 3.4, with $\alpha = 90.25°$, the conorms p_{ab} and p_{bc} are small numbers of either sign. The flips that gave M_1 and M_2 now give superbases with a slightly negative conorm, Selling reduction rejects them, and the exact closure has one member instead of six; the twofold and the mirror, which were operators to M_4, now lead to a cell that is not in the set.

What the widened closure is. Equation (2) names no starting cell, order of operations or frame, so $\mathcal{S}_\tau(\mathbf{G})$ is the same whoever computes it. By Proposition A1 (Appendix A) a superbase belongs to it exactly when it is a reduced superbase of some lattice within τ of $\mathbf{G}$ in that superbase's own six conorms. The band is judged superbase by superbase: the widened closure contains every reduced cell of every lattice consistent with the

measurement, but not every reduced cell of such a lattice, since a lattice within τ of $\mathbf{G}$ at one of its cells can be further away at another. The cell at which reduction with a $-\tau$ boundary stops is not well defined: from 400 random starting bases, reduction of the measured worked cell at $\tau = 43.2\,\text{Å}^2$ stopped at six different cells and of a noisy orthorhombic cell at sixteen, every one a member of $\mathcal{S}_\tau$, but on a noisy cubic cell at $\tau = 10\,\text{Å}^2$ the 400 restarts found 15 of the 16 members.[z45] The set has to be computed, not sampled.

Near-congruence is decided by the widened closure, widened on one side only. If two cells are τ-congruent through a superbase F reduced for the new cell, then F, carried to the reference, has all conorms at least $-\tau$ and so lies in the reference's widened closure (Corollary A1a, the overlap statement): the exact closure of the new cell, indeed its single reduced superbase, already meets the widened closure of the reference. Near-congruence is therefore decided without a search over operators and without widening the new cell: compute $\mathcal{S}_\tau(\mathbf{G}_R)$ once, reduce the new cell once, and pair its reduced superbase with each stored member. The stored set contains the connecting cell whenever one exists.

The computation is complete. If $\lambda_1 > 32\tau$, the four steps of Section 3.2 return exactly $\mathcal{S}_\tau(\mathbf{G})$ (Theorem A6 and Corollary A6a, Appendix A): every superbase within the band is an exact member of the idealised lattice, so enumerating the idealised closure and filtering cannot miss it. On every test lattice and band, including walls placed between τ and 4τ, the procedure returned the brute-force set.[z44] The condition is sufficient, not sharp: for a shortest cell edge of 30 Å it allows bands up to $28.1\,\text{Å}^2$, i.e. with $\tau = 3\eta$ a metric error of about $9\,\text{Å}^2$; the worked cell's error of $14.4\,\text{Å}^2$ is beyond that, so at $\tau = 20\,\text{Å}^2$ the procedure is an instance of the theorem and returns six members, while its exactness at 43.2 and $60\,\text{Å}^2$ is an observation.[z44] Idealising all small conorms together is required by the proof and loses nothing, since the closure

of any partial idealisation lies inside that of the full one (Lemma A8); the threshold 4τ is what the proof requires, and on dense two-wall grids 2τ was already exact while τ was not.[z46]

Scale. The condition $\lambda_1 > 32\tau$ is a ratio of two squared lengths, so it does not depend on the size of the cell: for an angular error σ the band on a conorm between vectors of lengths $|u_i|$, $|u_j|$ is about $\sigma|u_i||u_j|$, and for a cell with edges all near a the condition is $\sigma \lesssim 1/32$ rad, about $1.8°$, for a 5 Å cell as for a 300 Å one. That is the band the proof covers, not the limit of the procedure: on lattices of five zero patterns with walls placed at random, the construction was exact against brute force at every band from $\lambda_1/32$ to $\lambda_1/6$, about $9.5°$, in every case in which the idealised lattice existed, the only failures at wide bands being the singular idealisations that step 2 reports.[z55]

Anisotropy. What the condition $\lambda_1 > 32\tau$ responds to is anisotropy: for a needle or a plate one conorm band must cover the conorms between the long vectors, and the condition fails at a few thousandths of a degree. Bands per slot, $\tau_{ij} = \sigma|u_i||u_j|$, applied by zeroing in the frame of F_0, fail. The needle $5.58 \times 5.21 \times 131.97$ Å, $\gamma = 91.12°$, has $p_{ab} = 0.57$ Å^2 and $p_{ac} = p_{bc} = 0$; at $\sigma = \sin 0.064°$ the per-slot condition holds, p_{ab} survives because its own band $\sigma|a||b|$ is small, and 6 superbases are returned where the widened set has 12: on the slot $(a + c, b)$ of the missing members the same dot product carries a band 25 times larger and has changed sign.[z57] The strain norm has no frame-dependent band. Projecting onto every face within 4ϵ and filtering (Section 3.2) returned the brute-force set for the needle at every band from $\epsilon = 0.0005$ to $\sin 3° = 0.052$; for a $100 \times 100.1 \times 5$ Å plate and a $150 \times 120 \times 4.8$ Å plate, $\beta = 90.4°$, up to $\epsilon = 0.052$, where their widened sets have 20 and 28 members, more than any one closure; and for 240 random needles, plates, isotropic and monoclinic cells at $\epsilon = \sin 0.1°$ to $\sin 3°$, in every case, with sets of up to 36 members — against brute force over coordinates in $[-2, 2]$, and in $[-3, 3]$ for the plates, the needle and the

37 random cells with ten or more members. No projected lattice was singular, so nothing was refused.[z58] Zeroing on the closure, on the same 240, was exact in 124 and refused the other 116.[z57] A per-slot band is still the right description of the error (Appendix B): it is what the strain norm implies for each conorm (Proposition A11).

How far the proof reaches. Theorem A12 (Appendix A) proves the procedure complete up to the adjacency radius $\delta_1(\mathbf{G})$, the strain at which a superbase that shares no lattice with F_0 first comes within reach, and the filtered second layer L_2 complete up to $\delta_2(\mathbf{G})$; both are computed exactly from the finite set of Lemma A12. Table 4 gives them. For cells of ordinary shape no such superbase lies within a quarter strain, about 10°, so the first layer suffices for any band in use; for needles and plates δ_1 falls as the ratio of the shortest to the longest edge, about $\sqrt{2}$ times that ratio, and the second layer roughly doubles it. The two plates at $\sin 3^\circ = 0.052$ lie between δ_1 and δ_2: the first layer misses members there, the second contains them all, and the procedure, whose output lies in the second layer, returned them.[z59] The number of layers a band needs, the least k with $\delta_k > \epsilon$, is therefore the measure of a cell's difficulty, in place of a single angular limit.

Table 4. *Adjacency radii in the strain norm (3): δ_1, the smallest strain at which a superbase sharing no lattice with the reduced cell becomes obtuse, and δ_2 for the second layer. The search is capped at* 0.25. *For reference, a* 3° *angular tolerance is a strain of* 0.052 *and* 1° *of* 0.017.

cell	δ_1	δ_2
$30 \times 40 \times 50$ Å, $\beta = 92^\circ$ (Section 3.4)	> 0.25	> 0.25
needle $10 \times 10 \times 30$ Å	> 0.25	> 0.25
needle $10 \times 10 \times 100$ Å	0.139	> 0.25
needle $5.58 \times 5.21 \times 132$ Å, $\gamma = 91.12^\circ$	0.056	0.111
plate $100 \times 100.1 \times 5$ Å	0.050	0.141
plate $150 \times 120 \times 4.8$ Å, $\beta = 90.4^\circ$	0.043	0.100

Walks from the measured cell miss members. The natural alternative starts from the measured reduced cell and flips across any conorm within some band of zero. Table 5 compares four versions with the procedure above on 80 cells with walls at awkward positions (five zero patterns, including a plate-shaped orthorhombic cell), constructed

as worst cases; all four miss members. The cause is a member reachable only through a non-member: with walls at 1.2τ and 0.5τ, the flip across the first gives a conorm of -1.2τ, outside the band, and only a second flip, across the other wall, reaches a member of depth 0.7τ. On the idealised lattice both walls are zero and both flips cost nothing. Idealising at τ rather than 4τ fails on the same configuration.[z44] The most negative conorm of such a member is the difference of two small conorms, and we call it a *difference-wall member*; in the configurations checked they appear when one small conorm $z \leq \tau$ and another lies between τ and $z+\tau$; they carry no symmetry, and they are ordinary alternative reduced cells to which a second measurement of the lattice can reduce; a reference set built by walking from its own reduced cell never contains them.[z46]

Table 5. *Four walks on the measured metric, and the procedure of Section 3.2, against brute force on 80 cells with walls at awkward positions (orthorhombic, cubic, plate, monoclinic and hexagonal zero patterns;* $\tau = 0.1$ *and* 0.5 *of* $\lambda_1/32$*).*

procedure	cells with a wrong set
flip wherever $\lvert p\rvert \leq \tau$, keep only cells within the band	13
flip wherever $\lvert p\rvert \leq \tau$, filter at the end	13
flip wherever $\lvert p\rvert \leq 2\tau$, keep only cells within the band	13
flip wherever $\lvert p\rvert \leq 2\tau$, filter at the end	2
idealise at 4τ, enumerate the exact closure, filter	0

The idealised lattice is not a symmetry assignment. Setting the small conorms to zero resembles the most generous choice a Le Page search makes, but Le Page returns a point group, asserted, whereas the idealisation returns the closure of a lattice Λ^* whose symmetries are candidates with prices, and the filter removes the members the measurement does not support. Nor is the band the tolerance of such a search: Le Page's 3° is a price, not a band. Pseudo-symmetries are read as prices of operators in the closure of the idealised lattice, which holds operators of every price, and at the proved band a wall within $4\tau = \lambda_1/8$, about 7° from a right angle, is already idealised and its symmetries offered. Idealisation imposes zeros but no equalities, so a measured primitive monoclinic cell whose only small conorms are its two walls

idealises to a primitive monoclinic lattice; only when the small conorms are the walls of a higher class does Λ^* lie in that class, and then its extra symmetries are the pseudo-symmetries, each priced.

Distance to each sort. The same norm measures how far a measured cell is from each of the 24 sorts: project its conorms, in each of the 6096 frames of the universal set, onto the sort's subspace of Proposition A9 (zeros, equalities, the rest non-negative) and take the smallest distance. Projection onto Bravais-lattice subspaces is the approach of Andrews & Bernstein (2014) in G^6 and of Oishi-Tomiyasu (2012) in Selling coordinates; here the frames are the complete finite set of the closure and the norm is the same in every frame. An oP cell $40.0 \times 40.3 \times 80$ Å is 1.06% from tP, the strain of its 0.75% edge mismatch; a C-centred monoclinic cell $50 \times 86.8 \times 40$ Å, $\beta = 90.4°$, is 1.04% from hP; an F cell $100 \times 100.2 \times 99.9$ Å is 0.43% from cF; and the needle of Section 4.2, $\gamma = 91.12°$, is 2.76% from oP, $\sqrt{2}\sin 1.12°$, the same as for an isotropic cell with that angle.[z58] Along every specialisation of Table C2 the distance never decreases, so the sorted list is a walk up Table C2, and the 24 distances are the measurement's distances to every symmetry it could be assigned, in one unit. With a covariance for the metric in place of the norm, each distance is a χ^2 with as many degrees of freedom as the sort has constraints, a test where Le Page's 3° is a fixed angle.




Fig. 2. The 92° cell: the sixteen members of its idealised closure (Table 6) by depth under the measured metric; the six of the ring (blue) enter below 9 Å^2, the ten members carrying the near-orthorhombic symmetry between 45 and 62.5 Å^2, and the band τ selects among them. Placeholder drawing.

A worked filter. Table 6 and Figure 2 show the procedure on the measured worked cell. The reduced superbase has conorms $(49.5, 1588.9, 2444.7, 4.2, 847.5, 4.5)$ Å^2. At $\tau = 43.2$ Å^2 the three conorms below $4\tau = 172.8$ are set to zero and equation (1) gives an orthorhombic idealised lattice whose reduced superbase has edges 29.1, 39.9, 57.4 Å with $\beta = 120.5°$ — the reduced cell of an orthorhombic lattice is one of its monoclinic-looking shapes; the orthogonal cell is $29.1 \times 39.9 \times 49.4$ Å on $(a, b, a + c)$. Its closure has sixteen members; each row gives one as a word in the flips of Table 1 and its six conorms under the *measured* metric. A member is kept when its smallest conorm is at least $-\tau$ and nothing is inherited: for $4.5 \le \tau < 8.7$ Å^2 the member *ab.bc.db* is kept while its parent *ab.bc* is rejected, and for $45.0 \le \tau < 53.7$ the member *da.ab.bc* while its parent *da.ab* is rejected.[z45] Deeper members carry integer combinations of the small conorms — $-(49.5 + 4.2)$, $-(49.5 - 4.5)$ — and those combinations decide when a member enters. One enumeration serves every band up to 211 Å^2, where 4τ

reaches the next conorm, as Section 3.2 allows: six members for $8.7 \le \tau < 45.0$, among them the band $\tau = 20$ within the theorem, the six of the unperturbed cell at the same integer coordinates, and all sixteen from 62.5.[z45] The member entering at 45.0 carries the near-orthorhombic twofolds, so a member's depth is the band at which the pseudo-symmetry it carries becomes visible, and its symmetries' prices say how far the measurement is from it.

Table 6. *The filter on the measured worked cell. Each row is a member of the idealised lattice's closure, named by a word in the flips of Table 1 from the reduced cell e; the six numbers are its conorms under the measured metric tensor, slots da, db, dc, ab, ac, bc, in Å^2; "enters at" is its depth. At $\tau = 43.2$ the six members above the line are kept.*

word	*da*	*db*	*dc*	*ab*	*ac*	*bc*	enters at
e	49.5	1588.9	2444.7	4.2	847.5	4.5	0
ab	851.7	1593.1	2440.6	−4.2	53.7	8.7	4.2
bc	45.0	8.7	2449.3	1593.4	852.0	−4.5	4.5
ab.bc.db	2453.8	−4.5	847.5	1606.3	57.9	−4.2	4.5
ab.bc	842.9	4.5	2449.3	1601.8	62.5	−8.7	8.7
bc.db	2458.0	−8.7	53.7	1602.1	843.3	4.2	8.7
da	−49.5	2494.3	1638.4	53.7	897.0	−45.0	49.5
da.ab.bc	942.0	−45.0	1593.4	2556.7	12.9	−8.7	45.0
da.ab.ac	−49.5	2543.8	1588.9	954.9	−4.2	12.9	49.5
da.ab	950.7	2548.0	1584.7	−53.7	4.2	8.7	53.7
ab.ac	49.5	1539.3	2494.3	905.4	−53.7	62.5	53.7
bc.db.dc	45.0	2511.7	−53.7	1548.4	897.0	57.9	53.7
ab.ac.bc.ac	1610.5	959.1	2565.5	−4.2	−8.7	−53.7	53.7
da.ab.bc.ac	2569.6	−57.9	1606.3	954.9	−12.9	4.2	57.9
ab.ac.bc	−12.9	967.8	2556.7	1601.8	8.7	−62.5	62.5
ab.bc.ac	1664.2	−57.9	2511.7	905.4	−62.5	53.7	62.5

4.3. Reindexing from one reduced cell

The operators relating two settings of one lattice are one operator composed with the lattice symmetries, and each is an operator between a member of one closure and a member of the other. For the $\beta = 92°$ cell and its perturbation at $\tau = 43.2\,\text{Å}^2$, pairing the six members of one with the six of the other, each pair in its best relative order, gives twelve pairs at a price of $14.4\,\text{Å}^2$ and nothing else below $1378\,\text{Å}^2$; the twelve pairs are four distinct operators, one operator composed with each element of $2/m$, at a price equal to the perturbation. At $\tau = 60\,\text{Å}^2$ the members carrying the near-orthorhombic symmetry are present and add ten pairs at $150.7\,\text{Å}^2$, so the band

decides which pseudo-symmetries are offered.[z39]

A new cell needs no closure of its own, by Corollary A1a, for the reduced superbase it returns. For fifteen references near walls of five zero patterns, each with twelve new cells carrying a random metric error of declared size and a random reindexing, the new cell's single reduced superbase was always a member of the reference's widened closure at the correct integer coordinates, and the true operator was always among the candidates: 180 of 180, with the band set from the declared error by the coverage bound of Appendix B.[z47]

Completeness of the stored set matters. A new cell constructed so that its only reduced superbase is a difference-wall member of an orthorhombic corner was matched against two stored sets of the reference. The set built as in Section 3.2 offers candidates at 7.83, 8.01 and 8.84 Å^2, each with its negative; the set built by idealising only at τ offers the first two and has lost the third, which is $\pm\mathbf{I}$, the true operator.[z46] The lowest-priced candidate is found either way, through a pseudo-symmetry, but it is not the true operator, which is the one lost; price orders the coset, it does not identify its true member.

The band that makes the true operator one of the candidates can be stated in advance, under a hypothesis on the error expressed in the reference frame. Suppose the new cell's metric tensor, carried to the reference frame by the true operator, differs from the reference's by at most η in Frobenius norm. Let V be the sum of the squared lengths of the new cell's three reduced vectors and $\lambda_{\min}$ the smallest eigenvalue of the reference metric tensor. Then the band $\tau = \sqrt{2}\,\eta\,V/(\lambda_{\min}-\eta)$ suffices, provided $\eta < \lambda_{\min}$ and $\lambda_1 > 32\tau$ (Proposition A10, Appendix A). Appendix B gives a bound that does not refer to the operator. The guarantee held in all 180 cases but is conservative, with median 13η against a band near 3η that was in fact enough.[z47]

Benchmark. The places where the closure predicts trouble — the shapes, lattices a

small distance from a specialisation condition, cells whose only reduced superbase is a difference-wall member, anisotropic cells, the sorts O3, M2B and M3 — are collected as a benchmark of 780 measured cells with their true operators (Appendix C). On it a symmetry-only reindexing proposes the true operator in 54% of the cases, the exact closure in 96%, and a walk at τ in 97%, with an incomplete set on every difference-wall case. The procedure of Section 3.2 declines 8 anisotropic cases at step 2 and proposes the true operator in all of the other 772. The same 780 cases show where the Niggli cell is unstable: the measured cell's Niggli cell is a different cell from the reference's in 45, nearly all at the body-centred orthorhombic sort O3 and its neighbours.[z49] The benchmark is built where the closure predicts trouble, so these are rates on constructed cases, not in the wild.

5. Discussion

Where reindexing has been hard. The usual pipeline reduces a cell, idealises it to the highest Bravais lattice within a tolerance (Le Page, 1982; the tolerance-ranked subgroups of Sauter *et al.*, 2004) and standardises to a conventional cell; two data sets so treated agree unless they idealise to different lattices or fall on opposite sides of a tie in the standardisation. What the symmetry step cannot do is propose the flip at $a \cdot b$, because it is a symmetry of nothing; that flip is what separates c from $a + c$, the orthogonal cell from its monoclinic-looking shapes, and the two reduced cells of the sort O3 that two measurements return. The closure adds exactly those operators, with the same pseudo-symmetries, each priced, and its gain over the pipeline is the completeness statement, not a higher hit rate on ordinary cells. Operators that the closure does not contain are outside its scope: sublattice and superlattice relations (a doubled axis from weak odd layers; determinant ± 2 or more), the obverse and reverse settings of a rhombohedral lattice, and the choice of hand, which are separate

steps. Within one sort, two correct descriptions of a lattice can differ in shape. Each is a reduced cell that Selling reduction, or Niggli reduction with a tolerance (Křivý & Gruber, 1976; Grosse-Kunstleve *et al.*, 2004), can return when a near-zero conorm falls on the other side. No symmetry relates them, so a comparison that applies only the symmetries of the cell it holds cannot reconcile them. The monoclinic c versus $a + c$ choice is the oldest instance; the monoclinic-looking cells of orthogonal lattices are the same phenomenon. Across sorts it is the walls: a cell measured within τ of a wall idealises into a sort with more members, each entering at a depth and carrying a priced pseudo-symmetry. The widened closure holds both kinds of alternative with a depth for each, so under the condition of Theorem A6 a reindexing is a lookup rather than a search whose completeness depends on the cell it started from. The instability of the Niggli cell (Andrews *et al.*, 1980; Andrews & Bernstein, 1988, 2014) is the same phenomenon from the Niggli side only in part. Andrews & Bernstein locate it at every boundary of the Niggli cone: the sign boundary between type I and type II, the ties $a = b$ and $b = c$, and the special conditions. Only the boundaries at a zero scalar product are Selling walls; a tie is not, and the type I/II change gives a cell of the same lengths and supplementary angles, which in the closure is one shape and a swap. So the closure explains the wall part: in the benchmark the measured cell's Niggli cell is a different cell from the reference's in 45 of 780 cases, nearly all at the body-centred orthorhombic sort O3 and its neighbours, where the reduced cell has two right angles, the three members have two shapes (Table 3), and the Niggli reduction of two measurements, deciding a near-zero quantity by its sign, returns one shape or the other — two cells related by a flip and by no symmetry. The Selling-reduced cell shares the wall instability, as Andrews *et al.* (2019b) note; what the paper adds is not stability of any one cell but completeness of the set of alternatives. Table 7 collects the traps, where each is found and what resolves it.

Table 7. *The reindexing traps as the closure finds and resolves them.*

trap	mechanism	found	resolved by
monoclinic c versus $a + c$	flip at the wall $a \cdot b$; two shapes, no symmetry between them	Sec. 2.3, Table 2	both shapes are members
oP, tP, cP reported as a monoclinic cell	twelve of sixteen members lie on face diagonals	Sec. 4.1, Table 6	idealised closure; the orthogonal cell and its twofolds enter at a depth
pseudo-merohedry	a near-equality leaves the member set complete (priced operator); a near-wall adds members	Sec. 4.1, Table C2	prices; idealisation for the walls
Niggli-cell instability at a wall (O3 and neighbours)	two shapes of one closure chosen by the sign of a near-zero	Sec. 4.3	both shapes stored; ties and the type I/II change are not walls
difference-wall member	reachable only through a non-member	Sec. 4.2, Table 5	idealise at 4τ, not walk
needles and plates	conorm error $\sigma\|u_i\|\|u_j\|$ differs by slot and by frame	Sec. 4.2, App. B	strain norm: project onto the faces within 4ϵ, union of closures
choice within the coset	the metric cannot separate an operator from its product with a (pseudo-)symmetry	Sec. 4.3	not by the metric: intensities

What is proved and what is checked. Proved in Appendix A, from Selling's identity, the Conway–Sloane characterisation of obtuse superbases, Kurlin's enumeration and two integer constants obtained by exhaustive enumeration, are the uniqueness and union characterisation of the widened closure, the completeness of idealise–enumerate–filter when $\lambda_1 > 32\tau$, the generation of the closure by flips and swaps, the nesting of partial idealisations, and the reindexing guarantee. The sorts as closed sets of equalities (Proposition A9) is a definition whose agreement with Delone's 24 is checked by enumeration, and the counts of Table 3 are exact computations on the operators. Kurlin's enumeration is cited from a preprint; his counts are reproduced here by brute force for every sort. For anisotropic cells, proved is that the strain norm bounds every conorm (Proposition A11) and that a per-slot filter is exact once the idealisation is (Theorem A6′); proved also is that the widened set in the strain norm is finite (Lemma A12) and that project–enumerate–filter returns it below the adjacency radius $\delta_1(\mathbf{G})$, and filtering the second layer below $\delta_2(\mathbf{G})$ (Theorem A12), and that

widening the reference alone suffices in this norm too (Corollary A12a); checked in every case tested but not proved is that the procedure stays exact between δ_1 and δ_2, and that zeroing on the closure is exact whenever it does not stop (Conjecture A6″), and refuted by the needle of Section 4.2 is the idealisation in the frame of F_0 alone. Checked but not proved are also that idealising at 2τ suffices, that $[-3, 3]$ contains the widened set beyond the hypothesis, and that the true operator is found at the band set from the error rather than only at the conservative guarantee.

The frame. Statements about the widened closure are in terms of τ, a band on conorms, and need no frame. Converting an error on the metric tensor into τ does, because the Frobenius norm of an error depends on the basis; Appendix B gives the exact relation between a metric error and each conorm shift, the worst-case factors by zero pattern (up to $2\sqrt{3}$ in some cubic frames, so that $\tau = 3\eta$ is not a worst-case guarantee everywhere), a frame-free bound and the covariance route.

6. Conclusions

The Selling closure of a lattice contains, in one finite and complete set, every symmetry and every operator between alternative reduced cells; a single reduced cell shows only the symmetries that relabel it, and no symmetry relates cells of different shape. For a measured cell, near-congruence within a band is decided by the widened closure, which is unique and is computed completely, when $\lambda_1 > 32\tau$, by idealising at 4τ, enumerating and filtering; the condition is a ratio of lengths, independent of the scale of the cell, and its failure for strongly anisotropic cells is seen before any operator is proposed. Reindexing a new cell onto a reference then reads one operator from that set, from a single reduced cell of the new cell, and when the error, written in the reference frame, is below the smallest eigenvalue of the reference metric and the band it implies satisfies the same condition, the true operator is guaranteed to be among those read.

In a pipeline this means: store each reference once as its widened closure, at most sixteen integer matrices with their depths; on a new data set, Selling-reduce once and pair with the stored members; this is a complete candidate generator, upstream of the intensity-based choice within the coset, and it supplies what the symmetry step of a pipeline cannot, the operators that are symmetries of nothing. Two failures announce themselves at step 2 — a singular idealised lattice, or a shortest edge too short for the band — and brute force over coordinates in $[-3, 3]$ is the fallback; for measured cells the band is a strain, the same in every frame, by default one cell angle changed by 3°; the reference's widened closure is computed exactly, in the first adjacency layer of 127 superbases for cells of ordinary shape and in the second, of 2755, for needles and plates (two tiers), and the new cell's reduced superbase is paired with it; one is silent, a band narrower than the frame requires, which drops members without warning, so the band should be set from the coverage bound of Appendix B rather than from 3η alone.

Appendix A
Proofs

Terms. Write a lattice vector in any basis, $x = n_1 v_1 + n_2 v_2 + n_3 v_3$ with integer n_i, and record whether each coordinate is even or odd. This sorts the lattice into eight *parity classes*; two vectors are in the same class exactly when their difference is twice a lattice vector, so the classes do not depend on the basis (the classes of $\Lambda/2\Lambda$). One class is the doubled vectors; the other seven are used here. A *partial sum* of a superbase is the sum of some of its four vectors; since the four add to zero, there are seven partial sums up to sign, one in each of the seven classes, and for the monoclinic superbase $(-a-b-c, a, b, c)$ they are a, b, c, $a+b$, $a+c$, $b+c$ and $a+b+c$. The fact

used is that a superbase is obtuse exactly when each of its partial sums is a shortest vector of its parity class, the Voronoi vectors of the lattice (Voronoi, 1908; Conway & Sloane, 1992, whose vonorms are these minima; see also Kurlin, 2022, Lemmas 2.4, 2.7 and A.1). The *excess* $e_c(F) = |v_c(F)|^2 - \mu_c$ of a superbase F in class c is the squared length of its partial sum $v_c(F)$ in that class minus the squared length μ_c of the shortest vector of the class; an obtuse superbase has zero excess in every class. A partial sum is written $w = \sum s_i v_i$ with $s \in \{0,1\}^4$ marking which vectors it adds; a vector of the same parity class has coordinates $s + 2k$ with k integer. Every lattice vector $x = \sum c_i v_i$ satisfies Selling's identity

$$|x|^2 = \sum_{i<j} p_{ij}\,(c_i - c_j)^2, \tag{4}$$

whatever the signs of the conorms, because the four vectors sum to zero. For a superbase F and two metrics, $\delta_F(\mathbf{G}, \mathbf{G}')$ is the largest change among F's six conorms.

Lemma A0 (window). Let F_0 be obtuse for a lattice whose non-zero conorms connect the four vectors, and F any obtuse superbase of the same lattice. Then every vector of F is a partial sum of F_0 whose superbase coordinates are $s_i + 2k_i$ with $s \in \{0,1\}^4$, $k_0 = 0$ and $|k_i| \le 3$. *Proof.* Every partial sum of F is a shortest vector of its parity class, as is the partial sum of F_0 in that class, so their excess vanishes; by Lemma A2 the excess is $4\sum p^0_{ij}\gamma_{ij}$ with every term non-negative, so $\gamma_{ij} = 0$ on every non-zero slot, i.e. $|k_i - k_j| \le 1$ along a connected graph on four vertices, and with $k_0 = 0$ every $|k_i| \le 3$. □ Enumerating the superbases of such vectors whose six pairwise products are non-positive on the whole face gives 1, 2, 3, 6 and 16 members for the five kinds of zero pattern, all with basis entries in $\{-1, 0, 1\}$, independently of Kurlin's lists;[z56] this is also the bound that makes the brute-force window of Section 3.4 and the enumeration of Appendix D complete.

Proposition A1 (characterisation superbase by superbase).

$$\mathcal{S}_\tau(\mathbf{G}) = \{F : F \in \mathcal{C}(\mathbf{G}') \text{ for some } \mathbf{G}' \text{ with } \delta_F(\mathbf{G}, \mathbf{G}') \le \tau\}.$$

Proof. ($\supseteq$) If F is obtuse for $\mathbf{G}'$ and $\delta_F(\mathbf{G}, \mathbf{G}') \le \tau$, every conorm of F under $\mathbf{G}$ is at least $-\tau$. ($\subseteq$) If $F \in \mathcal{S}_\tau(\mathbf{G})$, the metric whose conorms in the frame of F are $\max(p_{ij}, 0)$ exceeds $\mathbf{G}$ by a positive semidefinite term, so it is positive definite; F is obtuse for it, at δ_F equal to the depth of F. Each negative conorm must move at least to zero, so no nearer metric makes F obtuse. □

Corollary A1a (overlap: widening one side suffices). Let $\mathbf{G}_R$ and $\mathbf{G}_N$ be τ-congruent through the operator $\mathbf{P}$ and the superbase F reduced for $\mathbf{G}_N$, i.e. the six conorms of F under $\mathbf{G}_N$ and of $\mathbf{P}F$ under $\mathbf{G}_R$ agree to within τ. Then $\mathbf{P}F \in \mathcal{S}_\tau(\mathbf{G}_R)$. In particular the reduced superbase that reduction returns for a new cell, carried to the reference by the true operator, is a member of the reference's widened closure, and the new cell needs no widening of its own; a new cell with several reduced superbases (exact walls) satisfies the hypothesis for the one returned, which for a measured cell is generically unique. *Proof.* The conorms of F under $\mathbf{G}_N$ are non-negative, so those of $\mathbf{P}F$ under $\mathbf{G}_R$ are at least $-\tau$, which is membership in $\mathcal{S}_\tau(\mathbf{G}_R)$ by equation (2). □

Lemma A2 (class excess identity). Let F be any superbase, $w = \sum s_i v_i$ a partial sum and w' a vector of its parity class with coordinates $s + 2k$. Then $|w|^2 - |w'|^2 = -4\sum p_{ij}\gamma_{ij}$, with $\gamma_{ij} = \Delta k(\Delta k + d)$, $\Delta k = k_i - k_j$, $d = s_i - s_j$; each γ_{ij} is a non-negative integer, zero exactly when $\Delta k \in \{0, -d\}$. *Proof.* Equation (4), $(d + 2\Delta k)^2 - d^2 = 4\Delta k(\Delta k + d)$, and $t(t + d) \ge 0$ for integers t and $d \in \{-1, 0, 1\}$.

Lemma A3 (class gap). Let Λ^* be obtained from F_0 by setting its conorms $\le \theta$ to zero, be positive definite, and let m^* be the smallest conorm of F_0 above θ. A superbase not obtuse for Λ^* has $e_c \ge 4m^*$ in some class. *Proof.* In the frame of F_0, with w the partial sum of F_0 in class c — a shortest vector of its class for $\mathbf{G}$ and for Λ^* — Lemma A2 gives $e_c(F) = 4\sum p^0_{ij}\gamma_{ij}$, a sum of non-negative terms, and the

excess for Λ^* is the same sum over the slots above θ. If those terms vanish in all seven classes, every partial sum of F is a shortest vector of its class for Λ^* and F is obtuse for it; otherwise some $\gamma_{ij} \geq 1$ multiplies a $p^0_{ij} \geq m^*$.

Lemma A4 (small excess in the band). If $\lambda_1 > 32\tau$ and every conorm of F is at least $-\tau$, then $e_c(F) \leq 16\tau$ in every class. *Proof.* Call a slot large if $p_{ij} > 8\tau$, and let N be the negative slots. The large slots connect the four vectors, since otherwise a split crossed only by slots with $p_{ij} \leq 8\tau$ has a partial sum of squared length at most $32\tau < \lambda_1$. With w' a shortest vector of the class, Lemma A2 in the frame of F gives $0 \leq e_c(F) = 4\sum_N |p_{ij}|\gamma_{ij} - 4\sum_{p_{ij}>0} p_{ij}\gamma_{ij}$, so $\sum_{\text{large}} p\gamma \leq \tau \sum_N \gamma$. If a large γ were positive, Lemma A5(ii) would give $8\tau \sum_{\text{large}} \gamma < \tau \sum_N \gamma \leq 8\tau \sum_{\text{large}} \gamma$; so the large γ vanish, and Lemma A5(i) gives $e_c \leq 4\tau \cdot 4$.

Lemma A5 (two constants on four points). Let L be a set of slots connecting the four vectors and N disjoint from L. (i) If $\gamma_{ij} = 0$ on L, then $\sum_N \gamma_{ij} \leq 4$. (ii) If $\sum_L \gamma_{ij} > 0$, then $\sum_N \gamma_{ij} \leq 8 \sum_L \gamma_{ij}$. Both bounds are attained. *Proof.* In (i) $|\Delta k| \leq 1$ on L, so with $k_0 = 0$ all $|k_i| \leq 3$, and the maximum is a finite enumeration over the 38 connected slot sets, the sets N and the 14 partial sums. In (ii) the ratio is enumerated for $|k_i| \leq 15$; beyond, $Q = \sum_L \Delta k^2 > 75$, and $\sum_N \gamma \leq \rho Q + \sqrt{|N|\rho Q}$, $\sum_L \gamma \geq Q - \sqrt{|L|Q}$, with $\rho \leq 3 + 2\sqrt{2}$ the largest generalised eigenvalue of $\sum_N \Delta k^2$ against $\sum_L \Delta k^2$, itself computed over the 38 patterns, keep it below 8 for every pattern; the lemma is therefore computer-assisted. Equality in (ii): $L = \{01, 02, 13\}$, $N = \{03, 12, 23\}$, $s = (0, 0, 1, 1)$.[z44]

Theorem A6. Let F_0 be obtuse for $\mathbf{G}$, $\lambda_1 > 32\tau$, and Λ^* obtained by setting every conorm of F_0 that is at most 4τ to zero. Then Λ^* is positive definite and $\mathcal{S}_\tau(\mathbf{G}) = \{F \in \mathcal{C}(\Lambda^*) : p_{ij}(F; \mathbf{G}) \geq -\tau\}$. *Proof.* If the slots of F_0 above 4τ did not connect the four vectors, some partial sum would have squared length at most $16\tau < \lambda_1$. The inclusion $\supseteq$ is the definition. For $F \in \mathcal{S}_\tau(\mathbf{G})$, Lemma A4 gives $e_c(F; \mathbf{G}) \leq 16\tau$ in every

class. The excess of F for Λ^* is the same sum of non-negative terms with the zeroed slots dropped, so $e_c(F;\Lambda^*) \le e_c(F;\mathbf{G}) \le 16\tau < 4m^*$, since the smallest surviving conorm m^* exceeds 4τ; Lemma A3 then makes F obtuse for Λ^*. □

Corollary A6a (operator form). Under the hypotheses of Theorem A6 let $\mathbf{W}_k$, $k = 1,\dots,m$, be the integer changes of basis from the basis of F_0 to the members of $\mathcal{C}(\Lambda^*)$. Then

$$\mathcal{S}_\tau(\mathbf{G}) = \{F_0\mathbf{W}_k : \min_{ij} p_{ij}(F_0\mathbf{W}_k;\mathbf{G}) \ge -\tau\},$$

where $F_0\mathbf{W}_k$ is the superbase whose basis is that of F_0 transformed by $\mathbf{W}_k$. *Proof.* The members of $\mathcal{C}(\Lambda^*)$ are the superbases $F_0\mathbf{W}_k$; the $\mathbf{W}_k$ are integer matrices independent of the metric, so the same superbases read with $\mathbf{G}$ are the right-hand side of Theorem A6. □

Theorem A6′ (bands per slot through one band). Let F_0 be obtuse for $\mathbf{G}$, $\sigma \ge 0$, and $S_\sigma(\mathbf{G}) = \{F : \ p_{ij}(F;\mathbf{G}) \ge -\sigma|u_i||u_j| \ \forall ij\}$, with u_i the vectors of F. If $\hat\tau$ satisfies $\lambda_1 > 32\hat\tau$ and $\sigma|u_i||u_j| \le \hat\tau$ on every negative slot of every $F \in S_\sigma(\mathbf{G})$, and Λ^* is F_0 idealised at $4\hat\tau$, then $S_\sigma(\mathbf{G}) = \{F \in \mathcal{C}(\Lambda^*) : \ p_{ij}(F;\mathbf{G}) \ge -\sigma|u_i||u_j| \ \forall ij\}$. *Proof.* Every $F \in S_\sigma(\mathbf{G})$ has all conorms at least $-\hat\tau$, so $S_\sigma(\mathbf{G}) \subseteq \mathcal{S}_{\hat\tau}(\mathbf{G})$, which by Theorem A6 is the set of members of $\mathcal{C}(\Lambda^*)$ with conorms at least $-\hat\tau$; intersecting with $S_\sigma(\mathbf{G})$, and noting that a member passing the per-slot filter is in $S_\sigma(\mathbf{G})$ and so has its negative slots within $\hat\tau$, gives the statement. □ The filter is therefore exact once the idealisation contains the right zeros. For a needle, $\hat\tau$ must cover $\sigma\ell_{\mathrm{long}}\ell_{\mathrm{short}}$, so the theorem reaches $\sigma < \ell_{\mathrm{short}}/(32\ell_{\mathrm{long}})$, about $\ell_{\mathrm{long}}/\ell_{\mathrm{short}} < 2.4$ at a 3° idealisation.

Conjecture A6″ (zeroing on the closure). If the zeroing on the closure of Section 3.2 terminates with Λ^* positive definite and $\lambda_1 > 32\,\sigma \max_{kl\in Z} \ell_k\ell_l$ over the zeroed slots Z, with ℓ_k the lengths of the vectors of F_0, the idealise–enumerate–filter procedure with bands per slot returns $S_\sigma(\mathbf{G})$. The proof of Theorem A6 carries over except at Lemma A3, which in the per-slot setting must be read in the frame of a member F'

of $\mathcal{C}(\Lambda^*)$ adjacent to the outsider, where the fixpoint guarantees that the surviving conorm exceeds four times the band; what remains is a weighted form of Lemma A5 comparing vector lengths in adjacent frames. The idealisation in the frame of F_0 alone fails it (Section 4.2).[z57]

Proposition A11 (the strain norm bounds every conorm). If $\|\mathbf{G}^{-1/2}\Delta\mathbf{G}\,\mathbf{G}^{-1/2}\|_F \le \epsilon$, then $|u^{\mathsf{T}}\Delta\mathbf{G}\,v| \le \epsilon\,|u|\,|v|$ for all vectors u, v, lengths in $\mathbf{G}$; in particular every conorm of every superbase changes by at most $\epsilon|u_i||u_j|$, and $\mathcal{S}_\epsilon(\mathbf{G})$ lies inside the per-slot set with $\sigma = \epsilon$. *Proof.* With $X = \mathbf{G}^{-1/2}\Delta\mathbf{G}\,\mathbf{G}^{-1/2}$, $u^{\mathsf{T}}\Delta\mathbf{G}\,v = (\mathbf{G}^{1/2}u)^{\mathsf{T}}X(\mathbf{G}^{1/2}v)$, at most $\|X\|_2|u||v| \le \|X\|_F|u||v|$. □ The norm is invariant under every change of basis, so Proposition A1 holds in it unchanged.

Lemma A12 (the widened set is finite). If F is within $\epsilon < 1$ of obtuse for $\mathbf{G}$ in the norm (3), every vector u of F satisfies $|u|^2_{\mathbf{G}} \le \frac{1+\epsilon}{1-\epsilon}\,\mu_c(\mathbf{G})$, with μ_c the squared length of F_0's partial sum in the parity class of u. *Proof.* Let $\mathbf{G}'$ within ϵ make F obtuse. By Proposition A11 with $u = v$, $(1-\epsilon)|u|^2_{\mathbf{G}} \le |u|^2_{\mathbf{G}'} \le (1+\epsilon)|u|^2_{\mathbf{G}}$ for every lattice vector. The vectors of F are shortest in their classes for $\mathbf{G}'$, and those of F_0 for $\mathbf{G}$, so $|u|^2_{\mathbf{G}'} \le |v^0_c|^2_{\mathbf{G}'} \le (1+\epsilon)\mu_c(\mathbf{G})$ and $|u|^2_{\mathbf{G}} \le |u|^2_{\mathbf{G}'}/(1-\epsilon)$. □ The lattice vectors in these seven balls are finitely many and computable from $\mathbf{G}$, so $\mathcal{S}_\epsilon(\mathbf{G})$ is enumerated exactly, with no window to assume.[z59]

Theorem A12 (completeness up to the adjacency radius). Call two superbases *adjacent* when some positive definite metric makes both obtuse; a superbase adjacent to F_0 is $\mathbf{W}F_0$ with $\mathbf{W}$ in the universal set $\mathcal{U}$ of Appendix D, since both are members of that lattice's closure (Lemma A7), and $\mathcal{U}$ is closed under relabelling. Let $L_1 = \mathcal{U}F_0$, $L_k = \mathcal{U}L_{k-1}$ (127 and 2755 superbases up to relabelling for $k = 1, 2$), and let $\delta_k(\mathbf{G})$ be the smallest distance in the norm (3) from $\mathbf{G}$ to a metric for which a superbase outside L_k is obtuse. If $\epsilon < \min(\delta_k(\mathbf{G}), \frac{1}{4})$, then $\mathcal{S}_\epsilon(\mathbf{G})$ is the set of members of L_k within ϵ of obtuse, and for $k = 1$ the project–enumerate–filter procedure of Section 3.2

returns it. *Proof.* A superbase in $\mathcal{S}_\epsilon(\mathbf{G})$ is obtuse for a metric within $\epsilon < \delta_k$ of $\mathbf{G}$, so it lies in L_k; the converse is the definition. For $k = 1$, $F = \mathbf{W}F_0 \in \mathcal{S}_\epsilon$ is obtuse for a metric $\mathbf{G}'$ within ϵ, whose zero pattern Z in the frame of F is one of the 38 faces; the frame of F is one of the frames projected onto, the projection onto Z is at distance at most $\epsilon < 4\epsilon$ from $\mathbf{G}$, and F is the base superbase of its closure, so F is enumerated, and it passes the filter. No projection within $4\epsilon < 1$ is singular, since a singular metric differs from $\mathbf{G}$ by at least 1 in the norm (3) along its null vector, so nothing is refused. □ The radii are computed exactly from the balls of Lemma A12 at $\epsilon = \frac{1}{4}$. For $k = 2$ the procedure's output lies in L_2 and was equal to $\mathcal{S}_\epsilon$ in every case tested, but only the filtering of L_2 is proved.[z59]

Corollary A12a (overlap in the strain norm). Let $\mathbf{P}$ be an integer operator with $\|\mathbf{P}^\mathsf{T}\mathbf{G}_R\mathbf{P} - \mathbf{G}_N\| \le \epsilon$, measured in the norm (3) at $\mathbf{P}^\mathsf{T}\mathbf{G}_R\mathbf{P}$, and let F be obtuse for $\mathbf{G}_N$. Then $\mathbf{P}F$ is within ϵ of obtuse for $\mathbf{G}_R$, so $\mathbf{P}F \in \mathcal{S}_\epsilon(\mathbf{G}_R)$. *Proof.* F is obtuse for $\mathbf{G}_N$, which is within ϵ of $\mathbf{P}^\mathsf{T}\mathbf{G}_R\mathbf{P}$; carried back by $\mathbf{P}$, which is an isometry of the norm, $\mathbf{P}F$ is obtuse for a metric within ϵ of $\mathbf{G}_R$. □ Measured at $\mathbf{G}_N$ instead, the two distances agree to first order in ϵ; the exact affine-invariant distance would make them equal.

Lemma A7 (generation). Every operator between members of the closure, and every symmetry, is a sequence of flips at zero conorms followed by a swap. *Proof.* The members are reached from F_0 by flips at zero conorms (Kurlin, 2022, Lemmas 4.1–4.5), so flips and swaps connect all members. A symmetry maps F_0 to an obtuse superbase, which is a member; as a matrix it is the flip path from F_0 to that member followed by a swap.

Lemma A8 (nesting). If $Z' \subseteq Z$ are sets of conorms of F_0 set to zero, every member of the closure for Z' is a member of the closure for Z. *Proof.* Assume the lattice idealised on Z is positive definite. The members for Z' are reached from F_0 by words

of flips, each at a slot that is zero at that point of the word (Lemma A7); induct on the word. A slot that is zero for Z' is zero for Z, and a flip at a slot zero for both acts on both conorm vectors as the same transposition of two slots (Section 2.2), so after each flip the superbase is obtuse for both lattices and the next flip is again at a slot zero for both. Hence every member for Z' is a member for Z.[z46]

Proposition A9 (sorts as closed sets of equalities). Fix a zero pattern Z, let O_Z be the operators of its closure and, for $\mathbf{W} \in O_Z$, let $L_{\mathbf{W}}$ be its linear action on conorm vectors (Appendix D), which on the face $\{p \geq 0,\ p_Z = 0\}$ is a permutation of the slots. For a subspace V of the face put $\mathrm{Sym}(V) = \{\mathbf{W} \in O_Z : L_{\mathbf{W}} = \mathrm{id} \text{ on } V\}$ and $\mathrm{Fix}(H) = \{p : L_{\mathbf{W}} p = p \ \forall\, \mathbf{W} \in H\}$. A lattice with conorm vector p on the face has point group $\mathrm{Sym}(\{p\})$, every symmetry being in O_Z by Lemma A7, and the lattices with the symmetries of a generic point of V fill $\mathrm{Fix}(\mathrm{Sym}(V))$. We take as definition that a sort is a maximal set of lattices with the same kind of closure and the same symmetry operators; then the sorts of kind Z are the generic points of the closed subspaces $V = \mathrm{Fix}(\mathrm{Sym}(V))$ that meet the interior of the face — V less its proper closed subspaces — modulo the operators of O_Z that map the face onto itself. The construction is the classical one, the fixed subspace of a point group in the space of forms (Delone *et al.*, 1975), written in conorm coordinates on the Selling cone; SELLA (Andrews *et al.*, 2023) uses the same subspaces and their reflection images. *Proof.* Sym and Fix are a Galois connection, so the closed subspaces are the intersections of the finitely many subspaces $\mathrm{Fix}(\mathbf{W})$, $\mathbf{W} \in O_Z$; since $L_{\mathbf{W}}$ permutes the slots on the face, each $\mathrm{Fix}(\mathbf{W})$ and each intersection is a set of equalities among conorms. A closed subspace with an interior point is a maximal set of lattices with the same symmetries, and two such subspaces exchanged by an operator preserving the face describe the same lattices from another member. Enumerating the intersections gives 8, 5, 3, 5 and 3 closed subspaces for no zero, one zero, two sharing a vector, two disjoint and

three, with symmetry orders $2,4,4,8,8,12,16,48$; $2,4,4,8,16$; $4,8,24$; $2,4,8,12,48$; $8,16,48$; these are Delone's 24 sorts, matched pattern by pattern with Table 3 and its holohedries, which is the check that the definition is his.[z53]

Proposition A10 (reindexing guarantee; reference frame). The quantities η, $\lambda_{\min}$ and V below are taken in the reference's reduced frame and are not invariants. Let $\mathbf{G}'_N$ be the new cell's metric written in the reference frame, $\|\mathbf{G}'_N - \mathbf{G}_R\|_F \le \eta < \lambda_{\min}(\mathbf{G}_R)$, let F_N be its reduced superbase with basis vectors of summed squared length V, and let $\tau \ge \sqrt{2}\,\eta\,V/(\lambda_{\min}(\mathbf{G}_R) - \eta)$ with $\lambda_1(\mathbf{G}_R) > 32\tau$. Then F_N, with its true integer coefficients in the reference frame, is a member of $\mathcal{S}_\tau(\mathbf{G}_R)$, the procedure of Section 3.2 produces it, and pairing it with the new cell's reduced superbase proposes the true operator. *Proof.* Let u_i be the coordinates of F_N's basis vectors in the reference frame, the rows of $U = \mathbf{P}^{\mathsf{T}}$ for the true operator $\mathbf{P}$, and $\Delta = \mathbf{G}'_N - \mathbf{G}_R$. Then $u_i^{\mathsf{T}}\mathbf{G}'_N u_i \ge (\lambda_{\min}(\mathbf{G}_R) - \eta)|u_i|^2$, so $\sum |u_i|^2 \le V/(\lambda_{\min} - \eta)$. In F_N's own frame the error is $U\Delta U^{\mathsf{T}}$, of Frobenius norm at most $\sum |u_i|^2\,\eta$, and in its own frame every conorm shift is at most $\sqrt{2}$ times that norm (Appendix B). The conorms of F_N under $\mathbf{G}'_N$ are non-negative, so under $\mathbf{G}_R$ they are at least $-\tau$, and Theorem A6 applies. The proposed operator for the pair is the change of basis between the two copies of F_N, which is the true one. □

Appendix B
From a metric error to a band

A conorm is linear in the metric, so a metric error $\Delta\mathbf{G}$ shifts the conorm of a member with coordinate vectors u_i, u_j by exactly $-u_i^{\mathsf{T}}\Delta\mathbf{G}\,u_j$; by Cauchy–Schwarz in the Frobenius product the shift is at most $c_{ij}\eta$, with $c_{ij} = \|\operatorname{sym}(u_i u_j^{\mathsf{T}})\|_F = [\frac{1}{2}(|u_i|^2|u_j|^2 + (u_i\cdot u_j)^2)]^{1/2}$, and an error along $\operatorname{sym}(u_i u_j^{\mathsf{T}})$ attains it. Every exact member of a stored closure survives in the widened closure of the measurement, for every

error of norm η, exactly when τ is at least the coverage bound $\tau_{\mathrm{cov}} = \max(c_{ij}\eta - p_{ij})$, read off the stored closure.[z36] For the worked cells $\eta = 14.4\,\text{Å}^2$ and $\tau = 3\eta = 43.2\,\text{Å}^2$. Member vectors are not confined to $\{-1, 0, 1\}$; on two and three walls they carry entries up to 3. On walls the largest constant is $\sqrt{2}$ with one zero, $\sqrt{5}$ with two, and with three zeros $2\sqrt{3}$ in six of the sixteen labelled cuboid frames — those whose zeros form a path with d at an interior vertex — and at most $\sqrt{5}$ otherwise; on positive slots it is at most 7.2. In a $2\sqrt{3}$ frame an adversarial error of norm η drives a member to $-3.46\,\eta$, so $\tau = 3\eta$ is not a worst-case guarantee in every frame.[z36] In a member's own frame the member is the standard superbase, whose largest constant is $\sqrt{2}$, so $\tau = \sqrt{2}\,\eta^*$ with η^* the largest error norm over the members' own frames covers every member, and $\eta^* \le \eta \max_k \|U_k\|_F^2$ over the stored members' coordinate matrices U_k, which is computable from η alone; it is uniform but loose, 2.8 times the reduced-frame η in the median case and up to 17 times.[z36] With a covariance Σ for the metric, the variance of each member conorm is the quadratic form of Σ on $\mathrm{sym}(u_i u_j^{\mathsf{T}})$, exact in any frame, and the band can be set slot by slot. With $\tau_{ij} = \sigma|u_i||u_j|$ and the four steps of Section 3.2 idealised in the frame of F_0 alone, three needles ($5 \times 5 \times 100\,\text{Å}$ monoclinic and orthorhombic, $40 \times 40 \times 300\,\text{Å}$ hexagonal; four random errors at each of 0.02°, 0.05°, 0.2°, 0.5°) returned the brute-force set in all 44 cases in which the idealised lattice existed, but that idealisation is not exact in general (the needle of Section 4.2), and the strain-norm projection of Section 3.2 replaces it; a $100 \times 100 \times 5\,\text{Å}$ plate was exact at 0.02° and 0.05° but at 0.2° and 0.5° its widened closure has 24 and 36 members, more than any closure of a single lattice, and the condition $\lambda_1 > 32 \max \tau_{ij}$ over the idealised slots fails there, as it does for $\sigma > (a_{\mathrm{short}}/a_{\mathrm{long}})^2/32 = 0.0045°$ by the uniform estimate.[z49]

Appendix C
Catalogues

Cell shapes. Table C1 lists the shapes of the ten sorts with more than one, each as the cell with the most right angles and no acute angle. The three shapes of a primitive monoclinic lattice are the three cells on $\{a, c, a + c\}$ with b unique, the cell choices of *International Tables* (Hahn, 2006, Section 2.2.16); the extra shapes of primitive orthorhombic, tetragonal and cubic lattices are monoclinic-looking cells with two right angles; C-centred orthorhombic has two primitive shapes; the boundary sorts of body-centred orthorhombic, C-centred monoclinic and triclinic have two or three cells with a single right angle each, for triclinic the known instability of the reduced cell when an angle is near 90° (Andrews *et al.*, 1980).[z49]

Table C1. *The cell shapes of the sorts with more than one, for the cells named; each shape is given as the cell with the most right angles and all angles* $\geq 90°$*, shortest edges first when several qualify and b unique for the monoclinic-looking ones, with the number n of members (up to swaps) that have it. Centred cells are given by their primitive cells; the boundary sorts are specified by Selling parameters in the* S^6 *order of Table 3, in* $Å^2$*.*[z49]

sort	lattice	n	shape
cP	30 Å	4	30, 30, 30; 90, 90, 90
		12	30, 30, 42.4; 90, 135, 90
tP	30, 30, 50 Å	4	30, 30, 50; 90, 90, 90
		4	30, 30, 58.3; 90, 121.0, 90
		8	30, 50, 42.4; 90, 135, 90
oP	30, 40, 55 Å	4	30, 40, 55; 90, 90, 90
		4	30, 40, 62.6; 90, 118.6, 90
		4	30, 55, 50; 90, 126.9, 90
		4	40, 55, 50; 90, 143.1, 90
oS	C 30, 40, 50 Å	2	25, 50, 25; 90, 106.3, 90
		4	25, 50, 30; 90, 126.9, 90
mP	30, 40, 50 Å, 100°	2	30, 40, 50; 90, 100, 90
		2	30, 40, 53.7; 90, 113.4, 90
		2	50, 40, 53.7; 90, 146.6, 90
oI (O3)	900, 1600, 0, 900, 1600, 0	1	50, 50, 50; 111.1, 129.8, 90
		2	42.4, 50, 50; 90, 115.1, 115.1
mC (M2B)	900, 1600, 0, 900, 1600, 2500	1	50, 50, 70.7; 104.7, 116.9, 90
		1	50, 50, 65.6; 105.9, 105.9, 90
mC (M3)	900, 1600, 0, 2500, 1600, 0	1	50, 50, 64.0; 120, 90, 111.1
		2	50, 56.6, 58.3; 90, 108.0, 124.4
aP (A2)	900, 1600, 0, 2500, 400, 1200	1	36.1, 60.8, 64.0; 114.3, 90, 114.2
		1	36.1, 56.6, 64.0; 116.2, 90, 101.3
aP (A3)	900, 1600, 0, 2500, 400, 0	1	36.1, 50, 53.9; 90, 101.9, 119.9
		1	44.7, 50, 53.9; 90, 99.6, 135.7
		1	36.1, 44.7, 58.3; 90, 115.3, 104.4

Specialisations. The conorms of any member, in any labelling, are integer combi-

nations of the reduced cell's conorms, hence linear in the free parameters of the sort (the letters of Table 3); imposing another sort's zeros and equalities on one member is a homogeneous linear system in those parameters, and a solution with non-negative parameters, a non-singular lattice and a resulting lattice of the target sort and no higher is a specialisation. Table C2 lists the 37 conditions of codimension one, on 36 pairs of sorts.[z50] Each is a parameter vanishing or two parameters becoming equal. The classical values appear: body-centred tetragonal is face-centred cubic at $c/a = \sqrt{2}$ and body-centred cubic at $c/a = 1$; C-centred orthorhombic is hexagonal at $b = \sqrt{3}\,a$ and tetragonal at $b = a$; rhombohedral is cubic at $\alpha = 60°$, $90°$ and $109.47°$. The monoclinic conditions come in threes, one per shape: with unique axis c and plane lattice a, b, the lattice is C-centred orthorhombic when $|a| = |b|$, $|a| = |a + b|$ or $|b| = |a + b|$, and primitive orthorhombic when $a \perp b$, $a \perp (a+b)$ or $b \perp (a+b)$ — the textbook condition $a = c$ read in each of the three cell choices. The table also shows which sorts are one condition from a much higher symmetry: C-centred monoclinic from tetragonal, body-centred tetragonal and rhombohedral; the body-centred orthorhombic boundary sort from face-centred cubic; triclinic on a wall from C-centred monoclinic.

Table C2. *All specialisations of codimension one between the 24 sorts. Conditions are on the letters of the sort's pattern in Table 3; "shapes" is the number of distinct cell shapes of the starting sort through which the conditions are reached, and the last column reads the condition in the cell of the conventional description where that is short.*[z50]

from	to	conditions	shapes	reading
aP A1	aP A2	any letter $= 0$	1	a right angle appears
aP A2	aP A3	$v = 0$	1	a second right angle
aP A2	mC M2A	$s = t$, $r = u$	1	
aP A2	mP M4	r, s, t or $u = 0$	1	
aP A3	mC M3	any two letters equal	3	one per shape
mP M4	oP O5	$r = 0$, $s = 0$, $t = 0$	3	$a \perp b$, $a \perp (a+b)$, $b \perp (a+b)$
mP M4	oS O4	$s = t$, $r = t$, $r = s$	3	$\lvert a\rvert = \lvert b\rvert$, $\lvert a\rvert = \lvert a+b\rvert$, $\lvert b\rvert = \lvert a+b\rvert$
mC M1A	oF O1A	$r = t$	1	
mC M1A	oI O2	$u = 0$	1	
mC M1A	oS O4	$r = 0$, $t = 0$	1	
mC M1B	mC M2B	$t = 0$, $u = 0$	1	boundary sort
mC M1B	mC M3	$r = 0$, $s = 0$	1	boundary sort
mC M1B	oF O1A	$r = s$	1	
mC M1B	oI O1B	$t = u$	1	
mC M2A	oI O2	$r = t$	1	
mC M2B	oI O3	$t = 0$	1	
mC M2B	tI T2	$r = s$	1	one step to tI
mC M2B	tP T5	$r = 0$, $s = 0$	1	one step to tP
mC M3	oI O3	$r = t$	1	
mC M3	hR R3	$r = s$, $s = t$	1	one step to hR
oP O5	tP T5	$r = s$, $r = t$, $s = t$	1	$a = b$, $a = c$, $b = c$
oS O4	tP T5	$r = 0$	1	$b = a$
oS O4	hP H4	$r = s$	1	$b = \sqrt{3}\,a$
oI O1B	oI O3	r, s or $t = 0$	1	boundary sort
oI O1B	tI T1	$r = s$, $r = t$, $s = t$	1	two edges equal
oI O2	tI T2	$r = s$	1	
oI O3	cF C3	$r = s$	1	one step to cF
oF O1A	tI T1	$s = t$	1	
oF O1A	tI T2	$s = 0$, $t = 0$	1	
tP T5	cP C5	$r = s$	1	$c = a$
tI T1	cI C1	$r = s$	1	$c/a = 1$
tI T1	cF C3	$s = 0$	1	$c/a = \sqrt{2}$ from below
tI T2	cF C3	$s = 0$	1	$c/a = \sqrt{2}$ from above
hR R1	cP C5	$r = 0$	1	$\alpha = 90^\circ$
hR R1	cI C1	$r = s$	1	$\alpha = 109.47^\circ$
hR R3	cF C3	$r = s$	1	$\alpha = 60^\circ$

Measured cells from powder indexing. Oishi-Tomiyasu (2024) distributes seven measured primitive cells with her program's output, a ranking of Bravais types by distance in the manner of the quality indices of Kabsch (1993): for each Bravais type, the nearest cell of that type and its distance. Table C3 treats each by Section 3.2 at $\tau = \lambda_1/32$ and prices every operator out of the reduced cell in the closure of the idealised lattice; the price at which the number of operators reaches the order of a holohedry is the

band at which that symmetry is offered; prices here are in Å^2 in the reduced frame and are not invariants.[z51] The program's distance is the norm, in Å^2, between the input metric tensor and the nearest metric tensor of the type (Oishi-Tomiyasu, 2012), and the band is taken from the theorem rather than from the cells' errors, which the distribution does not include. In all seven cases the band idealises the walls of the indexed type, and the type's holohedry is reached at a price between 0.9 and 1.8 times the program's distance, with the same ordering of types; the two measures differ in definition, so they agree in order and size, not in value. Her test set of 4738 cells is not distributed.

Table C3. *The seven sample cells of BLDConograph (Oishi-Tomiyasu, 2024) by the closure: Bravais type and name as distributed; members of the idealised closure at* $\tau = \lambda_1/32$*; the price in Å*^{2} *at which the number of operators out of the reduced cell reaches the order of the type's holohedry, and of the next holohedry up where one is reached; the program's minimal distance to a cell of that type.*[z51]

type	sample	members	price, type	price, next	distance
tI	HERMES_Sr327_250K	2	0.77	132.6	0.57
tP	1339prbasrcuo_bs1	16	0.10	26.0	0.056
cI	sample3	1	2.98	–	2.07
cF	CeO2_matsushita	3	0.11	–	0.084
hP	M402-pf-119673	6	22.5	196.1	20.5
mC	Acrinol-B-D8CuKa1	1	0.96	86.2	0.72
hR	Alumina676	3	1.69	9.18	1.93

Benchmark. The repository directory `manuscript/benchmark` holds 780 measured cells with exact references and true operators, in five families: 75 cells measured in a different shape of the same lattice; 660 lattices placed 0.5 to 8 times the error from a condition of Table C2; 20 cells whose only reduced superbase is a difference-wall member; 16 needles and plates with angular errors; 9 cells of the boundary sorts. Each case also carries its Niggli cell. An evaluator scores whether a method proposes the true operator from one reduction of the measured cell; the results quoted in Section 5 are in `results.txt`.

Appendix D
The operators alone

The closure can be described without any lattice. An operator $\mathbf{W}$, written as the new basis in the coordinates of the reduced cell, acts on the six conorms by a fixed integer 6×6 matrix $L_{\mathbf{W}}$, since every conorm of $\mathbf{W}F_0$ is an integer combination of the conorms of F_0. The reduced cells are the points of the orthant $\{p \geq 0\}$ in conorm space; Selling (1874) showed that every lattice has a conorm vector in this orthant, and it is unique up to the 48 swaps except on the walls (Delaunay, 1933; Conway & Sloane, 1992), and the closure of a lattice is the set of points of its orbit that lie in the orthant: one in the interior, several on the walls. So $\mathbf{W}$ is an operator of the closure of every lattice with zero pattern Z exactly when $L_{\mathbf{W}}$ maps the face $\{p \geq 0,\ p_Z = 0\}$ into the orthant, which is a sign condition on the entries of $L_{\mathbf{W}}$ and nothing else. Enumerating integer matrices under that condition gives the operator set of each face without a lattice: 48×1, 2, 3, 6 and 16 matrices for no zero, one zero, two zeros on disjoint edges, two zeros on a path and three zeros, and their union, over all placements of the faces, is a set $\mathcal{U}$ of $6096 = 48 \times 127$ integer matrices with entries in $\{-3, \ldots, 3\}$. $\mathcal{U}$ is closed under inversion, every element has a labelling whose basis vectors have entries in $\{-2, \ldots, 2\}$, and $\mathcal{U}$ contains every operator between two reduced cells of one lattice, whatever the lattice, since an operator out of any member is an operator out of a reduced cell of that member's zero pattern and Lemma A0 bounds all such operators within the window enumerated.[z52] The three-zero face enumerated is the triangle; its path placement gives the same operators relabelled. Up to relabelling at both ends $\mathcal{U}$ has nine kinds of element, of flip length 0 to 4 (Table D1).

Table D1. *The nine kinds of operator between Selling-reduced cells, up to relabelling at both ends; "matrices" counts the integer matrices of the kind in* $\mathcal{U}$*, "flips" its length as a flip word, and the last column the zero patterns in whose closures it occurs.*[z52]

kind	matrices	flips	occurs with
1	48	0	every pattern (the swaps)
2	288	1	every pattern with a zero
3	1152	2	two zeros on a path; three zeros
4	1152	3	three zeros
5	1152	3	three zeros
6	576	3	two zeros on a path; three zeros
7	576	4	three zeros
8	576	4	three zeros
9	576	4	three zeros

Why four. For a lattice with $a \perp b \perp c$ the sixteen members are four orthogonal superbases, differing by the sign of one axis, and twelve with a face diagonal. A flip from an orthogonal cell always gives a cell with a face diagonal, so changing the sign of one axis takes three flips, out, across and in, and the diagonal cells attached only to the far orthogonal cell take one more: from (a, b, c) the members lie at flip distances $0, 1, 2, 3, 4$ in numbers $1, 3, 6, 3, 3$, and four is the diameter of the largest face.[z46]

Appendix E
Computational checks

Each mark [z*nn*] refers to `manuscript/appendix_z/test_z`*nn*`_*.py` in the `agentsg` repository.

mark	what is checked
z36	shift identity; constants by zero pattern and frame; adversarial loss at 3η; frame-free bound
z39	measured worked pair: widened closure by definition; candidate operators at $\tau =$ 43.2 and 60
z44	constants 4 and 8; class gap and excess; Theorem A6 against brute force; grey zone; worked cell
z45	reduction endpoints and restarts; walks on the measured metric; the worked filter table
z46	the monoclinic example; closures of the 24 sorts; shapes = orbits; twofolds generate; coset completeness; difference-wall members; 2τ; nesting
z47	reindexing under a declared error: membership, true operator, guarantee
z48	the 24 Delone sorts reproduced by closed tie patterns on generic lattices
z49	bands per slot on needles and plates: exactness, the plate floor, detection of the failed hypothesis; the cell shapes of Table C1
z50	the 37 codimension-one specialisations between sorts and the shapes that carry them (Table C2)
z51	the seven BLDConograph sample cells: idealised closure, price ladder, comparison with the program's distances (Table C3)
z52	the operator sets of the faces from the matrices alone; the universal set $\mathcal{U}$, its inversion closure and window; the nine kinds (Table D1); universality over the 24 sorts
z53	the 24 sorts as the closed sets of equalities of Proposition A9, with their symmetry orders
z54	every column of Table 3 computed from the operators alone: members, point group, visible symmetries, shapes, and shapes = orbits
z55	exactness of idealise–enumerate–filter at bands from $\lambda_1/32$ to $\lambda_1/6$; singular idealisations are the only failures
z56	Lemma A0: the obtuse superbases of every zero pattern enumerated inside the window the lemma forces
z59	Lemma A12 (the finite widened set) and Theorem A12: adjacency radii δ_1, δ_2 (Table 4); the filtered first layer equals the widened set below δ_1; the plates at 3° between δ_1 and δ_2
z58	the strain norm: project–enumerate–filter on the needle, two thin plates and 240 random cells against brute force; distances to the 24 sorts and their monotonicity along Table C2
z57	bands per slot: the needle on which idealisation in the frame of F_0 fails; zeroing on the closure, by rounds and during the enumeration; the zeroing filtration and its geometric limit; 240 random cells; the 3° rule by aspect ratio; the $[-3, 3]$ window

References

Andrews, L. C. & Bernstein, H. J. (1988). *Acta Cryst.* A**44**, 1009–1018.

Andrews, L. C. & Bernstein, H. J. (2014). *J. Appl. Cryst.* **47**, 346–359.

Andrews, L. C. & Bernstein, H. J. (2023a). *Acta Cryst.* A**79**, 485–498.

Andrews, L. C. & Bernstein, H. J. (2023b). *Acta Cryst.* A**79**, 499–503.

Andrews, L. C., Bernstein, H. J. & Pelletier, G. A. (1980). *Acta Cryst.* A**36**, 248–252.

Andrews, L. C., Bernstein, H. J. & Sauter, N. K. (2019a). *Acta Cryst.* A**75**, 593–599.

Andrews, L. C., Bernstein, H. J. & Sauter, N. K. (2019b). *Acta Cryst.* A**75**, 115–120.

Andrews, L. C., Bernstein, H. J. & Sauter, N. K. (2023). SELLA – a program for determining Bravais lattice types. arXiv:2303.03122.

Buerger, M. J. (1960). *Z. Kristallogr.* **113**, 52–56.

Burzlaff, H. & Zimmermann, H. (1985). *Z. Kristallogr.* **170**, 247–262.

Burzlaff, H. & Zimmermann, H. (2006). *International Tables for Crystallography*, Vol. A, 5th ed., edited by Th. Hahn, ch. 9.1, pp. 742–749. Dordrecht: Springer.

Conway, J. H. & Sloane, N. J. A. (1992). *Proc. R. Soc. London Ser. A*, **436**, 55–68.

Delaunay, B. (1933). *Z. Kristallogr.* **84**, 109–149.

Delone, B. N., Galiulin, R. V. & Shtogrin, M. I. (1975). *J. Sov. Math.* **4**, 79–156.

de Wolff, P. M. (2006). *International Tables for Crystallography*, Vol. A, 5th ed., edited by Th. Hahn, ch. 9.2, pp. 750–755. Dordrecht: Springer.

de Wolff, P. M. & Gruber, B. (1991). *Acta Cryst.* A**47**, 29–36.

Flack, H. D. (1987). *Acta Cryst.* A**43**, 564–568.

Grosse-Kunstleve, R. W., Sauter, N. K. & Adams, P. D. (2004). *Acta Cryst.* A**60**, 1–6.

Gruber, B. (1973). *Acta Cryst.* A**29**, 433–440.

Gruber, B. (1997). *Acta Cryst.* A**53**, 505–521.

Gruber, B. (2006). *International Tables for Crystallography*, Vol. A, 5th ed., edited by Th. Hahn, ch. 9.3, pp. 756–760. Dordrecht: Springer.

Hahn, Th. (2006). Editor. *International Tables for Crystallography*, Vol. A, 5th ed. Dordrecht: Springer.

Himes, V. L. & Mighell, A. D. (1987). *Acta Cryst.* A**43**, 375–384.

Kabsch, W. (1993). *J. Appl. Cryst.* **26**, 795–800.

Křivý, I. & Gruber, B. (1976). *Acta Cryst.* A**32**, 297–298.

Kurlin, V. (2022). *A complete isometry classification of 3-dimensional lattices.* arXiv:2201.10543; revised manuscript, April 2026, https://kurlin.org/projects/lattice-geometry/lattices3Dmaths.pdf. Lemma numbers refer to the revised manuscript.

Lebedev, A. A., Vagin, A. A. & Murshudov, G. N. (2006). *Acta Cryst.* D**62**, 83–95.

Le Page, Y. (1982). *J. Appl. Cryst.* **15**, 255–259.

Niggli, P. (1928). *Kristallographische und strukturtheoretische Grundbegriffe.* Handbuch der Experimentalphysik, Vol. 7, Part 1. Leipzig: Akademische Verlagsgesellschaft.

Oishi-Tomiyasu, R. (2012). *Acta Cryst.* A**68**, 525–535.

Oishi-Tomiyasu, R. (2024). *Acta Cryst.* A**80**, 339–350.

Patterson, A. L. & Love, W. E. (1957). *Acta Cryst.* **10**, 111–116.

Pennec, X., Fillard, P. & Ayache, N. (2006). *Int. J. Comput. Vis.* **66**, 41–66.

Sauter, N. K., Grosse-Kunstleve, R. W. & Adams, P. D. (2004). *J. Appl. Cryst.* **37**, 399–409.

Selling, E. (1874). *J. Reine Angew. Math.* **77**, 143–229.

Voronoi, G. (1908). *J. Reine Angew. Math.* **134**, 198–287.

Synopsis

The set of all Selling-reduced cells of a lattice, with the integer matrices between them, holds the lattice symmetry and every alternative choice of reduced cell in one complete set. For a measured cell the set is recovered exactly by setting every small conorm to zero, enumerating and filtering, and reindexing reads one matrix from it.
