

COMPOSE: Pivotal phase III trial to compare ¹⁷⁷Lu-edotreotide with best standard of care for well-differentiated aggressive grade 2 and 3 gastroenteropancreatic neuroendocrine tumours

902TiP

Halfdanarson TR,¹ Halperin DM,² Reidy-Lagunes D,³ Kong G,⁴ Capdevila J,⁵ Mailman J,⁶ Herrmann K,⁷ Srirajaskanthan R,⁸ Leyden S,⁹ Sierras C,¹⁰ Harris P¹⁰

1. Mayo Clinic, Rochester, MN USA; 2. MD Anderson Cancer Center, Houston, Texas, USA; 3. Memorial Sloan Kettering Cancer Center, New York, USA; 4. Peter MacCallum Cancer Centre, Melbourne, VIC, Australia; 5. Vall d'Hebron University Hospital, Vall d'Hebron Institute of Oncology, Barcelona, Spain; 6. NorCal CarcINET Community, Oakland, CA; 7. University Hospital Essen, Germany; 8. Kings College Hospital, London, UK; 9. NeuroEndocrine Cancer Australia, Victoria, Australia; 10. ITM Oncologics GmbH, Garching/Munich, Germany


Background

Approximately 70% of neuroendocrine tumours are gastroenteropancreatic (GEP-NETs). These frequently develop metastatic disease with limited treatment options.¹

For well-differentiated high grade 2 and 3 GEP-NETs, current therapies include peptide receptor radionuclide therapy (PRRT), somatostatin analogues, chemotherapy, cytoreduction and molecular targeted therapies (everolimus, sunitinib) with no specified use sequence.

PRRT uses radiolabelled somatostatin analogues to selectively target somatostatin receptor expressing (SSTR+) tumour cells. Use may stabilise disease and induce objective tumour response.²

Figure 1. Kaplan-Meier estimates of progression free survival in the study population depending on number of no-carrier added ¹⁷⁷Lu-edotreotide PRRT cycles³



¹⁷⁷Lu edotreotide is an innovative radiolabelled somatostatin analogue with promising efficacy and a favourable safety profile.^{2,3} Retrospective data in metastatic GEP-NETs treated with ≥2 cycles of ¹⁷⁷Lu-edotreotide demonstrate progression free survival (PFS) of ≥30 months (Figure 1).³


The Phase III COMPETE trial, which has completed patient recruitment, is comparing ¹⁷⁷Lu edotreotide efficacy/safety versus everolimus in grade 1 and 2 GEP-NETs.⁴

Trial design

COMPOSE is a prospective, randomised, controlled, open-label, multi-center Phase III trial recruiting patients with well-differentiated aggressive grade 2 and 3 (Ki-67 index 15–55%), SSTR+, GEP-NETs.

COMPOSE will evaluate efficacy, safety and patient-reported outcomes of 1st- or 2nd-line treatment with ¹⁷⁷Lu edotreotide PRRT compared to best standard of care [investigator's choice of chemotherapy (capecitabine-temozolomide: CAPTEM or folinic acid, fluorouracil and oxaliplatin: FOLFOX) or everolimus] (Figure 2).

Figure 2. Summary schedule of treatments and follow-up consultation



COMPOSE aims to randomize 202 patients 1:1 to a defined number of cycles ¹⁷⁷Lu-edotreotide or an active comparator (Figure 2).

Conflicts of interest: Co-authors participated in the trial as Principal Investigators
Study sponsored by: ITM Solucin GmbH

Presented at: 2022 ESMO Congress; Paris, France; Sept 9–13

Primary endpoint

- Progression free survival, assessed every 12 weeks until disease progression (response evaluation criteria in solid tumors v1.1) or death, whichever occurs earlier

Key secondary endpoint

- Overall survival, assessed up to 2 years after disease progression

Current status

- Recruitment has opened in several countries with approximately 40 specialist centres in 10 countries scheduled to be open

COMPOSE results are expected to increase treatment options for patients with well-differentiated aggressive grade 2 and 3 GEP-NETs, including for 1st-line therapy



Correspondence to:
cristina.sierras@itm-radiopharma.com

Clinical Phase III Trial COMPOSE
NCT04919226: ClinicalTrials.gov

Find more trial information on
www.itm-gep-net-trials.com

References

- Severi S, et al. Onco Targets Ther 2017;10:551–557;
- Strosberg J, et al. N Engl J Med 2017;376:125–135; 3. Baum RP, et al. Theranostics 2016;6:501-510. 4. ClinicalTrials.gov Identifier: NCT03049189