

Supporting Information

Comprehensive Understanding of SiO₂-promoted Iron Fischer-Tropsch Synthesis

Catalysts: Fe-SiO₂ Interaction and Beyond

Yu Zhang^{a, b}, Ming Qing^{c, *}, Hong Wang^c, Xing-Wu Liu^c, Suyao Liu^c, Hongliu Wan^{a, b}, Linge Li^{a, b}, Xiang Gao^{a, b}, Yong Yang^{a, c, *}, Xiao-Dong Wen^{a, c, *}, and Yong-Wang Li^{a, c}

^a State Key Laboratory of Coal Conversion, Institute of Coal Chemistry Chinese Academy of Sciences Taiyuan 030001, People's Republic of China

^b University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China

^c National Energy Research Center for Clean Fuels, Synfuels China Co., Ltd, Beijing 101400, People's Republic of China

* Corresponding author.

E-mail address: qingming@synfuelschina.com.cn (M. Qing);

yyong@sxicc.ac.cn (Y. Yang)

wxd@sxicc.ac.cn (X. Wen).




Fig. S1. XRD patterns of fresh catalysts.




Fig. S2. Relationship between CO₂ selectivity and CO conversion of catalysts.




Fig. S3. Iron species particle distributions of fresh and spent Fe@Si catalysts.

Table S1. Quantitative results of Fe-O-Si and Si-O-Si bonds intensity in FT-IR.

Catalysts	Fe-O-Si bond (I)	Si-O-Si bond (II)	Ratio (I/II)
Fe@Si	4.8	62.4	0.077
Fe/Si	8.3	75.8	0.109
Fe-Si	12.3	64.5	0.191