# Classification of Finitely Often Self-Intersecting Loops in $\mathbb{R}^2$ under equivalence given by isomorphism of corresponding Region Graphs.

Adam Kurkiewicz School of Computer Science University of Glasgow United Kingdom

May 19, 2016

#### Abstract

Abstract placeholder

### 1 Introduction

**Definition 1** (Almost Injectivity). Let  $f: X \mapsto Y$  be a function. Let us say that f is almost injective, if  $f(x_1) = f(x_2) \implies x_1 = x_2$  holds for all but finitely many  $x_i \in X$ .

**Definition 2** (Crossing Set). Let  $f: X \mapsto Y$  be an almost injective function. Let the crossing set  $C_f$  of f be defined by

$$C_f = \{x_1, x_2 \mid f(x_1) = f(x_2) \land x_1 \neq x_2\}$$

**Definition 3** (Crossing Equivalence). Let  $C_f$  be the crossing set of of an almost injective function f. Let crossing equivalence  $\sim_C$  be a relation on  $C_f$  defined by  $c_1 \sim_C c_2 \iff f(c_1) = f(c_2)$  for any  $c_1, c_2 \in C_f$ .

**Proposition 1** (Crossing Equivalence Relation). Crossing equivalence  $\sim_C$  is an equivalence relation on the crossing set  $C_f$  of an almost injective function f.

*Proof.* Reflexivity, Symmetry and Transitivity follow easily from properties of set equality =, and that f is a function.

**Definition 4** (FOSIL). Let  $\gamma:[0,1]\mapsto \mathbb{R}^2$  be a continuous, almost injective function with  $\gamma(0)=\gamma(1)$ . We will say that  $\gamma$  is a finitely often self-intersecting loop in  $\mathbb{R}^2$ , or FOSIL in short.

**Definition 5** (Region Number). Let  $\gamma$  be a FOSIL. Let  $\sim_C$  be a crossing equivalence on  $C_{\gamma}$ , the crossing set of  $\gamma$ . Define region number of  $\gamma$ ,  $\rho_{\gamma}$  as

$$\rho_{\gamma} = 1 + \sum_{[c] \in C_{\gamma}/\sim_C} (|[c]| - 1)$$

.

**Definition 6** (Regions). Let  $\gamma$  be a FOSIL. Let  $R^{\gamma}$  denote a collection of open, connected, disjoint subsets of  $\mathbb{R}^2$ , whose union gives the complement of  $\gamma$ , that is  $\mathbb{R}^2 \setminus \gamma([0,1])$ . Then  $R^{\gamma}$  is called a collection of regions of  $\gamma$ , and  $R_i \in R^{\gamma}$  is called a region.

**Theorem 1** (Jordan Curve Theorem for FOSILs). Let  $\gamma$  be a FOSIL,  $R^{\gamma}$  be a collection of regions of  $\gamma$ , and  $\rho_{\gamma}$  be the region number of  $\gamma$ . Then the collection of regions is unique, and

$$|R^{\gamma}| = \rho_{\gamma}$$

This auxiliary result has a rather technical proof, to which we devote a separate chapter.

**Definition 7** (Edge-Connectedness). Let  $\gamma$  be a FOSIL. Let  $R_1, R_2 \in R^{\gamma}$  be regions of  $\gamma$ , and  $C_{\gamma}$  be the crossing set of  $\gamma$ . We will say that  $R_1$  and  $R_2$  are edge-connected if  $R_1 \neq R_2$  and there exists an open, connected subset of [0,1], say S, with the property that  $S \cap C_f = \emptyset$ , and such that  $R_1 \cup \gamma(S) \cup R_2$  is connected.

**Proposition 2** (Edge-Connectedness Reflexivity). Edge connectedness on the collection of regions of  $\gamma$  is a reflexive relation.

*Proof.* Reflexivity follows from commutativity of set union  $\cup$ .

**Definition 8** (Graph). Let V be a finite set. Let E be a subset of  $V \times V$ . Then G = (V, E) is called a graph if for any  $(v_1, v_2) \in E$  we have that  $(v_2, v_1) \in E$ .  $v \in V$  is called a vertex in G, and  $(v_1, v_2) \in E$  is called an edge in G.

**Definition 9** (Region Graph). Let  $R^{\gamma}$  be the collection of regions of a FOSIL  $\gamma$ . Let  $E = \{(R_1, R_2) \in R^{\gamma} \times R^{\gamma} \mid R_1 \text{ edge-connected } R_2\}$ . Then  $G = (R^{\gamma}, E)$  is called a region graph of a FOSIL  $\gamma$ .

**Proposition 3** (Region Graph is a Graph). The region graph G = (V, E) of a FOSIL  $\gamma$  is a graph.

*Proof.* By Jordan Curve Theorem for FOSILs, V is uniquely defined and finite. Let  $(R_1, R_2) \in E$ . Then  $(R_2, R_1) \in E$  by reflexivity of edge-connectedness.  $\square$ 

**Definition 10** (Graph Isomorphism). Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be graphs with  $|V_1| = |V_2|$  and  $|E_1| = |E_2|$ . We will say that  $G_1$  is graph isomorphic,  $\sim_G$ , to  $G_2$  if there exists an injective function  $\psi : V_1 \mapsto V_2$  such that for each  $(v_1, v_2) \in E_1$ , we have that  $(\psi(v_1), \psi(v_2)) \in E_2$ .

**Proposition 4** (Graph Isomorphism Equivalence Relation). Let  $\mathfrak{G}$  be a collection of graphs. Then  $\sim_G$  is an equivalence relation on  $\mathfrak{G}$ .

Proof. TODO Symmetry, transitivity, reflexivity.

**Definition 11** (Graph Path). Let G = (V, E) be a graph. Let  $P = \{v_k\}$  be a sequence of vertices in G, such that for any  $v_i, v_{i+1} \in P$ , we have that  $(v_i, v_{i+1}) \in E$ . Then P is called a path in graph G.

**Definition 12** (Connected Graph). Let G = (V, E) be a graph. Let us say that G is connected if there exists a path P in G, such that for any  $v \in V$  we have that  $v \in P$ .

**Definition 13** (Bipartite Graph). Let G = (V, E) be a graph. If there exist disjoint  $S_1, S_2 \subseteq V$  with  $S_1 \cup S_2 = V$  such that for any  $v_i, w_i \in S_i$  we have  $(v_i, w_i) \notin E$ , then G is called a bipartite graph.

**Theorem 2** (FOSIL Classification Theorem). Let  $\mathfrak{F}$  be the collection of all FOSILs. Let  $\mathfrak{R} = \{R^{\gamma} | \gamma \in \mathfrak{F}\}$  be the collection of corresponding region graphs. Let  $\mathfrak{B}$  be the collection of all connected bipartite graphs. The function

$$\Phi: \mathfrak{R}/\sim_G \mapsto \mathfrak{F}/\sim_G$$

given by

$$\Phi([R]_{\mathfrak{R}}) = [R]_{\mathfrak{B}}$$

is a well-defined bijection.

In other words,  $\mathfrak{R} \subseteq \mathfrak{B}$  and for any  $B \in \mathfrak{B}$  there is an  $R \in \mathfrak{R}$  with  $R \sim_G B$ .

We devote the next chapter to the proof of this theorem.

### 2 FOSIL Classification Theorem

The inclusion bit should follow from Hopf's degree theorem. Surjectivity is much trickier, but there seems to be an algorithm to go from any bipartite graph to a FOSIL, but I don't quite have the details yet.

### 3 Jordan Curve Theorem for FOSILs

The starting point should be the proof of Jordan-Holder. Two proofs in particular seem of interest [1], [2].

## References

- [1] T. Hales. Jordan's proof of the Jordan Curve Theorem. Studies in Logic, Grammar and Rhetoric, 10(23):116–131, 2007.
- [2] C. Thomassen. The Jordan-Schonflies theorem and the classification of surfaces. *The American Mathematical Monthly*, 99(2):116–131, 1992.