Rare decays at CMS

Jónatan Piedra* [⁺]
IFCA (CSIC - Universidad de Cantabria
E-mail: piedra@cern.ch

XIV International Conference on Heavy Quarks and Leptons (HQL2018) May 27- June 1, 2018 Yamagata Terrsa, Yamagata,Japan

^{*}Speaker.

[†]On behalf of the CMS Collaboration.

Rare decays at CMS Jónatan Piedra

- 1. Introduction
- 2. FCNC with top quarks
- **2.1** FCNC in $tZq \rightarrow 3\ell$

[1]

2.2 FCNC in $tH \rightarrow bb$

[2]

- 3. Angular analyses with bottom quarks
- 3.1 Angular observables in $B^+ \to K^+ \mu \mu$

[3]

3.2 Angular observables in $B^0 \to K^{*0} \mu \mu$

[4]

4. Conclusions

References

- [1] CMS Collaboration, Search for flavour changing neutral currents in top quark production and decays with three-lepton final state using the data collected at $\sqrt{s} = 13$ TeV, https://cds.cern.ch/record/2292045, CMS-PAS-TOP-17-017.
- [2] CMS Collaboration, Search for the flavor-changing neutral current interaction of the top quark and the Higgs boson which decays into a pair of b quarks at $\sqrt{s} = 13$ TeV, accepted for publication in JHEP https://cds.cern.ch/record/2296416, CERN-EP-2017-309 [hep-ex/1712.02399].
- [3] CMS Collaboration, Angular analysis of the decay $B^+ \to K^+ \mu^+ \mu^-$ at $\sqrt{s} = 8$ TeV, https://cds.cern.ch/record/2621370 CERN-EP-2018-125 [hep-ex/1806.00636].
- [4] CMS Collaboration, Measurement of angular parameters from the decay $B^0 \to K^{*0} \mu^+ \mu^-$ at $\sqrt{s} = 8$ TeV, https://cds.cern.ch/record/2287571, CERN-EP-2017-240 [hep-ex/1710.02846].