

## 1 Finite Difference Methods

### General System Formulas

The general system is a BVP set up as:  $x'' = F(t, x, x')$  with boundary values  $x(a) = \alpha$  and  $x(b) = \beta$ . Here  $t = a + ih$  and  $h = \frac{b-a}{n}$  where  $(0 \leq i \leq n)$

### Central Difference Formulas

For the frist derivative the formula is:

$$x'(t) = \frac{x(t+h) - x(t-h)}{2h}$$

$$\rightarrow x'_i = \frac{x_{i+1} - x_{i-1}}{2h}$$

For the second derivative the formula is:

$$x''(t) = \frac{x(t+h) - 2x(t) + x(t-h)}{h^2}$$

$$\rightarrow x''_i = \frac{x_{i+1} - 2x_i + x_{i-1}}{h^2}$$

central difference for PDEs are as follows:

$$\frac{\partial u}{\partial t} \approx \frac{u_i^{j+1} - u_i^j}{\Delta t} \text{ and } \dots$$

$$\frac{\partial^2 u}{\partial t^2} \approx \frac{(u_{i+1}^j - 2u_i^j + u_{i-1}^j) + (u_{i+1}^{j+1} - 2u_i^{j+1} + u_{i-1}^{j+1})}{2\Delta x^2}$$

### Matrix Formulas

The formulas also take a matrix form in a linear system  $Ax = b$ . The pricipal equation of the system is found in the form:

$$a_i x_{i-1} + d_i x_i + c_i x_{i+1} = b_i$$

In  $Ax = b$  form this looks like:

$$A = \begin{bmatrix} d_1 & c_1 & & \\ a_2 & d_2 & c_2 & \\ & & \ddots & \\ & & & a_n & d_n \end{bmatrix}$$

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad b = \begin{bmatrix} b_1 - a_1 \alpha \\ b_2 \\ \vdots \\ b_n - c_n \beta \end{bmatrix}$$

## 2 Shooting Method Formulas

### General System Formulas

Similar to finite methods. The general system is a BVP set up as:  $x'' = F(t, x, x')$  with boundary values  $x(a) = \alpha$  and  $x(b) = \beta$ . Here we may also have formulas for hypothesized terminal values at  $b$  in the form  $x(b) \approx \beta \approx \phi(z)$ .  $z$  values will be of an IVP form  $x'(a) = z_{IVP}$

The error formula for  $\phi(z)$  is given by:  $\epsilon = |\phi(z) - \beta|$

### Iterative z Formulas

The general form iterative z formula is as follows:

$$z_{n+1} = z_n + \frac{\beta - \phi(z_n)}{\phi(z_n) - \phi(z_{n-1})} (z_n - z_{n-1})$$

In the simple form the formula is:

$$z_3 = z_2 + \frac{\beta - \phi(z_2)}{\phi(z_n) - \phi(z_1)} (z_2 - z_1)$$

## 3 Linear Algebra Overview

### LU Decomposition Formulas

Here  $A = LU$ , this would look like the following:

$$A = LU, \quad A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$L = \begin{bmatrix} l_{11} & 0 & 0 \\ l_{21} & l_{22} & 0 \\ l_{31} & l_{32} & l_{33} \end{bmatrix}, U = \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix}$$

to make this fit on the cheatsheet I have broken  $A$  into several vectors,  $A = [a_1 | a_2 | a_3]$ , so these parts of  $A$  are:

$$a_1 = \begin{bmatrix} l_{11} u_{11} \\ l_{21} u_{11} \\ l_{31} u_{11} \end{bmatrix}, \quad a_2 = \begin{bmatrix} l_{11} u_{12} \\ l_{21} u_{12} + l_{22} u_{22} \\ l_{31} u_{12} + l_{32} u_{22} \end{bmatrix}$$

$$a_3 = \begin{bmatrix} l_{11} u_{13} \\ l_{21} u_{13} + l_{22} u_{23} \\ l_{31} u_{13} + l_{32} u_{23} + l_{33} u_{33} \end{bmatrix}$$

### Ax = b, LU Decomposition Formulas

Note that before  $Ax = b$  and we also have  $A = LU$ , thus  $LUx = b$ . With a substitution we get  $Ux = y$ , so  $Ly = b$ .

### LU Decomposition Inverse Formula

This is similar to normal  $LU$  decomposition. The formula takes the form of  $AX = I$  where all matrices are  $n$  by  $n$ , so  $X = [x_1 | x_2 | \dots | x_n]$  with the standard identity  $I = [I_1 | I_2 | \dots | I_n]$ .

### Gaussian Inverse Formula

The gaussian inverse does the following:

$$[A|I] \rightarrow [I|A^{-1}]$$

### Cholesky Factorization Formulas

This is similar to LU decomposition, only  $U = L^T$ . So:  $A = LL^T$  which means that:

$$L = \begin{bmatrix} l_{11} & 0 & 0 \\ l_{21} & l_{22} & 0 \\ l_{31} & l_{32} & l_{33} \end{bmatrix}, L^T = \begin{bmatrix} l_{11} & l_{21} & l_{31} \\ 0 & l_{22} & l_{32} \\ 0 & 0 & l_{33} \end{bmatrix}$$

$$A = \begin{bmatrix} l_{11}^2 & l_{11}l_{21} & l_{11}l_{31} \\ l_{21}l_{11} & l_{21}^2 + l_{22}^2 & l_{21}l_{31} + l_{22}l_{32} \\ l_{31}l_{11} & l_{31}l_{21} + l_{32}l_{22} & l_{31}^2 + l_{32}^2 + l_{33}^2 \end{bmatrix}$$

Then solving for the system and getting a cleaner  $L$  at the end gives the formula:

$$L = \begin{bmatrix} \sqrt{a_{11}} & 0 & 0 \\ \frac{a_{21}}{l_{11}} & \sqrt{a_{22} - l_{21}^2} & 0 \\ \frac{a_{31}}{l_{11}} & \frac{a_{32} - l_{31}l_{21}}{l_{22}} & \sqrt{A_{33} - l_{31}^2 - l_{32}^2} \end{bmatrix}$$

But the actual Cholesky factorization is just the matrix  $L$ .

### Gaussian Elimination Formulas

Gassian elimination is an  $Ax = b$  solving method. It makes an upper triangular matrix with the form:  $[A|b]$

## 4 Initial Value Problems

### RK4

This is the 4th order (RK4) Runge-Kutta method for the Initial Value Problem (IVP):

$$x(t+h) = x(t) + \frac{1}{6}(K_1 + 2K_2 + 2K_3 + K_4)$$

where the following are values of  $K_n$ :

$$K_1 = hf(t, x)$$

$$K_2 = hf(t + \frac{1}{2}h, x + \frac{1}{2}K_1)$$

$$K_3 = hf(t + \frac{1}{2}h, x + \frac{1}{2}K_2)$$

$$K_4 = hf(t + h, x + K_3)$$

first, the  $K_n$  values are calculated in succession. They the  $K_n$  values are filled into the first formula above.

### Modified Euler's Method

the Modified Euler's method is simply:

$$K_1 = f(t_n, x_n), K_2 = f(t_n + h, x_n + hK_1)$$

$$x_{n+1} = x(t_n) + \frac{h/2}{(} K_1 + K_2)$$

## 5 Generally Useful Maths

### Trig Properties

$$\sin^2 x + \cos^2 x = 1 \quad \sec x = \frac{1}{\cos x}$$

$$2 \sin x = \sin x \cos x$$

$$\tan x = \frac{\sin x}{\cos x} = \frac{1}{\cot x} \quad \csc x = \frac{1}{\sin x}$$

$$\frac{d}{dx} \sin x = \cos x \quad \frac{d}{dx} \cos x = -\sin x$$

$$\frac{d}{dx} \tan x = \sec^2 x \quad \frac{d}{dx} \cot x = -\csc^2 x$$

$$\frac{d}{dx} \arcsin x = \frac{1}{\sqrt{1-x^2}} \quad \frac{d}{dx} \arccos x = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \arctan x = \frac{1}{1+x^2} \quad \frac{d}{dx} \sec x = \sec x \tan x$$

### Log & Exp Properties

$$\log x^n = n \log x \quad \log\left(\frac{1}{x}\right) = -\log x$$

$$\log_a x = \frac{\log_b x}{\log_a b} \quad \frac{d}{dx} e^{ax} = ae^{ax}$$

$$x^0 = 1 \quad x^n \cdot x^m = x^{n+m} \quad x^{-n} = \frac{1}{x^n}$$

$$\log_a x^n = n \log_a x \quad \frac{e^{-nx}}{e^x} = e^{-(n+1)x}$$

$$\log_a\left(\frac{x}{y}\right) = \log_a x - \log_a y$$

$$\log_a(xy) = \log_a x + \log_a y$$

$$\frac{d}{dx} \ln x = \frac{1}{x}, \frac{d}{dx} a^{g(x)} = \ln(a) a^{g(x)} g'(x) \\ \frac{d}{dx} a^{g(x)} = \ln(a) a^{g(x)} g'(x), \frac{d}{dx} b^x = b^x \ln b \\ \frac{d}{dx} e^{g(x)} = g'(x) e^{g(x)}, \frac{d}{dx} a^x = a^x \ln a \\ \frac{d}{dx} \log_a(g(x)) = \frac{g'(x)}{\ln(a)g(x)}$$

### Other Derivative Rules

$$\frac{d}{dx} f(g(x)) = f'(g(x))g'(x)$$

$$\frac{d}{dx} f(x)/g(x) = \frac{(f'(x)g(x) - g'(x)f(x))}{g(x)^2}$$

## In Class Terminology

the relative error formula:  $\frac{|x-\hat{x}|}{\hat{x}}$

more generally, with  $\hat{x}, \hat{y}$  being rounded terms we get relative error as:

$$\frac{(x-y) - (\hat{x} - \hat{y})}{(x-y)} = \text{relative error}$$

## 6 Boundary Value Problems

Can be descretized with finite difference methods. The formula goes from  $t_0, \Delta t, \dots, t_n$ . The fromulas for the known boundary values are of the form:  $x(t_0) = \alpha$  and  $x(t_n) = \beta$ .

### Jacobi Method

Solves  $Ax = b$ . Solve the system of equations by guessing at  $x_n$  valeus. starts with initial guess  $x = [0 \ 0 \ 0]^T$ , converges on an answer after iteration. Uses the formula:

$$x_i^{(k+1)} = \frac{1}{a_{ii}} (b_i - \sum_{j \neq i} a_{ij} x_j^{(k)}), i = 1, \dots, n.$$

### Gauss-Seidel Method

Just like Jacobi, only it substitutions during iteration. Uses formula:  $x_i^{(k+1)} =$

$$\frac{1}{a_{ii}} (b_i - \sum_j^{i-1} a_{ij} x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)})$$

## 7 PDE's

Solved like BVPs.

### Classification Formulas

Discriminant gives classification:

$B^2 - 4AC$ , with form:

$$A \frac{\partial^2 u}{\partial x^2} + B \frac{\partial^2 u}{\partial x \partial t} + C \frac{\partial^2 u}{\partial t^2} + D \frac{\partial u}{\partial x} + E \frac{\partial u}{\partial t} + Fu = G$$

$B^2 - 4AC = 0$ , Parabolic

$B^2 - 4AC > 0$ , Hyperbolic

$B^2 - 4AC < 0$ , Elliptic

### Crank-Nicholson

the general form of the problem uses the equation  $s = \frac{\Delta x^2}{\Delta t}$ .