

Il leone delle caverne



Il bisonte della Steppa

A steppe bison on display at the University of Alaska Museum of the North in Fairbanks. The steppe bison is one of several extinct large mammals that roamed interior Alaska during the Wisconsinan glacial period, 100,000 to 10,000 years ago. This specimen died about 36,000 years ago and was found during the summer of 1979. It has a bluish color over the entire carcass, caused by the phosphorus in the animal tissue reacting with the iron in the soil to produce a mineral coating of vivianite - which became a brilliant blue when it was exposed to air. Hence the name Blue Babe.



Si salvarono le specie più piccole e quelle diffuse anche in Africa e/o in Asia

I daini (gen. *Dama*) sopravvissero in Asia Minore, camosci (gen. *Rupicapra*) e stambecchi (gen. *Capra*) nelle regioni alpine, le antilopi Saiga (*Saiga tatarica*) ed i cavalli di Przewalski (*Equus ferus*) in Asia centrale, i buoi muschiati (*Ovibos moschatus*) nell'estremo Nord (Groenlandia), renne e caribù (*Rangifer tarandus*) rispettivamente nel Nord dell'Eurasia e del Nordamerica, ippopotami (*Hippopotamus amphibius*), bufali d'acqua (*Bubalus murrensis*), leopardi (*Panthera pardus*) e leoni (*Panthera leo*) in Africa e questi ultimi anche in Asia

Il bue muschiato

- Allevato in Alaska per la lana



Diametro della fibra di lana in alcuni mammiferi

DIAMETRO DELLA FIBRA (μm)	RAZZA
4÷30	Bue muschiato
12÷13*	Changthangi (Kashmir Pashmina) cashmere
13,00÷>40,00	Gentile di Puglia
17,70÷19,14	Merino
18÷25	Alpaca
≤ 19	Cashmere
19,15÷20,59	Rambouillet
20,60÷22,04	Targhee, Southdown
22,05÷23,49	Corriedale, Columbia
23,50÷30,99	Panama, Romeldale
25÷45	Angora
26,40÷32,69	Shropshire, Hampshire, Suffolk, Oxford, Dorset, Cheviot
32,70÷38,09	Romney
36,20÷40,20	Costwold
32,70÷40,20	Lincoln
32,00÷50,00	Sambucana

L'alpaca

L'**alpaca** (*Vicugna pacos*) è un camelide originario del Sudamerica, discendente dalla **vigogna** (*Lama vicugna*), allevato soprattutto per utilizzarne la pregiata lana



La razza ovina merino

Merino wool is fine and soft. Staples are commonly 65÷100 mm long. A Saxon Merino produces 3÷6 kg of greasy wool a year, while a good quality Peppin Merino ram (ariete) produces up to **18 kg**. Merino wool is generally < 24 µm in diameter.

Basic Merino types include:

- **strong** (broad) wool (23÷24.5 µm)
- **medium** (21÷22.9 µm)
- **fine** (18.6÷20.9 µm)
- **superfine** (15÷18.5 µm)
- **ultra-fine** (11.5÷15 µm)



La razza ovina gentile di Puglia

The Gentile di Puglia derives from cross-breeding local ewes with Merino rams brought from Spain, first by Alfonso V of Aragon **in the fifteenth century**, and later, repeatedly, by the Bourbon kings of Naples, who had extensive estates near Foggia.

In the nineteenth century, after the Unification of Italy, there was cross-breeding with imported French Rambouillet and German Merinolandschaf animals, with the aim of further improving the quality of the wool.



La razza ovina gentile di Puglia

The collapse of the wool trade in the later twentieth century caused a sharp fall in number of the breed. Various indiscriminate attempts were made to improve the meat yield by cross-breeding with other types, some of them imported. In the 1980s a study of 10,000 head found 13% to be pure-bred stock. As a result, a controlled cross-breeding and selection programme was started, which led to the creation of the Trimeticcia di Segezia breed.

The Gentile di Puglia is one of the seventeen autochthonous Italian sheep breeds for which a genealogical herdbook is kept by the Associazione Nazionale della Pastorizia, the Italian national association of sheep-breeders. The herdbook was established in 1971. In 1983 total numbers for the breed were estimated at 500,000, of which 31,700 were registered in the herdbook; in 2013 the number recorded in the herdbook was 3532.

Il cavallo di Przewalski



L'antilope saiga




L'Olocene

The natural (mid- to late-Holocene) guild of large herbivores in Europe comprises **European bison** (*Bison bonasus*); **aurochs** (*Bos primigenius*), now globally extinct as a wild species; **wild horse** (*Equus ferus*), now globally extinct as a wild species; **Eurasian elk** (*Alces alces*), also known as moose; **red deer** (*Cervus elaphus*); **roe deer** (*Capreolus capreolus*); **fallow deer** (*Dama dama*); **reindeer** (*Rangifer tarandus*); **wild boar** (*Sus scrofa*); **muskox** (*Ovibos moschatus*); **chamois** (*Rupicapra rupicapra*); and **ibex** (*Capra sp.*).


Early Holocene sea level rise

The early Holocene sea level rise (EHSLR) was a significant jump in sea level by about 60 m during the early Holocene, between about 12,000 and 7,000 years ago, spanning the **Eurasian Mesolithic**.

Early Holocene sea level rise Post-Glacial Sea Level Rise



Doggerland and Doggerbank



Cosquer Cave

Cosquer Cave is located in the Calanque de Morgiou in Marseille, France, near Cap Morgiou. The entrance to the cave is located 37 m (121 ft) underwater, due to the Holocene sea level rise, and contains various prehistoric rock art engravings (incisioni). Its submarine entrance was discovered in 1985 by Henri Cosquer, a professional diver. The underwater passage leading to the cave was progressively explored until 1990 by cave divers, without the divers being aware of the archaeological character of the cave.

Only in the last period (1990-91) of the progressive underwater explorations did the cave divers emerge in the non-submerged part of the cave. The prehistoric paintings were not immediately discovered by the divers to first emerge from the other side of the sump. The cave was named after Henri Cosquer when its existence was made public in 1991, after three divers became lost in the cave and died.


Cosquer Cave

The cave can now be accessed by divers through a 175 m (574 ft) long tunnel; the entrance is located 37 m (121 ft) below sea level, which has risen since the cave was inhabited. During the glacial periods of the Pleistocene, the shore of the Mediterranean was several kilometers to the south and the sea level up to 100 m (330 ft) below the entrance of the cave.

Cosquer Cave



Cosquer Cave



La transizione dal Paleolitico al Mesolitico

Già in crisi per la progressiva scomparsa dei loro pascoli, i grandi erbivori pleistocenici furono sottoposti ad una persecuzione spietata ed implacabile da parte di sempre più numerose comunità di cacciatori-raccoglitori: l'opera di sterminio fu resa ancora più efficiente dall'affinarsi delle tecniche di lavorazione della pietra e dei palchi dei cervidi (localmente delle zanne di mammut), materiali con cui venivano fabbricate le punte delle lance e delle frecce.




Fig. 1 — Reconstruction of the methods of hafting and using microliths in the projectile weapons. 1: Pushkari Culture; 2: Aurignacian Culture; 3-6: Gravettian Culture; 7-10: Early Mesolithic Shan-Koba Culture; 11-17: Late Mesolithic and Neolithic Cultures of Ukraine.

La transizione dal Paleolitico al Mesolitico

The bow and arrow were much better adapted to the denser forest cover: This new weapon was precise, rapid and silent



La transizione dal Paleolitico al Mesolitico

Cacciatori mesolitici seguono le tracce di bovidi (camosci e stambecchi) in una valle alpina





Non tagliate i grandi alberi!

Sono la «casa» di una miriade di creature!




In questa foto: le cavità nel tronco di una farnia (*Quercus robur*) danno rifugio a molti animali.

Questa si trova sull'Isola di Öland, in Svezia, dai tempi di Linneo, che la vide nel 1741.

Pagina a lato, sopra: i corpi fruttiferi, tondi e arancioni, di funghi saprofitti; sotto: *Trametes versicolor*, fungo a mensola.

Non tagliate i grandi alberi!

- Chirotteri (Pipistrelli), diverse specie, alcune delle quali minacciate di estinzione e/o scoperte solo recentemente



Non tagliate i grandi alberi!

- Coleotteri Cerambicidi
- Strigiformi (rapaci notturni)
- Picchi...



Chi se ne avvantaggia?



Chi se ne avvantaggia?

In a Mediterranean area, where agricultural crops are not available and supplementary food is not provided, diet varies according to the availability of energy-rich foods such as acorns and olives; pine-seeds are actively consumed, even when their availability is low. When abundant, acorns and olives account for most of the diet, and when scarce are replaced by graminoids and juniper berries. In summer, graminoids and pine-seeds account for most of the diet.

Following a high production of acorns and olives, **wild boar exhibit higher body weight, more breeding females and a larger litter size** than in years of poor production of these foods.



Ma dopo un anno di pasciona...

Domenica 13 ottobre 2024

AREA RISERVATA



CUNEO PAESI CRONACA POLITICA CHIESA ECONOMIA SANITÀ SPORT SPETTACOLI

ABBONATI



EVENTI È Nino Haratischwili la vincitrice della XIV edizione del Premio Lattes Grinzane 2024 Borgo, i vincitori delle borse di studio "Sebastiano Grandi"

Castagne, un'annata pesantissima per i boschi del cuneese

Nei dati diffusi da Cia Cuneo, stagione in ritardo e cali di produzione (in certi casi vicina allo zero)

Cuneo

Giovedì 3 ottobre 2024



Il 2024 si avvia ad essere archiviato come l'annus horribilis delle castagne in provincia di Cuneo. La stagione non è conclusa, ma i primi dati sulla raccolta, in ritardo di un paio di

Sono tornati gli **ECOINCENTIVI**

RISPARMIA FINO A **10.000€**

PRENOTA SUBITO IL TUO!

AZZURRA CENTER

Il Consiglio per la scuola



