

Supplementary Data for “Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation”

Raw assemblies and polished results at <http://gembox.cbcb.umd.edu/shared/canu/index.html>

| | |
|--|----|
| Supplementary Note 1: Kmer Filtering..... | 3 |
| Table S1: Overlap Sensitivity With and Without Filtering | 3 |
| Supplementary Note 2: Sequencing Data | 5 |
| Supplementary Note 3: Low Coverage Assembly | 5 |
| Table S2: Continuity and QV statistics for <i>A. thaliana</i> hybrid versus hierachical. | 5 |
| Figure S1: Canu 20X+Quiver, Canu 20X+Pilon, and SPAdes 100X+20X and SPAdes 100X+150X <i>A. thaliana</i> assemblies..... | 6 |
| Supplementary Note 4: Assemblers | 6 |
| Falcon | 6 |
| SPAdes | 7 |
| Miniasm | 7 |
| Canu..... | 7 |
| Supplementary Note 5: Polishing | 8 |
| Supplementary Note 6: Validation | 8 |
| Supplementary Note 7: Assembler Runtimes..... | 9 |
| Table S3: Assembly Time Canu (CPU hours). | 9 |
| Table S4: Assembly Time Falcon (CPU hours). | 9 |
| Table S5. Assembly Time Miniasm (CPU hours). | 10 |
| Table S6. Assembly Time SPAdes (CPU hours). | 10 |
| Supplementary Note 8: PacBio Assembly Statistics with Quiver..... | 11 |
| Table S7 Quiver Statistics For Canu. | 11 |
| Table S8 Quiver Statistics For Falcon. | 11 |
| Table S9 Quiver Statistics For Miniasm | 12 |
| Table S10 Quiver Assembly Statistics For SPAdes | 12 |
| Figure S2 Canu, Falcon, Miniasm, and SPAdes <i>E. coli</i> K12 polished assemblies. | 12 |
| Figure S3 Canu, Falcon, and Miniasm <i>D. melanogaster</i> assemblies..... | 13 |
| Figure S4 Canu, Falcon, and Miniasm <i>A. thaliana</i> assemblies. | 13 |
| Figure S5 Canu, Falcon, and Miniasm <i>C. elegans</i> assemblies. | 13 |
| Figure S6 Canu and Falcon assemblies of <i>H. sapiens</i> CHM1. | 14 |
| Supplementary Note 9: Oxford Nanopore Assemblies with Nanopolish..... | 15 |
| Table S11 Nanopolish Statistics for Canu | 15 |
| Table S12 Nanopolish Statistics for Falcon | 15 |
| Table S13 Nanopolish Statistics for Miniasm | 16 |
| Figure S7 Canu, Falcon, and Miniasm assemblies of <i>E. coli</i> K12 MAP005. | 16 |
| Figure S8 Canu, Falcon, and Miniasm assemblies of <i>E. coli</i> K12 MAP006-1. | 16 |

| | |
|--|----|
| Figure S9 Canu, Falcon, and Miniasm assemblies of <i>E. coli</i> K12 MAP006-2. | 17 |
| Figure S10 Canu, Falcon, and Miniasm assemblies of <i>E. coli</i> K12 MAP006-PCR-1. | 17 |
| Figure S11 Canu, Falcon, and Miniasm assemblies of <i>E. coli</i> K12 MAP006-PCR-2. | 17 |
| Figure S12 Canu, Falcon, and Miniasm assemblies of <i>B. anthracis</i> | 18 |
| Figure S13 Canu, Falcon, and Miniasm assemblies of <i>Y. pestis</i> | 18 |
| Supplementary Note 10: Oxford Nanopore assemblies with Pilon | 19 |
| Table S14 Pilon Statistics for Miniasm after 300 rounds of pilon | 19 |
| Figure S14 Minimap remaining errors in <i>E. coli</i> K12 MAP006-1. | 19 |
| Table S15 Pilon Statistics for Canu | 20 |
| Table S16 Pilon Statistics for Falcon | 20 |
| Table S17 Pilon Statistics for Miniasm | 21 |
| Table S18 Pilon Statistics for Spades | 21 |
| Figure S15 Canu, Falcon, Miniasm, and SPAdes assemblies of <i>E. coli</i> K12 MAP005. | 21 |
| Figure S16 Canu, Falcon, Miniasm, and SPAdes assemblies of <i>E. coli</i> K12 MAP006-1. | 22 |
| Figure S17 Canu, Falcon, Miniasm, and SPAdes assemblies of <i>E. coli</i> K12 MAP006-2. | 22 |
| Figure S18 Canu, Falcon, Miniasm, and SPAdes assemblies of <i>E. coli</i> K12 MAP006-PCR-1. | 22 |
| Figure S19 Canu, Falcon, Miniasm, and SPAdes assemblies of <i>E. coli</i> K12 MAP006-PCR-2. | 22 |
| Figure S20 Canu, Falcon, Miniasm, and SPAdes assemblies of <i>B. anthracis</i> | 23 |
| Figure S21 Canu, Falcon, Miniasm, and SPAdes assemblies of <i>Y. pestis</i> | 23 |
| Supplementary Note 11: 1D Assembly | 23 |
| Figure S22: Canu 1D, Canu+three rounds of Nanopolish, Canu 1D+three rounds of Pilon, and Miniasm assemblies for <i>E. coli</i> K12 MAP006-1 1D. | 24 |
| Supplementary Note 12: <i>S. cerevisiae</i> assembly | 24 |
| Figure S23: Canu 1D, and Canu 1D+three rounds of Pilon, and Canu PacBio 115X assemblies. | 24 |
| Figure S24: Parallel overlap store construction | 25 |
| Figure S25: Suspicious read detection. | 25 |
| References | 26 |

Supplementary Note 1: Kmer Filtering

We used the *B. anthracis* Oxford Nanopore MinION dataset (Sequencing). All sequences were mapped to the reference *B. anthracis* str. ‘Ames Ancestor’ (GCF_000008445.1) with blasr (Chaisson and Tesler 2012):

```
blasr ba_filtered.fasta ref.fna -sa ref.sa -bestn 10 -maxAnchorsPerPosition 100 -  
advanceExactMatches 10 -affineAlign -affineOpen 100 -affineExtend 0 -insertion 5 -deletion 5 -  
extend -maxExtendDropoff 20 -nproc 8 -m 4 -out merged.blasr.m4
```

No significant mappings to pXO2 were found. K-mers predominantly coming from pXO1 were identified to have copy-number >40. Thus, all k-mers with >=40 occurrences were flagged as repeat and passed in a filter file to MHAP. We ran a sweep of suppression parameters including no suppression and full suppression and evaluated sensitivity on both pXO1 and the chromosome.

Table S1: Overlap Sensitivity With and Without Filtering

| Program | Filtering | Sensitivity on Chromosome | PPV on Chromosome | Sensitivity on pXO1 | PPV on pXO1 |
|----------------|---------------|---------------------------|-------------------|---------------------|-------------|
| MHAP | None | 90.32% | 99.52% | 89.69% | 99.55% |
| | 0.1 | 90.30% | 99.59% | 87.63% | 99.60% |
| | 0.2 | 90.30% | 99.62% | 87.82% | 99.66% |
| | 0.3 | 90.30% | 99.64% | 87.86% | 99.69% |
| | 0.4 | 90.30% | 99.61% | 88.00% | 99.65% |
| | 0.5 | 90.30% | 99.63% | 88.15% | 99.64% |
| | 0.6 | 90.30% | 99.56% | 88.23% | 99.65% |
| | 0.7 | 90.30% | 99.63% | 88.53% | 99.65% |
| | 0.8 | 90.30% | 99.57% | 88.77% | 99.60% |
| | 0.9 | 90.30% | 99.53% | 89.05% | 99.58% |
| | Full | 89.68% | 99.83% | 26.34% | 99.82% |
| MHAP sensitive | 0.9 | 96.37% | 92.96% | 95.65% | 93.02% |
| Daligner | Default | 75.74% | 100.00% | 59.68% | 100.00% |
| | -t1000 -M0 | 75.96% | 99.98% | 76.04% | 99.98% |
| Minimap | Default | 93.54% | 89.22% | 16.99% | 89.23% |
| | -f 0.00000001 | 95.22% | 84.63% | 94.49% | 84.64% |

Overlaps were evaluated as in Berlin et. al (Berlin et al. 2015) using the commands:

```
java -cp mhap-2.1test.jar edu.umd.marbl.mhap.main.EstimateROC chromosome.m4 raw_reads.ovls
ba_filtered.fasta 2000 10000 true false 0.7 0.2 true
java -cp mhap-2.1test.jar edu.umd.marbl.mhap.main.EstimateROC pX01.m4 raw_reads.ovls ba_filtered.fasta
2000 10000 true false 0.7 0.2 true
```

Minimap was run with the commands:

```
minimap -k 15 -Sw5 -L100 -m0 -t8 -I6G ba_filtered.fasta ba_filtered.fasta > minimap.paf
paf2mhap.pl ba_filtered.fasta minimap.paf > minimap.ovls
```

```
minimap -f 0.00000001 -k 15 -Sw5 -L100 -m0 -t8 -I6G ba_filtered.fasta ba_filtered.fasta >
minimap.sens.paf
paf2mhap.pl ba_filtered.fasta minimap.sens.paf > minimap.sens.ovls
```

Daligner was run with the commands:

```
daligner -v -t16 -H1000 -e0.7 -s1000 raw_reads raw_reads
LASort -v raw_reads.raw_reads.C0 raw_reads.raw_reads.N0 raw_reads.raw_reads.C1 raw_reads.raw_reads.N1
raw_reads.raw_reads.C2 raw_reads.raw_reads.N2 raw_reads.raw_reads.C3 raw_reads.raw_reads.N3 && LAmerge -v
raw_reads.1 raw_reads.raw_reads.C0.S raw_reads.raw_reads.N0.S raw_reads.raw_reads.C1.S
raw_reads.raw_reads.N1.S raw_reads.raw_reads.C2.S raw_reads.raw_reads.N2.S raw_reads.raw_reads.C3.S
raw_reads.raw_reads.N3.S && rm raw_reads.raw_reads.C0.S.las raw_reads.raw_reads.N0.S.las
raw_reads.raw_reads.C1.S.las raw_reads.raw_reads.N1.S.las raw_reads.raw_reads.C2.S.las
raw_reads.raw_reads.N2.S.las raw_reads.raw_reads.C3.S.las raw_reads.raw_reads.N3.S.las
LAsow raw_reads raw_reads.1.las > default.ovls
```

```
daligner -v -t1000 -H1000 -e0.7 -s1000 raw_reads raw_reads
LASort -v raw_reads.raw_reads.C0 raw_reads.raw_reads.N0 raw_reads.raw_reads.C1 raw_reads.raw_reads.N1
raw_reads.raw_reads.C2 raw_reads.raw_reads.N2 raw_reads.raw_reads.C3 raw_reads.raw_reads.N3 && LAmerge -v
raw_reads.1 raw_reads.raw_reads.C0.S raw_reads.raw_reads.N0.S raw_reads.raw_reads.C1.S
raw_reads.raw_reads.N1.S raw_reads.raw_reads.C2.S raw_reads.raw_reads.N2.S raw_reads.raw_reads.C3.S
raw_reads.raw_reads.N3.S && rm raw_reads.raw_reads.C0.S.las raw_reads.raw_reads.N0.S.las
raw_reads.raw_reads.C1.S.las raw_reads.raw_reads.N1.S.las raw_reads.raw_reads.C2.S.las
raw_reads.raw_reads.N2.S.las raw_reads.raw_reads.C3.S.las raw_reads.raw_reads.N3.S.las
LAsow raw_reads raw_reads.1.las > sens.ovls
```

Truth overlaps were obtained from a bwa mem (Li 2013) mapping to the reference genome:

```
bwa mem -x ont2d b_anthraxis.fna ba_filtered.fasta -t 16 samtools view -b -S - > mapping.bam
convertSam mapping.bam b_anthraxis.fna > mapping.m4
cat mapping.m4 |grep NC_007530.2 > chromosome.m4
cat mapping.m4 |grep NC_007322.2 > pX01.m4
```

Output was converted to BLASR's m4 format. Minimap overlaps were converted to MHAP format using the provided paf2mhap.pl script. Overlaps not found based on the reference mapping were confirmed using full Smith-Waterman alignment. If no alignment of at least 65% identity could be found, the overlap was considered a false-positive. While MHAP without filtering does not suffer the same PPV penalty as Minimap (due to its second-stage filter (Berlin et al. 2015) which checks the estimated mutation rate), performance suffers as the average number of sequences hit by a k-mer lookup increases from an average of 24.42 with filtering to 25.72 and the % of sequences which make it past the second-stage filter drops from an average of 78.29 to 77.06%. DALIGNER runtime increases 2-fold (from 1.28 CPU h to 2.58) and memory use increases 1.6-fold (from 38G to 61G).

Supplementary Note 2: Sequencing Data

Illumina MiSeq data for *E. coli* K12 was downloaded from the Illumina Scientific Data website (http://www.illumina.com/systems/miseq/scientific_data.html) and used for all analysis requiring Illumina data (Pilon polishing/SPAdes assembly). Illumina MiSeq *S. cerevisiae* W303 data was downloaded from Goodwin et. al. (Goodwin et al. 2015) (<http://schatzlab.cshl.edu/data/nanocorr/>). Illumina MiSeq *A. thaliana* Ler-0 data was downloaded from Lee et. al. (Lee et al. 2014) (<http://schatzlab.cshl.edu/data/ectools>). All datasets were downsampled to 100X before assembly.

Ba + Yp from NBACC, trimmed with Trimmomatic:

```
java -jar trimmomatic-0.36.jar PE -phred33 input.r1.fastq.gz input.r2.fastq.gz output.r1.fastq
output.r1.bad.fastq output.r2.fastq output.r2.bad.fastq ILLUMINACLIP:NexteraPE-PE.fa:2:30:10
LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:36
```

Only the surviving paired-end sequences were used for analysis.

PacBio SMRT sequencing was downloaded from DevNet (<https://github.com/PacificBiosciences/DevNet/wiki/Datasets>) and is available from NCBI SRA. *E. coli* K12 [https://s3.amazonaws.com/files.pacb.com/datasets/secondary-analysis/e-coli-k12-P6C4/p6c4_ecoli_RSII_DDR2_with_15kb_cut_E01_1.tar.gz], *C. elegans* [http://datasets.pacb.com.s3.amazonaws.com/2014/c_elegans/list.html], *A. thaliana* [SRX533607], *D. melanogaster* [SRX499318], *H. sapiens* [PRJNA246220]. In all cases the raw H5 files and fastq sequences were downloaded. Fastq sequences were used for assembly.

Oxford Nanopore data from Loman et. al. (Loman et al. 2015) was downloaded from EBI (ERX708228, ERX708229, ERX708230, ERX708231) sequencing *E. coli* K12 MG1665 and co-assembled as one dataset (MAP005). Four additional runs were generated for *E. coli* K12 by the same lab and available at (<http://lab.loman.net/2015/09/24/first-sqk-map-006-experiment/>). Raw sequences were downloaded from EBI (ERR1147227 (MAP006-1), ERR1147228 (MAP006-2), ERR1147229 (MAP006-PCR-1), ERR1147230 (MAP006-PCR-2) and assembled individually.

Supplementary Note 3: Low Coverage Assembly


Table S2: Continuity and QV statistics for *A. thaliana* hybrid versus hierarchical.

SPAdes 100X+20X and SPAdes 100X+150X contained 79,651 and 79,025 contigs, respectively. Only contigs greater than 1000 bp (2,696 and 2,428 respectively) were included for the analysis. The filtered contig set did not decrease the percentage of the reference covered by the full set, indicating the shorter contigs are likely redundant or assembly artifacts. SPAdes 100X+20X has a smaller max and NG50 than Canu but a higher accuracy than the initial assembly. Polishing with Quiver or Pilon increases the QV to one that is comparable with SPAdes. Despite having a higher max contig, the SPAdes 100X+150X NG50 is almost equivalent to Canu with comparable error statistics, despite having over 10-fold more sequencing data. Same columns as table S7.

| Assembler | Assembly | | | | | Polishing | | | | |
|---------------------------------|----------|----------|--------|---------------|-------|-----------|----------|--------|---------------|-------|
| | Max (mb) | N50 (mb) | Errors | % ref covered | QV | Max (mb) | N50 (mb) | Errors | % ref covered | QV |
| Canu 20X + Quiver | 4.54 | 1.16 | 185 | 79.07% | 19.23 | 4.54 | 1.17 | 183 | 81.83% | 20.11 |
| Canu 20X + Pilon 100X | 4.54 | 1.16 | 185 | 79.07% | 19.23 | 4.54 | 1.16 | 201 | 81.91% | 20.31 |
| SPAdes 100X+20X + Pilon 100X | 3.71 | 0.84 | 156 | 82.23% | 20.32 | 3.71 | 0.84 | 155 | 82.24% | 20.32 |
| SPAdes 100X + 150X + Pilon 100X | 5.69 | 1.23 | 163 | 82.40% | 20.32 | 5.69 | 1.23 | 164 | 82.43% | 20.32 |

Figure S1: Canu 20X+Quiver, Canu 20X+Pilon, and SPAdes 100X+20X and SPAdes 100X+150X *A. thaliana* assemblies.

The plot shows the best (1-to-1) alignments between the reference (x-axis) and each assembly (y-axis). Red lines indicate forward-strand matches while blue lines indicate reverse-complement matches. Dashed vertical lines delineate chromosome ends while dashed horizontal lines delineate contigs. A diagonal indicates concordant matches while off-diagonal matches indicate assembly errors or differences versus the reference.



Supplementary Note 4: Assemblers

Falcon

Falcon v0.4.1 was checked out on 3/16/2016 (commit c602aad3667b3fd49263028dac44da8e42caa17c). Each test was run using the configurations provided in examples, when available. The LG configuration was used for CHM1. No configuration was available for *C. elegans* so the same one as *D. melanogaster* was used. For Oxford Nanopore datasets, read names were altered using the <https://github.com/jts/nanocorrect/blob/master/nanocorrect-preprocess.pl> script to match DALIGNER expectation.

SPAdes

SPAdes v3.7.1 was used for all experiments and run with the command:

```
spades.py -t 48 -m 128 -o asm -1 <illumina.1.fastq> -2 <illumina.2.fastq> --nanopore  
<ont.fasta>
```

for Oxford Nanopore datasets and

```
spades.py -t 48 -m 128 -o asm -1 <illumina.1.fastq> -2 <illumina.2.fastq> --pacbio <pac.fasta>
```

For PacBio SMRT datasets.

Miniasm

Minimap/miniasm was checked out of github on 3/16/2016 (commit 1cd6ae3bc7c7a6f9e7c03c0b7a93a12647bba244 minimap, 17d5bd12290e0e8a48a5df5afaeaeef4d171aa133 miniasm). Minimap/miniasm ran as specified in the misc/demo-worm-pacbio.sh script with the commands:

```
minimap -k 15 -Sw5 -L100 -m0 -t48 -I6G reads.fasta reads.fasta |gzip -1 > reads.paf.gz  
miniasm -f reads.fasta reads.paf.gz > reads.gfa
```

GFA primary sequences were converted to a fasta file for downstream analysis.

Canu

Canu version 1.3 was used for all experiments. The default Canu error rates, 0.05 for Oxford Nanopore data and 0.025 for PacBio data, are designed to work on a variety of datasets. The error rate is an upper bound on overlaps used in assembly and runtime can be improved by decreasing it. MAP005 *E. coli* ran with the default parameters. Newer Oxford Nanopore datasets, MAP006 *E. coli* datasets and *Y. pestis*, which are more similar to PacBio sequence error rates, ran with errorRate=0.025. *B. anthracis* used sensitive parameters (corMhapSensitivity=high corOutCoverage=80 errorRate=0.04). The *Y. pestis* had one high-coverage unassembled unitig with 15 reads which was included with the assembled contigs for downstream analysis and polishing. PacBio experiments used errorRate=0.013, as suggested for high-coverage PacBio datasets on the online FAQ (<http://canu.readthedocs.io/en/latest/faq.html>), matching the corrected read error rate used by Falcon.

Supplementary Note 5: Polishing

For PacBio data raw H5 bax.h5 were input to Quiver from SMRTportal 2.3.0 patch4 using the SGE pipeline available at <https://github.com/skoren/QuiverGrid>

For Oxford Nanopore data, two alternative polishing strategies were used. First using Nanopolish (Loman et al. 2015) checked out on 03/16/2016 (commit aba1b3201f46b4a00ae7463da3d2e3142366fd0e) using the SGE pipeline available at <https://github.com/skoren/NanoGrid>.

Second, using complementary Illumina data, polishing was performed with Pilon 1.13 with the commands:

```
bwa index <asm.fasta>
bwa map <asm.fasta> <illumina.1.fastq> <illumina.2.fastq>
java -jar pilon.jar --fix bases,local --genome <asm.fasta> --output <asm.pilon.fasta> --
changes --vcf --diploid
using the SGE pipeline available at https://github.com/skoren/PilonGrid.
```

Supplementary Note 6: Validation

Validation was performed using MUMer3.23 (Kurtz et al. 2004) and dnadiff (Phillippy et al. 2008). For *E. coli*, *B. anthracis*, and *Y. pestis* the command used was

```
dnadiff reference.fasta asm.fasta
```

For the other genomes to improve runtime, the commands were

```
nucmer -l 100 -c 1000 reference.fasta asm.fasta
dnadiff -d out.delta
```

For the unpolished 1D *E. coli* and the 1D+2D *S. cerevisiae* assemblies reference alignments were run with relaxed MUMmer parameters, to identify low-identity matches, for both Canu and Miniasm:

```
nucmer -l 10 -c 100 reference.fasta
dnadiff -d out.delta
```

Percent reference covered and identity was reported from 1-to-1 alignments and errors were calculated as the sum of Relocations, Translocations, and Inversions. Validation was performed after assembly and after each round of polishing.

For *E. coli* K12 the reference used was NC_000913.3. For *B. anthracis* the reference was GCF_000008445.1. For *Y. pestis* the reference was GCF_000009065.1. For *D. melanogaster* the reference was Release 6 (GCF_000001215.4). For *A. thaliana* the reference was TAIR10 (GCF_000001735.3). For *C. elegans* the reference was GCF_000002985.6. for *H. sapiens*, the reference was GRCh38 (GCF_000001405.34).

Supplementary Note 7: Assembler Runtimes

All tests ran on an SGE grid composed of 24 hosts, each with dual Intel(R) Xeon(R) CPU E5-2680 v3 @ 2.50GHz CPUs (24-cores, 48 hyperthreads) and 128GB of ram. Canu and Falcon were allowed to use SGE to submit processes to the grid using built-in grid management code. Since Miniasm and SPAdes are designed to run on a single shared-memory machines, each was submitted to a single machine and allowed exclusive use of a machine.

Table S3: Assembly Time Canu (CPU hours).

Times were broken out to multiple steps, including the time to build MHAP indices (Raw MHAP index), the time to compute overlaps given the index (Raw MHAP overlapping), time to compute alignments and generate corrected read consensus sequences (read correction), and time to compute correct read overlaps for both the trimming and unitigging Canu stages. In all cases for raw data, Canu computed the all-vs-all overlaps for the input raw data.

| Genome | Raw MHAP Index | Raw MHAP Overlapping | Read Correction | Trim overlapInCore Overlapping | Unitigging overlapInCore Overlapping | Total Assembly Time |
|------------------------|----------------|----------------------|-----------------|--------------------------------|--------------------------------------|---------------------|
| <i>E. coli</i> | 1.45 | 0.35 | 1.95 | 0.12 | 0.19 | 4.26 |
| <i>D. melanogaster</i> | 28.29 | 36.85 | 192.25 | 88.68 | 215.71 | 573.65 |
| <i>A. thaliana</i> | 46.32 | 116.65 | 80.90 | 151.74 | 157.87 | 577.49 |
| <i>C. elegans</i> | 17.99 | 9.11 | 36.30 | 14.64 | 88.22 | 170.14 |
| CHM1 | 455.81 | 2340.66 | 1,624.57 | 6,945.50 | 6,401.53 | 18,019.70 |
| MAP005 | 0.60 | 0.15 | 1.97 | 3.05 | 3.21 | 9.38 |
| MAP006-1 | 0.80 | 0.17 | 2.09 | 0.55 | 1.00 | 4.85 |
| MAP006-2 | 0.56 | 0.08 | 1.27 | 0.29 | 0.43 | 2.78 |
| MAP006-PCR-1 | 0.57 | 0.52 | 1.47 | 0.12 | 0.19 | 3.04 |
| <i>B. anthracis</i> | 2.63 | 0.84 | 7.61 | 23.77 | 26.15 | 63.13 |
| <i>Y. pestis</i> | 1.59 | 0.76 | 3.38 | 1.07 | 5.53 | 12.55 |

Table S4: Assembly Time Falcon (CPU hours).

Times were broken out as for Canu, with time to generate raw overlaps (raw DALIGNER overlapping), time to align and generate corrected consensus sequences (read correction), and time to overlap the corrected sequences. In all cases DALIGNER uses a seed read length threshold (set in the Falcon spec) to limit overlaps to only the longest sequences.

| Genome | Raw DALIGNER Overlapping | Read Correction | Corrected DALIGNER Overlapping | Total Assembly Time |
|-----------------|--------------------------|-----------------|--------------------------------|---------------------|
| E. coli | 2.10 | 7.10 | 0.86 | 10.08 |
| D. melanogaster | 564.91 | 273.94 | 89.29 | 930.76 |
| A. thaliana | 654.47 | 140.18 | 39.92 | 836.61 |
| C. elegans | 70.55 | 99.21 | 32.68 | 202.74 |
| CHM1 | 51,282.2 | 2,252.99 | 10,472.40 | 64,087.80 |
| MAP005 | 0.18 | 2.63 | 0.16 | 2.98 |
| MAP006-1 | 0.41 | 4.14 | 0.36 | 4.93 |
| MAP006-2 | 0.14 | 1.31 | 0.11 | 1.56 |
| MAP006-PCR-1 | 0.14 | 1.47 | 0.12 | 1.74 |
| MAP006-PCR-2 | 0.53 | 6.44 | 0.43 | 7.42 |
| B. anthracis | 2.8 | 9.67 | 0.26 | 12.75 |
| Y. pestis | 0.48 | 3.86 | 0.22 | 4.57 |

Table S5. Assembly Time Miniasm (CPU hours).

The time for both steps in miniasm, overlapping (minimap) and assembly (unitigging with miniasm) are reported. In all cases, minimap computed all-vs-all overlaps for the input raw data.

| Genome | Raw minimap overlapping | Unitigging | Total Assembly Time |
|---------------------|-------------------------|------------|---------------------|
| E. coli | 0.17 | 0.01 | 0.18 |
| D. melanogaster | 19.03 | 0.44 | 19.47 |
| A. thaliana | 35.03 | 1.20 | 36.23 |
| C. elegans | 8.08 | 0.15 | 8.23 |
| CHM1 | 3,109.20 | N/A | N/A |
| MAP005 | 0.02 | 0.00 | 0.02 |
| MAP006-1 | 0.04 | 0.01 | 0.05 |
| MAP006-2 | 0.02 | 0.00 | 0.02 |
| MAP006-PCR-1 | 0.02 | 0.00 | 0.02 |
| MAP006-PCR-2 | 0.06 | 0.00 | 0.06 |
| <i>B. anthracis</i> | 0.1 | 0.01 | 0.11 |
| Y. pestis | 0.05 | 0.05 | 0.10 |

Table S6. Assembly Time SPAdes (CPU hours)

| Genome | Total Assembly Time |
|---------------------|---------------------|
| E. coli | 4.08 |
| MAP005 | 3.61 |
| MAP006-1 | 3.65 |
| MAP006-2 | 3.56 |
| MAP006-PCR-1 | 3.57 |
| MAP006-PCR-2 | 4.00 |
| <i>B. anthracis</i> | 8.47 |
| <i>Y. pestis</i> | 17.08 |

Supplementary Note 8: PacBio Assembly Statistics with Quiver

Quiver (Chin et al. 2013) relies on the signal-level quality values from PacBio and a model trained on a specific chemistry to correct errors. The tables show quality for the initial assembly and after multiple rounds of Quiver. The figures show an alignment of the final assembly for each assembler to the reference genome. The final assembly is after one round of Quiver (for Canu and Falcon) and four rounds (for miniasm).

Table S7 Quiver Statistics For Canu.

Each assembly was aligned to the reference genome using MUMmer and errors identified using dnadiff (Supplementary Note 5). Summary assembly consensus accuracy statistics are reported for the initial assembly as well as multiple rounds of polishing. QV was computed from total SNPs versus the reference and the total aligned bases in the assembly

| Genome | Assembly | | | Quiver1 | | | Quiver2 | | |
|-----------------|---------------|-------|-----------|---------------|-------|----------|---------------|-------|----------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| E. coli | 100.00% | 46.81 | 4.26 | 100% | 58.90 | 7.99 | 100% | 58.90 | 7.54 |
| D. melanogaster | 97.22% | 31.72 | 573.65 | 97.47% | 36.88 | 822.87 | 97.48% | 37.21 | 824.25 |
| A. thaliana | 82.90% | 20.31 | 577.49 | 82.94% | 20.32 | 347.82 | 82.93% | 20.32 | 376.72 |
| C. elegans | 99.61% | 33.46 | 170.14 | 99.70% | 35.93 | 239.93 | 99.7% | 35.76 | 229.93 |
| CHM1 | 86.48% | 25.14 | 18,019.70 | 86.84% | 27.17 | 4,730.01 | 86.86% | 27.19 | 4,796.51 |

Table S8 Quiver Statistics For Falcon.
Same columns as table S7.

| Genome | Assembly | | | Quiver1 | | | Quiver2 | | |
|-----------------|---------------|-------|-----------|---------------|-------|----------|---------------|-------|----------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| E. coli | 100% | 30.22 | 10.08 | 100% | 58.22 | 7.67 | 100% | 58.89 | 7.39 |
| D. melanogaster | 95.52% | 27.14 | 930.76 | 96.12% | 37.27 | 1375.16 | 96.15% | 37.34 | 1359.23 |
| A. thaliana | 82.06% | 19.87 | 836.61 | 82.72% | 20.32 | 295.64 | 82.73% | 20.32 | 335.51 |
| C. elegans | 98.70% | 26.59 | 202.74 | 98.82% | 35.76 | 194.66 | 98.82% | 35.69 | 175.32 |
| CHM1 | 86.03% | 23.25 | 64,087.80 | 86.58% | 27.18 | 4,701.18 | 86.62% | 27.19 | 4,702.71 |

Table S9 Quiver Statistics For Miniasm
Same columns as table S7.

| Genome | Assembly | | | Quiver1 | | | Quiver2 | | | Quiver3 | | | Quiver4 | | |
|-----------------|---------------|------|-------|---------------|-------|--------|---------------|-------|--------|---------------|-------|--------|---------------|-------|--------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| E. coli | 95.20% | 9.36 | 0.18 | 99.98% | 37.67 | 8.19 | 99.99% | 57.65 | 7.78 | 99.99% | 57.12 | 8.38 | 99.99% | 57.12 | 7.40 |
| D. melanogaster | 0.00% | 0 | 19.47 | 96.42% | 35.05 | 282.83 | 96.48% | 36.95 | 383.80 | 96.50% | 37.20 | 395.93 | 96.51% | 37.27 | 402.30 |
| A. thaliana | 0.00% | 0 | 36.23 | 82.79% | 20.26 | 197.45 | 82.88% | 20.32 | 255.95 | 82.87% | 20.32 | 250.71 | 82.88% | 20.32 | 236.09 |
| C. elegans | 0.00% | 0 | 8.23 | 99.24% | 32.14 | 111.35 | 99.37% | 34.61 | 127.54 | 99.43% | 35.14 | 133.16 | 99.44% | 35.32 | 145.88 |

Table S10 Quiver Assembly Statistics For SPAdes
Same columns as table S7.

| Genome | Assembly | | | Quiver1 | | | Quiver2 | | |
|---------|---------------|-------|------|---------------|-------|------|---------------|-------|------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| E. coli | 100% | 44.55 | 4.09 | 99.97% | 47.97 | 9.82 | 100% | 45.05 | 7.44 |

Figure S2 Canu, Falcon, Miniasm, and SPAdes *E. coli* K12 polished assemblies.
See caption for Figure S1.




Figure S3 Canu, Falcon, and Miniasm *D. melanogaster* assemblies.
See caption for Figure S1.




Figure S4 Canu, Falcon, and Miniasm *A. thaliana* assemblies.
See caption for Figure S1.




Figure S5 Canu, Falcon, and Miniasm *C. elegans* assemblies.
See caption for Figure S1.





Figure S6 Canu and Falcon assemblies of *H. sapiens* CHM1.
See caption for Figure S1.



Supplementary Note 9: Oxford Nanopore Assemblies with Nanopolish

Nanopolish (Loman et al. 2015) relies on the events underlying the basecall for a Nanopore sequence and a model of the sequencing error to correct errors. The tables show quality for the initial assembly and after multiple rounds of Nanopolish. The figures show an alignment of the final assembly for each assembler to the reference genome. The final assembly is after one round of Nanopolish (for Canu and Falcon) and three rounds (for miniasm).

Table S11 Nanopolish Statistics for Canu
Same columns as table S7.

| Genome | Assembly | | | Nanopolish 1 | | | Nanopolish 2 | | |
|--------------|---------------|-------|-------|---------------|-------|--------|---------------|-------|--------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| MAP005 | 99.97% | 14.59 | 9.38 | 99.98% | 22.44 | 367.49 | 99.98% | 22.91 | 202.82 |
| MAP006-1 | 99.79% | 21.53 | 4.85 | 99.80% | 27.11 | 162.19 | 99.81% | 27.19 | 154.01 |
| MAP006-2 | 99.85% | 20.91 | 2.78 | 99.91% | 26.61 | 165.91 | 99.92% | 26.86 | 151.57 |
| MAP006-PCR-1 | 99.95% | 22.28 | 3.04 | 99.95% | 27.92 | 161.04 | 99.95% | 27.93 | 142.63 |
| MAP006-PCR-2 | 99.99% | 22.42 | 5.25 | 99.99% | 28.27 | 200.84 | 100.00% | 28.41 | 171.34 |
| B. anthracis | 99.77% | 15.69 | 63.13 | 99.77% | 20.65 | 831.27 | 99.77% | 21.04 | 468.11 |
| Y. pestis | 99.97% | 21.35 | 12.55 | 99.97% | 26.13 | 241.70 | 99.97% | 26.20 | 192.00 |

Table S12 Nanopolish Statistics for Falcon
Same columns as table S7.

| Genome | Assembly | | | Nanopolish 1 | | | Nanopolish 2 | | |
|--------------|---------------|-------|------|---------------|-------|--------|---------------|-------|--------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| MAP005 | 22.89% | 13.54 | 2.98 | 23.03% | 22.29 | 103.22 | 23.00% | 23.07 | 38.55 |
| MAP006-1 | 99.86% | 19.38 | 4.93 | 99.86% | 26.64 | 202.52 | 99.86% | 27.16 | 174.34 |
| MAP006-2 | 99.94% | 18.68 | 1.56 | 99.94% | 26.19 | 194.63 | 99.94% | 26.89 | 153.68 |
| MAP006-PCR-1 | 99.80% | 20.16 | 1.74 | 99.80% | 27.46 | 166.63 | 99.80% | 27.88 | 143.49 |
| MAP006-PCR-2 | 100.00% | 20.31 | 7.42 | 100.00% | 27.83 | 205.47 | 100.00% | 28.35 | 179.51 |

| | | | | | | | | | |
|--------------|--------|-------|-------|--------|-------|--------|--------|-------|--------|
| B. anthracis | 86.27% | 14.73 | 12.75 | 86.29% | 20.81 | 783.18 | 86.30% | 21.41 | 368.58 |
| Y. pestis | 99.97% | 19.16 | 4.57 | 99.97% | 25.51 | 290.44 | 99.96% | 25.95 | 188.91 |

Table S13 Nanopolish Statistics for Miniasm
Same columns as table S7.

| Genome | Assembly | | | Nanopolish 1 | | | Nanopolish 2 | | | Nanopolish 3 | | |
|--------------|---------------|------|------|---------------|-------|---------|---------------|-------|--------|---------------|-------|--------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| MAP005 | 79.81% | 7.81 | 0.02 | 99.62% | 15.58 | 1801.51 | 99.93% | 20.31 | 320.75 | 99.96% | 21.94 | 221.74 |
| MAP006-1 | 91.76% | 9.24 | 0.05 | 99.71% | 16.94 | 1363.27 | 99.94% | 22.75 | 257.37 | 99.97% | 25.56 | 180.33 |
| MAP006-2 | 92.89% | 9.18 | 0.02 | 99.48% | 16.78 | 1130.5 | 99.68% | 22.35 | 233.74 | 99.70% | 25.15 | 118.69 |
| MAP006-PCR-1 | 93.83% | 9.52 | 0.02 | 99.79% | 17.62 | 1014.35 | 99.93% | 23.58 | 208.51 | 99.96% | 26.28 | 115.40 |
| MAP006-PCR-2 | 93.77% | 9.31 | 0.06 | 99.89% | 18.10 | 1457.47 | 99.97% | 24.74 | 252.86 | 99.98% | 27.18 | 159.24 |
| B. anthracis | 66.46% | 7.75 | 0.11 | 97.07% | 14.36 | 4013.98 | 97.19% | 18.67 | 676.67 | 97.21% | 20.23 | 404.14 |
| Y. pestis | 89.94% | 9.20 | 0.10 | 99.65% | 16.56 | 1502.52 | 99.90% | 21.99 | 295.81 | 99.91% | 24.61 | 201.73 |

Figure S7 Canu, Falcon, and Miniasm assemblies of *E. coli* K12 MAP005.
See caption for Figure S1.




Figure S8 Canu, Falcon, and Miniasm assemblies of *E. coli* K12 MAP006-1.
See caption for Figure S1.




Figure S9 Canu, Falcon, and Miniasm assemblies of *E. coli* K12 MAP006-2.
See caption for Figure S1.




Figure S10 Canu, Falcon, and Miniasm assemblies of *E. coli* K12 MAP006-PCR-1.
See caption for Figure S1.




Figure S11 Canu, Falcon, and Miniasm assemblies of *E. coli* K12 MAP006-PCR-2.
See caption for Figure S1.




Figure S12 Canu, Falcon, and Miniasm assemblies of *B. anthracis*.
See caption for Figure S1.





Figure S13 Canu, Falcon, and Miniasm assemblies of *Y. pestis*.
See caption for Figure S1.



Supplementary Note 10: Oxford Nanopore assemblies with Pilon

Pilon (Walker et al. 2014) uses Illumina sequences aligned to an assembly to correct errors. The tables show quality for the initial assembly and after multiple rounds of Pilon. The figures show an alignment of the final assembly for each assembler to the reference genome. The final assembly is after three round of Pilon (for Canu, Falcon, and Miniasm) and unpolished (for SPAdes).

Table S14 Pilon Statistics for Miniasm after 300 rounds of pilon
Same columns as table S7.

| Genome | Assembly | | | Pilon 300 | | |
|--------------|---------------|------|------|---------------|-------|------------|
| | % ref covered | QV | Time | % ref covered | QV | Total Time |
| MAP005 | 79.81% | 7.81 | 0.02 | 91.80% | 16.79 | 182.57 |
| MAP006-1 | 91.76% | 9.24 | 0.05 | 97.07% | 24.45 | 114.18 |
| MAP006-2 | 92.89% | 9.18 | 0.02 | 98.03% | 24.55 | 96.82 |
| MAP006-PCR-1 | 93.83% | 9.52 | 0.02 | 98.46% | 25.19 | 83.28 |
| MAP006-PCR-2 | 93.77% | 9.31 | 0.06 | 98.62% | 25.18 | 94.96 |
| B. anthracis | 65.34% | 7.75 | 0.11 | 79.88% | 11.51 | 842.89 |
| Y. pestis | 89.94% | 9.20 | 0.10 | 93.92% | 19.49 | 543.77 |

Figure S14 Minimap remaining errors in *E. coli* K12 MAP006-1.

Using dnadiff after 300 rounds of pilon on miniasm we tabulated GAP entries in the rdiff file. These indicate unaligned regions between the reference and the assembly as well as the size of the discrepancy. The average discrepancy size was 569.17 bp in six discrepant regions. The figure highlights several gap regions which show a) an expansion in the assembly with respect to the reference b) a collapse in the reference with respect to the assembly.




Table S15 Pilon Statistics for Canu
Same columns as table S7.

| Genome | Assembly | | | Pilon 1 | | | Pilon 2 | | | Pilon 3 | | |
|--------------|---------------|-------|-------|---------------|-------|------|---------------|-------|------|---------------|-------|------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| MAP005 | 99.97% | 14.59 | 9.38 | 99.97% | 29.99 | 0.96 | 99.99% | 38.62 | 0.32 | 99.99% | 38.95 | 0.32 |
| MAP006-1 | 99.79% | 21.53 | 4.85 | 99.79% | 40.98 | 0.50 | 99.82% | 48.40 | 0.28 | 99.82% | 53.44 | 0.26 |
| MAP006-2 | 99.85% | 20.91 | 2.78 | 99.89% | 37.37 | 0.56 | 99.94% | 42.90 | 0.29 | 99.94% | 48.90 | 0.29 |
| MAP006-PCR-1 | 99.95% | 22.28 | 3.04 | 99.95% | 43.72 | 0.56 | 99.95% | 50.54 | 0.28 | 99.95% | 51.75 | 0.27 |
| MAP006-PCR-2 | 99.99% | 22.42 | 5.25 | 100% | 43.79 | 0.38 | 100% | 50.44 | 0.27 | 100% | 50.99 | 0.26 |
| B. anthracis | 99.77% | 15.69 | 63.13 | 99.77% | 25.56 | 0.83 | 99.77% | 27.95 | 0.53 | 99.77% | 28.17 | 0.52 |
| Y. pestis | 99.97% | 21.35 | 12.55 | 99.87% | 28.78 | 2.26 | 99.83% | 29.62 | 1.52 | 99.83% | 29.77 | 1.59 |

Table S16 Pilon Statistics for Falcon
Same columns as table S7.

| Genome | Assembly | | | Pilon 1 | | | Pilon 2 | | | Pilon 3 | | |
|----------|---------------|-------|------|---------------|-------|------|---------------|-------|------|---------------|-------|------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| MAP005 | 22.89% | 13.54 | 2.98 | 23.06% | 27.72 | 0.65 | 23.07% | 33.31 | 0.41 | 23.07% | 33.47 | 0.32 |
| MAP006-1 | 99.86% | 19.38 | 4.93 | 99.86% | 33.77 | 1.30 | 99.86% | 43.04 | 0.57 | 99.86% | 44.43 | 0.50 |
| MAP006-2 | 99.94% | 18.68 | 1.56 | 99.94% | 32.67 | 1.27 | 99.94% | 40.24 | 0.58 | 99.94% | 41.77 | 0.52 |

| | | | | | | | | | | | | |
|--------------|--------|-------|-------|--------|-------|------|--------|-------|------|--------|-------|------|
| MAP006-PCR-1 | 99.80% | 20.16 | 1.74 | 99.80% | 35.44 | 0.76 | 99.80% | 43.31 | 0.54 | 99.80% | 45.13 | 0.51 |
| MAP006-PCR-2 | 100% | 20.31 | 7.42 | 100% | 35.80 | 0.75 | 100% | 42.17 | 0.54 | 100% | 44.33 | 0.51 |
| B. anthracis | 86.27% | 14.73 | 12.75 | 86.31% | 24.95 | 0.45 | 86.31% | 29.08 | 0.19 | 86.31% | 29.54 | 1.47 |
| Y. pestis | 99.97% | 19.16 | 4.57 | 99.71% | 26.94 | 1.54 | 99.65% | 28.73 | 2.74 | 99.65% | 28.91 | 1.78 |

Table S17 Pilon Statistics for Miniasm
Same columns as table S7.

| Genome | Assembly | | | Pilon 1 | | | Pilon 2 | | | Pilon 3 | | |
|--------------|---------------|------|------|---------------|-------|------|---------------|-------|------|---------------|-------|------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| MAP005 | 79.81% | 7.81 | 0.02 | 85.62% | 10.87 | 1.43 | 88.46% | 13.97 | 1.01 | 90.63% | 15.83 | 0.69 |
| MAP006-1 | 91.76% | 9.24 | 0.05 | 93.09% | 16.42 | 2.01 | 96.32% | 22.23 | 0.74 | 96.97% | 24.11 | 0.34 |
| MAP006-2 | 92.89% | 9.18 | 0.02 | 94.12% | 16.05 | 1.87 | 97.47% | 22.75 | 0.50 | 97.98% | 24.36 | 0.34 |
| MAP006-PCR-1 | 93.83% | 9.52 | 0.02 | 95.17% | 17.02 | 1.36 | 97.93% | 23.20 | 0.49 | 98.41% | 24.86 | 0.28 |
| MAP006-PCR-2 | 93.77% | 9.31 | 0.06 | 94.85% | 16.45 | 1.83 | 98.15% | 23.20 | 0.46 | 98.57% | 24.86 | 0.34 |
| B. anthracis | 66.46% | 7.75 | 0.11 | 77.41% | 9.38 | 0.29 | 78.54% | 10.45 | 2.19 | 79.36% | 11.12 | 2.31 |
| Y. pestis | 89.94% | 9.20 | 0.10 | 91.89% | 14.09 | 2.18 | 93.16% | 18.05 | 3.99 | 93.76% | 19.16 | 2.41 |

Table S18 Pilon Statistics for Spades
Same columns as table S7.

| Genome | Assembly | | | Pilon 1 | | | Pilon 2 | | | Pilon 3 | | |
|--------------|---------------|-------|-------|---------------|-------|------|---------------|-------|------|---------------|-------|------|
| | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time | % ref covered | QV | Time |
| MAP005 | 100% | 44.55 | 3.61 | 100% | 48.75 | 0.33 | 100% | 49.27 | 0.37 | 100% | 49.59 | 0.29 |
| MAP006-1 | 100% | 44.55 | 3.65 | 100% | 48.74 | 0.39 | 100% | 49.26 | 0.46 | 100% | 49.59 | 0.30 |
| MAP006-2 | 100% | 44.55 | 3.56 | 100% | 48.74 | 0.36 | 100% | 49.26 | 0.28 | 100% | 49.59 | 0.28 |
| MAP006-PCR-1 | 100% | 44.60 | 3.57 | 100% | 48.54 | 0.28 | 100% | 49.03 | 0.28 | 100% | 49.34 | 0.27 |
| MAP006-PCR-2 | 100% | 44.60 | 4.00 | 100% | 48.74 | 0.31 | 100% | 49.26 | 0.29 | 100% | 49.59 | 0.27 |
| B. anthracis | 100% | 42.83 | 8.47 | 100% | 44.82 | 1.06 | 100% | 45.20 | 0.99 | 100% | 45.28 | 0.90 |
| Y. pestis | 95.99% | 33.56 | 17.08 | 96.00% | 33.51 | 1.29 | 96.00% | 33.53 | 1.29 | 96.00% | 33.53 | 1.24 |

Figure S15 Canu, Falcon, Miniasm, and SPAdes assemblies of *E. coli* K12 MAP005.
See caption for Figure S1.




Figure S16 Canu, Falcon, Miniasm, and SPAdes assemblies of *E. coli* K12 MAP006-1.
See caption for Figure S1.




Figure S17 Canu, Falcon, Miniasm, and SPAdes assemblies of *E. coli* K12 MAP006-2.
See caption for Figure S1.




Figure S18 Canu, Falcon, Miniasm, and SPAdes assemblies of *E. coli* K12 MAP006-PCR-1.
See caption for Figure S1.




Figure S19 Canu, Falcon, Miniasm, and SPAdes assemblies of *E. coli* K12 MAP006-PCR-2.
See caption for Figure S1.




Figure S20 Canu, Falcon, Miniasm, and SPAdes assemblies of *B. anthracis*.
See caption for Figure S1.





Figure S21 Canu, Falcon, Miniasm, and SPAdes assemblies of *Y. pestis*.
See caption for Figure S1.



Supplementary Note 11: 1D Assembly


We ran 10 rounds of correction followed by assembly. For 1D:

```
NAME=1d.fasta
COUNT=0
for i in `seq 1 10`; do
  canu -correct -p asm -d round$i \
    corOutCoverage=500 corMinCoverage=0 corMhapSensitivity=high \
    genomeSize=4.8m -nanopore-raw $NAME
  NAME="round$i/asm.correctedReads.fasta.gz"
  COUNT=`expr $COUNT + 1`
done
canu -p asm -d asm genomeSize=4.8m -nanopore-corrected $NAME \
  errorRate=0.1 utgGraphDeviation=50 batOptions="-ca 500 -cp 50"
```

The initial Canu assembly had 0 errors, covering 89.38% of the reference at 85.52% identity. We ran Nanopolish 1d-workflows with modifications available from

<https://github.com/skoren/nanopolish>. Compiler optimization had to be disabled for Nanopolish to run to completion. We ran a total of 10 rounds of polishing, resulting in 96.67%, 97.78%, 98.04%, 98.18%, 98.25%, 98.31%, 98.34%, 98.37%, 98.38%, and 98.40% identity. Post three rounds of polishing the assembly has 1 errors, covering 93.32% of the reference at 98.04% identity. Post three Pilon polishing rounds it has 10 errors (all relocations due to collapsed sequences which could not be mapped pre-polishing), covering 93.84% of the reference at 99.70% identity. The initial Minimap assembly had 3 errors, covering only 9.19% of the reference at 75.76% identity.

Figure S22: Canu 1D, Canu+three rounds of Nanopolish, Canu 1D+three rounds of Pilon, and Miniasm assemblies for *E. coli* K12 MAP006-1 1D.
See caption for Figure S1.



Supplementary Note 12: *S. cerevisiae* assembly

As above, we ran 10 rounds of correction followed by assembly:

For *S. cerevisiae*:

```
COUNT=0
NAME=input.fasta
for i in `seq 1 10`; do
    canu -correct -p asm -d round$i \
        corOutCoverage=500 corMinCoverage=0 corMhapSensitivity=high \
        genomeSize=12.1m -nanopore-raw $NAME
    NAME="round$i/asm.correctedReads.fasta.gz"
    COUNT=`expr $COUNT + 1`
done
canu -p asm -d asm genomeSize=12.1m -nanopore-corrected $NAME utgGraphDeviation=50
batOptions="-ca 500 -cp 50"
done
```

The initial Canu assembly had 2 errors, covering 94.93% of the reference at 94.38% identity. Post polishing it has 14 errors, covering 96.86% of the reference at 99.83% identity. The reference (S288c) is not identical to the sequenced strain (W303) and the number of errors is similar to that from a high-coverage PacBio only assembly (29 ctg, 99.12% of ref, 99.88% idy, 13 errors) and could be true variation between the strains.

Figure S23: Canu 1D, and Canu 1D+three rounds of Pilon, and Canu PacBio 115X assemblies.
See caption for Figure S1.




Figure S24: Parallel overlap store construction

Canu constructs the overlap store by a parallel bucket sort. Each overlap file is first bucketized based on their read range, with each bucket sized to be approximately the same size. Once bucketized, parallel processes collect all overlaps from a bucket and sort them independently. Finally, a merged index is written to disk.




Figure S25: Suspicious read detection.

A read A has an anomaly at position X which prevents the A-C overlap from being discovered. Read A's best overlap is to read B, but read B's best overlap is to read C. If read A ends in a repeat, the best overlap could be to a diverged repeat, leading to a misassembly.

A -----X-----
 C -----
 B -----

References

- Berlin K, Koren S, Chin C-S, Drake JP, Landolin JM, Phillippy AM. 2015. Assembling large genomes with single-molecule sequencing and locality-sensitive hashing. *Nat Biotech* **33**: 623-630.
- Chaisson MJ, Tesler G. 2012. Mapping single molecule sequencing reads using basic local alignment with successive refinement (BLASR): application and theory. *BMC Bioinformatics* **13**: 238.
- Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE et al. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nature methods* **10**: 563-569.
- Goodwin S, Gurtowski J, Ethe-Sayers S, Deshpande P, Schatz MC, McCombie WR. 2015. Oxford Nanopore sequencing, hybrid error correction, and de novo assembly of a eukaryotic genome. *Genome Res* **25**: 1750-1756.
- Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile and open software for comparing large genomes. *Genome biology* **5**: R12-R12.
- Lee H, Gurtowski J, Yoo S, Marcus S, McCombie WR, Schatz M. 2014. Error correction and assembly complexity of single molecule sequencing reads. *bioRxiv* doi:10.1101/006395.
- Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv preprint arXiv:13033997*.
- Loman NJ, Quick J, Simpson JT. 2015. A complete bacterial genome assembled de novo using only nanopore sequencing data. *Nature methods* **12**: 733-735.
- Phillippy AM, Schatz MC, Pop M. 2008. Genome assembly forensics: finding the elusive mis-assembly. *Genome biology* **9**: R55-R55.
- Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK et al. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* **9**: e112963.