

Systemic risk and contagion effects in Australian financial institutions and sectors

Piet de Jong, Geoff Loudon and Weihao Choo

*Department of Applied Finance and Actuarial Studies
Macquarie University, Sydney, NSW 2109.*

piet.dejong@mq.edu.au

1. Literature review

The starting point for the proposed research is the recent literature and the CIFR targeted areas and APRA aims and functions. This recent literature includes the following Adrian and Brunnermeier (2011), Acharya et al. (2012), Acharya et al. (2012) and Brownlees and Engle (2010). The proposed research aims to extend and apply these techniques particularly in relation to the entities regulated by APRA. Thus our broad aim is to develop, implement and bring to bear recent developments in stress testing on the aims of APRA and the CIFR targeted research areas detailed above.

2. Improved measures of contagion and systematic risk

CoVaR_q as proposed in Adrian and Brunnermeier (2011) is a basis for proposed measures contagion, exposure and systemic risk. It suffers from a number of drawbacks:

- Couched in terms of VaR_q containing the scale of the original measurements. It is worthwhile to have measures and techniques robust to scale.
- Conditioning on VaR_{0.5} is undesirable and relatively intractable. In our proposal we reference stress with respect to the unconditional VaR_q. This permits a more transparent analysis and estimation.
- Our proposed approach separates out scale effects and interdependence effects and aims to relates these separately to external variables including shocks and drivers of systemic risk. Thus VaR_q movements due to scale are disentangled from movements due to codependence with separate driver responses.

3. Significance of the project and policy implications

Understanding the impact of external shocks and their propagation through the financial system is vital for managing and remediating systemic risk. Effective regulation is dependent upon the development of a robust and reliable set of appropriate risk measures. We propose new measures of systemic risk that relate marginal and joint distributions separately to external drivers. This allows for more cogent and coherent stress testing as it includes the estimation of contagion effects, exposure effects and systemic risk across related entities and different financial sectors. Improved stress testing, estimation of risk effects and transmission of shocks through the financial system will make for more cogent prudential policy, prudential margin setting and better identify sources of risk to the financial system.

4. Percentile sensitivity and contagion

4.1. Theorem

Suppose x is a random vector with marginal distributions¹

$$F(x) \equiv \{F(x_1), \dots, F(x_p)\} \equiv (u_1, \dots, u_p) \equiv u .$$

Further suppose $0 \leq q \leq 1$ is given and $Q(x)$ is the vector of q -quantiles

$$F\{Q(x)\} = q1 = Q(u) ,$$

where 1 is a vector of p ones. Define the stress vector with respect to x_j as²

$$\frac{dQ(x)}{dx_j} \equiv Q(x|u_j > q) - Q(x) , \quad (1)$$

where $Q(x|u_j > q)$ is the vector of q -quantiles of x given $u_j > q$. Then if $Q(x)$ is linear in q then

$$\frac{dQ(x_i)}{dQ(x_j)} \equiv \frac{dQ(x_i)/dx_j}{dQ(x_j)/dx_j} = \frac{f_j}{f_i} \frac{q_{ij}}{q(1-q)} , \quad q_{ij} = C_{ij}(q + q_{ij}, q) - q^2 , \quad (2)$$

where f_i and f_j are the densities of x_i and x_j evaluated at $Q(x_i)$ and $Q(x_j)$ and C_{ij} is the copula of (u_i, u_j) . Further if

$$s_{ij} \equiv \frac{q_{ij}}{q(1-q)} ,$$

then $-1 \leq s_{ij} \leq 1$ with $s_{ij} = \pm 1$ if x_i and x_j are comonotonic and counter monotonic, respectively. If x_i and x_j are independent then $s_{ij} = 0$. If u_i and u_j are exchangeable then $s_{ji} = s_{ij}$.

¹To economise on notation, write $F_j(x) \equiv F_j(x_j) \equiv F(x_j)$ and similar for other vector functions.

²In Adrian and Brunnermeier (2011) $\Delta CoVar_q \equiv q_{y|x=q_x} - q_y$. Variable y is generally the “financial system” and hence considered is the change in the VaR_q of the system when institution x stressed, with stress interpreted as $x = q_x$. On page 10 of their paper they incorrectly state “... *CoVaR* conditions on the event that [an] institution is at its VaR level, which occurs with probability q .”

4.2. Proof

By definition

$$\frac{dQ(u_i)}{du_j} \equiv Q(u_i|u_j > q) - q \equiv q_{ij} ,$$

where q_{ij} is such that

$$q = \frac{P(u_i \leq q + q_{ij}, u_j > q)}{1 - q} = \frac{q + q_{ij} - C_{ij}(q + q_{ij}, q)}{1 - q} , \quad (3)$$

Rearranging yields the second equation in (2). Now

$$\frac{dQ(x_i)}{dx_j} \equiv Q_{q+q_{ij}}(x_i) - Q(x_i) \approx Q'(x_i)q_{ij} = \frac{q_{ij}}{f_i} , \quad (4)$$

where $'$ denotes differentiation with respect to q and the subscript on Q indicates the revised q for the quantile calculation. The approximation follows from a first order Taylor expansion and is exact if the quantile is linear in q . Similarly

$$\frac{dQ(x_j)}{dx_j} \equiv Q_{q+q(1-q)}(x_j) - Q(x_j) \approx \frac{q(1-q)}{f_j} . \quad (5)$$

Dividing (4) by (5) yields the first equation in (2) and completes the proof.

4.3. Discussion

The critical result is that sensitivities factor into contributions from the ratios of the marginal densities and quantities calculated from the pairwise copulas. The via the implicit equation for q_{ij} in (2), solved using a root finding algorithm. The quantities q_{ij} are implicitly defined from the pairwise copulas. In summary the “sensitivity” matrix

$$S \equiv \frac{dQ(x)}{dQ'(x)} ,$$

where $'$ denotes transposition has ones on the diagonal quantities between ± 1 off the diagonal. The matrix S is called the sensitivity matrix and displays the sensitivity of the q -quantile of each component of x to stress in the same or other components.

Since f_i and f_j are the densities evaluated at $Q(x_i)$ and $Q(x_j)$ it follows that the ratio f_i/f_j can be replaced by the ratio of the hazards. This is useful in cases were it is practical to model the hazard rather than the density.

4.4. Implementation

Figure 1 displays the empirical copulas calculated from four weekly closing stock prices labelled anz, cba, mcq and wbc for $n = 761$ weeks from 2000 April 12 through to 2014 October 29. The empirical copulas are calculated by converting each observation to an empirical percentile and plotting the same against each of the other series percentiles.




Figure 1: Pairwise copulas of bank stocks cba, anz, mqg, wbc, and the overall bank index. Red lines plot sensitivity s_{ij} as a function of q .

To estimate q_{ij} at a particular q , the second equation in (2) is iterated³ starting from $q_{ij} = 0$ where copulas are estimated as

$$\hat{C}_{ij}(u_i, u_j) \equiv \hat{E}\{(p_{ik} \leq u_i)(p_{jk} \leq u_j)\} .$$

Here \hat{E} computes the empirical mean over the cases $k = 1, \dots, n$ and p_{ik} and p_{jk} are the empirical percentiles of case k of x_i and x_j , respectively.

4.5. Generalisations

Similar results apply when $Q(x)$ is replaced by other risk measures such as $R(x) \equiv E\{xr(u)\}$ where r is a given function which acts componentwise and $xr(u)$ denotes componentwise multiplication. For example if $r(u) = mu^{m-1}$ then $R(x) = E\{\max(x^1, \dots, x^m)\}$ where x^1, \dots, x^m are m independent copies of x and the risk measure is the expected worst outcome in m independent trials.

With $R(x)$, the analogue of (1) is

$$\frac{dR(x)}{dx_j} \equiv R(x|u_j > r_j) - R(x_j) , \quad r_j \equiv P\{x_j \leq R(x_j)\} . \quad (6)$$

This differs from (1) in that a different risk measure is used and r_j replaces q . If the components of $R(x)$ are linear in the r_i then

$$\frac{dR(x_i)}{dR(x_j)} = \frac{f_j}{f_i} \frac{q_{ij}}{q(1-q)} , \quad q_{ij} = C_{ij}(r_i + q_{ij}, r_j) - r_i r_j , \quad (7)$$

where f_i and f_j are the x_i and x_j densities at the r_i and r_j quantiles, respectively.

5. Partially Gaussian copulas

Suppose $(u, v) = F_*(x, y)$ where x and y are scalar random variables. Suppose $u_1 < \dots < u_n$ are ordered observed values of u and $q_t = \Phi^-(u_t)$, $t = 1, \dots, n$. The correspondingly ordered values of v and $z = \Phi^-(v)$ are denoted v_t and z_t . Note $\mathcal{E}(z_t) = \mathcal{E}(q_t) = 0$ and $\mathcal{V}(z_t) = \mathcal{V}(q_t) = 1$ where \mathcal{E} and \mathcal{V} evaluate the mean and variance assuming t is uniform on the integers $1, \dots, n$.

Smoothing methods are here proposed to smooth and simulate y values using a copula on (u, v) constructed via a cubic spline model linking z and u . Of particular concern are simulated z when u is near 0 or 1 corresponding to x extremes. Given a simulated u and in turn z , $(x, y) = F_*^{-1}\{u, \Phi(z)\}$, a draw from the joint. Constraining u draws to say the upper tail, yields corresponding upper tail values for y . The challenge is to produce cogent z 's or y 's that properly reflect dependence in different parts of the distribution unencumbered by inappropriate constraints, and in particular tail constraints, inherent in say the Gaussian copula setup. Practical methods admit fitting on the basis of observed

³The second equation in (2) is a contraction mapping and hence has a fixed point.

data, and testing procedures for departures or otherwise from Gaussianity. The method outlined below achieves this.

The partially Gaussian (PG) copula for (u, v) is stated in terms of $q \equiv \Phi^-(u)$ and $z \equiv \Phi^-(v)$ on an arbitrarily fine u grid $0 < u_1 < \dots < u_n < 1$:

$$z_t = \alpha_t + \beta q_t + s\epsilon_t, \quad \alpha_{t+1} = \alpha_t + \delta_t + 1.268\eta_t, \quad \delta_{t+1} = \delta_t + \eta_t. \quad (8)$$

Here the ϵ_t and η_t are serially and contemporaneously uncorrelated zero mean disturbances with common variance σ^2 . The q_t term is the Gaussian component and α_t a nonparametric deviation. If α_t is identically zero then (z_t, q_t) is bivariate normal with correlation $\rho = \pm\sqrt{1 - (s\sigma)^2} = \beta$.

The second and third equations in (8) spell out the model for the nonparametric deviation α_t with α_0 and δ_0 assumed unknown parameters. If $\sigma \rightarrow 0$, equivalent to $s \rightarrow \infty$, the first equation in (8) is $z_t = \alpha_0 + \delta_0 t + \beta q_t + s\epsilon_t$. With this linear formulation for α_t , $\beta = \rho$ if $\alpha_0 = \delta_0 = 0$. Model (8) with $\beta = 0$ is the nonparametric model since z_t varies smoothly with t , unrelated to q_t . If $\beta = 0$ and $\sigma \rightarrow 0$ there is a probit relationship between z_t and u_t . Finally $s = 0$ implies $a(t) + \beta q_t = z_t$ for any β . The full range of possibilities is set out in Table 1 discussed in more detail below.

The $\alpha_t = 0$ and $\alpha_t = \alpha_0 + \delta_0 t$ specifications are generalised using function $a(t)$ minimising

$$\sum_{t=1}^n \{z_t - a(t) - \beta q_t\}^2 + s^2 \int_{-\infty}^{\infty} \{a''(t)\}^2 dt. \quad (9)$$

The interpolates of $a(t)$ equal $E(\alpha_t | z_1, \dots, z_n)$ computed with (8) where the ϵ_t and η_t are assumed normal and q_t fixed. From (9), $a(t)$ is piecewise twice differentiable passing closest, in a mean square error sense, to $z_t - \beta q_t$ with a penalty s^2 for roughness measured in terms of the second derivative $a''(t)$. Using observed percentile data (u_t, v_t) , $t = 1, \dots, n$, the aim is to estimate $a(t)$ and β and associated confidence bounds and assess goodness of fit. The function $a(t)$ is thought of as an intercept function.

6. Special cases of the PG model

Special cases of the general setup (8) or (9) are set out in Table 1. In this table, the notation $s \rightarrow \infty$ means s becomes large and σ small in such a way that $s\sigma$, the measurement noise in the first equation in (8), stays finite. In fact $s\sigma \leq 1$ since the variance of z_t is 1. Thus $s \rightarrow \infty$ implies $\sigma \rightarrow 0$ and states that the noise to signal ratio is becoming arbitrarily large. With the model the squared correlation between z_t and the signal $\alpha_t + \beta q_t$ is, since the variance of z_t is 1:

$$\rho^2 = 1 - (s\sigma)^2 = 1 - \sum_{t=1}^n \{z_t - a(t) - \beta q_t\}^2$$

The first row in Table 1 is the standard Gaussian copula setup. Here there is no α_t signal or component. The additional restriction $\beta = 0$ yields independence. Co (or counter) monotonicity results in the Gaussian framework if

$\rho = \pm 1$ as displayed in the third row. If $s = 0$ then there is no measurement noise and the combined signal $\alpha_t + \beta q_t$ must pass through the points z_t resulting in a perfect fit. In terms of (9), $s = 0$ implies there is no roughness penalty and hence the minimisation is solely focussed on minimising the first, squared error, term, achieved by setting $\alpha_t + \beta q_t = z_t$.

Table 1: Special cases of the PG copula; in all cases $\rho = \pm\sqrt{1 - (s\sigma)^2}$

case	$a(t)$	β	s	σ	α_0	δ_0	ρ_q	other
Gaussian	0		∞	0	0	0	ρ	$\rho = \beta$
independence	0	0	∞	0	0	0	0	$\rho = 0$
\pm comonotonic	0	± 1	0	0	0	0	± 1	$z_t = \pm q_t$
empirical	z_t		0				0	$\rho = 1$
linear probit	$\alpha_0(1 - 2u_t)$	0	∞	0		$\frac{-2\alpha_0}{n+1}$	0	indep. if $\alpha_0 = 0$
nonparametric	smooth	0					0	non Gaussian
PG	smooth						(10)	

The linear probit copula specification in Table 1 eliminates the Gaussian component q_t by setting $\beta = 0$, and giving maximum penalty for roughness. The resulting prediction is a straight line $\hat{z}_t = \alpha_0 + \delta_0 t$. However the mean of the z_t and hence \hat{z}_t is zero implying $0 = \alpha_0 + \delta_0(n+1)/2$. Solving for δ_0 gives the δ_0 constraint displayed in Table 1 and the resulting straight line signal $a(t)$.

The bottom row of Table 1 displays the general partial Gaussian specification absent any constraints on the parameters. Eliminating the Gaussian component from the signal by setting $\beta = 0$ leads to the nonparametric, non Gaussian specification in the second last row of Table 1.

7. Fitting PG copulas

The PG is readily fit to percentile data, and provides a practical platform for copula simulation. This is discussed, implemented, and used in subsequent sections. The parameters defining PG are s , β , σ and starting conditions α_0 and δ_0 . Given s , the other parameters have closed form, readily ‘computable’ expressions calculated with the Kalman filter.

The parameter s can be estimated by minimising least squares, closely related to maximum likelihood. In particular suppose the ϵ_t and η_t are normal.

8. Correlational properties of the PG copula

The column headed ρ_q in Table 1 is the partial correlation between z_t and α_t after removing the effect of the Gaussian component q_t . If $\beta = 0$, there is no Gaussian component and $\rho_q = \rho$.

This section considers alternative correlations. If $s \rightarrow \infty$ and $\alpha_0 = \delta_0 = 0$ then $a(t) = 0$ and (8) reduces to $z_t = \beta q_t + s\epsilon_t$ where $s\epsilon_t$ has mean zero and variance $0 \leq (s\sigma)^2 \leq 1$ and $\rho = \pm \sqrt{1 - (s\sigma)^2} = \beta$ is the correlation, called the overall correlation. The overall correlation is used when using a Gaussian copula. As many authors have pointed out Gaussian copula modelling can be misleading especially when considering correlations in the tails. Use of model (8) permits a more nuanced view of correlation without completely abandoning the Gaussian setup.

8.1. Partial correlation after allowing for semiparametric component

The partial correlation between z_t and q_t after allowing for $a(t)$ is

$$\rho_q \equiv \frac{\text{cov}(z_t - \alpha_t, q_t)}{\sqrt{\text{cov}(z_t - \alpha_t)\text{cov}(q_t)}} = \frac{\beta}{\sqrt{\beta^2 + (s\sigma)^2}} = \pm \left\{ 1 + \left(\frac{s\sigma}{\beta} \right)^2 \right\}^{-1/2}, \quad (10)$$

where the sign is that of β . Equation (10) is equivalent to $\mathcal{O}(\rho_q) = \beta/(s\sigma)$ where $\mathcal{O}(x) \equiv \pm \sqrt{x^2/(1-x^2)}$. If $s \rightarrow 0$ then $\rho_q \rightarrow 1$. If $a(t) \rightarrow 0$ then $\rho_q \rightarrow \rho$.

8.2. Local correlation

The local correlation between z_t and q_t is

$$\rho_t \equiv \text{cor}(\Delta z_t, \Delta q_t | t) = \frac{\beta \text{cov}(\Delta q_t)}{\sqrt{\text{cov}(\Delta z_t | t) \text{cov}(\Delta q_t)}} = \pm \left\{ \frac{\text{cov}(\Delta z_t | t)}{\beta^2 \text{cov}(\Delta q_t)} \right\}^{-1/2}$$

where “ $|t$ ” indicates conditioning on information up to and including time t . A calculation shows writing $c = 4 - \sqrt{3}$,

$$\text{cov}(\Delta z_t | t) = (2s^2 + c)\sigma^2 + \beta^2 \text{cov}(\Delta q_t), \quad \text{cov}(\Delta q_t) \approx (q'_t)^2 \text{cov}(u_t) = \frac{\sigma^2 \pi e^{q_t^2}}{6}.$$

Hence

$$\mathcal{O}(\rho_t) = \frac{\beta}{s_t \sigma}, \quad s_t = \sqrt{\frac{6(2s^2 + c)}{\pi}} e^{-q_t^2/2}.$$

Divisor s_t is small if $|q_t|$ is large indicating the model posits higher tail correlation than normal if $\beta \neq 0$. The ratio

$$\frac{\mathcal{O}(\rho_t)}{\mathcal{O}(\rho)} = \frac{s}{s_t} = \sqrt{\frac{\pi}{6(2 + c/s^2)}} e^{q_t^2/2} \rightarrow \frac{\sqrt{2\pi} e^{q_t^2/2}}{\sqrt{24}},$$

compares local to overall correlation. If $s \rightarrow \infty$ local correlation in the tails will far exceed overall correlation. This is obvious given $s \rightarrow \infty$ implies the model will exactly reproduce the data. If $s \rightarrow 0$

8.3. Alternative view of local correlation

In (8), a large $|\beta|$ suggests more extreme tail behaviour than in the Gaussian copula. To see this note

$$\Delta E(\alpha_t + \beta q_t) \approx \frac{-2\delta_0}{n+1} + \frac{\beta}{(n+1)\Phi'(q_t)} = \frac{\sqrt{2\pi}\beta e^{q_t^2/2} - 2\delta_0}{n+1}, \quad (11)$$

where Δ denotes change with respect to t . Change (11) is relatively large if q_t^2 is large, that is t is near 1 or n . Higher than normal tail dependence implies $|\beta|$ large. For a Gaussian copula, $\beta = \pm\sqrt{1-(s\sigma)^2}$ and comparing the ratio of estimates of the two sides to 1 is a test for Gaussianity.

8.4. Correlation between z_t and $a(t)$

The correlation between z_t and $a(t)$ after allowing for q_t is

$$\rho_a \equiv \frac{\text{cov}(z_t - \beta q_t, \alpha_t)}{\sqrt{\text{cov}(z_t - \beta q_t)\text{cov}(\alpha_t)}} = \left\{ 1 + \frac{(s\sigma)^2}{\text{cov}(\alpha_t)} \right\}^{1/2}. \quad (12)$$

This is BS – has to be done properly –. The overall correlation between z_t and $a(t)$ of

$$\frac{\text{cov}(z_t, \alpha_t)}{\sqrt{\text{cov}(z_t)\text{cov}(\alpha_t)}} = \frac{\text{cov}(z_t - \beta q_t, \alpha_t)}{\sqrt{1 - \beta^2 - (s\sigma)^2}} = \sqrt{1 - \beta^2 - (s\sigma)^2}$$

while the nonparametric correlation is the correlation between the nonparametric component α_t and z_t after removing the the effect of the Gaussian component:

The term $\alpha_t + \beta q_t$ is the signal and ϵ_t , noise. The signal is relatively smooth and contains both a nonparametric and Gaussian component q_t . Departures of z_t from the Gaussian component are the sum of a persistent component α_t and transient component, $s\epsilon_t$. Since ϵ_t and the driver of persistence η_t share σ , the smoothing parameter s measures relative importance of transience as opposed to persistence. The scale parameter σ relates to the degree of roughness since it is the variance of the changes in the slope of α_t .

Since $\mathcal{E}(q_t) = \mathcal{E}(z_t) = \mathcal{E}(\epsilon_t) = \mathcal{E}(\eta_t) = 0$ it follows

$$0 = \mathcal{E}(\alpha_t) = \alpha_0 + \delta_0 \mathcal{E}(t) = \alpha_0 + \delta_0 \mathcal{E}(t) = \alpha_0 + \delta_0 \frac{n+1}{2} \quad \Rightarrow \quad \delta_0 = \frac{-2\alpha_0}{n+1}.$$

Further $\mathcal{V}(q_t) = \mathcal{V}(z_t) = 1$ implies an analysis of variance type decomposition

$$1 = \mathcal{V}(\alpha_t) + \beta^2 + (s\sigma)^2. \quad (13)$$

Decomposition (13) splits the total variance of 1 into that due to the nonparametric component α_t , Gaussian component β^2 and residual noise, $(s\sigma)^2$. Alternatively (13) states the variance of the nonparametric component is $1 - \beta^2 - (s\sigma)^2$.

An example PG copula with $s\sigma = xx$, $\sigma = 0$, $\beta = xx$ and $\alpha_0 = xx$ is displayed in Figure ??.

9. Correlations with the PG copula

With (8) and t uniform on $1, \dots, n$

$$\mathcal{V} \begin{pmatrix} z_t \\ \alpha_t \\ q_t \end{pmatrix} = \begin{pmatrix} 1 & 1 - (s\sigma)^2 & \beta \\ 1 - (s\sigma)^2 & 1 - \beta^2 - (s\sigma)^2 & 0 \\ \beta & 0 & 1 \end{pmatrix}.$$

This covariance matrix serves to define a variety of ordinary and partial correlations.

The correlation between z_t and q_t is β . The partial correlation between z_t and the Gaussian component q_t , after removing the effect of the nonparametric component α_t , is called the Gaussian correlation:

$$\rho_q \equiv \frac{\text{cov}(q_t, z_t - \alpha_t)}{\sqrt{\text{cov}(q_t)\text{cov}(z_t - \alpha_t)}} = \frac{\beta}{\sqrt{\beta^2 + (s\sigma)^2}} = \pm \left\{ 1 + \left(\frac{s\sigma}{\beta} \right)^2 \right\}^{-1/2}, \quad (14)$$

where the sign is that of β . Hence $\rho_q = 1$ or 0 if $s\sigma = 0$ or $\beta = 0$, respectively. Using (13), ρ_q^2 equals the proportion of residual variance $\mathcal{V}(z_t) - \mathcal{V}(\alpha_t) = \beta^2 + (s\sigma)^2$ explained by the Gaussian component q_t .

and ρ_a is 1 or 0 according to whether $s\sigma$ equals 0 or $\sqrt{1 - \beta^2}$, respectively. Finally the overall correlation is the correlation between z_t and the signal, consisting of both nonparametric and Gaussian component:

$$\rho \equiv \text{cor}(z_t, \alpha_t + \beta q_t) = \text{cor}(z_t, z_t - s\epsilon_t) = \frac{1 - (s\sigma)^2}{\sqrt{\text{cov}(z_t - s\epsilon_t)}} = \sqrt{1 - (s\sigma)^2}.$$

with $\rho = \rho_a$ if $\beta = 0$. If $s = 0$, there is no measurement error in (8) or, equivalently, there is no roughness penalty in (9) and $\rho_q = \rho_a = \rho = 1$.

10. Special cases of the PG copula

Table ?? sets out possible choices for s and β . These combinations are discussed in the further subsections. Note that $s = 0$ does not imply $s\sigma = 0$ as $s \rightarrow 0$ may imply $s\sigma$ converges to a positive number.

10.1. Empirical copula

The empirical copula results in (8) when $s = 0$. In this case $z_t = \alpha_t + \beta q_t$. The deviation of $z_t - (\alpha_t + \beta q_t)$ can be made zero by selecting appropriate η_t . Hence $s = 0$ implies the fitted copula is the empirical copula. Note $\mathcal{V}(\alpha_t) = 1 - \beta^2$ and since β is arbitrary, the correlation between z_t and the nonparametric component is 1.

10.2. Large local variation

The other extreme $s \rightarrow \infty$ is displayed in the bottom rows of Table ??. Large s forces negligible σ and straight line behaviour for α_t yielding

$$z_t = \alpha_0(1 - 2u_t) + \beta q_t + s\epsilon_t,$$

where $s\epsilon_t$ has finite variance $(s\sigma)^2$.

10.3. Gaussian copula

If $s \rightarrow \infty$ and $\alpha_0 = 0$ then (q_t, z_t) is bivariate Gaussian with correlation β and (8) reduces to a Gaussian copula with correlation β and $\rho = \rho_q$. In this case $s\sigma = \sqrt{1 - \beta^2}$.

10.4. Probit copula

If $s \rightarrow \infty$ and $\beta = 0$ then (8) defines a probit copula. From (??), $\alpha_0 = \pm\sqrt{3\{1 - (s\sigma)^2\}}$.

10.4.1. Independence

If $s \rightarrow \infty$ and $\alpha_0 = \beta = 0$ then the variables are independent. The situation is achieved either letting $\beta \rightarrow 0$ with a Gaussian copula or letting $\alpha_0 \rightarrow 0$ with a Probit copula. Under independence $s\sigma = 1$.

10.5. Conditionally Gaussian copula

This is the intermediate situation where $0 < s < \infty$. Testing $\alpha_0 = 1/s = 0$ is a test for a Gaussian copula. Goodness of fit is measured with

$$\hat{\rho}^2 = 1 - (s\hat{\sigma})^2 = 1 - \mathcal{E}\{(z_t - \hat{z}_t)^2\}, \quad (15)$$

where \hat{z}_t is the estimate of z_t based on (8) using estimates of α_0 , β and $s\sigma$. Values of $\hat{\rho}^2$ lie between 0 and 1 and have the usual interpretation: near 1 indicates the percentiles of u well explain z values of v . Note $R \rightarrow \beta$ if $\alpha_0 = 0$ and $s \rightarrow \infty$.

11. Estimation of the PG model

A convenient feature of the GC model is the straightforward estimation. The unknown parameters are α_0 , β , σ , and s . Given s , estimates of the other parameters are in closed form – generalised least squares (GLS) estimators of α_0 and β , while σ is estimated from the fitted residuals. GLS estimates are derived using the Kalman filter (KF). The KF and the associated Smoothing filter (SF) also deliver diagnostics for assessing goodness of fit and estimates of sampling error. In these calculations s is either given or alternatively is iterated over to produce a quasi maximum likelihood estimate.

From (15)

$$\mathcal{E}\left\{\left(\frac{z_t - \hat{z}_t}{s\hat{\sigma}}\right)^2\right\} = 1$$

suggesting the comparison of standardised residuals to ± 2 and perhaps plotting the percentile rank of the standardised residuals against u_t to identify lack of fit.

The implied correlation between z_t and q_t is

$$\beta = \frac{\beta}{\sqrt{\beta^2 + (s\sigma)^2}} = \pm \left\{1 + \left(\frac{s\sigma}{\beta}\right)^2\right\}^{-1/2}.$$

The term $\beta/(s\sigma)$ is the signal to noise ratio. If $z_t, q_t \sim (0, 1)$ then $\beta^2 + (s\sigma)^2 = 1$ and

$$\beta = \pm \sqrt{1 - (s\sigma)^2} = R = \beta, \quad \frac{\beta}{s\sigma} = \frac{\pm \sqrt{1 - (s\sigma)^2}}{s\sigma} = \frac{\beta}{\sqrt{1 - \beta^2}}$$

Thus the empirical correlation between z_t and q_t is

$$\sum_{t=1}^n z_t q_t = \beta + \delta_0 \sum_{t=1}^n t q_t$$

By construction both z_t and q_t have mean zero and variance 1 implying the correlation between z_t and q_t is

If $\beta = 0$ then

$$z_t = \alpha_0 + n\delta_0 u_t + \beta\Phi^-(u_t) + s\epsilon_t,$$

Testing $\delta_0 = 0$ amounts to testing for independence. If $\beta = \pm 1$ then $z_t - (\pm q_t) = \alpha_0 + t\delta_0$ implying $\alpha_0 = n\delta_0 = 0$ and $z_t = \pm q_t$, that is the variables are co or counter monotonic. Finally if $\beta \neq 0, \pm 1$ then $z_t = \alpha_0 + t\delta_0 + \beta q_t + s\epsilon_t$ since $\alpha_0 = \delta_0 = 0$, implied by the fact that Since both z_t and q_t are, by construction, standard normal, this amounts to prescribing a Gaussian copula with correlation indicated in Table ???. The ratio $\beta/(s\sigma)$ is the signal to noise ratio.

The middle column of the body of Table ?? displays alternatives for finite $s \neq 0$. If $\beta = 0$ the implied copula is nonparametric with α_t evolving smoothly as a function of t with the actual degree of smoothness increasing with s . A constraint is $\alpha_0 = \alpha_n = \sum_t \delta_t = 0$ and hence α_t forms a bridge. If $\beta = \pm 1$ then the signed difference $z_t - (\pm q_t)$ is modelled as a smooth. In a fit, the bridge structure is expected to be enforced by the data.

The scale σ quantifies roughness in the signal α_t but its size is determined by s with $s\sigma \leq 1$ implied by the standard normality of z_t . In fitting, s is either chosen a-priori, or estimated together with the other parameters. Given s , estimates of β , α_0 , δ_0 and σ are closed form generalised least squares calculations. Thus s can be estimated using a one dimensional search. Given the values of the parameters, standard diagnostics are used to assess the fit and reveal areas of inadequate fit. These techniques are used and explained in the applications.

Use of q_t permits a partially Gaussian explanation of the copula with Gaussianity increasing with s . In the limit the enforced Gaussian copula has correlation R signed according to $\hat{\beta}$. Co or countermonotonicity implies $R = \pm 1$ with the sign indicating the direction of the relationship. Setting $\beta = 0$ excludes the Gaussian copula component.

The fitted values \hat{z}_t stacked in vector \hat{z} provide a basis for simulating all percentiles including extreme percentiles. In particular suppose $z \sim N(\hat{z}, \Sigma)$ where Σ is the covariance matrix implied by the model. Then $v \sim \Phi_*(z)$ is a simulation of the the v_t . Bringing v into the original order yields (u_t, v_t) , a simulation from the copula. Simulated percentiles are mapped back into the original scale as $F_*^-(u_t, v_t)$ to provide simulated (x_t, y_t) outcomes.

Table 2: Predicted z_t with conormal copula model

	$s = 0$	$0 < s < \infty$	$s \rightarrow \infty$
$\beta = 0$	z_t empirical	α_t non parametric probit	$\alpha_0(1 - 2u_t)$ linear probit independence: $\alpha_0 = 0$
$\beta \neq 0, \pm 1$		$\alpha_t + \beta q_t$ conormal	$\alpha_0(1 - 2u_t) + \beta q_t$ Gaussian if $\alpha_0 = 0$
$\beta = \pm 1$		$\alpha_t \pm q_t$ cointegrated departure	$\alpha_0(1 - 2u_t) \pm q_t$ co/counter monotonic if $\alpha_0 = 0$

12. Copula forcing

Even if data suggests a strict Gaussian copula it may be desirable to force stronger dependence, especially in the tails. This forcing is achieved with

13. Application to copula fitting

The following example illustrates the copula fitting technique.

14. Financial sensitivity and contagion

Suppose the singular value decomposition of the matrix S of sensitivities at a particular q is $S = UDV'$. Here D a diagonal matrix of singular values, arranged in decreasing order along the diagonal and U and V are orthonormal: $U'U = V'V = I$. If the series are independent then $U = D = V = I$. If the series are comonotonic then $U = V = (1, 0)$ while $D = \text{diag}(1, 0, \dots, 0)$. Hence in the comonotonic case $S = 11'$.

So the best thing is to compare

$$Q = s_{i1}c_{j1} + \dots + s_{ip}c_{jp}$$

If $Q = I$ then $S = C = I$. If $Q = 11'$, a matrix of ones, then $S = \sqrt{p}(1, 0)$ and $C = (c, 0)$ both have rank 1: here s and c are the leading columns of S and C , respectively and 0 a matrix of zeros.

The approximation $Q \approx sc'$ is exact if Q has rank 1 implying $Q = 11'$. The approximation $Q \approx sc'$ is inadequate if $Q = I$. Inadequacy can be judged from the largest singular value in D , as a proportion of p , the number of variables or rows in Q .

$$Q_* \equiv \frac{1}{p-1}(Q - I) = UDV', \quad s \equiv Q_*1, \quad c \equiv Q'_*1.$$

$$Q = I + s1' + UDV' , \quad s \equiv \frac{1}{p-1}(Q - I)1$$

Then s is the vector of average VaR_q sensitivity of each variable to all others. Further c is the average VaR_q impact of each variable on all others. The matrices U , D and V define the singular value (svd) decomposition of Q_* , arranged so that diagonal matrix D has the singular values on the diagonal in descending order. If $b = d_1 u_1$ and $c = v_1$ where u_1 , d_1 and v_1 are the first column, top diagonal entry, and first column of U , D and V respectively, then for two variables $y \neq x$ in Q ,

$$\Delta_x q_y = s_y + b_y c_x + \epsilon_{yx}$$

This states that percentage sensitivities are, apart from the “error” ϵ_{yx} , an average sensitivity plus a scaled response to the contagious effect of the x variable. The contagious effects contained in c are estimated by maximising the explanation of Q .

The vector $1'Q$ sums the changes in VaR_q when each of the column variables is stressed, and writes this as a proportion of the change in the variable being stressed. These proportional sums, subtracting 1 and divided by $p - 1$ where p is the number of variables, measures the average contagion of each variable on all others: $c' = (p - 1)^{-1}1'(Q - I)$ or $c = (p - 1)^{-1}(Q - I)1$

Alternatively the vector $Q1$ sums the changes in VaR_q of each row variable when all the column variables are stressed. Again it is appropriate to remove the effect of a variable on itself and consider the average over the remaining variables: $s \equiv (p - 1)^{-1}(Q - I)1$. If $Q_* \equiv (p - 1)^{-1}(Q - I)$ then $s = Q_*1$ and $c = Q_*'1$ are the vectors of sensitivities and contagions, respectively. If $Q = I$ then $s = c = 0$. If $Q = 11'$ then $s = c = 1$.

15. Systemic risk and causal chains

A rank one approximation to the matrix Q is $Q \approx sc'$. Vector c is an index of the contagious impact of each of variables on the others while s measures the sensitivity of each variable to each of the others. The vectors s and c are derived from the singular value decomposition $Q = UDV'$ where $U'U = V'V = 1$ and where s and c are the first column of UD and V respectively, assuming the svd is organised so that the singular values in the diagonal matrix D are organised from largest to smallest. The appropriateness of the summarisation sc' is measured with $\text{tr}(Q - sc')$.

If the variables are independent then $Q = I$ and both s and c equal a column of the identity matrix with sc' a matrix of 0's except in a single diagonal position where it is 1. Then all but one variable has a contagion effect and only that variable is sensitive to the contagion provided by the variable.

If the variables are comonotonic then $Q = 11'$ and $s = 1$ and $c = 1$ where 1 denotes a vector of ones. Thus the rank 1 approximation is exact and each variable is equally contagious and equally sensitive.

Note that

$$Q^n \approx (s'c)^{n-1}sc' = ,$$

If the random variables are independent the $Q - I = 0$ and $a = b = k = 0$ and there is no error in the first order svd approximation. If the random variables are comonotonic then $Q = 11'$ and $Q - I$ has ones everywhere except on the diagonal where it is zero. The vector of row means is then $p^{-1}(p - 1)1$

Systemic risk in the system is measured with $b'k = d_1(u_1'v_1)$. In the case of comonotonic random variables $Q = 11'$, $a = 1$ and x , where p is the number of variables, $d_1 = 1$.

Furthermore we may define quantities such as $u^- \equiv \text{VaR}_q(v|u \leq q)$ measuring the impact of a non distressed state in v . For brevity we do not dwell on these constructs in this writeup although the ramifications and potential uses of these constructs will be investigated in the research.

16. Generalising Sklar's Theorem

Sklar's Theorem states that if $F(x)$ is a joint then there exists a C such that $F = C \circ F_*$ where $C_F = F$

17. Literature

We propose a measure for systemic risk: CoVaR, the value at risk (VaR) of financial institutions conditional on other institutions being in distress. We define an institution's (marginal) contribution to systemic risk as the difference between CoVaR and the financial system's VaR. From our estimates of CoVaR for characteristic-sorted portfolios of publicly traded financial institutions, we quantify the extent to which characteristics such as leverage, size, and maturity mismatch predict systemic risk contribution. We argue for macro-prudential regulation based on the degree to which such characteristics forecast systemic risk contribution.

18. Econometric implementation

The above development sets out our proposed broad framework for linking bivariate copulas and marginals to external variables and shocks study the impact of the same on stresses within the system and the contagious effects of crises. Proposed econometric analysis will implement and extend Brownlees and Engle (2010).

19. Data

We will employ publicly available data as published by APRA and other regulators.

References

- Acharya, V., R. Engle, and M. Richardson (2012). Capital shortfall: A new approach to ranking and regulating systemic risks. *The American Economic Review* 102(3), 59–64.
- Acharya, V., L. Pedersen, T. Philippon, and M. Richardson (2012). Measuring systemic risk.
- Adrian, T. and M. K. Brunnermeier (2011). Covar. Technical report, National Bureau of Economic Research.
- Brownlees, C. T. and R. Engle (2010). Volatility, correlation and tails for systemic risk measurement. *New York University, mimeo*.