Thesis progress

Pietro Daniele

UNIVERSITÀ
DEGLI STUDI
DI MILANO

Index

1 1° week

2 2° week

GitHub

The whole code is available at this GitHib link

1 ° week

Uploading file .root

The file uploaded are:

- PowhegPy8_NNLOPS_ggH110.root;
- PowhegPy8_NNLOPS_ggH125.root;
- PowhegPy8_NNLOPS_ggH130.root;
- PowhegPy8_NNLOPS_ggH140.root;

Data

In each file, there is a `TTree myCatNtuple`, which contains all the variables. Only three of them were used:

- **m_yy**: mass distribution;
- **EventWeight**: it contains all type of weights.
- **cutFlow**



$m_{\gamma\gamma}$ distibution

Using all entries the distributions at different masses are:









Data analysis and fits

The distributions are fitted using two type of fit:

- **gaussian fit**: there are 2 parameters:
  - mean μ;
  - sigma σ;

$$f(x, \mu, \sigma) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

- **DCB fit**: there are six parameters:
  - mean μ;
  - sigma σ
  - a_1;
  - p_1;
  - a_2;
  - p_2;

$$f(x, \mu, \sigma, a_1, p_1, a_2, p_2) = \cdots$$

Gaussian and DCB fits

Applying the fits to all entries:









CutFlow

Events with at least two isolated γ with tight identification could be selected, where the transverse momentum of the leading photon is such that $p_T/m_{\gamma\gamma} > 0.35$, whereas that of the sub-leading photon is $p_T/m_{\gamma\gamma} > 0.25$. Then there are also other demands on the "quality" of data collection and triggers.

$\Longrightarrow$ Using the `cutFlow` greater than 13, the events are selected.

$m_{\gamma\gamma}$ distribution with cutFlow > 13

Using `cutFlow`>13 entries the distributions at different masses are:









Gaussian and DCB fits with cutFlow > 13

Applying the fits to all entries:









DCB fit and ratio plot [110 GeV]

The `ratio plot` between the fit and the distribution is added to each DCB fit. Using the 110 GeV mass distribution:



DCB fit and ratio plot [125 GeV]

The `ratio plot` between the fit and the distribution is added to each DCB fit. Using the 125 GeV mass distribution:



DCB fit and ratio plot [130 GeV]

The `ratio plot` between the fit and the distribution is added to each DCB fit. Using the 130 GeV mass distribution:



DCB fit and ratio plot [140 GeV]

The `ratio plot` between the fit and the distribution is added to each DCB fit. Using the 140 GeV mass distribution:



DCB fit+ratioplot [110 GeV] (cutFlow > 13)

The `ratio plot` between the fit and the distribution is added to each DCB fit. Using the 110 GeV mass distribution with `cutFlow` > 13:



DCB fit+ratioplot [125 GeV] (cutFlow > 13)

The `ratio plot` between the fit and the distribution is added to each DCB fit. Using the 125 GeV mass distribution with `cutFlow` > 13:



DCB fit+ratioplot [130 GeV] (cutFlow > 13)

The `ratio plot` between the fit and the distribution is added to each DCB fit. Using the 130 GeV mass distribution with `cutFlow` > 13:



DCB fit+ratioplot [140 GeV] (cutFlow > 13)

The `ratio plot` between the fit and the distribution is added to each DCB fit. Using the 140 GeV mass distribution with `cutFlow` > 13:



2° week

DSCB(m_h)

The DSCD fit are studied with 3 different methods, in addiction to all free parameters fit:

1. multi-fit;
2. μ and σ are functions of m_h;
3. all parameters are functions of m_h;

DSCB multifit

A DSCB multifit consists of a series of consecutive fits:

1. fit on μ and σ, tails params fixed;
2. fit on tails params, μ and σ fixed;
3. fit on all parameters;

DSCB multifit [110 GeV]



DSCB multifit [125 GeV]



DSCB multifit [130 GeV]



DSCB multifit [140 GeV]



$\mu(m_h)$

Fitting the μ values at different m_h with a linear fit:

$$\mu(m_h) = A + B \cdot m_h$$



$\sigma(m_h)$

Fitting the σ values at different m_h with a linear fit:

$$\sigma(m_h) = A + B \cdot m_h$$



$a_1(m_h)$

Fitting the a_1 values at different m_h with a linear fit:

$$a_1(m_h) = A + B \cdot m_h$$



$p_1(m_h)$

Fitting the p_1 values at different m_h with a linear fit:

$$p_1(m_h) = A + B \cdot m_h$$



$a_2(m_h)$

Fitting the a_2 values at different m_h with a linear fit:

$$a_2(m_h) = A + B \cdot m_h$$



$p_2(m_h)$

Fitting the p_2 values at different m_h with a linear fit:

$$p_2(m_h) = A + B \cdot m_h$$



DSCB(m_h) with $\mu(m_h)$ and $\sigma(m_h)$ [110 GeV]



DSCB(m_h) with $\mu(m_h)$ and $\sigma(m_h)$ [125 GeV]



DSCB(m_h) with $\mu(m_h)$ and $\sigma(m_h)$ [130 GeV]



DSCB(m_h) with $\mu(m_h)$ and $\sigma(m_h)$ [140 GeV]



DSCB(m_h) with all params(m_h) [110 GeV]



DSCB(m_h) with all params(m_h) [125 GeV]



DSCB(m_h) with all params(m_h) [130 GeV]



DSCB(m_h) with all params(m_h) [140 GeV]



Global fit [110 GeV]



Global fit [125 GeV]



Global fit [130 GeV]



Global fit [140 GeV]



Single vs Global fit params: μ



Single vs Global fit params: σ



Single vs Global fit params: a_1



Single vs Global fit params: p_1



Single vs Global fit params: a_2



Single vs Global fit params: p_2



Efficiencies(m_h)

$$eff = \frac{\#\ of\ events\ with\ cutFlow > 13}{\#\ of\ all\ events}$$