

# The Arithmetic of Elliptic Curves

Luca NOTARNICOLA

PhD Away Days 2017

September, 2017



# Elliptic curve

## Definition

An **elliptic curve**  $E$  over a field  $K$  of  $\text{char}(K) \neq 2, 3$  is a non-singular algebraic plane curve given by an equation of the form

$$Y^2 = X^3 - AX - B \quad ; \quad A, B \in K .$$


# Elliptic curve

## Definition


An **elliptic curve**  $E$  over a field  $K$  of  $\text{char}(K) \neq 2, 3$  is a non-singular algebraic plane curve given by an equation of the form

$$Y^2 = X^3 - AX - B \quad ; \quad A, B \in K .$$

$$E : Y^2 = X^3 - X + 1$$



$$E : Y^2 = X^3 - 3X$$



# Elliptic curve

- *Discriminant  $\Delta$  of  $E$* : discriminant of the cubic polynomial
- $E$  elliptic curve  $\iff \Delta \neq 0$
- Homogeneous equation of *projective curve*  $E$  in  $\mathbf{P}^2(K)$ :

$$y^2z = x^3 - Axz^2 - Bz^3$$

- For  $z \neq 0$ ,  $[x : y : z] \in \mathbf{P}^2(K) \longleftrightarrow (X, Y) = (x/z, y/z)$
- Unique point with  $z = 0$ : *point at infinity*  $\mathcal{O} = [0 : 1 : 0]$

## Other definition encountered


An *elliptic curve* over a field  $K$  is a smooth projective curve of genus 1 together with a distinguished point  $\mathcal{O}$ .

# Group structure

- Elliptic curves define **group varieties**: The set of points on  $E$  is an abelian group for  $+$  with neutral element  $\mathcal{O}$


# Group structure

- Elliptic curves define **group varieties**: The set of points on  $E$  is an abelian group for  $+$  with neutral element  $\mathcal{O}$




# Group structure

- Elliptic curves define **group varieties**: The set of points on  $E$  is an abelian group for  $+$  with neutral element  $\mathcal{O}$



# Group structure


- Elliptic curves define **group varieties**: The set of points on  $E$  is an abelian group for  $+$  with neutral element  $\mathcal{O}$





# Group structure

- Elliptic curves define **group varieties**: The set of points on  $E$  is an abelian group for  $+$  with neutral element  $\mathcal{O}$



# Torsion points on elliptic curves

Consider:

- $E$  elliptic curve over  $K = \mathbb{Q}$
- $E(\mathbb{Q})$  group of points of  $E$  with coordinates in  $\mathbb{Q}$

# Torsion points on elliptic curves

Consider:

- $E$  elliptic curve over  $K = \mathbb{Q}$
- $E(\mathbb{Q})$  group of points of  $E$  with coordinates in  $\mathbb{Q}$

## Definition

For  $n \in \mathbb{N}_{\geq 2}$ , the  $n$ -th torsion group of  $E$  is defined by

$$E[n] = \{P \in E(\mathbb{Q}) : [n]P := \underbrace{P + \dots + P}_{n \text{ times}} = \mathcal{O}\}$$

# Torsion points on elliptic curves

Consider:

- $E$  elliptic curve over  $K = \mathbb{Q}$
- $E(\mathbb{Q})$  group of points of  $E$  with coordinates in  $\mathbb{Q}$

## Definition

For  $n \in \mathbb{N}_{\geq 2}$ , the  $n$ -th torsion group of  $E$  is defined by

$$E[n] = \{P \in E(\mathbb{Q}) : [n]P := \underbrace{P + \dots + P}_{n \text{ times}} = \mathcal{O}\}$$

- **Important fact:**  $E[n] \simeq \mathbb{Z}/n\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$  as abelian groups

# Elliptic curves and Galois representations

Consider  $E$  elliptic curve over  $K = \mathbb{Q}$

# Elliptic curves and Galois representations

Consider  $E$  elliptic curve over  $K = \mathbb{Q}$

- $\overline{\mathbb{Q}}$  field of algebraic numbers

# Elliptic curves and Galois representations

Consider  $E$  elliptic curve over  $K = \mathbb{Q}$

- $\overline{\mathbb{Q}}$  field of algebraic numbers
- Galois group of the field extension  $\overline{\mathbb{Q}}/\mathbb{Q}$

$$\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) = \{\sigma \in \mathrm{Aut}(\overline{\mathbb{Q}}) : \sigma(x) = x, \forall x \in \mathbb{Q}\}$$

# Elliptic curves and Galois representations

Consider  $E$  elliptic curve over  $K = \mathbb{Q}$

- $\overline{\mathbb{Q}}$  field of algebraic numbers
- Galois group of the field extension  $\overline{\mathbb{Q}}/\mathbb{Q}$

$$\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) = \{\sigma \in \mathrm{Aut}(\overline{\mathbb{Q}}) : \sigma(x) = x, \forall x \in \mathbb{Q}\}$$

- $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  acts on  $E[n] \simeq \mathbb{Z}/n\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$



# Elliptic curves and Galois representations

Consider  $E$  elliptic curve over  $K = \mathbb{Q}$

- $\overline{\mathbb{Q}}$  field of algebraic numbers
- Galois group of the field extension  $\overline{\mathbb{Q}}/\mathbb{Q}$

$$\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) = \{\sigma \in \mathrm{Aut}(\overline{\mathbb{Q}}) : \sigma(x) = x, \forall x \in \mathbb{Q}\}$$

- $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  acts on  $E[n] \simeq \mathbb{Z}/n\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$
- Obtain a Galois representation

$$\rho_n : \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \mathrm{Aut}(E[n]) \simeq \mathrm{GL}_2(\mathbb{Z}/n\mathbb{Z})$$

# The Tate module

- Consider a prime number  $\ell$  and the projective system

$$\dots \rightarrow E[\ell^n] \rightarrow \dots \rightarrow E[\ell^2] \rightarrow E[\ell] \quad (1)$$

with transition maps given by  $P \mapsto [l]P$

# The Tate module

- Consider a prime number  $\ell$  and the projective system

$$\dots \rightarrow E[\ell^n] \rightarrow \dots \rightarrow E[\ell^2] \rightarrow E[\ell] \quad (1)$$

with transition maps given by  $P \mapsto [l]P$

## Definition

The **Tate module** is defined as the projective limit of (1)

$$T_\ell E = \varprojlim E[\ell^n]$$

# The Tate module

- Recall the ring of  $\ell$ -adic integers  $\mathbb{Z}_\ell$

$$\mathbb{Z}_\ell = \varprojlim \mathbb{Z}/\ell^n \mathbb{Z}$$

- From  $E[\ell^n] \simeq \mathbb{Z}/\ell^n \mathbb{Z} \times \mathbb{Z}/\ell^n \mathbb{Z}$  it follows

$$T_\ell E \simeq \mathbb{Z}_\ell \times \mathbb{Z}_\ell$$

- $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  acts on  $T_\ell E$
- Obtain a **Galois representation**

$$\rho_{\ell^\infty} : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{Z}_\ell)$$

# Elliptic curves over finite fields

Consider an elliptic curve  $E$  over  $\mathbb{Q}$  given by  $Y^2 = X^3 - AX - B$

- Without loss of generality coefficients  $A, B$  lie in  $\mathbb{Z}$
- Reduction of  $E$  modulo prime number  $q$

# Elliptic curves over finite fields

Consider an elliptic curve  $E$  over  $\mathbb{Q}$  given by  $Y^2 = X^3 - AX - B$

- Without loss of generality coefficients  $A, B$  lie in  $\mathbb{Z}$
- Reduction of  $E$  modulo prime number  $q$
- Obtain a cubic curve  $\hat{E}$  over finite field  $\mathbb{F}_q$

# Elliptic curves over finite fields

Consider an elliptic curve  $E$  over  $\mathbb{Q}$  given by  $Y^2 = X^3 - AX - B$

- Without loss of generality coefficients  $A, B$  lie in  $\mathbb{Z}$
- Reduction of  $E$  modulo prime number  $q$
- Obtain a cubic curve  $\hat{E}$  over finite field  $\mathbb{F}_q$  – not necessarily elliptic curve, may have singular points !

# Elliptic curves over finite fields

Consider an elliptic curve  $E$  over  $\mathbb{Q}$  given by  $Y^2 = X^3 - AX - B$

- Without loss of generality coefficients  $A, B$  lie in  $\mathbb{Z}$
- Reduction of  $E$  modulo prime number  $q$
- Obtain a cubic curve  $\hat{E}$  over finite field  $\mathbb{F}_q$  – not necessarily elliptic curve, may have singular points !

## Definition

- $q$  is called a **good prime** if  $\hat{E}$  is non-singular
- $q$  is called a **bad prime** otherwise



# Elliptic curves over finite fields

- For good primes  $q \neq \ell$ ,  $\text{Gal}(\overline{\mathbb{F}}_q/\mathbb{F}_q)$  acts on  $T_\ell \hat{E}$ ,  $\ell$ -adic Tate module of  $E$  modulo  $q$
- Galois representation  $\text{Gal}(\overline{\mathbb{F}}_q/\mathbb{F}_q) \rightarrow \text{GL}_2(\mathbb{Z}_\ell)$
- We define integers  $a_q \in \mathbb{Z}$  by the relation

$$|E(\mathbb{F}_q)| = q + 1 - a_q \quad (2)$$

- Theorem (Hasse):  $|a_q| \leq 2\sqrt{q}$

# The $L$ -series of an elliptic curve

## Definition

For  $s \in \mathbb{C}$ , the  $L$ -series of an elliptic curve  $E$  is defined by

$$L(E, s) = \prod_{q \text{ good}} \frac{1}{1 - a_q q^{-s} + q^{1-2s}} \prod_{q \text{ bad}} \frac{1}{1 - a_q q^{-s}} = \sum_{n \geq 1} \frac{a_n}{n^s}$$

# The $L$ -series of an elliptic curve

## Definition

For  $s \in \mathbb{C}$ , the  $L$ -series of an elliptic curve  $E$  is defined by

$$L(E, s) = \prod_{q \text{ good}} \frac{1}{1 - a_q q^{-s} + q^{1-2s}} \prod_{q \text{ bad}} \frac{1}{1 - a_q q^{-s}} = \sum_{n \geq 1} \frac{a_n}{n^s}$$

- Compute  $a_q$  as follows
  - If  $q$  good prime, use (2)
  - If  $q$  bad prime, look at the unique singular point  $P$  of  $E$  modulo  $q$

$$a_q = \begin{cases} \pm 1 & \text{if } P \text{ is an ordinary double point (node)} \\ 0 & \text{if } P \text{ is not an ordinary double point (cusp)} \end{cases}$$

# A worked example

Consider  $E : y^2 = x^3 + 3$  over  $\mathbb{Q}$

- Look for reduction modulo prime  $q$

# A worked example

Consider  $E : y^2 = x^3 + 3$  over  $\mathbb{Q}$

- Look for reduction modulo prime  $q$
- discriminant  $\Delta = -3^5 \cdot 2^4 \implies \begin{cases} q \geq 5 & \text{good primes} \\ q = 2, 3 & \text{bad primes} \end{cases}$

# A worked example

Consider  $E : y^2 = x^3 + 3$  over  $\mathbb{Q}$

- Look for reduction modulo prime  $q$
- discriminant  $\Delta = -3^5 \cdot 2^4 \implies \begin{cases} q \geq 5 & \text{good primes} \\ q = 2, 3 & \text{bad primes} \end{cases}$
- $E$  modulo 7 defines an elliptic curve  $\hat{E}$  over  $\mathbb{F}_7$ 
  - $|\hat{E}(\mathbb{F}_7)| = 13$
  - Compute  $a_7$  using (2):  $a_7 = 7 + 1 - 13 = -5$

# A worked example

Consider  $E : y^2 = x^3 + 3$  over  $\mathbb{Q}$

- Look for reduction modulo prime  $q$
- discriminant  $\Delta = -3^5 \cdot 2^4 \implies \begin{cases} q \geq 5 & \text{good primes} \\ q = 2, 3 & \text{bad primes} \end{cases}$
- $E$  modulo 7 defines an elliptic curve  $\hat{E}$  over  $\mathbb{F}_7$ 
  - $|\hat{E}(\mathbb{F}_7)| = 13$
  - Compute  $a_7$  using (2):  $a_7 = 7 + 1 - 13 = -5$
- $E$  modulo 3 given by  $y^2 = x^3$  is not an elliptic curve
  - $(0, 0)$  is a cusp  $\implies a_3 = 0$

Thank you for your attention.