

Strogatz

Malcolm

Started 20th November 2025

Contents

1	Flows on a line	2
1.1	Introduction	2
1.2	Geometric intuition	3
1.2.1	Instructive example	3
1.2.2	Fixed points and stability	6

Chapter 1

Flows on a line

1.1 Introduction

We have the general system as

$$\begin{aligned}\dot{x}_1 &= f_1(x_1, \dots, x_n) \\ &\vdots \\ \dot{x}_n &= f_n(x_1, \dots, x_n)\end{aligned}$$

its solutions can be intuitively visualised as trajectories flowing through an n -dimensional phase space. In these early sections we start with the simple case where $n = 1$; then we get a single equation of the form

$$\dot{x} = f(x)$$

We call such systems one-dimensional or first-order systems.

We clarify the following:

1. The word *system* is being used here in the context of a dynamical system, not in the classical sense of a collection of two or more equations. Thus a single equation can be called a ‘system’.
2. We do not allow f to depend explicitly on time. Time-dependent or ‘nonautonomous’ equations of the form $\dot{x} = f(x, t)$ are more complicated because they require two variables. Such a system in this case would be regarded as a two-dimensional system, to be discussed later.

1.2 Geometric intuition

1.2.1 Instructive example

Consider the following nonlinear differential equation:

$$\dot{x} = \sin x$$


This particular equation can be solved in closed form; using separation of variables then integration we get

$$t = \ln \left| \frac{\csc x_0 + \cot x_0}{\csc x + \cot x} \right|$$

where $x = x_0$ at $t = 0$. Although the result is exact, it isn't really possible to interpret. For instance consider the following

1. Suppose $x_0 = \pi/4$. Describe the qualitative features of the solution $x(t)$ for all $t > 0$. What happens as $t \rightarrow \infty$?
2. For an arbitrary initial condition x_0 , what is the behaviour of $x(t)$ as $t \rightarrow \infty$?


The formula above is not transparent. We consider instead a graphical analysis; the differential equation $\dot{x} = \sin x$ represents a *vector field* on the line: it dictates the velocity vector \dot{x} at each x . To sketch this vector field, it is convenient to plot \dot{x} vs x , drawing arrows on the x -axis to indicate the corresponding velocity vector at each x . The arrows point to the right when $\dot{x} > 0$ and to the left when $\dot{x} < 0$:



The *flow* is to the right when $\dot{x} > 0$ and to the left when $\dot{x} < 0$; at points where $\dot{x} = 0$, there is no flow, and such points are therefore called *fixed points*. Solid dots represent *stable* fixed points (often called *attractors/sinks*, because the flow is toward them). Open dots represent *unstable* fixed points (also known as *repellers/sources*).


(next page)

Cont.
We had




We start at x_0 and watch how x is carried along by the flow. Consider now the questions proposed earlier.

1. Starting at $x_0 = \pi/4$, the flow moves to the right faster and faster until x crosses $\pi/2$ (where $\sin x$ reaches its maximum). Then the flow slows down and eventually approaches the stable fixed point $x = \pi$ from the left. As such the qualitative form of the solution is as follows



See that the curve is concave up at first, then concave down; this change in concavity corresponds to the initial acceleration for $x < \pi/2$, followed by the deceleration toward $x = \pi$.
(next page)


2. The same reasoning applies to any initial condition x_0 . If $\dot{x} > 0$ initially, the flow is to the right and x asymptotically approaches the nearest stable point, and similarly to the left for the other case where $\dot{x} < 0$. If $\dot{x} = 0$, then x remains constant. The qualitative form of the solution for any initial condition can be sketched:



Note that the picture can't tell us certain *quantitative* things: for instance we don't know the time at which the speed $|\dot{x}|$ is greatest. But in many cases the *qualitative* information is what we care about. In this case these sketches are useful.

1.2.2 Fixed points and stability

The ideas discussed can be extended to any one-dimensional system $\dot{x} = f(x)$. We draw the graph of $f(x)$.



As before, the flow is to the right where $f(x) > 0$ and to the left when $f(x) < 0$. To find the solution to $\dot{x} = f(x)$ starting from an arbitrary initial condition x_0 , we start at $x = x_0$ and watch how it is carried along by the flow.

The point moves along the x axis according to some function $x(t)$. This function is called the *trajectory* based at x_0 , and it represents the solution of the differential equation starting from the initial condition x_0 . A figure like the one above, which shows all the different trajectories of the system, is called a *phase portrait*.

The appearance of the phase portrait is controlled by the fixed points x^* , defined by $f(x^*) = 0$; they correspond to stagnation points of flow. As shown earlier the solid dots represent stable fixed points while open dots represent unstable fixed points. In terms of trajectories, fixed points represent *equilibrium* solutions (sometimes called steady/constant/rest solutions), since if $x = x^*$ initially it stays there for all of time.

An equilibrium is defined to be stable if all sufficiently small disturbances away from it damp out in time, so stable equilibria are represented geometrically by stable fixed points. Conversely, unstable equilibria, in which disturbances grow over time, are represented by unstable points.

(next page)

Instructive examples

Say we want to find all the fixed points for $\dot{x} = x^2 - 1$, and then classify their stability.