

Bondy & Murty

Malcolm

Started 13th October 2025

Contents

1	Graphs	2
1.1	Definitions and examples	2

Chapter 1

Graphs

1.1 Definitions and examples

A *graph* is an ordered pair $(V(G), E(G))$ consisting of a set $V(G)$ of *vertices* and a set $E(G)$, disjoint from $V(G)$, of *edges*, together with an *incidence function* ψ_G that associates with each edge of G an unordered pair of (not necessarily distinct) vertices of G .

If e is an edge and u and v are vertices such that $\psi_G(e) = \{u, v\}$ then e is said to *join* u and v , and the vertices u and v are called the *ends* of e .

We denote the numbers of vertices and edges in G by $v(G)$ and $e(G)$; these two basic parameters are called the *order* and *size* of G respectively.

Example 1

These examples serve to clarify the definition. For notational simplicity, we write uv for the unordered pair $\{u, v\}$.

$$G = (V(G), E(G))$$

where

$$\begin{aligned} V(G) &= \{u, v, w, x, y\} \\ E(G) &= \{a, b, c, d, e, f, g, h\} \end{aligned}$$

and ψ_G is defined by

$$\begin{aligned} \psi_G(a) &= uv & \psi_G(b) &= uu & \psi_G(c) &= vw & \psi_G(d) &= wx \\ \psi_G(e) &= vx & \psi_G(f) &= wx & \psi_G(g) &= ux & \psi_G(h) &= xy \end{aligned}$$

(next page)

Example 2

$$H = (V(H), E(H))$$

where

$$V(H) = \{v_0, v_1, v_2, v_3, v_4, v_5\}$$

$$E(H) = \{e_1, e_2, e_3, e_4, e_5, e_6, e_7, e_8, e_9, e_{10}\}$$

and ψ_H is defined by

$$\begin{array}{llll} \psi_H(e_1) = v_1v_2 & \psi_H(e_2) = v_2v_3 & \psi_H(e_3) = v_3v_4 & \psi_H(e_4) = v_4v_5 \\ \psi_H(e_5) = v_5v_1 & \psi_H(e_6) = v_0v_1 & \psi_H(e_7) = v_0v_2 & \psi_H(e_8) = v_0v_3 \\ & \psi_H(e_9) = v_0v_4 & \psi_H(e_{10}) = v_0v_5 & \end{array}$$

Drawings

Graphs are so named because