

Bloch

Malcolm

Started 15th June 2026

Contents

1	Construction of the Real Numbers	2
1.1	Axioms for the Natural numbers	2
1.1.1	Peano Postulates	2
1.1.2	Consequences of Peano Postulates	3
1.1.3	Addition on $\mathbb{N}$	6
1.1.4	Multiplication on $\mathbb{N}$	8
1.1.5	Properties of addition and multiplication on the naturals	11
1.1.6	Ordering on $\mathbb{N}$	18
1.1.7	Well-ordering	22
1.2	Construction of the integers	23
1.2.1	Construction	23
1.2.2	Operations on the integers	24

Chapter 1

Construction of the Real Numbers

Definition 1.1. Let S be a set. A **binary operation** on S is a function $S \times S \rightarrow S$. A **unary operation** on S is a function $S \rightarrow S$.

Let S be a set and let $*$: $S \times S \rightarrow S$ be a binary operation. If $x, y \in S$, the correct way to write the result of the operation to the pair $\langle x, y \rangle$ would be $*(\langle x, y \rangle)$. However since this is a bit cumbersome one may write $x * y$ instead. Similarly if $\neg$: $S \rightarrow S$ is a unary operation and $x \in S$ one may choose to write $\neg x$ over $\neg(x)$.

1.1 Axioms for the Natural numbers

1.1.1 Peano Postulates

Axiom: Peano Postulates

There exists a set $\mathbb{N}$ with an element $1 \in \mathbb{N}$ and a function $s : \mathbb{N} \rightarrow \mathbb{N}$ that satisfy the following three properties.

- There is no $n \in \mathbb{N}$ such that $s(n) = 1$.
- The function s is injective.
- Let $G \subseteq \mathbb{N}$ be a set. Suppose that $1 \in G$, and that if $g \in G$ then $s(g) \in G$. Then $G = \mathbb{N}$.

It will be shown that the set $\mathbb{N}$ is unique. We can therefore give it a name.

Definition 1.2. The set of **natural numbers** denoted $\mathbb{N}$, is the set the existence of which is given in the Peano Postulates.

1.1.2 Consequences of Peano Postulates

Lemma 1.3. *Let $a \in \mathbb{N}$. Suppose that $a \neq 1$. Then there is a unique $b \in \mathbb{N}$ such that $a = s(b)$. (where existence of s and $\mathbb{N}$ are outlined by the peano axioms)*

Proof. First we prove existence. Let

$$G = \{1\} \cup \{c \in \mathbb{N} \mid \exists b \in \mathbb{N}(s(b) = c)\}$$

(which exists by the subset, pairing, and union axioms) We will prove that $G = \mathbb{N}$. To do this we need to show $1 \in G$, $G \subseteq \mathbb{N}$ and $\forall g \in G(s(g) \in G)$. The first two are straightforward. For the third suppose $g \in G$; we have two cases. First suppose $g \in \{1\}$; then $g = 1 \in \mathbb{N}$. Since $s : \mathbb{N} \rightarrow \mathbb{N}$, we then have $s(1) \in \mathbb{N}$. So by definition $s(1) \in \{c \in \mathbb{N} \mid \exists b \in \mathbb{N}(s(b) = c)\}$ and $s(g) = s(1) \in G$. Now suppose $g \in \{c \in \mathbb{N} \mid \exists b \in \mathbb{N}(s(b) = c)\}$; then by definition $g \in \mathbb{N}$ and $s(g) \in \mathbb{N}$ exists. By definition $s(g) \in \{c \in \mathbb{N} \mid \exists b \in \mathbb{N}(s(b) = c)\}$ so $s(g) \in G$. By the third peano postulate we then can assert that $G = \mathbb{N}$. So $a \in \mathbb{N} = G$. Since $a \neq 1$ we must then have $a \in \{c \in \mathbb{N} \mid \exists b \in \mathbb{N}(s(b) = c)\}$ and by definition, $\exists b \in \mathbb{N}(s(b) = a)$.

Now for uniqueness. Let m and n be arbitrary elements of $\mathbb{N}$ such that $a = s(m)$ and $a = s(n)$. Then $\langle m, a \rangle \in s$ and $\langle n, a \rangle \in s$. Since s is injective we must have $n = m$. $\square$

Theorem 1.4. *Let H be a set, Let $e \in H$ and let $k : H \rightarrow H$ be a function. Then there is a unique function $f : \mathbb{N} \rightarrow H$ such that $f(1) = e$ and $f \circ s = k \circ f$.*

Proof. Letting H, e , and k be arbitrary such that $e \in H$ and $k : H \rightarrow H$. We will show existence and uniqueness separately. We start with uniqueness.

Let g and h be arbitrary such that $g : \mathbb{N} \rightarrow H$, $g(1) = e$, and $g \circ s = k \circ g$ and $h : \mathbb{N} \rightarrow H$, $h(1) = e$, and $h \circ s = k \circ h$. We will first define V as the unique set (guaranteed by the subset and extensionality axioms) such that

$$\forall a(a \in V \leftrightarrow a \in \mathbb{N} \wedge g(a) = h(a))$$

and show that $V = \mathbb{N}$. By peano's postulates, we need to show $V \subseteq \mathbb{N}$, $1 \in V$, and $\forall n \in V(s(n) \in V)$. The first is straightforward. For the second requirement, we know by peano's postulates that $1 \in \mathbb{N}$, so by our assumptions $g(1)$ and $h(1)$ exist and $g(1) = e = h(1)$. So $1 \in V$.

For the third requirement, let n be arbitrary and suppose that $n \in V$. Then by definition $n \in \mathbb{N}$ and $g(n) = h(n)$. Since $n \in \mathbb{N} = \text{dom}(h \circ s)$, we have, by our assumptions,

$$h(s(n)) = h \circ s(n) = k \circ h(n) = k(h(n)) = k(g(n)) = k \circ g(n) = g \circ s(n) = g(s(n))$$

we know by definition $\text{ran}(s) \subseteq \mathbb{N}$, so $s(n) \in \mathbb{N}$ and again by definition $s(n) \in V$. We can conclude that $V = \mathbb{N}$.

Suppose $\langle x, y \rangle \in g$, then $x \in \text{dom} g = \mathbb{N} = V$. So by definition $y = g(x) = h(x)$, and $\langle x, y \rangle \in h$. Suppose $\langle x, y \rangle \in h$, then $x \in \text{dom} h = \mathbb{N} = V$. So by definition $y = h(x) = g(x)$, and $\langle x, y \rangle \in g$.

(next page)

Now for existence. We need to come up with a function from $\mathbb{N}$ into H such that $f(1) = e$ and $f \circ s = k \circ f$. We first define C , where

$$\forall X (X \in C \leftrightarrow X \subseteq \mathbb{N} \times H \wedge \langle 1, e \rangle \in X \wedge \forall n \forall y (\langle n, y \rangle \in X \rightarrow \langle s(n), k(y) \rangle \in X))$$

(guaranteed by the subset and extensionality axioms.) We propose that $f = \bigcap C$ is the desired function. To ensure its existence, we need to show that C is nonempty. We will show that $\mathbb{N} \times H \in C$. For the first requirement, clearly $\mathbb{N} \times H \subseteq \mathbb{N} \times H$. Next, since $1 \in \mathbb{N}$ and $e \in H$ then $\langle 1, e \rangle \in \mathbb{N} \times H$. Finally, let n and y be arbitrary such that $\langle n, y \rangle \in \mathbb{N} \times H$. Since $s : \mathbb{N} \rightarrow \mathbb{N}$ and $k : H \rightarrow H$, we must then have $s(n) \in \mathbb{N}$ and $k(y) \in H$. So $\langle s(n), k(y) \rangle \in \mathbb{N} \times H$. $f = \bigcap C$ is therefore a valid candidate.

Before we prove the functionality of f , we first state a few facts about f . First we note that $f \in C$. Let arbitrary $x \in \bigcap C$. Since C is nonempty there exists $X \in C$, and by definition $x \in X \subseteq \mathbb{N} \times H$. So $\bigcap C \subseteq \mathbb{N} \times H$. Next let arbitrary $X \in C$. By definition $\langle 1, e \rangle \in X$, so $\langle 1, e \rangle \in \bigcap C$. Finally let n and y be arbitrary and suppose $\langle n, y \rangle \in \bigcap C$. We need to show that $\langle s(n), k(y) \rangle \in \bigcap C$. Let arbitrary $X \in C$. We have $\langle n, y \rangle \in X$. Since $X \in C$, by definition we must then have $\langle s(n), k(y) \rangle \in X$. We are done.

Next we note that $\forall X \in C (f \subseteq X)$. This is straightforward to prove. Finally we show

$$\begin{aligned} \forall n \forall y ([\langle n, y \rangle \in f \wedge \langle n, y \rangle \neq \langle 1, e \rangle] \\ \rightarrow \exists m \exists u (\langle m, u \rangle \in f \wedge \langle n, y \rangle = \langle s(m), k(u) \rangle)) \end{aligned}$$

Let n and y be arbitrary such that $\langle n, y \rangle \in f$ and $\langle n, y \rangle \neq \langle 1, e \rangle$. Suppose, for contradiction, that the statement after the implication is false. Then $\forall m \forall u (\langle m, u \rangle \in f \rightarrow \langle n, y \rangle \neq \langle s(m), k(u) \rangle)$. The set $f - \{\langle n, y \rangle\}$ exists by the subset axiom; we will show that this set is a member of C . We know that $f \in C$, so $f \subseteq \mathbb{N} \times H$, and it is straightforward to show that $(f - \{\langle n, y \rangle\}) \subseteq \mathbb{N} \times H$. Since $\langle 1, e \rangle \neq \langle n, y \rangle$ and $\langle 1, e \rangle \in f$, we also have $\langle 1, e \rangle \in f - \{\langle n, y \rangle\}$. Finally, let arbitrary $\langle a, b \rangle \in f - \{\langle n, y \rangle\}$. Then $\langle a, b \rangle \in f \in C$ and $\langle s(a), k(b) \rangle \in f$. Since $\langle a, b \rangle \in f$, we must have, by our assumption, $\langle n, y \rangle \neq \langle s(a), k(b) \rangle$ and $\langle s(a), k(b) \rangle \in f - \{\langle n, y \rangle\}$. So $f - \{\langle n, y \rangle\} \in C$. It is straightforward to show that $f \not\subseteq f - \{\langle n, y \rangle\}$, contradicting the fact that $\forall X \in C (f \subseteq X)$.

We now prove functionality. We will do this by defining G :

$$\forall z (z \in G \leftrightarrow z \in \mathbb{N} \wedge \exists! x \in H (\langle z, x \rangle \in f))$$

(which exists by the subset axiom) and using the peano postulates to show that $G = \mathbb{N}$.

(next page)

We defined G :

$$\forall z(z \in G \leftrightarrow z \in \mathbb{N} \wedge \exists! x \in H(\langle z, x \rangle \in f))$$

To show that $G = \mathbb{N}$, we need to show that $G \subseteq \mathbb{N}$, $1 \in G$, and $\forall n \in G(s(n) \in G)$. The first requirement is straightforward. For the second, we already know by definition that $1 \in \mathbb{N}$. Since $f \in C$ we know by definition that $\langle 1, e \rangle \in f$, where $e \in H$. Let arbitrary $x' \in H$ such that $\langle 1, x' \rangle \in f$. Suppose for contradiction that $e \neq x'$. Then we have $\langle 1, e \rangle \neq \langle 1, x' \rangle$; so by the third statement proven earlier there exists m, u such that $\langle m, u \rangle \in f$ and $\langle 1, x' \rangle = \langle s(m), k(u) \rangle$. This means that $1 = s(m)$, where since $f \subseteq \mathbb{N} \times H$, $m \in \mathbb{N}$. This contradicts the peano postulates, which assert that $\neg \exists n \in \mathbb{N}(s(n) = 1)$.

For the third and final requirement, let n be an arbitrary element of G . Then by definition $n \in \mathbb{N}$ and there exists a unique x such that $\langle n, x \rangle \in f$. By definition we know $s(n) \in \mathbb{N}$, and since $f \in C$, we have $\langle s(n), k(x) \rangle \in f$, this verifies existence. Now let x' be arbitrary such that $\langle s(n), x' \rangle \in f$. By peano's postulates we know that $s(n) \neq 1$, so $\langle s(n), x' \rangle \neq \langle 1, e \rangle$. By the third statement proven earlier there exists m, u such that $\langle m, u \rangle \in f$ and $\langle s(n), x' \rangle = \langle s(m), k(u) \rangle$. So $s(n) = s(m)$ and $x' = k(u)$. Since s is injective we must have $m = n$. So $\langle n, u \rangle \in f$. By uniqueness of x we then have $u = x$. So $x' = k(u) = k(x)$, proving uniqueness. $s(n) \in G$.

We have shown that $G = \mathbb{N}$. It follows from this that $f : \mathbb{N} \rightarrow H$. For functionality, let $x \in \text{dom}f$. Then there exists y such that $\langle x, y \rangle \in f$. Since $f \subseteq \mathbb{N} \times H$ we have $x \in \mathbb{N} = G$, and by definition y is unique. Next to show $\text{dom}f = \mathbb{N}$, first let arbitrary $x \in \text{dom}f$, similarly it immediately follows that $x \in \mathbb{N}$. Now let $x \in \mathbb{N}$. Then $x \in G$ and by definition there exists $y \in H$ such that $\langle x, y \rangle \in f$, meaning $x \in \text{dom}f$. Finally to show $\text{ran}f \subseteq H$, let $y \in \text{ran}f$. Then there exists x such that $\langle x, y \rangle \in f \subseteq \mathbb{N} \times H$, so $y \in H$.

The fact that $\langle 1, e \rangle \in \bigcap C = f$ was already shown earlier. Finally we show $f \circ s = k \circ f$. Let $\langle x, y \rangle \in f \circ s$. Then $y = f \circ s(x) = f(s(x))$. Since $s(x) \neq 1$, $\langle s(x), y \rangle \neq \langle 1, e \rangle$ and as proven earlier there then exists m, u such that $\langle m, u \rangle \in f$ and $\langle s(x), y \rangle = \langle s(m), k(u) \rangle$. So $s(x) = s(m)$ and $y = k(u)$. Since s is injective $x = m$. We then have $\langle x, u \rangle \in f$ and $y = k(u) = k(f(x)) = k \circ f(x)$. so $\langle x, y \rangle \in k \circ f$. Next, suppose $\langle x, y \rangle \in k \circ f$. By definition there exists z such that $\langle x, z \rangle \in f$ and $\langle z, y \rangle \in k$. Since $f \in C$ we have $\langle s(x), k(z) \rangle \in f$. We then have $y = k(z) = f(s(x)) = f \circ s(x)$ and $\langle x, y \rangle \in f \circ s$. We are done. $\square$

1.1.3 Addition on $\mathbb{N}$

Theorem 1.5. *There is a unique binary operation $+$: $\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ that satisfies the following two properties for all $n, m \in \mathbb{N}$:*

- $n + 1 = s(n)$
- $n + s(m) = s(n + m)$

Proof. We start with uniqueness. Let $+$ and $\oplus$ be arbitrary such that $+, \oplus : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$, $\forall m \in \mathbb{N} \forall n \in \mathbb{N} (+(\langle n, 1 \rangle) = s(n) \wedge +(\langle n, s(m) \rangle) = s(+(\langle n, m \rangle)))$, and $\forall m \in \mathbb{N} \forall n \in \mathbb{N} (\oplus(\langle n, 1 \rangle) = s(n) \wedge \oplus(\langle n, s(m) \rangle) = s(\oplus(\langle n, m \rangle)))$. We need to show $+$ = $\oplus$. We define G as

$$\forall x (x \in G \leftrightarrow x \in \mathbb{N} \wedge \forall n \in \mathbb{N} (+(\langle n, x \rangle) = \oplus(\langle n, x \rangle)))$$

(guaranteed by the subset axiom.) We will show that $G = \mathbb{N}$. To do this we need to show $G \subseteq \mathbb{N}$, $1 \in G$, and $\forall g \in G (s(g) \in G)$. The first requirement is straightforward. For the second requirement, we already know $1 \in \mathbb{N}$, and for arbitrary $n \in \mathbb{N}$, $+(\langle n, 1 \rangle) = s(n) = \oplus(\langle n, 1 \rangle)$.

For the third requirement, let arbitrary $g \in G$. Then by definition $g \in \mathbb{N}$ and $\forall n \in \mathbb{N} (+(\langle n, g \rangle) = \oplus(\langle n, g \rangle))$. Since $s : \mathbb{N} \rightarrow \mathbb{N}$ we know that $s(g) \in \mathbb{N}$ exists. Let n be arbitrary. By definition $+(\langle n, s(g) \rangle) = s(+(\langle n, g \rangle))$ and $\oplus(\langle n, s(g) \rangle) = s(\oplus(\langle n, g \rangle))$. By assumption we can assert that $+(\langle n, g \rangle) = \oplus(\langle n, g \rangle)$, so $+(\langle n, s(g) \rangle) = s(+(\langle n, g \rangle)) = s(\oplus(\langle n, g \rangle)) = \oplus(\langle n, s(g) \rangle)$. We conclude that $s(g) \in G$ and $G = \mathbb{N}$.

Let $\langle \langle x, y \rangle, z \rangle \in +$. By definition $y \in \mathbb{N} = G$ and $x \in \mathbb{N}$. So $+(\langle x, y \rangle) = \oplus(\langle x, y \rangle)$ and $z = \oplus(\langle x, y \rangle)$. So $\langle \langle x, y \rangle, z \rangle \in \oplus$. Now let $\langle \langle x, y \rangle, z \rangle \in \oplus$. A similar argument shows that $\langle \langle x, y \rangle, z \rangle \in +$.

Now for existence. We want to show

$$\begin{aligned} \exists + (+ : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N} \wedge \forall m \in \mathbb{N} \forall n \in \mathbb{N} (&+(\langle n, 1 \rangle) = s(n) \\ &\wedge +(\langle n, s(m) \rangle) = s(+(\langle n, m \rangle)))) \end{aligned}$$

Theorem 1.4 allows us to assert

$$\forall k \forall H \forall e ([e \in H \wedge k : H \rightarrow H] \rightarrow \exists ! f (f : \mathbb{N} \rightarrow H \wedge f(1) = e \wedge f \circ s = k \circ f))$$

It immediately follows that

$$\forall p \in \mathbb{N} \exists ! f (f : \mathbb{N} \rightarrow \mathbb{N} \wedge f(1) = p \wedge f \circ s = s \circ f)$$

We will define $+$ as the set where

$$\begin{aligned} \forall z (z \in + \leftrightarrow \exists a \in \mathbb{N} \exists b \in \mathbb{N} (\exists f (f : \mathbb{N} \rightarrow \mathbb{N} \wedge f(1) = s(a) \wedge f \circ s = s \circ f \\ \wedge \exists c (\langle b, c \rangle \in f \wedge z = \langle \langle a, b \rangle, c \rangle)))) \end{aligned}$$

Existence follows from the subset axiom, where it can be shown that $+$ $\subseteq (\mathbb{N} \times \mathbb{N}) \times \mathbb{N}$.

(next page)

We defined $+$ as

$$\forall z(z \in + \leftrightarrow \exists a \in \mathbb{N} \exists b \in \mathbb{N} (\exists f(f : \mathbb{N} \rightarrow \mathbb{N} \wedge f(1) = s(a) \wedge f \circ s = s \circ f \wedge \exists c(\langle b, c \rangle \in f \wedge z = \langle \langle a, b \rangle, c \rangle))))$$

First we prove functionality. Let x be an arbitrary element of $\text{dom}(+)$. Then there exists y such that $\langle x, y \rangle \in +$. Let y' be arbitrary such that $\langle x, y' \rangle \in +$. $\langle x, y \rangle \in +$ by definition means that there exists $a, b \in \mathbb{N}$, and $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $f(1) = s(a)$ and $f \circ s = s \circ f$, and that there exists c such that $\langle b, c \rangle \in f$ and $\langle x, y \rangle = \langle \langle a, b \rangle, c \rangle$. $\langle x, y' \rangle \in +$ similarly implies the existence of $a_0, b_0 \in \mathbb{N}$ and $f_0 : \mathbb{N} \rightarrow \mathbb{N}$ such that $f_0(1) = s(a_0)$ and $f_0 \circ s = s \circ f_0$, and the existence of c_0 such that $\langle b_0, c_0 \rangle \in f_0$ and $\langle x, y' \rangle = \langle \langle a_0, b_0 \rangle, c_0 \rangle$. Then $\langle a, b \rangle = x = \langle a_0, b_0 \rangle$, so $a_0 = a$ and $b_0 = b$. It follows that $s(a_0) = s(a) \in \mathbb{N}$. This and

$$\forall p \in \mathbb{N} \exists! f(f : \mathbb{N} \rightarrow \mathbb{N} \wedge f(1) = p \wedge f \circ s = s \circ f)$$

implies that $f = f_0$. So we have $y = c = f(b) = f(b_0) = f_0(b_0) = c_0 = y'$.

Next we show that $\text{dom}(+) = \mathbb{N} \times \mathbb{N}$. First let x be an arbitrary element of $\text{dom}(+)$. Then there exists y such that $\langle x, y \rangle \in +$. By definition there then exists $a, b \in \mathbb{N}$ and there exists c such that $\langle x, y \rangle = \langle \langle a, b \rangle, c \rangle$. So $x = \langle a, b \rangle \in \mathbb{N} \times \mathbb{N}$.

Now suppose $x \in \mathbb{N} \times \mathbb{N}$. Then by definition there exists $a, b \in \mathbb{N}$ such that $x = \langle a, b \rangle$. $s(a) \in \mathbb{N}$ exists, so by theorem 1.4 there exists f such that $f : \mathbb{N} \rightarrow \mathbb{N}$, $f(1) = s(a)$, and $f \circ s = s \circ f$. Since $b \in \mathbb{N} = \text{dom} f$, $f(b)$ exists and $\langle b, f(b) \rangle \in f$. Then by definition, $\langle \langle a, b \rangle, f(b) \rangle \in +$, so $x = \langle a, b \rangle \in \text{dom}(+)$.

Next we show that $\text{ran}(+) \subseteq \mathbb{N}$. Let arbitrary $y \in \text{ran}(+)$. Then there exists x such that $\langle x, y \rangle \in +$. By definition there then exists $a, b \in \mathbb{N}$, $f : \mathbb{N} \rightarrow \mathbb{N}$, and c such that $\langle b, c \rangle \in f$ and $\langle x, y \rangle = \langle \langle a, b \rangle, c \rangle$. So we have $y = c = f(b) \in \mathbb{N}$.

Now we verify the two properties of addition, that is,

$$\forall m \in \mathbb{N} \forall n \in \mathbb{N} (+(\langle n, 1 \rangle) = s(n) \wedge +(\langle n, s(m) \rangle) = s(+(\langle n, m \rangle)))$$

Let arbitrary $n, m \in \mathbb{N}$. We know that $+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$, so $\langle n, 1 \rangle \in \mathbb{N} \times \mathbb{N} = \text{dom}(+)$ and there exists y such that $\langle \langle n, 1 \rangle, y \rangle \in +$. By definition there then exists $a, b \in \mathbb{N}$ and $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $f(1) = s(a)$, and there exists c such that $\langle b, c \rangle \in f$ and $\langle \langle n, 1 \rangle, y \rangle = \langle \langle a, b \rangle, c \rangle$. Then $n = a$, $1 = b$, and $y = c$. We then have $\langle 1, y \rangle \in f$ and $y = f(1) = s(a) = s(n)$. We conclude that $\langle \langle n, 1 \rangle, s(n) \rangle \in +$ and $+(\langle n, 1 \rangle) = s(n)$.

(next page)

Now for the second property. We know that $s(m) \in \mathbb{N}$. So $\langle n, s(m) \rangle \in \mathbb{N} \times \mathbb{N} = \text{dom}(+)$ and there exists y such that $\langle \langle n, s(m) \rangle, y \rangle \in +$. By definition there then exists $a, b \in \mathbb{N}$ such that $f : \mathbb{N} \rightarrow \mathbb{N}$, $f(1) = s(a)$, and $f \circ s = s \circ f$, and there exists c such that $\langle b, c \rangle \in f$ and $\langle \langle n, s(m) \rangle, y \rangle = \langle \langle a, b \rangle, c \rangle$. We then have $n = a$, $s(m) = b$, and $y = c = f(b) = f(s(m)) = s(f(m))$. We know that $\langle n, m \rangle \in \mathbb{N} \times \mathbb{N} = \text{dom}(+)$, so there exists y_0 such that $\langle \langle n, m \rangle, y_0 \rangle \in +$. Again by definition there exists $a_0, b_0 \in \mathbb{N}$ and $f_0 : \mathbb{N} \rightarrow \mathbb{N}$ such that $f_0(1) = s(a_0)$ and $f_0 \circ s = s \circ f_0$ and there exists c_0 such that $\langle b_0, c_0 \rangle \in f_0$ and $\langle \langle n, m \rangle, y_0 \rangle = \langle \langle a_0, b_0 \rangle, c_0 \rangle$. Since it follows that $a_0 = n = a$ and $s(a_0) = s(a)$, theorem 1.4 allows us to assert $f = f_0$. Putting it all together, we have $+(\langle n, m \rangle) = y_0 = c_0 = f_0(b_0) = f_0(m) = f(m)$. We conclude that $+(\langle n, s(m) \rangle) = y = s(f(m)) = s(+(\langle n, m \rangle))$. $\square$

1.1.4 Multiplication on $\mathbb{N}$

Theorem 1.6. *There is a unique binary operation $\cdot : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ that satisfies the following two properties for all $n, m \in \mathbb{N}$:*

- $n \cdot 1 = n$
- $n \cdot s(m) = (n \cdot m) + n$

Proof. We start with uniqueness. Let $\cdot, \odot$ be arbitrary. Suppose $\cdot : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ and $\forall m \in \mathbb{N} \forall n \in \mathbb{N} (\cdot(\langle n, 1 \rangle) = n \wedge \cdot(\langle n, s(m) \rangle) = +(\cdot(\langle n, m \rangle), n))$. Similarly suppose $\odot : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ and $\forall m \in \mathbb{N} \forall n \in \mathbb{N} (\odot(\langle n, 1 \rangle) = n \wedge \odot(\langle n, s(m) \rangle) = +(\odot(\langle n, m \rangle), n))$. We want to show that $\cdot = \odot$. We do this by defining G such that

$$\forall z (z \in G \leftrightarrow z \in \mathbb{N} \wedge \forall n \in \mathbb{N} (\cdot(\langle n, z \rangle) = \odot(\langle n, z \rangle)))$$

(where existence is assured by the subset axiom) and showing that $G = \mathbb{N}$.

We need to show $G \subseteq \mathbb{N}$, $1 \in G$, and $\forall g \in G (s(g) \in G)$. The first requirement is straightforward. For the second, we already know $1 \in \mathbb{N}$, and see that for arbitrary $n \in \mathbb{N}$ we have $\cdot(\langle n, 1 \rangle) = n = \odot(\langle n, 1 \rangle)$.

For the third requirement let arbitrary $g \in G$. Then by definition $g \in \mathbb{N}$ and $\forall n \in \mathbb{N} (\cdot(\langle n, g \rangle) = \odot(\langle n, g \rangle))$. We know that $s(g) \in \mathbb{N}$ exists. So since $\langle n, s(g) \rangle \in \mathbb{N} \times \mathbb{N} = \text{dom}(\cdot)$, we have, by our assumptions, $\cdot(\langle n, s(g) \rangle) = +(\cdot(\langle n, g \rangle), n) = +(\odot(\langle n, g \rangle), n) = \odot(\langle n, s(g) \rangle)$. So $s(g) \in G$ and $G = \mathbb{N}$.

To prove the desired result, let arbitrary $z \in \cdot \subseteq (\mathbb{N} \times \mathbb{N}) \times \mathbb{N}$. Then there exists $a_0, a_1, b \in \mathbb{N}$ such that $\langle \langle a_0, a_1 \rangle, b \rangle = z \in \cdot$. So $b = \cdot(\langle a_0, a_1 \rangle)$. Since $a_1 \in \mathbb{N} = G$, we can then conclude that $b = \cdot(\langle a_0, a_1 \rangle) = \odot(\langle a_0, a_1 \rangle)$. So $z = \langle \langle a_0, a_1 \rangle, b \rangle \in \odot$. Next suppose $z \in \odot$. It then follows from a similar argument that $z \in \cdot$. We are done.

(next page)

Now for existence. We start by defining a function h for all q , where

$$\forall q \exists h \forall z (z \in h \leftrightarrow \exists m \in \mathbb{N} \exists x \in \mathbb{N} (z = \langle m, x \rangle \wedge \langle \langle m, q \rangle, x \rangle \in +))$$

(which exists by the subset axiom, where $\mathbb{N} \times \mathbb{N}$ is a bounding set. Uniqueness for each q follows from extensionality.) We show that for all $q \in \mathbb{N}$, this h is a function from $\mathbb{N}$ into $\mathbb{N}$. That is

$$\forall q \in \mathbb{N} [\forall z (z \in h \leftrightarrow \exists m \in \mathbb{N} \exists x \in \mathbb{N} (z = \langle m, x \rangle \wedge \langle \langle m, q \rangle, x \rangle \in +)) \rightarrow h : \mathbb{N} \rightarrow \mathbb{N}]$$

Let arbitrary $q \in \mathbb{N}$. Suppose that the statement before the implication is true. First to prove functionality. Let $a \in \text{dom}h$. Then there exists b such that $\langle a, b \rangle \in h$. Let b' be arbitrary such that $\langle a, b' \rangle \in h$. By our assumptions we must have $a, b, b' \in \mathbb{N}$, $+(\langle a, q \rangle) = b$, and $+(\langle a, q \rangle) = b'$. Since $+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ we must have $b = b'$. Next for the domain bound suppose arbitrary $a \in \text{dom}h$. Then there exists b such that $\langle a, b \rangle \in h$. It immediately follows that $a \in \mathbb{N}$. Suppose instead that $a \in \mathbb{N}$. Then $\langle a, q \rangle \in \mathbb{N} \times \mathbb{N} = \text{dom}(+)$. So $+(\langle a, q \rangle) \in \mathbb{N}$ exists and by definition, $\langle a, +(\langle a, q \rangle) \rangle \in h$. So $a \in \text{dom}h$. Finally for the range bound, let arbitrary $b \in \text{ran}h$. Then there exists a such that $\langle a, b \rangle \in h$. It immediately follows by definition that $b \in \mathbb{N}$. We have therefore shown

$$\begin{aligned} \forall q \in \mathbb{N} \exists! h (\forall z (z \in h \leftrightarrow \exists m \in \mathbb{N} \exists x \in \mathbb{N} (z = \langle m, x \rangle \wedge \langle \langle m, q \rangle, x \rangle \in +)) \\ \wedge h : \mathbb{N} \rightarrow \mathbb{N}) \end{aligned}$$

Theorem 1.4 states

$$\forall H \forall q \forall h ([q \in H \wedge h : H \rightarrow H] \rightarrow \exists! g (g : \mathbb{N} \rightarrow H \wedge g(1) = q \wedge g \circ s = h \circ g))$$

which allows us to assert

$$\forall q \forall h ([q \in \mathbb{N} \wedge h : \mathbb{N} \rightarrow \mathbb{N}] \rightarrow \exists! g (g : \mathbb{N} \rightarrow \mathbb{N} \wedge g(1) = q \wedge g \circ s = h \circ g))$$

(next page)

We showed

$$\forall q \in \mathbb{N} \exists! h (\forall z (z \in h \leftrightarrow \exists m \in \mathbb{N} \exists x \in \mathbb{N} (z = \langle m, x \rangle \wedge \langle \langle m, q \rangle, x \rangle \in +)) \wedge h : \mathbb{N} \rightarrow \mathbb{N})$$

and

$$\forall q \forall h ([q \in \mathbb{N} \wedge h : \mathbb{N} \rightarrow \mathbb{N}] \rightarrow \exists! g (g : \mathbb{N} \rightarrow \mathbb{N} \wedge g(1) = q \wedge g \circ s = h \circ g))$$

Now we construct $\cdot$ as the set

$$\begin{aligned} \forall z (z \in \cdot \leftrightarrow \exists a \in \mathbb{N} \exists b \in \mathbb{N} (\\ \exists h (\forall z_0 (z_0 \in h \leftrightarrow \exists m \in \mathbb{N} \exists x \in \mathbb{N} (z_0 = \langle m, x \rangle \wedge \langle \langle m, a \rangle, x \rangle \in +)) \wedge h : \mathbb{N} \rightarrow \mathbb{N} \wedge \exists g (g : \mathbb{N} \rightarrow \mathbb{N} \wedge g(1) = a \wedge g \circ s = h \circ g \\ \wedge \exists c (\langle b, c \rangle \in g \wedge z = \langle \langle a, b \rangle, c \rangle)))))) \end{aligned}$$

(existence follows from the subset axiom, where $(\mathbb{N} \times \mathbb{N}) \times \mathbb{N}$ is a bounding set.)
We want to verify

$$\cdot : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N} \wedge \forall m \in \mathbb{N} \forall n \in \mathbb{N} (\cdot(\langle n, 1 \rangle) = n \wedge \cdot(\langle n, s(m) \rangle) = +(\cdot(\langle n, m \rangle), n))$$

We start with functionality. Let arbitrary $p \in \text{dom}(\cdot)$. Then by definition there exists q such that $\langle p, q \rangle \in \cdot$. This means that there exists $a, b \in \mathbb{N}$ and $h : \mathbb{N} \rightarrow \mathbb{N}$ such that $\forall z_0 (z_0 \in h \leftrightarrow \exists m \in \mathbb{N} \exists x \in \mathbb{N} (z_0 = \langle m, x \rangle \wedge \langle \langle m, a \rangle, x \rangle \in +))$. There also exists $g : \mathbb{N} \rightarrow \mathbb{N}$ such that $g(1) = a$ and $g \circ s = h \circ g$, and c such that $\langle b, c \rangle \in g$ and $\langle p, q \rangle = \langle \langle a, b \rangle, c \rangle$. Now suppose there exists $\langle p, q' \rangle \in \cdot$. Similarly there exists $a', b' \in \mathbb{N}$, $h' : \mathbb{N} \rightarrow \mathbb{N}$ such that $\forall z_0 (z_0 \in h' \leftrightarrow \exists m \in \mathbb{N} \exists x \in \mathbb{N} (z_0 = \langle m, x \rangle \wedge \langle \langle m, a' \rangle, x \rangle \in +))$, $g' : \mathbb{N} \rightarrow \mathbb{N}$ such that $g'(1) = a'$ and $g' \circ s = h' \circ g'$, and c' such that $\langle b', c' \rangle \in g'$ and $\langle p, q' \rangle = \langle \langle a', b' \rangle, c' \rangle$. We have $\langle a', b' \rangle = p = \langle a, b \rangle$. So $a' = a$ and $b' = b$. Uniqueness, as proven, implies that $h = h'$ and $g = g'$. So $q' = c' = g'(b') = g(b') = g(b) = c = q$.

Next we show $\text{dom}(\cdot) = \mathbb{N} \times \mathbb{N}$. Let arbitrary $p \in \text{dom}(\cdot)$. Then by definition there exists q such that $\langle p, q \rangle \in \cdot$. So there exists $a, b \in \mathbb{N}$ and c such that $\langle p, q \rangle = \langle \langle a, b \rangle, c \rangle$. So $p = \langle a, b \rangle \in \mathbb{N} \times \mathbb{N}$. Now let $p \in \mathbb{N} \times \mathbb{N}$. Then by definition there exists $a, b \in \mathbb{N}$ such that $p = \langle a, b \rangle$. Since $a \in \mathbb{N}$, as proven there exists $h : \mathbb{N} \rightarrow \mathbb{N}$ such that $\forall z (z \in h \leftrightarrow \exists m \in \mathbb{N} \exists x \in \mathbb{N} (z = \langle m, x \rangle \wedge \langle \langle m, a \rangle, x \rangle \in +))$, and there exists $g : \mathbb{N} \rightarrow \mathbb{N}$ such that $g(1) = a$ and $g \circ s = h \circ g$. By definition we then have $\langle \langle a, b \rangle, g(b) \rangle \in \cdot$ and $p = \langle a, b \rangle \in \text{dom}(\cdot)$.

For the range bound, let $q \in \text{ran}(\cdot)$. Then there exists p such that $\langle p, q \rangle \in \cdot$. So there exists $a, b \in \mathbb{N}$, $g : \mathbb{N} \rightarrow \mathbb{N}$, and c such that $\langle b, c \rangle \in g$ and $\langle p, q \rangle = \langle \langle a, b \rangle, c \rangle$. So $q = c = g(b) \in \mathbb{N}$.

Finally we verify the two properties of multiplication. Let arbitrary $m, n \in \mathbb{N}$. We first show $\cdot(\langle n, 1 \rangle) = n$. We know that $\langle \langle n, 1 \rangle, \cdot(\langle n, 1 \rangle) \rangle \in \cdot$. By definition there then exists $a, b \in \mathbb{N}$, $g : \mathbb{N} \rightarrow \mathbb{N}$ such that $g(1) = a$, and c such that $\langle b, c \rangle \in g$ and $\langle \langle n, 1 \rangle, \cdot(\langle n, 1 \rangle) \rangle = \langle \langle a, b \rangle, c \rangle$. So $\cdot(\langle n, 1 \rangle) = c = g(b) = g(1) = a = n$.
(next page)

Now we show $\cdot(\langle n, s(m) \rangle) = +(\cdot(\langle n, m \rangle), n)$. $s(m) \in \mathbb{N}$ exists, so we have $\langle \langle n, s(m) \rangle, \cdot(\langle n, s(m) \rangle) \rangle \in \cdot$. Then there exists $a, b \in \mathbb{N}$, $h : \mathbb{N} \rightarrow \mathbb{N}$ such that $\forall z_0 (z_0 \in h \leftrightarrow \exists m_0 \in \mathbb{N} \exists x \in \mathbb{N} (z_0 = \langle m_0, x \rangle \wedge \langle \langle m_0, a \rangle, x \rangle \in +))$, $g : \mathbb{N} \rightarrow \mathbb{N}$ such that $g(1) = a$ and $g \circ s = h \circ g$, and c such that $\langle b, c \rangle \in g$ and $\langle \langle n, s(m) \rangle, \cdot(\langle n, s(m) \rangle) \rangle = \langle \langle a, b \rangle, c \rangle$. We then have $\cdot(\langle n, s(m) \rangle) = c = g(b) = g(s(m)) = h(g(m))$ and $a = n$. Similarly we know that $\langle \langle n, m \rangle, \cdot(\langle n, m \rangle) \rangle \in \cdot$. So there exists $a_0, b_0 \in \mathbb{N}$, $h_0 : \mathbb{N} \rightarrow \mathbb{N}$ such that $\forall z_0 (z_0 \in h_0 \leftrightarrow \exists m_0 \in \mathbb{N} \exists x \in \mathbb{N} (z_0 = \langle m_0, x \rangle \wedge \langle \langle m_0, a_0 \rangle, x \rangle \in +))$, $g_0 : \mathbb{N} \rightarrow \mathbb{N}$ such that $g_0(1) = a_0$ and $g_0 \circ s = h_0 \circ g_0$, and c_0 such that $\langle b_0, c_0 \rangle \in g_0$ and $\langle \langle n, m \rangle, \cdot(\langle n, m \rangle) \rangle = \langle \langle a_0, b_0 \rangle, c_0 \rangle$. This means that $a_0 = n = a$, and that by our earlier statements we can assert that $h = h_0$, and $g = g_0$. So $\cdot(\langle n, m \rangle) = c_0 = g_0(b_0) = g(m)$. We then have $\cdot(\langle n, s(m) \rangle) = h(g(m)) = h(\cdot(\langle n, m \rangle))$. From the definition of h , there then exists $m_0, x \in \mathbb{N}$ such that $\langle \cdot(\langle n, m \rangle), h(\cdot(\langle n, m \rangle)) \rangle = \langle m_0, x \rangle$ and $\langle \langle m_0, a \rangle, x \rangle \in +$. This and the fact that $a = n$ allows us to conclude $+(\cdot(\langle n, m \rangle), n) = h(\cdot(\langle n, m \rangle)) = \cdot(\langle n, s(m) \rangle)$. $\square$

1.1.5 Properties of addition and multiplication on the naturals

Theorem 1.7. *Let $a, b, c \in \mathbb{N}$.*

1. *If $a + c = b + c$, then $a = b$ (Cancellation Law for Addition)*
2. *$(a + b) + c = a + (b + c)$ (Associative Law for Addition)*
3. *$1 + a = s(a) = a + 1$*
4. *$a + b = b + a$ (Commutative Law for Addition)*
5. *$a + b \neq 1$*
6. *$a + b \neq a$*
7. *$a \cdot 1 = a = 1 \cdot a$ (Identity Law for Multiplication)*
8. *$(a + b)c = ac + bc$ (Distributive Law)*
9. *$ab = ba$ (Commutative Law for Multiplication)*
10. *$c(a + b) = ca + cb$ (Distributive Law)*
11. *$(ab)c = a(bc)$ (Associative Law for Multiplication)*
12. *If $ac = bc$ then $a = b$ (Cancellation Law for Multiplication)*
13. *$ab = 1$ iff $a = 1 = b$*

(next page)

Proof. (1) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} \forall c \in \mathbb{N} (a + c = b + c \rightarrow a = b)$$

To do this we define the set G as the set where

$$\forall c (c \in G \leftrightarrow c \in \mathbb{N} \wedge \forall a \in \mathbb{N} \forall b \in \mathbb{N} (a + c = b + c \rightarrow a = b))$$

(where existence follows from the subset axiom using $\mathbb{N}$ as the bounding set. Uniqueness follows from extensionality.) Then we show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward; we need to show $1 \in G$ and $\forall g \in G (s(g) \in G)$. For the former, we already know $1 \in \mathbb{N}$. Let arbitrary $a, b \in \mathbb{N}$. Suppose that $a + 1 = b + 1$. By definition we then have $s(a) = s(b)$, so by injectivity $a = b$ and $1 \in G$.

Now the latter. Let arbitrary $g \in G$. As defined we then have $g \in \mathbb{N}$ and $\forall a \in \mathbb{N} \forall b \in \mathbb{N} (a + g = b + g \rightarrow a = b)$. $s(g) \in \mathbb{N}$ exists. Let arbitrary $a, b \in \mathbb{N}$ and suppose that $a + s(g) = b + s(g)$. Then by the definition of addition we have $s(a + g) = s(b + g)$ and, by injectivity, $a + g = b + g$. Since $g \in G$ we then have $a = b$. So $G = \mathbb{N}$. To prove the initial statement, let arbitrary $c \in \mathbb{N}$. Then from $\mathbb{N} = G$ the desired conclusion immediately follows. $\square$

Proof. (2) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} \forall c \in \mathbb{N} ((a + b) + c = a + (b + c))$$

To do this we define G as the set where

$$\forall c (c \in G \leftrightarrow c \in \mathbb{N} \wedge \forall a \in \mathbb{N} \forall b \in \mathbb{N} ((a + b) + c = a + (b + c)))$$

(where existence and uniqueness can be easily shown) and show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward. We need to show $1 \in G$ and $\forall g \in G (s(g) \in G)$.

For the former, we already know $1 \in \mathbb{N}$. Let arbitrary $a, b \in \mathbb{N}$. By the definition of addition we have

$$(a + b) + 1 = s(a + b) = a + s(b) = a + (b + 1)$$

Now the latter. Let arbitrary $g \in G$. Then by definition $g \in \mathbb{N}$ and $\forall a \in \mathbb{N} \forall b \in \mathbb{N} ((a + b) + g = a + (b + g))$. We know $s(g) \in \mathbb{N}$ exists. Let arbitrary $a, b \in \mathbb{N}$. By definition we then have

$$(a + b) + s(g) = s((a + b) + g) = s(a + (b + g)) = a + s(b + g) = a + (b + s(g))$$

So $\mathbb{N} = G$. To prove the initial statement, let arbitrary $c \in \mathbb{N} = G$. The desired conclusion immediately follows. $\square$

(next page)

Proof. (3) We want to show

$$\forall a \in \mathbb{N}(1 + a = s(a) = a + 1)$$

To do this we define G as the set where

$$\forall a(a \in G \leftrightarrow a \in \mathbb{N} \wedge 1 + a = s(a) = a + 1)$$

(Showing existence and uniqueness is straightforward.) We show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward. We need to show that $1 \in G$ and $\forall g \in G(s(g) \in G)$. For the former we already know that $1 \in \mathbb{N}$, and by definition we have $1 + 1 = s(1)$.

Now the latter. Let arbitrary $g \in G$. Then by definition $g \in \mathbb{N}$ and $1 + g = s(g) = g + 1$. We know $s(g) \in \mathbb{N}$ exists. We then have

$$1 + s(g) = s(1 + g) = s(s(g)) = s(g + 1) = (g + 1) + 1 = s(g) + 1$$

So $G = \mathbb{N}$. Proving the initial statement is straightforward. Let arbitrary $a \in \mathbb{N} = G$ and the desired conclusion immediately follows. $\square$

Proof. (4) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N}(a + b = b + a)$$

To do this we define G as the set such that

$$\forall a(a \in G \leftrightarrow a \in \mathbb{N} \wedge \forall b \in \mathbb{N}(a + b = b + a))$$

(Showing existence and uniqueness is straightforward.) We show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward. We need to show that $1 \in G$ and $\forall g \in G(s(g) \in G)$. For the former we already know $1 \in \mathbb{N}$. Letting arbitrary $b \in \mathbb{N}$, by (3) we have $1 + b = b + 1$.

Now the latter. Let arbitrary $g \in G$. Then by definition $g \in \mathbb{N}$ and $\forall b \in \mathbb{N}(g + b = b + g)$. $s(g) \in \mathbb{N}$ exists. Let arbitrary $b \in \mathbb{N}$. Then we have

$$s(g) + b = (g + 1) + b \stackrel{(2)}{=} g + (1 + b) \stackrel{(3)}{=} g + s(b) = s(g + b) = s(b + g) = b + s(g)$$

So $G = \mathbb{N}$. Showing the initial statement is straightforward. Let arbitrary $a \in \mathbb{N} = G$ and the desired conclusion immediately follows. $\square$

(next page)

Proof. (5) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} (a + b \neq 1)$$

Let arbitrary $a, b \in \mathbb{N}$. Suppose for contradiction that $a + b = 1$. We have 2 cases. First suppose $b = 1$. Then $a + b = a + 1 = s(a) = 1$, and the last equality is a contradiction of the peano postulates. Now suppose $b \neq 1$. Then by lemma 1.3 there exists $x \in \mathbb{N}$ such that $b = s(x)$. We then have $a + b = a + s(x) = s(a + x) = 1$, which is again a contradiction. $\square$

Proof. (6) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} (a + b \neq a)$$

We do this by defining G as the set such that

$$\forall a (a \in G \leftrightarrow a \in \mathbb{N} \wedge \forall b \in \mathbb{N} (a + b \neq a))$$

(Showing existence and uniqueness is straightforward.) We show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward. We need to show $1 \in G$ and $\forall g \in G (s(g) \in G)$. For the former, we already know $1 \in \mathbb{N}$. Let arbitrary $b \in \mathbb{N}$. By (5) $1 + b \neq 1$.

Now the latter. Let arbitrary $g \in G$. Then $g \in \mathbb{N}$ and $\forall b \in \mathbb{N} (g + b \neq g)$. $s(g) \in \mathbb{N}$ exists. Let arbitrary $b \in \mathbb{N}$. For contradiction, suppose $s(g) + b = s(g)$. We then have

$$s(g) = s(g) + b \stackrel{(4)}{=} b + s(g) = s(b + g)$$

By injectivity of s and (4) we then have $g = b + g = g + b$. This contradicts $\forall b \in \mathbb{N} (g + b \neq g)$. So $G = \mathbb{N}$. Showing the initial statement is straightforward. Let arbitrary $a \in \mathbb{N} = G$ and the desired conclusion immediately follows. $\square$

Proof. (7) We want to show

$$\forall a \in \mathbb{N} (a \cdot 1 = a = 1 \cdot a)$$

to do this we define G as the set such that

$$\forall a (a \in G \leftrightarrow a \in \mathbb{N} \wedge a \cdot 1 = a = 1 \cdot a)$$

(where existence and uniqueness can be easily shown.) We show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward. We need to show $1 \in G$ and $\forall g \in G (s(g) \in G)$. For the former we already know $1 \in \mathbb{N}$, and $1 \cdot 1 = 1$.

Now the latter. Let arbitrary $g \in G$. Then $g \in \mathbb{N}$ and $g \cdot 1 = g = 1 \cdot g$. $s(g) \in \mathbb{N}$ exists. We then have

$$s(g) \cdot 1 = s(g) = s(g \cdot 1) = s(1 \cdot g) = (1 \cdot g) + 1 = 1 \cdot s(g)$$

so $G = \mathbb{N}$. Proving the initial statement is straightforward. Let arbitrary $a \in \mathbb{N} = G$ and the desired conclusion immediately follows. $\square$

(next page)

Proof. (8) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} \forall c \in \mathbb{N} ((a + b)c = (ac) + (bc))$$

To do this we define G as the set such that

$$\forall c (c \in G \leftrightarrow c \in \mathbb{N} \wedge \forall a \in \mathbb{N} \forall b \in \mathbb{N} ((a + b)c = (ac) + (bc)))$$

(where proving existence and uniqueness is straightforward.) We show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward. We need to show $1 \in G$ and $\forall g \in G (s(g) \in G)$. For the former, we already know $1 \in \mathbb{N}$. Let arbitrary $a, b \in \mathbb{N}$. We have

$$(a + b) \cdot 1 = a + b = (a \cdot 1) + (b \cdot 1)$$

Now the latter. Let arbitrary $g \in G$. Then by definition $g \in \mathbb{N}$ and $\forall a \in \mathbb{N} \forall b \in \mathbb{N} ((a + b)g = (ag) + (bg))$. $s(g) \in \mathbb{N}$ exists. Let arbitrary $a, b \in \mathbb{N}$. We have

$$\begin{aligned} (a + b)s(g) &= ((a + b)g) + (a + b) = ((ag) + (bg)) + (a + b) \\ &\stackrel{(2)}{=} (ag) + ((bg) + (a + b)) \stackrel{(4,2)}{=} (ag) + ((bg + b) + a) \\ &= (ag) + ((bs(g)) + a) \stackrel{(4,2)}{=} ((ag) + a) + (bs(g)) \\ &= (as(g)) + (bs(g)) \end{aligned}$$

So $G = \mathbb{N}$. Proving the initial statement is straightforward. Let arbitrary $c \in \mathbb{N} = G$ and the desired conclusion immediately follows. $\square$

Proof. (9) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} (ab = ba)$$

To do this we define G as the set such that

$$\forall a (a \in G \leftrightarrow a \in \mathbb{N} \wedge \forall b \in \mathbb{N} (ab = ba))$$

(where proving existence and uniqueness is straightforward.) We show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward. We need to show $1 \in G$ and $\forall g \in G (s(g) \in G)$. For the former, we know $1 \in \mathbb{N}$. Let arbitrary $b \in \mathbb{N}$. Then by (7) we have $1 \cdot b = b \cdot 1$.

Now the latter. Let arbitrary $g \in G$. Then $g \in \mathbb{N}$ and $\forall b \in \mathbb{N} (gb = bg)$. $s(g) \in \mathbb{N}$ exists. Let arbitrary $b \in \mathbb{N}$. We have

$$s(g)b = (g + 1)b \stackrel{(8)}{=} (gb) + (1 \cdot b) \stackrel{(7)}{=} (bg) + b = bs(g)$$

So $G = \mathbb{N}$. Proving the initial statement is straightforward. Let arbitrary $a \in \mathbb{N} = G$ and the desired conclusion immediately follows. $\square$

(next page)

Proof. (10) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} \forall c \in \mathbb{N} (c(a + b) = (ca) + (cb))$$

Let arbitrary $a, b, c \in \mathbb{N}$. We have

$$c(a + b) \stackrel{(9)}{=} (a + b)c \stackrel{(8)}{=} (ac) + (bc) \stackrel{(9)}{=} (ca) + (cb) \quad \square$$

Proof. (11) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} \forall c \in \mathbb{N} ((ab)c = a(bc))$$

To do this we define G as the set such that

$$\forall b (b \in G \leftrightarrow b \in \mathbb{N} \wedge \forall a \in \mathbb{N} \forall c \in \mathbb{N} ((ab)c = a(bc)))$$

(where proving existence and uniqueness is straightforward.) We show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward. We need to show $1 \in G$ and $\forall g \in G (s(g) \in G)$. For the former, we already know $1 \in \mathbb{N}$. Let arbitrary $a, c \in \mathbb{N}$. We have

$$(a \cdot 1)c = ac \stackrel{(7)}{=} a(1 \cdot c)$$

Now the latter. Let arbitrary $g \in G$. Then $g \in \mathbb{N}$ and $\forall a \in \mathbb{N} \forall c \in \mathbb{N} ((ag)c = a(gc))$. $s(g) \in \mathbb{N}$ exists. Let arbitrary $a, c \in \mathbb{N}$. We have

$$\begin{aligned} (as(g))c &= ((ag) + a)c \stackrel{(8)}{=} ((ag)c) + (ac) = (a(gc)) + (ac) \\ &\stackrel{(10)}{=} a((gc) + c) \stackrel{(9)}{=} a((cg) + c) = a(cs(g)) \\ &\stackrel{(9)}{=} a(s(g)c) \end{aligned}$$

So $G = \mathbb{N}$. Proving the initial statement is straightforward. Let arbitrary $b \in \mathbb{N} = G$ and the desired conclusion immediately follows. $\square$

(next page)

Proof. (12) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} \forall c \in \mathbb{N} (ac = bc \rightarrow a = b)$$

To do this we define G as the set such that

$$\forall a (a \in G \leftrightarrow a \in \mathbb{N} \wedge \forall b \in \mathbb{N} \forall c \in \mathbb{N} (ac = bc \rightarrow a = b))$$

(where proving existence and uniqueness is straightforward.) We show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward. We need to show $1 \in G$ and $\forall g \in G (s(g) \in G)$.

For the former, we already know $1 \in \mathbb{N}$. Let arbitrary $b, c \in \mathbb{N}$ and suppose that $1 \cdot c = bc$. Suppose for contradiction that $b \neq 1$. Then by lemma 1.3 there exists $x \in \mathbb{N}$ such that $b = s(x)$. We then have

$$c \stackrel{(7)}{=} 1 \cdot c = bc = s(x)c = (x+1)c \stackrel{(8,7)}{=} (xc) + c \stackrel{(4,9)}{=} c + (cx)$$

which contradicts (6).

Now the latter. Let arbitrary $g \in G$. Then $g \in \mathbb{N}$ and $\forall b \in \mathbb{N} \forall c \in \mathbb{N} (gc = bc \rightarrow g = b)$. $s(g) \in \mathbb{N}$ exists. Let arbitrary $b, c \in \mathbb{N}$. Suppose $s(g)c = bc$. For contradiction, suppose $b = 1$. Then

$$c \stackrel{(7)}{=} 1 \cdot c = bc = s(g)c \stackrel{(9)}{=} cs(g) = (cg) + c \stackrel{(4)}{=} c + (cg)$$

contradicting (6). So we must have $b \neq 1$ and by lemma 1.3 there exists $x \in \mathbb{N}$ such that $b = s(x)$. Then

$$(cg) + c = cs(g) \stackrel{(9)}{=} s(g)c = bc = s(x)c = (x+1)c \stackrel{(8,7)}{=} (xc) + c$$

and by (1), $gc = cg = xc$. Since $g \in G$ we then have $g = x$ and $s(g) = s(x) = b$. So $G = \mathbb{N}$. Proving the initial statement is straightforward. Let arbitrary $a \in \mathbb{N} = G$ and the desired conclusion immediately follows. $\square$

Proof. (13) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} (ab = 1 \leftrightarrow a = 1 = b)$$

Let arbitrary $a, b \in \mathbb{N}$. First suppose $ab = 1$. For contradiction suppose $b \neq 1$. Then by lemma 1.3 there exists $x \in \mathbb{N}$ such that $b = s(x)$. So

$$1 = ab = as(x) = a(x+1) \stackrel{(10)}{=} (ax) + (a \cdot 1) = (ax) + a$$

which contradicts (5). So $b = 1$. We immediately have $1 = ab = a \cdot 1 = a$ and $a = 1 = b$. Now suppose $a = 1 = b$. We immediately have $ab = 1 \cdot 1 = 1$. $\square$

1.1.6 Ordering on $\mathbb{N}$

Definition 1.8. The relation $<$ on $\mathbb{N}$ is defined by $a < b$ if and only if there is some $p \in \mathbb{N}$ such that $a + p = b$ for all $a, b \in \mathbb{N}$.

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} (a < b \leftrightarrow \exists p \in \mathbb{N} (a + p = b))$$

The relation $\leq$ on $\mathbb{N}$ is defined by $a \leq b$ if and only if $a < b$ or $a = b$ for all $a, b \in \mathbb{N}$.

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} (a \leq b \leftrightarrow (a < b \vee a = b))$$

We will write $a > b$ to mean the same thing as $b < a$. The same applies to $a \geq b$.

Theorem 1.9. Let $a, b \in \mathbb{N}$. Suppose that $a < b$. Then there is a unique $p \in \mathbb{N}$ such that $a + p = b$.

Proof. By definition there exists $p \in \mathbb{N}$ such that $a + p = b$. Let $p' \in \mathbb{N}$ be arbitrary such that $a + p' = b$. Then by part 4 of theorem 1.7 we have $p + a = b$ and $p' + a = b$, so $p + a = p' + a$. By part 1 of theorem 1.7 we have $p = p'$. $\square$

Theorem 1.10. Let $a, b, c \in \mathbb{N}$.

1. $a \leq a$, and $a \not< a$, and $a < a + 1$
2. $1 \leq a$
3. If $a < b$ and $b < c$, then $a < c$; if $a \leq b$ and $b < c$, then $a < c$; if $a < b$ and $b \leq c$, then $a < c$; if $a \leq b$ and $b \leq c$, then $a \leq c$
4. $a < b$ if and only if $a + c < b + c$
5. Precisely one of $a < b$ or $a = b$ or $a > b$ holds (Trichotomy Law)
6. $a < b$ if and only if $ac < bc$
7. $a \leq b$ or $b \leq a$
8. If $a \leq b$ and $b \leq a$, then $a = b$
9. It cannot be that $b < a < b + 1$
10. $a \leq b$ if and only if $a < b + 1$
11. $a < b$ if and only if $a + 1 \leq b$

(next page)

Proof. (1) We want to show

$$\forall a \in \mathbb{N}(a \leq a \wedge a \not\leq a \wedge a < a + 1)$$

Let arbitrary $a \in \mathbb{N}$. Since $a = a$, by definition $a \leq a$. Next suppose for contradiction that $a < a$. Then there would exist $p \in \mathbb{N}$ such that $a + p = a$, contradicting part 6 of theorem 1.7. Finally since $1 \in \mathbb{N}$ and $a + 1 = a + 1$, by definition $a < a + 1$. $\square$

Proof. (2) We want to show

$$\forall a \in \mathbb{N}(1 \leq a)$$

Let arbitrary $a \in \mathbb{N}$. We have 2 cases. First suppose $a = 1$; then by definition $1 \leq a$. Next suppose $a \neq 1$. Then by lemma 1.3 there exists $x \in \mathbb{N}$ such that $a = s(x) = x + 1$. By theorem 1.7 part 4 we have $a = 1 + x$. So by definition $1 < a$. $\square$

Proof. (3) We want to show

$$\begin{aligned} \forall a \in \mathbb{N} \forall b \in \mathbb{N} \forall c \in \mathbb{N} ([(a < b \wedge b < c) \rightarrow a < c] \wedge [(a \leq b \wedge b < c) \rightarrow a < c] \\ \wedge [(a < b \wedge b \leq c) \rightarrow a < c] \wedge [(a \leq b \wedge b \leq c) \rightarrow a \leq c]) \end{aligned}$$

Let arbitrary $a, b, c \in \mathbb{N}$. First suppose $a < b \wedge b < c$. Then there exists $x, y \in \mathbb{N}$ such that $a + x = b$ and $b + y = c$. We then have $(a + x) + y = b + y = c$. By theorem 1.7 part 2 we have $a + (x + y) = c$ and therefore $a < c$.

Next suppose $a \leq b \wedge b < c$. There exists $y \in \mathbb{N}$ such that $b + y = c$. We have 2 cases. First suppose $a < b$. Then as proven earlier we have $a < c$. Next suppose $a = b$. Then we immediately have $a + y = c$ and $a < c$.

Next suppose $a < b \wedge b \leq c$. There exists $x \in \mathbb{N}$ such that $a + x = b$. We have 2 cases. First suppose $b < c$. Then again we immediately have $a < c$. Next suppose $b = c$. Then $a + x = c$ and $a < c$.

Finally suppose $a \leq b \wedge b \leq c$. We have 4 cases. First suppose $a < b \wedge b < c$. As proven $a < c$. Next suppose $a = b \wedge b < c$. As proven $a < c$. Next suppose $a < b \wedge b = c$. As proven $a < c$. Finally suppose $a = b$ and $b = c$. It immediately follows that $a = c$ and $a \leq c$. $\square$

Proof. (4) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} \forall c \in \mathbb{N}(a < b \leftrightarrow a + c < b + c)$$

Let arbitrary $a, b, c \in \mathbb{N}$. First suppose $a < b$. Then there exists $x \in \mathbb{N}$ such that $a + x = b$. Then $(a + x) + c = b + c$ and by theorem 1.7 parts 2 and 4 we have $(a + c) + x = b + c$. So $a + c < b + c$.

Next suppose $a + c < b + c$. Then there exists $x \in \mathbb{N}$ such that $(a + c) + x = b + c$. By theorem 1.7 parts 2 and 4 we have $(a + x) + c = b + c$, and by part 1 of that same theorem we have $a + x = b$. So $a < b$. $\square$

(next page)

Proof. (5) We want to prove trichotomy

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} ((a < b \vee a = b \vee b < a) \\ \wedge \neg((a < b) \wedge (a = b)) \wedge \neg((a < b) \wedge (b < a)) \wedge \neg((a = b) \wedge (b < a)))$$

Let arbitrary $a, b \in \mathbb{N}$. We start by proving the last three statements. First suppose for contradiction that $a < b$ and $a = b$. Then $a < a$, contradicting (1). Next suppose for contradiction that $a = b$ and $b < a$. A similar argument shows that this contradicts (1). Finally suppose for contradiction that $a < b \wedge b < a$. By (3) we have $a < a$, a contradiction.

Now for the first statement. We define G as the set such that

$$\forall a(a \in G \leftrightarrow a \in \mathbb{N} \wedge \forall b \in \mathbb{N}(a < b \vee a = b \vee b < a))$$

(proving existence and uniqueness of G is straightforward.) We show that $G = \mathbb{N}$. Showing $G \subseteq \mathbb{N}$ is straightforward. We need to show that $1 \in G$ and $\forall g \in G(s(g) \in G)$. For the former, we already know $1 \in \mathbb{N}$. Let arbitrary $b \in \mathbb{N}$. By (2) we have $1 \leq b$, meaning $1 < b$ or $1 = b$. Either case allows us to conclude that $1 \in G$.

Now the latter. Let arbitrary $g \in G$. Then $g \in \mathbb{N}$ and $\forall b \in \mathbb{N}(g < b \vee g = b \vee b < g)$. $s(g) \in \mathbb{N}$ exists. Let arbitrary $b \in \mathbb{N}$. We have 3 cases. First suppose $g < b$. Then there exists $x \in \mathbb{N}$ such that $g + x = b$. Suppose $x = 1$; then $s(g) = g + 1 = b$. Next suppose $x \neq 1$; then by lemma 1.3 there exists $y \in \mathbb{N}$ such that $x = s(y)$. So $b = g + x = g + s(y) = g + (y + 1)$. By theorem 1.7 parts 2 and 4 we have $(g + 1) + y = b$ and $s(g) < b$. For the second case suppose $g = b$. Then by (1) we have $b = g < g + 1 = s(g)$. Finally suppose $b < g$. Then since $g < g + 1$ by (3) we have $b < g + 1 = s(g)$. We then have $G = \mathbb{N}$.

Showing the desired result is straightforward. For all arbitrary $a \in \mathbb{N}$, $a \in G$. The result immediately follows. $\square$

Proof. (6) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} \forall c \in \mathbb{N}(ac < bc \leftrightarrow a < b)$$

Let arbitrary $a, b, c \in \mathbb{N}$. First suppose $a < b$. Then there exists $x \in \mathbb{N}$ such that $a + x = b$. Then by theorem 1.7 part 8 we have $bc = (a + x)c = (ac) + (xc)$; so $ac < bc$.

Next suppose $ac < bc$. Suppose for contradiction that $a \not< b$. Then by (5) we have $a = b \vee b < a$. First suppose $a = b$. Then we immediately have $ac < ac$, which contradicts (1). Next suppose $b < a$. There exists $x, y \in \mathbb{N}$ such that $(ac) + x = bc$ and $b + y = a$. By theorem 1.7 parts 8 and 2 we have $bc = ((b + y)c) + x = ((bc) + (yc)) + x = (bc) + ((yc) + x)$, contradicting part 6 of that same theorem. $\square$

(next page)

Proof. (7) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} (a \leq b \vee b \leq a)$$

Let arbitrary $a, b \in \mathbb{N}$. First suppose $a \not\leq b$. Then by (5) $a = b \vee b < a$, meaning $b \leq a$. Next suppose $a < b$. Then immediately it follows that $a \leq b$. $\square$

Proof. (8) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} ((a \leq b \wedge b \leq a) \rightarrow a = b)$$

Let arbitrary $a, b \in \mathbb{N}$. Suppose that $a \leq b \wedge b \leq a$. We have four cases. Three of the cases: $a < b \wedge b < a$, $a = b \wedge b < a$, and $a < b \wedge b = a$ all contradict (5). The only possibility is the last case $a = b \wedge b = a$. $\square$

Proof. (9) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} (\neg(b < a \wedge a < b + 1))$$

Let arbitrary $a, b \in \mathbb{N}$. Suppose for contradiction that $b < a \wedge a < b + 1$. Then there exists $x, y \in \mathbb{N}$ such that $b + x = a$ and $a + y = b + 1$. Then $(b + x) + y = b + 1$. By theorem 1.7 parts 2 and 4 we have $(x + y) + b = 1 + b$. By part 1 of that theorem we then have $x + y = 1$, contradicting part 5 of that theorem. $\square$

Proof. (10) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} (a \leq b \leftrightarrow a < b + 1)$$

Let arbitrary $a, b \in \mathbb{N}$. First suppose $a \leq b$. Suppose for contradiction that $b + 1 \leq a$. Then by (3) we would have $b + 1 \leq b$. The case where $b + 1 = b$ is a contradiction to theorem 1.7 part 6. The case where $b + 1 < b$ implies the existence of $x \in \mathbb{N}$ such that $(b + 1) + x = b$; by theorem 1.7 part 2 we have $b + (1 + x) = b$, again contradicting part 6 of that theorem. We can then conclude that $\neg(b + 1 \leq a)$, meaning $\neg(b + 1 = a \vee b + 1 < a)$. By (5) we must then have $a < b + 1$.

Now for the other direction. Suppose $a < b + 1$. Supposing that $b < a$ immediately contradicts (9). Therefore by (5) we must have $a \leq b$. $\square$

Proof. (11) We want to show

$$\forall a \in \mathbb{N} \forall b \in \mathbb{N} (a < b \leftrightarrow a + 1 \leq b)$$

Let arbitrary $a, b \in \mathbb{N}$. Suppose $a < b$. Supposing $b < a + 1$ immediately contradicts (9), so we must have, by (5), $a + 1 \leq b$.

Next suppose $a + 1 \leq b$. Suppose for contradiction that $b \leq a$, which would imply by (3) that $a + 1 \leq a$. The case where $a + 1 = a$ immediately contradicts theorem 1.7 part 6. The case where $a + 1 < a$ implies the existence of $x \in \mathbb{N}$ such that $(a + 1) + x = a$; theorem 1.7 part 2 then implies $a + (1 + x) = a$, again contradicting part 6 of that same theorem. We conclude that $\neg(b = a \vee b < a)$, so by (5) we have $a < b$. $\square$

1.1.7 Well-ordering

Theorem 1.11. (*Well-Ordering Principle*) Let $G \subseteq \mathbb{N}$ be a non-empty set. Then there is some $m \in G$ such that $m \leq g$ for all $g \in G$.

Proof. We want to show

$$\forall G(G \subseteq \mathbb{N} \wedge G \neq \emptyset \rightarrow \exists m \in G(\forall g \in G(m \leq g)))$$

Let G be arbitrary; assume that $G \subseteq \mathbb{N}$ and $G \neq \emptyset$. Assume for contradiction that the implication is false. That is

$$\neg \exists m \in G(\forall g \in G(m \leq g))$$

We define H as the set such that

$$\forall a(a \in H \leftrightarrow a \in \mathbb{N} \wedge \forall n \in \mathbb{N}(n \leq a \rightarrow n \notin G))$$

(where existence and uniqueness can be easily shown). We can show that $H \cap G = \emptyset$. For contradiction suppose there exists x such that $x \in H$ and $x \in G$. By definition then $x \in \mathbb{N}$ and $\forall n \in \mathbb{N}(n \leq x \rightarrow n \notin G)$. Then since $x \leq x$ we must have $x \notin G$, a contradiction. We will show that $H = \mathbb{N}$. Showing $H \subseteq \mathbb{N}$ is straightforward. We need to show $1 \in H$ and $\forall h \in H(s(h) \in H)$.

We start with the former. Suppose for contradiction that $1 \notin H$. We know $1 \in \mathbb{N}$, so we must have $\neg \forall n \in \mathbb{N}(n \leq 1 \rightarrow n \notin G)$. This is logically equivalent to

$$\exists n \in \mathbb{N}(n \leq 1 \wedge n \in G)$$

That is, there exists $n \in \mathbb{N}$ such that $n \leq 1$ and $n \in G$. $n < 1$ is a contradiction as it would imply the existence of $x \in \mathbb{N}$ such that $n + x = 1$, contradicting theorem 1.7 part 5. We therefore must have $1 = n \in G$. Let arbitrary $g \in G \subseteq \mathbb{N}$. By theorem 1.10 part 2 we must have $n = 1 \leq g$. This contradicts our assumption that such an element does not exist, so $1 \in H$.

Now the latter. Let arbitrary $h \in H$. Then $h \in \mathbb{N}$ and $\forall n \in \mathbb{N}(n \leq h \rightarrow n \notin G)$. We know $s(h) \in \mathbb{N}$ exists. Suppose for contradiction that $s(h) \notin H$. Then as before there then exists $n \in \mathbb{N}$ such that $n \leq s(h)$ and $n \in G$. By the contrapositive of one of the consequences of $h \in H$, since $n \in G$ we then have $\neg(n \leq h)$, which by theorem 1.10 part 5 implies $h < n$. Since $n < s(h) = h + 1$ would then lead to a contradiction of theorem 1.10 part 9, we must have $n = s(h)$. Let arbitrary $g \in G$. Suppose for contradiction that $g < n = s(h) = h + 1$. Then by theorem 1.10 part 10 we have $g \leq h$. Since $h \in H$ we then have $g \notin G$, a contradiction. So we must have $\neg(g < n)$, which by theorem 1.10 part 5 implies $n \leq g$. We know that $n \in G$, so this contradicts our earlier assumption that such an element must not exist. So $s(h) \in H$ and $H = \mathbb{N}$.

We know $H \cap G = \emptyset$. So $\mathbb{N} \cap G = \emptyset$. Suppose for contradiction that G is nonempty, meaning that there exists some $x \in G$. Then since $G \subseteq \mathbb{N}$ we would have $x \in \mathbb{N}$ and therefore $x \in \mathbb{N} \cap G = \emptyset$, a contradiction. So we must have $G = \emptyset$, contradicting a key assumption. $\square$

1.2 Construction of the integers

1.2.1 Construction

Definition 1.12. The relation $\sim$ on $\mathbb{N} \times \mathbb{N}$ is defined by $\langle a, b \rangle \sim \langle c, d \rangle$ if and only if $a + d = b + c$ for all $\langle a, b \rangle, \langle c, d \rangle \in \mathbb{N} \times \mathbb{N}$. That is, the set such that

$$\forall x (x \in \sim \leftrightarrow \exists \langle a, b \rangle \in \mathbb{N} \times \mathbb{N} \exists \langle c, d \rangle \in \mathbb{N} \times \mathbb{N} (x = \langle \langle a, b \rangle, \langle c, d \rangle \rangle \wedge a + d = b + c))$$

(Existence follows from using $(\mathbb{N} \times \mathbb{N}) \times (\mathbb{N} \times \mathbb{N})$ as a bounding set. Uniqueness from extensionality)

Lemma 1.13. The relation $\sim$ is an equivalence relation on $\mathbb{N} \times \mathbb{N}$.

Proof. Showing that $\sim \subseteq (\mathbb{N} \times \mathbb{N}) \times (\mathbb{N} \times \mathbb{N})$ is straightforward. We need to show reflexivity on $\mathbb{N} \times \mathbb{N}$, symmetry, and transitivity. We start with reflexivity; let arbitrary $\langle a, b \rangle \in \mathbb{N} \times \mathbb{N}$. We immediately have $a + b = b + a$ and $\langle a, b \rangle \sim \langle a, b \rangle$. For symmetry, let $\langle a, b \rangle, \langle c, d \rangle$ be arbitrary and suppose $\langle a, b \rangle \sim \langle c, d \rangle$. Then by definition both pairs are members of $\mathbb{N} \times \mathbb{N}$ and $a + d = b + c$. We then have $c + b = b + c = a + d = d + a$, which then means $\langle c, d \rangle \sim \langle a, b \rangle$.

Finally transitivity. Let $\langle a, b \rangle, \langle c, d \rangle, \langle e, f \rangle$ be arbitrary, suppose $\langle a, b \rangle \sim \langle c, d \rangle$ and $\langle c, d \rangle \sim \langle e, f \rangle$. This means that all three pairs are members of $\mathbb{N} \times \mathbb{N}$, $a + d = c + b$, and $c + f = e + d$. We then have

$$\begin{aligned} (a + f) + c &= a + (c + f) = a + (e + d) = (a + d) + e \\ &= (c + b) + e = (b + e) + c \end{aligned}$$

By cancellation we then have $a + f = b + e$. So $\langle a, b \rangle \sim \langle e, f \rangle$. $\square$

Definition 1.14. The set of *integers*, denoted $\mathbb{Z}$, is the quotient set $(\mathbb{N} \times \mathbb{N}) / \sim$. That is,

$$\forall x (x \in \mathbb{Z} \leftrightarrow \exists \langle a, b \rangle \in \mathbb{N} \times \mathbb{N} (x = [\langle a, b \rangle]))$$

where

$$\forall \langle a, b \rangle \in \mathbb{N} \times \mathbb{N} \forall x (x \in [\langle a, b \rangle] \leftrightarrow \langle a, b \rangle \sim x)$$

Each integer is an equivalence class. Existence of each equivalence class can be shown using $\text{ran}(\sim)$ as a bounding set. The same can be done for $\mathbb{Z}$ using $\mathcal{P}(\text{ran}(\sim))$. Uniqueness for both follows from extensionality.

Definition 1.15. We define $\hat{0}, \hat{1}$ as $\hat{0} = [\langle 1, 1 \rangle]$ and $\hat{1} = [\langle 1 + 1, 1 \rangle]$.

1.2.2 Operations on the integers

Definition 1.16. The binary operation $+$ on $\mathbb{Z}$ is defined as $+: \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$ where

$$\forall x(x \in + \leftrightarrow \exists \langle a, b \rangle \in \mathbb{N} \times \mathbb{N} \exists \langle c, d \rangle \in \mathbb{N} \times \mathbb{N} (x = \langle \langle [a, b], [c, d] \rangle, [a + c, b + d] \rangle))$$

put simply, for all $[a, b], [c, d] \in \mathbb{Z}$,

$$[\langle a, b \rangle] + [\langle c, d \rangle] = [\langle a + c, b + d \rangle]$$

(A bounding set is $(\mathbb{Z} \times \mathbb{Z}) \times \mathbb{Z}$. Uniqueness follows from extensionality.)

Proof. We start by showing the domain bound $\text{dom}(+) = \mathbb{Z} \times \mathbb{Z}$. First suppose $x \in \text{dom}(+)$. Then there exists y such that $\langle x, y \rangle \in +$ and $\langle a, b \rangle, \langle c, d \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x = \langle \langle [a, b], [c, d] \rangle \in \mathbb{Z} \times \mathbb{Z}$. Next suppose $x \in \mathbb{Z} \times \mathbb{Z}$. Then there exists $x_0, x_1 \in \mathbb{Z}$ such that $x = \langle x_0, x_1 \rangle$. There exists $\langle a, b \rangle, \langle c, d \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x_0 = [a, b]$ and $x_1 = [c, d]$. We know $\langle a + c, b + d \rangle \in \mathbb{N} \times \mathbb{N}$ and $\langle \langle [a, b], [c, d] \rangle, [a + c, b + d] \rangle \in +$, so $x = \langle x_0, x_1 \rangle = \langle \langle [a, b], [c, d] \rangle \in \text{dom}(+)$.

Now the range bound $\text{ran}(+) \subseteq \mathbb{Z}$. Let arbitrary $y \in \text{ran}(+)$. Then there exists x such that $\langle x, y \rangle \in +$ and $a, b, c, d \in \mathbb{N}$ such that $y = [a + c, b + d] \in \mathbb{Z}$.

Now we prove well definedness. That is, that addition between equivalence classes is independent of the particular elements used to represent each equivalence class:

$$\begin{aligned} \forall \langle a, b \rangle \in \mathbb{N} \times \mathbb{N} \forall \langle c, d \rangle \in \mathbb{N} \times \mathbb{N} \forall \langle a', b' \rangle \in \mathbb{N} \times \mathbb{N} \forall \langle c', d' \rangle \in \mathbb{N} \times \mathbb{N} \\ [\langle a, b \rangle] = [\langle a', b' \rangle] \wedge [\langle c, d \rangle] = [\langle c', d' \rangle] \\ \rightarrow [\langle a + c, b + d \rangle] = [\langle a' + c', b' + d' \rangle] \end{aligned}$$

Let arbitrary $\langle a, b \rangle, \langle c, d \rangle, \langle a', b' \rangle, \langle c', d' \rangle \in \mathbb{N} \times \mathbb{N}$ and suppose that $[\langle a, b \rangle] = [\langle a', b' \rangle]$ and $[\langle c, d \rangle] = [\langle c', d' \rangle]$. Then $\langle a, b \rangle \sim \langle a', b' \rangle$ and $a + b' = b + a'$. Similarly $c + d' = d + c'$. We then have, by associativity and commutativity of natural arithmetic:

$$\begin{aligned} (a + c) + (b' + d') &= (a + b') + (c + d') \\ &= (b + a') + (d + c') = (b + d) + (a' + c') \end{aligned}$$

and we can conclude that $\langle a + c, b + d \rangle \sim \langle a' + c', b' + d' \rangle$ and $[\langle a + c, b + d \rangle] = [\langle a' + c', b' + d' \rangle]$.

Finally we show functionality. Let arbitrary $x \in \text{dom}(+)$. Then by definition there exists y such that $\langle x, y \rangle \in +$. There then exists $\langle a, b \rangle, \langle c, d \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x = \langle \langle [a, b], [c, d] \rangle$ and $y = [a + c, b + d]$. Let y' be arbitrary such that $\langle x, y' \rangle \in +$. Then similarly there exists $\langle a', b' \rangle, \langle c', d' \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x = \langle \langle [a', b'], [c', d'] \rangle$ and $y' = [a' + c', b' + d']$. We know $\langle \langle [a, b], [c, d] \rangle = \langle \langle [a', b'], [c', d'] \rangle$. So since addition is well defined we have $y = [a + c, b + d] = [a' + c', b' + d'] = y'$. $\square$

(next page)

Definition 1.17. The binary operation $\cdot$ on $\mathbb{Z}$ is defined as $\cdot : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$ where

$$\forall x(x \in \cdot \leftrightarrow \exists \langle a, b \rangle \in \mathbb{N} \times \mathbb{N} \exists \langle c, d \rangle \in \mathbb{N} \times \mathbb{N} (\\ x = \langle \langle [a, b], [c, d] \rangle, [ac + bd, ad + bc] \rangle))$$

(A bounding set is $(\mathbb{Z} \times \mathbb{Z}) \times \mathbb{Z}$. Uniqueness follows from extensionality.)

Proof. We start with the domain bound $\text{dom}(\cdot) = \mathbb{Z} \times \mathbb{Z}$. First suppose $x \in \text{dom}(\cdot)$. Then there exists y such that $\langle x, y \rangle \in \cdot$, and $\langle a, b \rangle, \langle c, d \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x = \langle \langle [a, b], [c, d] \rangle, [ac + bd, ad + bc] \rangle \in \mathbb{Z} \times \mathbb{Z}$. Next suppose $x \in \mathbb{Z} \times \mathbb{Z}$. Then there exists $x_0, x_1 \in \mathbb{Z}$ such that $x = \langle x_0, x_1 \rangle$. There exists $\langle a, b \rangle, \langle c, d \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x_0 = [a, b]$ and $x_1 = [c, d]$. We have $\langle \langle [a, b], [c, d] \rangle, [ac + bd, ad + bc] \rangle \in \cdot$. So $x = \langle x_0, x_1 \rangle = \langle \langle [a, b], [c, d] \rangle, [ac + bd, ad + bc] \rangle \in \text{dom}(\cdot)$.

Next the range bound $\text{ran}(\cdot) \subseteq \mathbb{Z}$. Let arbitrary $y \in \text{ran}(\cdot)$. Then there exists x such that $\langle x, y \rangle \in \cdot$ and $a, b, c, d \in \mathbb{N}$ such that $y = [ac + bd, ad + bc] \in \mathbb{Z}$.

Next we show well definedness. That is

$$\begin{aligned} \forall \langle a, b \rangle \in \mathbb{N} \times \mathbb{N} \forall \langle c, d \rangle \in \mathbb{N} \times \mathbb{N} \forall \langle a', b' \rangle \in \mathbb{N} \times \mathbb{N} \forall \langle c', d' \rangle \in \mathbb{N} \times \mathbb{N} (\\ ([a, b] = [a', b'] \wedge [c, d] = [c', d']) \\ \rightarrow [ac + bd, ad + bc] = [a'c' + b'd', a'd' + b'c']) \end{aligned}$$

Let arbitrary $\langle a, b \rangle, \langle c, d \rangle, \langle a', b' \rangle, \langle c', d' \rangle \in \mathbb{N} \times \mathbb{N}$ and suppose $[a, b] = [a', b']$ and $[c, d] = [c', d']$. Then we have $\langle a, b \rangle \sim \langle a', b' \rangle$ and $a + b' = b + a'$. Similarly we have $c + d' = d + c'$. We want to show

$$(ac + bd) + (a'd' + b'c') = (ad + bc) + (a'c' + b'd')$$

See that

$$\begin{aligned} & ((a + b')d + (b + a')c) + ((c + d')b' + (d + c')a') \\ &= ((ad + b'd) + (bc + a'c)) + ((cb' + d'b') + (da' + c'a')) \\ &= ((ad + bc) + (b'd + a'c)) + ((a'c' + b'd') + (da' + cb')) \\ &= ((ad + bc) + (a'c' + b'd')) + ((b'd + a'c) + (da' + cb')) \end{aligned}$$

Also see that

$$\begin{aligned} & ((a + b')d + (b + a')c) + ((c + d')b' + (d + c')a') \\ &= ((b + a')d + (a + b')c) + ((d + c')b' + (c + d')a') \\ &= ((bd + a'd) + (ac + b'c)) + ((db' + c'b') + (ca' + d'a')) \\ &= ((ac + bd) + (da' + cb')) + ((a'd' + c'b') + (a'c + b'd)) \\ &= ((ac + bd) + (a'd' + c'b')) + ((b'd + a'c) + (da' + cb')) \end{aligned}$$

By cancellation we then have our desired result, which means

$$\langle ac + bd, ad + bc \rangle \sim \langle a'c' + b'd', a'd' + b'c' \rangle$$

and $[ac + bd, ad + bc] = [a'c' + b'd', a'd' + b'c']$.

(next page)

Finally functionality. Let arbitrary $x \in \text{dom}(\cdot)$. Then there exists y such that $\langle x, y \rangle \in \cdot$ and $\langle a, b \rangle, \langle c, d \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x = \langle [\langle a, b \rangle], [\langle c, d \rangle] \rangle$ and $y = [\langle ac + bd, ad + bc \rangle]$. Let y' be arbitrary such that $\langle x, y' \rangle \in \cdot$. Then similarly there exists $\langle a', b' \rangle, \langle c', d' \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x = \langle [\langle a', b' \rangle], [\langle c', d' \rangle] \rangle$ and $y' = [\langle a'c' + b'd', a'd' + b'c' \rangle]$. We have $\langle [\langle a, b \rangle], [\langle c, d \rangle] \rangle = \langle [\langle a', b' \rangle], [\langle c', d' \rangle] \rangle$, by well definedness we then have $y = [\langle ac + bd, ad + bc \rangle] = [\langle a'c' + b'd', a'd' + b'c' \rangle] = y'$. $\square$

Definition 1.18. The unary operation $-$ on $\mathbb{Z}$ is defined as $- : \mathbb{Z} \rightarrow \mathbb{Z}$ where

$$\forall x (x \in - \leftrightarrow \exists \langle a, b \rangle \in \mathbb{N} \times \mathbb{N} (x = \langle [\langle a, b \rangle], [\langle b, a \rangle] \rangle))$$

(A bounding set is $\mathbb{Z} \times \mathbb{Z}$. Uniqueness follows from extensionality.)

Proof. We start with the domain bound $\text{dom}(-) = \mathbb{Z}$. First let arbitrary $x \in \text{dom}(-)$. Then there exists y such that $\langle x, y \rangle \in -$, and $\langle a, b \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x = [\langle a, b \rangle] \in \mathbb{Z}$. Next suppose $x \in \mathbb{Z}$. Then there exists $\langle a, b \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x = [\langle a, b \rangle]$. We have $\langle [\langle a, b \rangle], [\langle b, a \rangle] \rangle \in -$. So $x = [\langle a, b \rangle] \in \text{dom}(-)$.

Next the range bound $\text{ran}(-) \subseteq \mathbb{Z}$. Suppose $y \in \text{ran}(-)$. Then there exists x such that $\langle x, y \rangle \in -$ and $a, b \in \mathbb{N}$ such that $y = [\langle b, a \rangle] \in \mathbb{Z}$.

Next we show well-definedness. That is

$$\forall \langle a, b \rangle \in \mathbb{N} \times \mathbb{N} \forall \langle a', b' \rangle \in \mathbb{N} \times \mathbb{N} ([\langle a, b \rangle] = [\langle a', b' \rangle] \rightarrow [\langle b, a \rangle] = [\langle b', a' \rangle])$$

Letting arbitrary $\langle a, b \rangle, \langle a', b' \rangle \in \mathbb{N} \times \mathbb{N}$ and supposing that $[\langle a, b \rangle] = [\langle a', b' \rangle]$, we have $a + b' = b + a'$. See that this also means $\langle b, a \rangle \sim \langle b', a' \rangle$ and therefore $[\langle b, a \rangle] = [\langle b', a' \rangle]$.

Finally functionality. Let arbitrary $x \in \text{dom}(-)$. Then there exists y such that $\langle x, y \rangle \in -$, and $\langle a, b \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x = [\langle a, b \rangle]$ and $y = [\langle b, a \rangle]$. Let y' be arbitrary such that $\langle x, y' \rangle \in -$. Then there exists $\langle a', b' \rangle \in \mathbb{N} \times \mathbb{N}$ such that $x = [\langle a', b' \rangle]$ and $y' = [\langle b', a' \rangle]$. We have $[\langle a, b \rangle] = [\langle a', b' \rangle]$; by well-definedness we have $y = [\langle b, a \rangle] = [\langle b', a' \rangle] = y'$. $\square$

(next page)

Definition 1.19.