

Mearurable circle squaring

Oleg Pikhurko
University of Warwick

Joint with Łukasz Grabowski and András Máthé

DIAMANT Symposium, 25 November 2016

Equidecomposability

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) :=
 $\exists A = A_1 \sqcup \cdots \sqcup A_n, B = B_1 \sqcup \cdots \sqcup B_n$, isometries
 $\gamma_1, \dots, \gamma_n$ s.t. $\forall i A_i = \gamma_i(B_i)$

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) :=
 $\exists A = A_1 \sqcup \dots \sqcup A_n, B = B_1 \sqcup \dots \sqcup B_n$, isometries
 $\gamma_1, \dots, \gamma_n$ s.t. $\forall i A_i = \gamma_i(B_i)$
- ▶ **Example:** triangle and rectangle

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) :=
 $\exists A = A_1 \sqcup \dots \sqcup A_n, B = B_1 \sqcup \dots \sqcup B_n$, isometries
 $\gamma_1, \dots, \gamma_n$ s.t. $\forall i A_i = \gamma_i(B_i)$
- ▶ **Example:** triangle and rectangle
- ▶ **Wallace 1807, Bolyai 1832, Gerwien 1832:**
equi-area polygons \Rightarrow dissection congruent

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) :=
 $\exists A = A_1 \sqcup \dots \sqcup A_n, B = B_1 \sqcup \dots \sqcup B_n$, isometries
 $\gamma_1, \dots, \gamma_n$ s.t. $\forall i A_i = \gamma_i(B_i)$
- ▶ **Example:** triangle and rectangle
- ▶ **Wallace 1807, Bolyai 1832, Gerwien 1832:**
equi-area polygons \Rightarrow dissection congruent
 - ▶ **Tarski'24:** equidecomposable

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) :=
 $\exists A = A_1 \sqcup \dots \sqcup A_n, B = B_1 \sqcup \dots \sqcup B_n$, isometries
 $\gamma_1, \dots, \gamma_n$ s.t. $\forall i A_i = \gamma_i(B_i)$
- ▶ **Example:** triangle and rectangle
- ▶ **Wallace 1807, Bolyai 1832, Gerwien 1832:**
equi-area polygons \Rightarrow dissection congruent
 - ▶ **Tarski'24:** equidecomposable
 - ▶ **Exercise:** $[0, 1] \sim [0, 1)$

Banach-Tarski paradox ($k \geq 3$)

Banach-Tarski paradox ($k \geq 3$)

- ▶ **Banach-Tarski'24:** $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$,

Banach-Tarski paradox ($k \geq 3$)

- ▶ **Banach-Tarski'24:** $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$,
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball

Banach-Tarski paradox ($k \geq 3$)

- ▶ **Banach-Tarski'24:** $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$,
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff'14:** paradoxes on sphere \mathbb{S}^{k-1}

Banach-Tarski paradox ($k \geq 3$)

- ▶ **Banach-Tarski'24:** $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$,
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff'14:** paradoxes on sphere \mathbb{S}^{k-1}
 - ▶ $SO(k) \supseteq$ free group $\mathbb{F}_2 = \langle a, b \rangle$

Banach-Tarski paradox ($k \geq 3$)

- ▶ **Banach-Tarski'24:** $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$,
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff'14:** paradoxes on sphere \mathbb{S}^{k-1}
 - ▶ $SO(k) \supseteq$ free group $\mathbb{F}_2 = \langle a, b \rangle$
 - ▶ $S_a := \{\text{irreducible words starting with } a\}$, etc

Banach-Tarski paradox ($k \geq 3$)

- ▶ **Banach-Tarski'24:** $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$,
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff'14:** paradoxes on sphere \mathbb{S}^{k-1}
 - ▶ $SO(k) \supseteq$ free group $\mathbb{F}_2 = \langle a, b \rangle$
 - ▶ $S_a := \{\text{irreducible words starting with } a\}$, etc
 - ▶ $a^{-1}.S_a = S_a \cup S_b \cup S_{b^{-1}} \cup \{\emptyset\}$

Banach-Tarski paradox ($k \geq 3$)

- ▶ **Banach-Tarski'24:** $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$,
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff'14:** paradoxes on sphere \mathbb{S}^{k-1}
 - ▶ $SO(k) \supseteq$ free group $\mathbb{F}_2 = \langle a, b \rangle$
 - ▶ $S_a := \{\text{irreducible words starting with } a\}$, etc
 - ▶ $a^{-1} \cdot S_a = S_a \cup S_b \cup S_{b^{-1}} \cup \{\emptyset\}$
- ▶ \Rightarrow Non non-trivial finitely-additive isometry-invariant mean on all bounded subsets of \mathbb{R}^k (or \mathbb{S}^{k-1})

Banach-Tarski paradox ($k \geq 3$)

- ▶ **Banach-Tarski'24:** $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$,
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff'14:** paradoxes on sphere \mathbb{S}^{k-1}
 - ▶ $SO(k) \supseteq$ free group $\mathbb{F}_2 = \langle a, b \rangle$
 - ▶ $S_a := \{\text{irreducible words starting with } a\}$, etc
 - ▶ $a^{-1} \cdot S_a = S_a \cup S_b \cup S_{b^{-1}} \cup \{\emptyset\}$
- ▶ \Rightarrow Non non-trivial finitely-additive isometry-invariant mean on all bounded subsets of \mathbb{R}^k (or \mathbb{S}^{k-1})
- ▶ **Tomkowicz-Wagon'16:** *"The Banach-Tarski Paradox"*

\mathbb{R}^k for $k \leq 2$

\mathbb{R}^k for $k \leq 2$

- ▶ **Banach'23:** Lebesgue measure can be extended to isometry-invariant finitely-additive mean on $2^{\mathbb{R}^k}$

\mathbb{R}^k for $k \leq 2$

- ▶ **Banach'23:** Lebesgue measure can be extended to isometry-invariant finitely-additive mean on $2^{\mathbb{R}^k}$
 - ▶ Lebesgue measurable $A \sim B \Rightarrow \lambda(A) = \lambda(B)$

\mathbb{R}^k for $k \leq 2$

- ▶ **Banach'23:** Lebesgue measure can be extended to isometry-invariant finitely-additive mean on $2^{\mathbb{R}^k}$
 - ▶ Lebesgue measurable $A \sim B \Rightarrow \lambda(A) = \lambda(B)$
 - ▶ \mathbb{B}^k cannot be doubled

\mathbb{R}^k for $k \leq 2$

- ▶ **Banach'23:** Lebesgue measure can be extended to isometry-invariant finitely-additive mean on $2^{\mathbb{R}^k}$
 - ▶ Lebesgue measurable $A \sim B \Rightarrow \lambda(A) = \lambda(B)$
 - ▶ \mathbb{B}^k cannot be doubled
- ▶ **von Neumann'28:** amenable groups

\mathbb{R}^k for $k \leq 2$

- ▶ **Banach'23:** Lebesgue measure can be extended to isometry-invariant finitely-additive mean on $2^{\mathbb{R}^k}$
 - ▶ Lebesgue measurable $A \sim B \Rightarrow \lambda(A) = \lambda(B)$
 - ▶ \mathbb{B}^k cannot be doubled
- ▶ **von Neumann'28:** amenable groups
 - ▶ $\text{Iso}(\mathbb{R}^k)$ is amenable $\Leftrightarrow k \leq 2$

\mathbb{R}^k for $k \leq 2$

- ▶ **Banach'23**: Lebesgue measure can be extended to isometry-invariant finitely-additive mean on $2^{\mathbb{R}^k}$
 - ▶ Lebesgue measurable $A \sim B \Rightarrow \lambda(A) = \lambda(B)$
 - ▶ \mathbb{B}^k cannot be doubled
- ▶ **von Neumann'28**: amenable groups
 - ▶ $\text{Iso}(\mathbb{R}^k)$ is amenable $\Leftrightarrow k \leq 2$
- ▶ Γ is **paradoxical** $:= \exists$ disjoint $A, B \subseteq \Gamma$ with $A, B \sim \Gamma$
(for $\Gamma \curvearrowright \Gamma$)

\mathbb{R}^k for $k \leq 2$

- ▶ **Banach'23**: Lebesgue measure can be extended to isometry-invariant finitely-additive mean on $2^{\mathbb{R}^k}$
 - ▶ Lebesgue measurable $A \sim B \Rightarrow \lambda(A) = \lambda(B)$
 - ▶ \mathbb{B}^k cannot be doubled
- ▶ **von Neumann'28**: amenable groups
 - ▶ $\text{Iso}(\mathbb{R}^k)$ is amenable $\Leftrightarrow k \leq 2$
- ▶ Γ is **paradoxical** $:= \exists$ disjoint $A, B \subseteq \Gamma$ with $A, B \sim \Gamma$ (for $\Gamma \curvearrowright \Gamma$)
- ▶ **Tarski'38**: no invariant mean \Leftrightarrow paradoxical

Tarski's circle squaring

Tarski's circle squaring

- ▶ Tarski'25: Is \mathbb{B}^2 equidecomposable to a square?

Tarski's circle squaring

- ▶ Tarski'25: Is \mathbb{B}^2 equidecomposable to a square?
- ▶ von Neumann'29: Yes, with affine transformations

Tarski's circle squaring

- ▶ Tarski'25: Is \mathbb{B}^2 equidecomposable to a square?
- ▶ von Neumann'29: Yes, with affine transformations
- ▶ Dubins-Hirsh-Karush'63: No, with topological disks

Tarski's circle squaring

- ▶ Tarski'25: Is \mathbb{B}^2 equidecomposable to a square?
- ▶ von Neumann'29: Yes, with affine transformations
- ▶ Dubins-Hirsh-Karush'63: No, with topological disks
- ▶ Gardner'85: No, with a discrete subgroup of $\text{Iso}(\mathbb{R}^2)$

Tarski's circle squaring

- ▶ Tarski'25: Is \mathbb{B}^2 equidecomposable to a square?
- ▶ von Neumann'29: Yes, with affine transformations
- ▶ Dubins-Hirsh-Karush'63: No, with topological disks
- ▶ Gardner'85: No, with a discrete subgroup of $\text{Iso}(\mathbb{R}^2)$
- ▶ Laczkovich'90: YES!

General result by Laczkovich

General result by Laczkovich

- ▶ Box (or upper Minkowski's) dimension

$$\dim_{\square}(A) := \lim_{\varepsilon \rightarrow 0} \varepsilon \log(\# \varepsilon\text{-grid boxes intersecting } A)$$

General result by Laczkovich

- ▶ Box (or upper Minkowski's) dimension

$$\dim_{\square}(A) := \lim_{\varepsilon \rightarrow 0} \varepsilon \log(\# \varepsilon\text{-grid boxes intersecting } A)$$

- ▶ $A \overset{\text{Tr}}{\sim} B$:= pieces are translated: $A_i = B_i + \mathbf{v}_i$

General result by Laczkovich

- ▶ **Box (or upper Minkowski's) dimension**

$$\dim_{\square}(A) := \lim_{\varepsilon \rightarrow 0} \varepsilon \log(\# \varepsilon\text{-grid boxes intersecting } A)$$

- ▶ $A \overset{\text{Tr}}{\sim} B$:= pieces are translated: $A_i = B_i + \mathbf{v}_i$
- ▶ **Laczkovich'90,92:** $A, B \subseteq \mathbb{R}^k$, $\lambda(A) = \lambda(B) > 0$,
bounded, $\dim_{\square}(\partial A), \dim_{\square}(\partial B) < k \Rightarrow A \overset{\text{Tr}}{\sim} B$

“Nice” pieces?

“Nice” pieces?

- ▶ Above results crucially use the Axiom of Choice

“Nice” pieces?

- ▶ Above results crucially use the Axiom of Choice
 - ▶ Pieces are non-measurable

“Nice” pieces?

- ▶ Above results crucially use the Axiom of Choice
 - ▶ Pieces are non-measurable
- ▶ **Borel** := σ -algebra generated by open sets

“Nice” pieces?

- ▶ Above results crucially use the Axiom of Choice
 - ▶ Pieces are non-measurable
- ▶ **Borel** := σ -algebra generated by open sets
- ▶ **Baire** := Borel \triangle meager

“Nice” pieces?

- ▶ Above results crucially use the Axiom of Choice
 - ▶ Pieces are non-measurable
- ▶ **Borel** := σ -algebra generated by open sets
- ▶ **Baire** := Borel \triangle meager
- ▶ **Lebesgue (measurable)** := Borel \triangle null-set

“Nice” pieces?

- ▶ Above results crucially use the Axiom of Choice
 - ▶ Pieces are non-measurable
- ▶ **Borel** := σ -algebra generated by open sets
- ▶ **Baire** := Borel \triangle meager
- ▶ **Lebesgue (measurable)** := Borel \triangle null-set
- ▶ **Dougherty-Foreman'92**: Banach-Tarski with Baire pieces

“Nice” pieces?

- ▶ Above results crucially use the Axiom of Choice
 - ▶ Pieces are non-measurable
- ▶ **Borel** := σ -algebra generated by open sets
- ▶ **Baire** := Borel \triangle meager
- ▶ **Lebesgue (measurable)** := Borel \triangle null-set
- ▶ **Dougherty-Foreman'92**: Banach-Tarski with Baire pieces
 - ▶ Solved **Ruziewicz Problem (1930)**

“Nice” pieces?

- ▶ Above results crucially use the Axiom of Choice
 - ▶ Pieces are non-measurable
- ▶ **Borel** := σ -algebra generated by open sets
- ▶ **Baire** := Borel \triangle meager
- ▶ **Lebesgue (measurable)** := Borel \triangle null-set
- ▶ **Dougherty-Foreman'92**: Banach-Tarski with Baire pieces
 - ▶ Solved **Ruziewicz Problem (1930)**
 - ▶ **Marks-Unger'16**: elegant proof

Measurable equidecompositions

Measurable equidecompositions

- ▶ $\mathcal{L} := \{\text{Lebesgue sets}\}$

Measurable equidecompositions

- ▶ $\mathcal{L} := \{\text{Lebesgue sets}\} = \{\text{Borel} \triangle \text{null-set}\}$

Measurable equidecompositions

- ▶ $\mathcal{L} := \{\text{Lebesgue sets}\} = \{\text{Borel} \triangle \text{null-set}\}$
- ▶ $A \sim_{\mathcal{L}} B := \text{equidecomposable with Lebesgue pieces}$

Measurable equidecompositions

- ▶ $\mathcal{L} := \{\text{Lebesgue sets}\} = \{\text{Borel} \triangle \text{null-set}\}$
- ▶ $A \sim_{\mathcal{L}} B := \text{equidecomposable with Lebesgue pieces}$
 - ▶ $A, B \in \mathcal{L}$

Measurable equidecompositions

- ▶ $\mathcal{L} := \{\text{Lebesgue sets}\} = \{\text{Borel} \triangle \text{null-set}\}$
- ▶ $A \sim_{\mathcal{L}} B := \text{equidecomposable with Lebesgue pieces}$
 - ▶ $A, B \in \mathcal{L}$
 - ▶ $\lambda(A) = \lambda(B)$

Measurable equidecompositions

- ▶ $\mathcal{L} := \{\text{Lebesgue sets}\} = \{\text{Borel} \triangle \text{null-set}\}$
- ▶ $A \sim_{\mathcal{L}} B :=$ equidecomposable with Lebesgue pieces
 - ▶ $A, B \in \mathcal{L}$
 - ▶ $\lambda(A) = \lambda(B)$
- ▶ **Banach-Tarski'24:** Banach-Tarski with Lebesgue pieces (but countably many)

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

- ▶ Necessary conditions:

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

► Necessary conditions:

► $A \in \mathcal{L}$

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

► Necessary conditions:

- $A \in \mathcal{L}$
- A is bounded

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

► Necessary conditions:

- $A \in \mathcal{L}$
- A is bounded
- $\lambda(A) = 1$

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

► Necessary conditions:

- $A \in \mathcal{L}$
- A is bounded
- $\lambda(A) = 1$
- $\bigcup \{\text{finitely many copies of } A\} \supseteq [0, 1]^k$

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

► Necessary conditions:

- $A \in \mathcal{L}$
- A is bounded
- $\lambda(A) = 1$
- $\bigcup \{\text{finitely many copies of } A\} \supseteq [0, 1]^k$

► Grabowski-Máthé-P. \geq '16: sufficient for $k \geq 3$

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

► Necessary conditions:

- $A \in \mathcal{L}$
- A is bounded
- $\lambda(A) = 1$
- $\bigcup \{\text{finitely many copies of } A\} \supseteq [0, 1]^k$

► Grabowski-Máthé-P. \geq '16: sufficient for $k \geq 3$

► Prove for spheres and then “lift” to \mathbb{R}^k

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

- ▶ **Necessary conditions:**
 - ▶ $A \in \mathcal{L}$
 - ▶ A is bounded
 - ▶ $\lambda(A) = 1$
 - ▶ $\bigcup \{\text{finitely many copies of } A\} \supseteq [0, 1]^k$
- ▶ **Grabowski-Máthé-P. \geq '16:** sufficient for $k \geq 3$
- ▶ **Prove** for spheres and then “lift” to \mathbb{R}^k
- ▶ **Grabowski-Máthé-P. \geq '16:** For sphere \mathbb{S}^k , $k \geq 2$:
 $A \sim_{\mathcal{L}}$ some cap iff

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

- ▶ **Necessary conditions:**
 - ▶ $A \in \mathcal{L}$
 - ▶ A is bounded
 - ▶ $\lambda(A) = 1$
 - ▶ $\bigcup \{\text{finitely many copies of } A\} \supseteq [0, 1]^k$
- ▶ **Grabowski-Máthé-P. \geq '16:** sufficient for $k \geq 3$
- ▶ **Prove** for spheres and then “lift” to \mathbb{R}^k
- ▶ **Grabowski-Máthé-P. \geq '16:** For sphere \mathbb{S}^k , $k \geq 2$:
 $A \sim_{\mathcal{L}}$ some cap iff
 - ▶ $A \in \mathcal{L}$

When is $A \sim_{\mathcal{L}} [0, 1]^k$?

- ▶ **Necessary conditions:**
 - ▶ $A \in \mathcal{L}$
 - ▶ A is bounded
 - ▶ $\lambda(A) = 1$
 - ▶ $\bigcup \{\text{finitely many copies of } A\} \supseteq [0, 1]^k$
- ▶ **Grabowski-Máthé-P. $\geq '16$:** sufficient for $k \geq 3$
- ▶ **Prove** for spheres and then “lift” to \mathbb{R}^k
- ▶ **Grabowski-Máthé-P. $\geq '16$:** For sphere \mathbb{S}^k , $k \geq 2$:
 $A \sim_{\mathcal{L}}$ some cap iff
 - ▶ $A \in \mathcal{L}$
 - ▶ $\bigcup \{\text{finitely many copies of } A\} \supseteq \mathbb{S}^k$

Measurable circle squaring?

Measurable circle squaring?

- ▶ Laczkovich'90: with Lebesgue pieces ?

Measurable circle squaring?

- ▶ Laczkovich'90: with Lebesgue pieces ?
- ▶ Marks-Unger'16: with Baire pieces ?

Measurable circle squaring?

- ▶ Laczkovich'90: with Lebesgue pieces ?
- ▶ Marks-Unger'16: with Baire pieces ?
- ▶ $\mathcal{T} := \{\text{Baire}\} = \{\text{Borel} \triangle \text{meager}\}$

Measurable circle squaring?

- ▶ Laczkovich'90: with Lebesgue pieces ?
- ▶ Marks-Unger'16: with Baire pieces ?
- ▶ $\mathcal{T} := \{\text{Baire}\} = \{\text{Borel} \triangle \text{meager}\}$
- ▶ Grabowski-Máthé-P. \geq '16: Yes

Measurable circle squaring?

- ▶ Laczkovich'90: with Lebesgue pieces ?
- ▶ Marks-Unger'16: with Baire pieces ?
- ▶ $\mathcal{T} := \{\text{Baire}\} = \{\text{Borel} \triangle \text{meager}\}$
- ▶ Grabowski-Máthé-P. \geq '16: Yes
- ▶ Grabowski-Máthé-P. \geq '16: $A, B \subseteq \mathbb{R}^k$, bounded,
 $\lambda(A) = \lambda(B) > 0$, $\dim_{\square}(\partial A), \dim_{\square}(\partial B) < k \Rightarrow$
 $A \overset{\text{Tr}}{\sim}_{\mathcal{L} \cap \mathcal{T}} B$

Decompositions via graph matching

Decompositions via graph matching

- ▶ $A, B \in \mathcal{L}$

Decompositions via graph matching

- ▶ $A, B \in \mathcal{L}$
- ▶ Isometries $\gamma_1, \dots, \gamma_n$

Decompositions via graph matching

- ▶ $A, B \in \mathcal{L}$
- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Infinite graph \mathcal{G} :

Decompositions via graph matching

- ▶ $A, B \in \mathcal{L}$
- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Infinite graph \mathcal{G} :
 - ▶ $V := A \sqcup B$

Decompositions via graph matching

- ▶ $A, B \in \mathcal{L}$
- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Infinite graph \mathcal{G} :
 - ▶ $V := A \sqcup B$
 - ▶ $E = \{xy \in A \times C : \exists i \gamma_i x = y\}$

Decompositions via graph matching

- ▶ $A, B \in \mathcal{L}$
- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Infinite graph \mathcal{G} :
 - ▶ $V := A \sqcup B$
 - ▶ $E = \{xy \in A \times B : \exists i \gamma_i x = y\}$
- ▶ Matching $\mathcal{M} \subseteq E \leftrightarrow$ disjoint $A_1, \dots, A_n \subseteq A$ st $\gamma_1 A_1, \dots, \gamma_n A_n \subseteq B$ are disjoint

Decompositions via graph matching

- ▶ $A, B \in \mathcal{L}$
- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Infinite graph \mathcal{G} :
 - ▶ $V := A \sqcup B$
 - ▶ $E = \{xy \in A \times B : \exists i \gamma_i x = y\}$
- ▶ Matching $\mathcal{M} \subseteq E \leftrightarrow$ disjoint $A_1, \dots, A_n \subseteq A$ st $\gamma_1 A_1, \dots, \gamma_n A_n \subseteq B$ are disjoint
- ▶ \exists perfect matching $\Rightarrow A \sim B$

Decompositions via graph matching

- ▶ $A, B \in \mathcal{L}$
- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Infinite graph \mathcal{G} :
 - ▶ $V := A \sqcup B$
 - ▶ $E = \{xy \in A \times B : \exists i \gamma_i x = y\}$
- ▶ Matching $\mathcal{M} \subseteq E \leftrightarrow$ disjoint $A_1, \dots, A_n \subseteq A$ st $\gamma_1 A_1, \dots, \gamma_n A_n \subseteq B$ are disjoint
- ▶ \exists perfect matching $\Rightarrow A \sim B$
- ▶ \mathcal{M} is measurable := each $A_i \in \mathcal{L}$

Measurable maximal matching

Measurable maximal matching

- ▶ Greedy Algorithm:

Measurable maximal matching

- ▶ Greedy Algorithm:
 - ▶ $A_1 := A \cap \gamma_1^{-1}(B)$

Measurable maximal matching

- ▶ Greedy Algorithm:
 - ▶ $A_1 := A \cap \gamma_1^{-1}(B)$
 - ▶ $B_1 := \gamma_1(A_1)$

Measurable maximal matching

- ▶ Greedy Algorithm:

- ▶ $A_1 := A \cap \gamma_1^{-1}(B)$
- ▶ $B_1 := \gamma_1(A_1)$
- ▶ $A_2 := (A \setminus A_1) \cap \gamma_2^{-1}(B \setminus B_1)$

Measurable maximal matching

- ▶ Greedy Algorithm:

- ▶ $A_1 := A \cap \gamma_1^{-1}(B)$
- ▶ $B_1 := \gamma_1(A_1)$
- ▶ $A_2 := (A \setminus A_1) \cap \gamma_2^{-1}(B \setminus B_1)$
- ▶ $B_2 := \gamma_2(A_2)$

Measurable maximal matching

- ▶ Greedy Algorithm:

- ▶ $A_1 := A \cap \gamma_1^{-1}(B)$
- ▶ $B_1 := \gamma_1(A_1)$
- ▶ $A_2 := (A \setminus A_1) \cap \gamma_2^{-1}(B \setminus B_1)$
- ▶ $B_2 := \gamma_2(A_2)$
- ▶ ...

Matchings without short augmenting paths

Matchings without short augmenting paths

- ▶ Greedy Algorithm $\rightarrow \mathcal{M}_1$

Matchings without short augmenting paths

- ▶ Greedy Algorithm $\rightarrow \mathcal{M}_1$
- ▶ To define \mathcal{M}_2 :

Matchings without short augmenting paths

- ▶ Greedy Algorithm $\rightarrow \mathcal{M}_1$
- ▶ To define \mathcal{M}_2 :
 - ▶ For all triples $(\gamma_i, \gamma_j, \gamma_h)$

Matchings without short augmenting paths

- ▶ Greedy Algorithm $\rightarrow \mathcal{M}_1$
- ▶ To define \mathcal{M}_2 :
 - ▶ For all triples $(\gamma_i, \gamma_j, \gamma_h)$
 - ▶ In parallel, **flip** all augmenting paths labelled $(\gamma_i, \gamma_j, \gamma_h)$

Matchings without short augmenting paths

- ▶ Greedy Algorithm $\rightarrow \mathcal{M}_1$
- ▶ To define \mathcal{M}_2 :
 - ▶ For all triples $(\gamma_i, \gamma_j, \gamma_h)$
 - ▶ In parallel, **flip** all augmenting paths labelled $(\gamma_i, \gamma_j, \gamma_h)$
 - ▶ Vertex-disjoint

Matchings without short augmenting paths

- ▶ Greedy Algorithm $\rightarrow \mathcal{M}_1$
- ▶ To define \mathcal{M}_2 :
 - ▶ For all triples $(\gamma_i, \gamma_j, \gamma_h)$
 - ▶ In parallel, **flip** all augmenting paths labelled $(\gamma_i, \gamma_j, \gamma_h)$
 - ▶ Vertex-disjoint
 - ▶ Measurability is preserved

Matchings without short augmenting paths

- ▶ Greedy Algorithm $\rightarrow \mathcal{M}_1$
- ▶ To define \mathcal{M}_2 :
 - ▶ For all triples $(\gamma_i, \gamma_j, \gamma_h)$
 - ▶ In parallel, **flip** all augmenting paths labelled $(\gamma_i, \gamma_j, \gamma_h)$
 - ▶ Vertex-disjoint
 - ▶ Measurability is preserved
- ▶ **Elek-Lippner'10**: \exists measurable \mathcal{M}_i without augmenting paths of length $\leq 2i - 1$

Borel-Cantelli-type argument

Borel-Cantelli-type argument

- ▶ $\mathcal{M}_1, \mathcal{M}_2, \dots$

Borel-Cantelli-type argument

- ▶ $\mathcal{M}_1, \mathcal{M}_2, \dots$
- ▶ $\mathcal{M} := \bigcup_{i=1}^{\infty} \bigcap_{j=1}^n M_j \subseteq E$

Borel-Cantelli-type argument

- ▶ $\mathcal{M}_1, \mathcal{M}_2, \dots$
 - ▶ $\mathcal{M} := \bigcup_{i=1}^{\infty} \bigcap_{j=1}^n \mathcal{M}_j \subseteq E$
 - ▶ If
 1. $\lambda(\{a \in A : \text{missed by } \mathcal{M}_i\}) \rightarrow 0$
 2. $\sum_{i=1}^{\infty} \lambda(\{a \in A : \mathcal{M}_i, \mathcal{M}_{i+1} \text{ differ on } a\}) < \infty$
- then \mathcal{M} is a perfect matching a.e.

Borel-Cantelli-type argument

- ▶ $\mathcal{M}_1, \mathcal{M}_2, \dots$
- ▶ $\mathcal{M} := \bigcup_{i=1}^{\infty} \bigcap_{j=1}^n \mathcal{M}_j \subseteq E$
- ▶ If
 1. $\lambda(\{a \in A : \text{missed by } \mathcal{M}_i\}) \rightarrow 0$
 2. $\sum_{i=1}^{\infty} \lambda(\{a \in A : \mathcal{M}_i, \mathcal{M}_{i+1} \text{ differ on } a\}) < \infty$

then \mathcal{M} is a perfect matching a.e.

- ▶ **Lyons-Nazarov'11:** Automatically holds if \mathcal{G} is a bipartite expander.

Open Problems

Open Problems

- ▶ Circle squaring with **Borel** pieces?

Open Problems

- ▶ Circle squaring with **Borel** pieces?
- ▶ **Minimum** number of pieces for circle squaring?

Open Problems

- ▶ Circle squaring with **Borel** pieces?
- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92**: “*a rough estimate*” is 10^{40}

Open Problems

- ▶ Circle squaring with **Borel** pieces?
- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92**: “*a rough estimate*” is 10^{40}
 - ▶ **Lower bound**:

Open Problems

- ▶ Circle squaring with **Borel** pieces?
- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92**: “*a rough estimate*” is 10^{40}
 - ▶ **Lower bound**:
 - ▶ 3 (if rotations allowed)

Open Problems

- ▶ Circle squaring with **Borel** pieces?
- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92**: “*a rough estimate*” is 10^{40}
 - ▶ **Lower bound**:
 - ▶ 3 (if rotations allowed)
 - ▶ 4 (for translations only)

Thank you!