

Measurable Combinatorics

Oleg Pikhurko
University of Warwick

"Interactions with Combinatorics", Birmingham, 30 June 2017

Motivation

Motivation

- ▶ Scientific curiosity

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$
- ▶ All degrees uniformly bounded

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$
- ▶ All degrees uniformly bounded
- ▶ Axiom of Choice

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$
- ▶ All degrees uniformly bounded
- ▶ Axiom of Choice \rightsquigarrow finite graphs

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$
- ▶ All degrees uniformly bounded
- ▶ Axiom of Choice \rightsquigarrow finite graphs
 - ▶ $\chi(G) \leq k$

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$
- ▶ All degrees uniformly bounded
- ▶ Axiom of Choice \rightsquigarrow finite graphs
 - ▶ $\chi(G) \leq k$ iff \forall finite $U \subseteq V$ $\chi(G[U]) \leq k$

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$
- ▶ All degrees uniformly bounded
- ▶ Axiom of Choice \rightsquigarrow finite graphs
 - ▶ $\chi(G) \leq k$ iff \forall finite $U \subseteq V$ $\chi(G[U]) \leq k$
- ▶ Extra structure

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$
- ▶ All degrees uniformly bounded
- ▶ Axiom of Choice \rightsquigarrow finite graphs
 - ▶ $\chi(G) \leq k$ iff \forall finite $U \subseteq V$ $\chi(G[U]) \leq k$
- ▶ Extra structure
 - ▶ End compactification (Freudenthal'31, ...)

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$
- ▶ All degrees uniformly bounded
- ▶ Axiom of Choice \rightsquigarrow finite graphs
 - ▶ $\chi(G) \leq k$ iff \forall finite $U \subseteq V$ $\chi(G[U]) \leq k$
- ▶ Extra structure
 - ▶ End compactification (Freudenthal'31, ...)
 - ▶ Martin boundary (Martin'41, Doob'59, Hung'60, ...)

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$
- ▶ All degrees uniformly bounded
- ▶ Axiom of Choice \rightsquigarrow finite graphs
 - ▶ $\chi(G) \leq k$ iff \forall finite $U \subseteq V$ $\chi(G[U]) \leq k$
- ▶ Extra structure
 - ▶ End compactification (Freudenthal'31, ...)
 - ▶ Martin boundary (Martin'41, Doob'59, Hung'60, ...)
- ▶ V : measure space

Motivation

- ▶ Scientific curiosity
- ▶ Extend known results to infinite graphs $G = (V, E)$
- ▶ All degrees uniformly bounded
- ▶ Axiom of Choice \rightsquigarrow finite graphs
 - ▶ $\chi(G) \leq k$ iff \forall finite $U \subseteq V$ $\chi(G[U]) \leq k$
- ▶ Extra structure
 - ▶ End compactification (Freudenthal'31, ...)
 - ▶ Martin boundary (Martin'41, Doob'59, Hung'60, ...)
- ▶ V : measure space
 - ▶ Allow only measurable $c : V \rightarrow [k]$

Finite graphs

Finite graphs

- ▶ Finite graph G

Finite graphs

- ▶ Finite graph G

- ▶ **Bijections** $\phi_1, \dots, \phi_k : V \rightarrow V$ st

$$E = \{\{x, y\} : x \neq y \text{ \& } \exists i \phi_i(x) = y\}$$

Finite graphs

- ▶ Finite graph G

- ▶ **Bijections** $\phi_1, \dots, \phi_k : V \rightarrow V$ st

$$E = \{\{x, y\} : x \neq y \text{ \& } \exists i \phi_i(x) = y\}$$

- ▶ Edge \leftrightarrow transposition

Finite graphs

- ▶ Finite graph G

- ▶ **Bijections** $\phi_1, \dots, \phi_k : V \rightarrow V$ st

$$E = \{\{x, y\} : x \neq y \text{ \& } \exists i \phi_i(x) = y\}$$

- ▶ Edge \leftrightarrow transposition
 - ▶ Matching \leftrightarrow involution

Finite graphs

- ▶ Finite graph G

- ▶ **Bijections** $\phi_1, \dots, \phi_k : V \rightarrow V$ st

$$E = \{\{x, y\} : x \neq y \text{ \& } \exists i \phi_i(x) = y\}$$

- ▶ Edge \leftrightarrow transposition
 - ▶ Matching \leftrightarrow involution
 - ▶ Cycles (with orientations) \leftrightarrow bijection

Finite graphs

- ▶ Finite graph G

- ▶ **Bijections** $\phi_1, \dots, \phi_k : V \rightarrow V$ st

$$E = \{\{x, y\} : x \neq y \text{ \& } \exists i \phi_i(x) = y\}$$

- ▶ Edge \leftrightarrow transposition
 - ▶ Matching \leftrightarrow involution
 - ▶ Cycles (with orientations) \leftrightarrow bijection
 - ▶ $E =$ **union** of edges

Finite graphs

- ▶ Finite graph G

- ▶ **Bijections** $\phi_1, \dots, \phi_k : V \rightarrow V$ st

$$E = \{\{x, y\} : x \neq y \text{ \& } \exists i \phi_i(x) = y\}$$

- ▶ Edge \leftrightarrow transposition
 - ▶ Matching \leftrightarrow involution
 - ▶ Cycles (with orientations) \leftrightarrow bijection
 - ▶ $E =$ **union** of edges / matchings

Finite graphs

- ▶ Finite graph G

- ▶ **Bijections** $\phi_1, \dots, \phi_k : V \rightarrow V$ st

$$E = \{\{x, y\} : x \neq y \text{ \& } \exists i \phi_i(x) = y\}$$

- ▶ Edge \leftrightarrow transposition
 - ▶ Matching \leftrightarrow involution
 - ▶ Cycles (with orientations) \leftrightarrow bijection
 - ▶ $E =$ **union** of edges / matchings / cycles

Finite graphs

- ▶ Finite graph G

- ▶ **Bijections** $\phi_1, \dots, \phi_k : V \rightarrow V$ st

$$E = \{\{x, y\} : x \neq y \text{ \& } \exists i \phi_i(x) = y\}$$

- ▶ Edge \leftrightarrow transposition
 - ▶ Matching \leftrightarrow involution
 - ▶ Cycles (with orientations) \leftrightarrow bijection
 - ▶ $E =$ **union** of edges / matchings / cycles
 - ▶ μ : uniform measure on V

Finite graphs

- ▶ Finite graph G

- ▶ **Bijections** $\phi_1, \dots, \phi_k : V \rightarrow V$ st

$$E = \{\{x, y\} : x \neq y \text{ \& } \exists i \phi_i(x) = y\}$$

- ▶ Edge \leftrightarrow transposition
 - ▶ Matching \leftrightarrow involution
 - ▶ Cycles (with orientations) \leftrightarrow bijection
 - ▶ $E =$ **union** of edges / matchings / cycles
 - ▶ μ : uniform measure on V
 - ▶ Measure-preserving invertible maps $\phi_i : V \rightarrow V$

Graphings

Graphings

- ▶ **Graphing** (V, E, \mathcal{A}, μ) :

Graphings

- ▶ **Graphing** (V, E, \mathcal{A}, μ) :
 - ▶ (V, E) : graph

Graphings

- ▶ **Graphing** (V, E, \mathcal{A}, μ) :
 - ▶ (V, E) : graph
 - ▶ (V, \mathcal{A}, μ) : standard probability space

Graphings

- ▶ **Graphing** (V, E, \mathcal{A}, μ) :
 - ▶ (V, E) : graph
 - ▶ (V, \mathcal{A}, μ) : standard probability space
 - ▶ eg unit interval $[0, 1]$

Graphings

- ▶ **Graphing** (V, E, \mathcal{A}, μ) :
 - ▶ (V, E) : graph
 - ▶ (V, \mathcal{A}, μ) : standard probability space
 - ▶ eg unit interval $[0, 1]$
 - ▶ \exists measure-preserving maps $\phi_1, \dots, \phi_k : V \rightarrow V$

Graphings

- ▶ **Graphing** (V, E, \mathcal{A}, μ) :
 - ▶ (V, E) : graph
 - ▶ (V, \mathcal{A}, μ) : standard probability space
 - ▶ eg unit interval $[0, 1]$
 - ▶ \exists measure-preserving maps $\phi_1, \dots, \phi_k : V \rightarrow V$
 $E = \{ \{x, y\} : x \neq y \text{ \& } \exists i \phi_i x = y \}$

Graphings

- ▶ **Graphing** (V, E, \mathcal{A}, μ) :
 - ▶ (V, E) : graph
 - ▶ (V, \mathcal{A}, μ) : standard probability space
 - ▶ eg unit interval $[0, 1]$
 - ▶ \exists measure-preserving maps $\phi_1, \dots, \phi_k : V \rightarrow V$
 $E = \{ \{x, y\} : x \neq y \text{ \& } \exists i \ \phi_i x = y \}$
- ▶ $\Delta(G) \leq 2k$

More motivation

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence
- ▶ Finite graph:

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence
- ▶ Finite graph:
 - ▶ Uniform vertex x

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence
- ▶ Finite graph:
 - ▶ Uniform vertex x
 - ▶ Output radius- r neighbourhood of x

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence
- ▶ Finite graph:
 - ▶ Uniform vertex x
 - ▶ Output radius- r neighbourhood of x
 - ▶ Rooted graph, up to isomorphism

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence
- ▶ Finite graph:
 - ▶ Uniform vertex x
 - ▶ Output radius- r neighbourhood of x
 - ▶ Rooted graph, up to isomorphism
- ▶ Graphing (V, E, \mathcal{A}, μ) :

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence
- ▶ Finite graph:
 - ▶ Uniform vertex x
 - ▶ Output radius- r neighbourhood of x
 - ▶ Rooted graph, up to isomorphism
- ▶ Graphing (V, E, \mathcal{A}, μ) :
 - ▶ Sample $x \in V$ using μ

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence
- ▶ Finite graph:
 - ▶ Uniform vertex x
 - ▶ Output radius- r neighbourhood of x
 - ▶ Rooted graph, up to isomorphism
- ▶ Graphing (V, E, \mathcal{A}, μ) :
 - ▶ Sample $x \in V$ using μ
 - ▶ Output radius- r neighbourhood of x in (V, E)

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence
- ▶ Finite graph:
 - ▶ Uniform vertex x
 - ▶ Output radius- r neighbourhood of x
 - ▶ Rooted graph, up to isomorphism
- ▶ Graphing (V, E, \mathcal{A}, μ) :
 - ▶ Sample $x \in V$ using μ
 - ▶ Output radius- r neighbourhood of x in (V, E)
- ▶ **Property testing**

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence
- ▶ Finite graph:
 - ▶ Uniform vertex x
 - ▶ Output radius- r neighbourhood of x
 - ▶ Rooted graph, up to isomorphism
- ▶ Graphing (V, E, \mathcal{A}, μ) :
 - ▶ Sample $x \in V$ using μ
 - ▶ Output radius- r neighbourhood of x in (V, E)
- ▶ **Property testing**
- ▶ **Not** unique

More motivation

- ▶ **Limit objects** for Benjamini-Schramm convergence
- ▶ Finite graph:
 - ▶ Uniform vertex x
 - ▶ Output radius- r neighbourhood of x
 - ▶ Rooted graph, up to isomorphism
- ▶ Graphing (V, E, \mathcal{A}, μ) :
 - ▶ Sample $x \in V$ using μ
 - ▶ Output radius- r neighbourhood of x in (V, E)
- ▶ **Property testing**
- ▶ **Not** unique
- ▶ **Aldous-Lyons'01**: Is every graphing approximable by finite graphs ?

Example: irrational rotation \mathcal{R}_α

Example: irrational rotation \mathcal{R}_α

- Fix **irrational** $\alpha \in \mathbb{R}$

Example: irrational rotation \mathcal{R}_α

- ▶ Fix **irrational** $\alpha \in \mathbb{R}$
- ▶ $V := [0, 1)$

Example: irrational rotation \mathcal{R}_α

- ▶ Fix **irrational** $\alpha \in \mathbb{R}$
- ▶ $V := [0, 1)$
- ▶ $T : V \rightarrow V, \quad x \mapsto x + \alpha \pmod{1}$

Example: irrational rotation \mathcal{R}_α

- ▶ Fix **irrational** $\alpha \in \mathbb{R}$
- ▶ $V := [0, 1)$
- ▶ $T : V \rightarrow V, \quad x \mapsto x + \alpha \pmod{1}$
- ▶ $E := \{\{x, Tx\} : x \in V\}$

Example: irrational rotation \mathcal{R}_α

- ▶ Fix **irrational** $\alpha \in \mathbb{R}$
- ▶ $V := [0, 1)$
- ▶ $T : V \rightarrow V, \quad x \mapsto x + \alpha \pmod{1}$
- ▶ $E := \{\{x, Tx\} : x \in V\}$
- ▶ Graphing **\mathcal{R}_α**

Properties of \mathcal{R}_α

Properties of \mathcal{R}_α

- ▶ Each component: infinite 2-regular path

Properties of \mathcal{R}_α

- ▶ Each component: infinite 2-regular path
- ▶ **Axiom of Choice:** $\chi(\mathcal{R}_\alpha) = 2$

Properties of \mathcal{R}_α

- ▶ Each component: infinite 2-regular path
- ▶ **Axiom of Choice**: $\chi(\mathcal{R}_\alpha) = 2$
- ▶ **Measurable chromatic number** $\chi_\mu(\mathcal{R}_\alpha)$

Properties of \mathcal{R}_α

- ▶ Each component: infinite 2-regular path
- ▶ **Axiom of Choice**: $\chi(\mathcal{R}_\alpha) = 2$
- ▶ **Measurable chromatic number** $\chi_\mu(\mathcal{R}_\alpha) \geq 3$:

Properties of \mathcal{R}_α

- ▶ Each component: infinite 2-regular path
- ▶ **Axiom of Choice**: $\chi(\mathcal{R}_\alpha) = 2$
- ▶ **Measurable chromatic number** $\chi_\mu(\mathcal{R}_\alpha) \geq 3$:
 - ▶ $V = A \cup B$

Properties of \mathcal{R}_α

- ▶ Each component: infinite 2-regular path
- ▶ **Axiom of Choice**: $\chi(\mathcal{R}_\alpha) = 2$
- ▶ **Measurable chromatic number** $\chi_\mu(\mathcal{R}_\alpha) \geq 3$:
 - ▶ $V = A \cup B$
 - ▶ $T(A) = B$

Properties of \mathcal{R}_α

- ▶ Each component: infinite 2-regular path
- ▶ **Axiom of Choice**: $\chi(\mathcal{R}_\alpha) = 2$
- ▶ **Measurable chromatic number** $\chi_\mu(\mathcal{R}_\alpha) \geq 3$:
 - ▶ $V = A \cup B$
 - ▶ $T(A) = B$
 - ▶ $\mu(A) = \mu(B) = 1/2$

Properties of \mathcal{R}_α

- ▶ Each component: infinite 2-regular path
- ▶ **Axiom of Choice**: $\chi(\mathcal{R}_\alpha) = 2$
- ▶ **Measurable chromatic number** $\chi_\mu(\mathcal{R}_\alpha) \geq 3$:
 - ▶ $V = A \cup B$
 - ▶ $T(A) = B$
 - ▶ $\mu(A) = \mu(B) = 1/2$
 - ▶ Contradicts **ergodicity** of T

Properties of \mathcal{R}_α

- ▶ Each component: infinite 2-regular path
- ▶ **Axiom of Choice**: $\chi(\mathcal{R}_\alpha) = 2$
- ▶ **Measurable chromatic number** $\chi_\mu(\mathcal{R}_\alpha) \geq 3$:
 - ▶ $V = A \cup B$
 - ▶ $T(A) = B$
 - ▶ $\mu(A) = \mu(B) = 1/2$
 - ▶ Contradicts **ergodicity** of T
- ▶ $\chi_\mu(\mathcal{R}_\alpha) = 3$

Properties of \mathcal{R}_α

- ▶ Each component: infinite 2-regular path
- ▶ **Axiom of Choice**: $\chi(\mathcal{R}_\alpha) = 2$
- ▶ **Measurable chromatic number** $\chi_\mu(\mathcal{R}_\alpha) \geq 3$:
 - ▶ $V = A \cup B$
 - ▶ $T(A) = B$
 - ▶ $\mu(A) = \mu(B) = 1/2$
 - ▶ Contradicts **ergodicity** of T
- ▶ $\chi_\mu(\mathcal{R}_\alpha) = 3$
- ▶ $\chi_{\text{Borel}}(\mathcal{R}_\alpha) = 3$

Borel and measurable set

Borel and measurable set

- ▶ $\{\text{Borel}\} \stackrel{\text{def}}{=} \text{smallest } \sigma\text{-algebra} \supseteq \text{open sets}$

Borel and measurable set

- ▶ $\{\text{Borel}\} \stackrel{\text{def}}{=} \text{smallest } \sigma\text{-algebra} \supseteq \text{open sets}$
- ▶ Some permitted operations:

Borel and measurable set

- ▶ $\{\text{Borel}\} \stackrel{\text{def}}{=} \text{smallest } \sigma\text{-algebra} \supseteq \text{open sets}$
- ▶ Some permitted operations:
 - ▶ Boolean operations

Borel and measurable set

- ▶ $\{\text{Borel}\} \stackrel{\text{def}}{=} \text{smallest } \sigma\text{-algebra} \supseteq \text{open sets}$
- ▶ Some permitted operations:
 - ▶ Boolean operations
 - ▶ Countably many iterations

Borel and measurable set

- ▶ $\{\text{Borel}\} \stackrel{\text{def}}{=} \text{smallest } \sigma\text{-algebra} \supseteq \text{open sets}$
- ▶ Some permitted operations:
 - ▶ Boolean operations
 - ▶ Countably many iterations
 - ▶ Colour of every vertex eventually stabilises \Rightarrow Borel

Borel and measurable set

- ▶ $\{\text{Borel}\} \stackrel{\text{def}}{=} \text{smallest } \sigma\text{-algebra} \supseteq \text{open sets}$
- ▶ Some permitted operations:
 - ▶ Boolean operations
 - ▶ Countably many iterations
 - ▶ Colour of every vertex eventually stabilises \Rightarrow Borel
- ▶ $(\text{Lebesgue measurable}) \stackrel{\text{def}}{=} \text{Borel} \triangle \text{measure-0}$

Borel and measurable set

- ▶ $\{\text{Borel}\} \stackrel{\text{def}}{=} \text{smallest } \sigma\text{-algebra} \supseteq \text{open sets}$
- ▶ Some permitted operations:
 - ▶ Boolean operations
 - ▶ Countably many iterations
 - ▶ Colour of every vertex eventually stabilises \Rightarrow Borel
- ▶ $(\text{Lebesgue measurable}) \stackrel{\text{def}}{=} \text{Borel} \triangle \text{measure-0}$
 - ▶ Stabilises a.e. \Rightarrow measurable

Greedy colourings Borel way

Greedy colourings Borel way

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$

Greedy colourings Borel way

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Proof:

Greedy colourings Borel way

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Proof:
 - ▶ WLOG $V = [0, 1]$

Greedy colourings Borel way

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Proof:
 - ▶ WLOG $V = [0, 1]$
 - ▶ Borel vertex colouring $f : V \rightarrow \{0, 1\}^{<\omega}$

Greedy colourings Borel way

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Proof:
 - ▶ WLOG $V = [0, 1]$
 - ▶ Borel vertex colouring $f : V \rightarrow \{0, 1\}^{<\omega}$
 - ▶ $f(x) := \min \text{ prefix } b_1 \dots b_k \text{ separating } x = 0.b_1 b_2 b_3 \dots$
from $N(x)$

Greedy colourings Borel way

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Proof:
 - ▶ WLOG $V = [0, 1]$
 - ▶ Borel vertex colouring $f : V \rightarrow \{0, 1\}^{<\omega}$
 - ▶ $f(x) := \min \text{ prefix } b_1 \dots b_k \text{ separating } x = 0.b_1 b_2 b_3 \dots$
from $N(x)$
 - ▶ Define $c : V \rightarrow [\Delta + 1]$:

Greedy colourings Borel way

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Proof:
 - ▶ WLOG $V = [0, 1]$
 - ▶ Borel vertex colouring $f : V \rightarrow \{0, 1\}^{<\omega}$
 - ▶ $f(x) := \min \text{ prefix } b_1 \dots b_k \text{ separating } x = 0.b_1 b_2 b_3 \dots$
from $N(x)$
 - ▶ Define $c : V \rightarrow [\Delta + 1]$:
 - ▶ For $i \in \{0, 1\}^{<\omega}$

Greedy colourings Borel way

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Proof:
 - ▶ WLOG $V = [0, 1]$
 - ▶ Borel vertex colouring $f : V \rightarrow \{0, 1\}^{<\omega}$
 - ▶ $f(x) := \min \text{ prefix } b_1 \dots b_k \text{ separating } x = 0.b_1 b_2 b_3 \dots$ from $N(x)$
 - ▶ Define $c : V \rightarrow [\Delta + 1]$:
 - ▶ For $i \in \{0, 1\}^{<\omega}$
Colour each $x \in f^{-1}(i)$ with smallest allowed colour

Greedy colourings Borel way

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Proof:
 - ▶ WLOG $V = [0, 1]$
 - ▶ Borel vertex colouring $f : V \rightarrow \{0, 1\}^{<\omega}$
 - ▶ $f(x) := \min \text{ prefix } b_1 \dots b_k \text{ separating } x = 0.b_1 b_2 b_3 \dots$ from $N(x)$
 - ▶ Define $c : V \rightarrow [\Delta + 1]$:
 - ▶ For $i \in \{0, 1\}^{<\omega}$
Colour each $x \in f^{-1}(i)$ with smallest allowed colour
- ▶ Descriptive set theory

Some known results

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Corollary: Borel chromatic index $\chi'_{\text{Borel}} \leq 2\Delta - 1$

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Corollary: Borel chromatic index $\chi'_{\text{Borel}} \leq 2\Delta - 1$
- ▶ Marks'16: $2\Delta - 1$ is attainable

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Corollary: Borel chromatic index $\chi'_{\text{Borel}} \leq 2\Delta - 1$
- ▶ Marks'16: $2\Delta - 1$ is attainable
 - ▶ bipartite & acyclic

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Corollary: Borel chromatic index $\chi'_{\text{Borel}} \leq 2\Delta - 1$
- ▶ Marks'16: $2\Delta - 1$ is attainable
 - ▶ bipartite & acyclic
- ▶ Vizing'64, Gupta'66: $\chi' \leq \Delta + 1$

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Corollary: Borel chromatic index $\chi'_{\text{Borel}} \leq 2\Delta - 1$
- ▶ Marks'16: $2\Delta - 1$ is attainable
 - ▶ bipartite & acyclic
- ▶ Vizing'64, Gupta'66: $\chi' \leq \Delta + 1$
- ▶ Abert'10, Marks'13: Measurable Vizing's theorem?

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Corollary: Borel chromatic index $\chi'_{\text{Borel}} \leq 2\Delta - 1$
- ▶ Marks'16: $2\Delta - 1$ is attainable
 - ▶ bipartite & acyclic
- ▶ Vizing'64, Gupta'66: $\chi' \leq \Delta + 1$
- ▶ Abert'10, Marks'13: Measurable Vizing's theorem?
- ▶ Csóka, Lippner & P'16:

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Corollary: Borel chromatic index $\chi'_{\text{Borel}} \leq 2\Delta - 1$
- ▶ Marks'16: $2\Delta - 1$ is attainable
 - ▶ bipartite & acyclic
- ▶ Vizing'64, Gupta'66: $\chi' \leq \Delta + 1$
- ▶ Abert'10, Marks'13: Measurable Vizing's theorem?
- ▶ Csóka, Lippner & P'16:
 - ▶ $\chi'_\mu \leq \Delta + O(\sqrt{\Delta})$

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Corollary: Borel chromatic index $\chi'_{\text{Borel}} \leq 2\Delta - 1$
- ▶ Marks'16: $2\Delta - 1$ is attainable
 - ▶ bipartite & acyclic
- ▶ Vizing'64, Gupta'66: $\chi' \leq \Delta + 1$
- ▶ Abert'10, Marks'13: Measurable Vizing's theorem?
- ▶ Csóka, Lippner & P'16:
 - ▶ $\chi'_\mu \leq \Delta + O(\sqrt{\Delta})$
 - ▶ Bipartite $\Rightarrow \chi'_\mu \leq \Delta + 1$

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Corollary: Borel chromatic index $\chi'_{\text{Borel}} \leq 2\Delta - 1$
- ▶ Marks'16: $2\Delta - 1$ is attainable
 - ▶ bipartite & acyclic
- ▶ Vizing'64, Gupta'66: $\chi' \leq \Delta + 1$
- ▶ Abert'10, Marks'13: Measurable Vizing's theorem?
- ▶ Csóka, Lippner & P'16:
 - ▶ $\chi'_\mu \leq \Delta + O(\sqrt{\Delta})$
 - ▶ Bipartite $\Rightarrow \chi'_\mu \leq \Delta + 1$
- ▶ Marks'16: Acyclic \mathcal{G} with $\chi_{\text{Borel}} = \Delta + 1$

Some known results

- ▶ Kechris, Solecky & Todorcevic'99: $\chi_{\text{Borel}} \leq \Delta + 1$
- ▶ Corollary: Borel chromatic index $\chi'_{\text{Borel}} \leq 2\Delta - 1$
- ▶ Marks'16: $2\Delta - 1$ is attainable
 - ▶ bipartite & acyclic
- ▶ Vizing'64, Gupta'66: $\chi' \leq \Delta + 1$
- ▶ Abert'10, Marks'13: Measurable Vizing's theorem?
- ▶ Csóka, Lippner & P'16:
 - ▶ $\chi'_\mu \leq \Delta + O(\sqrt{\Delta})$
 - ▶ Bipartite $\Rightarrow \chi'_\mu \leq \Delta + 1$
- ▶ Marks'16: Acyclic \mathcal{G} with $\chi_{\text{Borel}} = \Delta + 1$
- ▶ Conley, Marks & Tucker-Drob'16: Measurable Brooks' theorem for $\Delta \geq 3$

Example: group actions

Example: group actions

- ▶ Group Γ

Example: group actions

- ▶ Group Γ
- ▶ Measure-preserving action $\Gamma \curvearrowright (V, \mathcal{A}, \mu)$

Example: group actions

- ▶ Group Γ
- ▶ **Measure-preserving action** $\Gamma \curvearrowright (V, \mathcal{A}, \mu)$
- ▶ Finite generating set $S \subseteq \Gamma$, $S = S^{-1}$

Example: group actions

- ▶ Group Γ
- ▶ **Measure-preserving action** $\Gamma \curvearrowright (V, \mathcal{A}, \mu)$
- ▶ Finite generating set $S \subseteq \Gamma$, $S = S^{-1}$
- ▶ $E := \{ \{x, \gamma.x\} : x \in V, \gamma \in S, x \neq \gamma.x \}$

Example: group actions

- ▶ Group Γ
- ▶ **Measure-preserving action** $\Gamma \curvearrowright (V, \mathcal{A}, \mu)$
- ▶ Finite generating set $S \subseteq \Gamma$, $S = S^{-1}$
- ▶ $E := \{ \{x, \gamma.x\} : x \in V, \gamma \in S, x \neq \gamma.x \}$
- ▶ Graphing (V, E, \mathcal{A}, μ)

Example: group actions

- ▶ Group Γ
- ▶ **Measure-preserving action** $\Gamma \curvearrowright (V, \mathcal{A}, \mu)$
- ▶ Finite generating set $S \subseteq \Gamma$, $S = S^{-1}$
- ▶ $E := \{ \{x, \gamma.x\} : x \in V, \gamma \in S, x \neq \gamma.x \}$
- ▶ Graphing (V, E, \mathcal{A}, μ)
- ▶ $\mathcal{R}_\alpha: \mathbb{Z} \cong \langle T \rangle$ acting on $[0, 1)$

Example: group actions

- ▶ Group Γ
- ▶ **Measure-preserving action** $\Gamma \curvearrowright (V, \mathcal{A}, \mu)$
- ▶ Finite generating set $S \subseteq \Gamma$, $S = S^{-1}$
- ▶ $E := \{ \{x, \gamma.x\} : x \in V, \gamma \in S, x \neq \gamma.x \}$
- ▶ Graphing (V, E, \mathcal{A}, μ)
- ▶ $\mathcal{R}_\alpha: \mathbb{Z} \cong \langle T \rangle$ acting on $[0, 1)$
- ▶ \forall graphing is realisable this way

Example: group actions

- ▶ Group Γ
- ▶ **Measure-preserving action** $\Gamma \curvearrowright (V, \mathcal{A}, \mu)$
- ▶ Finite generating set $S \subseteq \Gamma$, $S = S^{-1}$
- ▶ $E := \{ \{x, \gamma.x\} : x \in V, \gamma \in S, x \neq \gamma.x \}$
- ▶ Graphing (V, E, \mathcal{A}, μ)
- ▶ $\mathcal{R}_\alpha: \mathbb{Z} \cong \langle T \rangle$ acting on $[0, 1)$
- ▶ \forall graphing is realisable this way
- ▶ **Measurable group theory**

Equidecomposability

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** $(A \sim B) \stackrel{\text{def}}{=}$

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) $\stackrel{\text{def}}{=}$
 - ▶ $\exists A = A_1 \sqcup \cdots \sqcup A_n$

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) $\stackrel{\text{def}}{=}$
 - ▶ $\exists A = A_1 \sqcup \cdots \sqcup A_n$
 - ▶ $\exists B = B_1 \sqcup \cdots \sqcup B_n$


Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) $\stackrel{\text{def}}{=}$
 - ▶ $\exists A = A_1 \sqcup \cdots \sqcup A_n$
 - ▶ $\exists B = B_1 \sqcup \cdots \sqcup B_n$
- st $\forall i \exists$ isometry γ_i with $A_i = \gamma_i(B_i)$

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) $\stackrel{\text{def}}{=}$
 - ▶ $\exists A = A_1 \sqcup \cdots \sqcup A_n$
 - ▶ $\exists B = B_1 \sqcup \cdots \sqcup B_n$st $\forall i \exists$ isometry γ_i with $A_i = \gamma_i(B_i)$


- ▶ **Example:**



Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) $\stackrel{\text{def}}{=}$
 - ▶ $\exists A = A_1 \sqcup \cdots \sqcup A_n$
 - ▶ $\exists B = B_1 \sqcup \cdots \sqcup B_n$st $\forall i \exists$ isometry γ_i with $A_i = \gamma_i(B_i)$

▶ **Example:**




- ▶ **Wallace 1807, Bolyai 1832, Gerwien 1832:**
equi-area polygons are dissection congruent

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) $\stackrel{\text{def}}{=}$
 - ▶ $\exists A = A_1 \sqcup \cdots \sqcup A_n$
 - ▶ $\exists B = B_1 \sqcup \cdots \sqcup B_n$st $\forall i \exists$ isometry γ_i with $A_i = \gamma_i(B_i)$

▶ **Example:**




- ▶ **Wallace 1807, Bolyai 1832, Gerwien 1832:**
equi-area polygons are dissection congruent
 - ▶ **Tarski'24:** equidecomposable

Equidecomposability

- ▶ $A, B \subseteq \mathbb{R}^k$ are **equidecomposable** ($A \sim B$) $\stackrel{\text{def}}{=}$
 - ▶ $\exists A = A_1 \sqcup \cdots \sqcup A_n$
 - ▶ $\exists B = B_1 \sqcup \cdots \sqcup B_n$st $\forall i \exists$ isometry γ_i with $A_i = \gamma_i(B_i)$

▶ **Example:**



- ▶ **Wallace 1807, Bolyai 1832, Gerwien 1832:**
equi-area polygons are dissection congruent
 - ▶ **Tarski'24:** equidecomposable
 - ▶ **Exercise:** $[0, 1] \sim [0, 1)$

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

- ▶ $k \geq 3$:

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

- ▶ $k \geq 3$:
 - ▶ **Banach-Tarski'24:** $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

- ▶ $k \geq 3$:
 - ▶ **Banach-Tarski'24**: $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k\text{-dim ball}$

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

- ▶ $k \geq 3$:
 - ▶ **Banach-Tarski'24**: $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff 1914**: paradoxes on sphere \mathbb{S}^{k-1}

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

- ▶ $k \geq 3$:
 - ▶ **Banach-Tarski'24**: $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff 1914**: paradoxes on sphere \mathbb{S}^{k-1}
- ▶ $k \leq 2$:

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

- ▶ $k \geq 3$:
 - ▶ **Banach-Tarski'24**: $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff 1914**: paradoxes on sphere \mathbb{S}^{k-1}
- ▶ $k \leq 2$:
 - ▶ **Banach'23**: Measurable $A, B \Rightarrow \mu(A) = \mu(B)$

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

- ▶ $k \geq 3$:
 - ▶ **Banach-Tarski'24**: $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff 1914**: paradoxes on sphere \mathbb{S}^{k-1}
- ▶ $k \leq 2$:
 - ▶ **Banach'23**: Measurable $A, B \Rightarrow \mu(A) = \mu(B)$
 - ▶ **Tarski'25**: Is \mathbb{B}^2 equidecomposable to a square ?

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

- ▶ $k \geq 3$:
 - ▶ **Banach-Tarski'24**: $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff 1914**: paradoxes on sphere \mathbb{S}^{k-1}
- ▶ $k \leq 2$:
 - ▶ **Banach'23**: Measurable $A, B \Rightarrow \mu(A) = \mu(B)$
 - ▶ **Tarski'25**: Is \mathbb{B}^2 equidecomposable to a square ?
 - ▶ **von Neumann'29**: Yes, with affine transformations

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

- ▶ $k \geq 3$:
 - ▶ **Banach-Tarski'24**: $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff 1914**: paradoxes on sphere \mathbb{S}^{k-1}
- ▶ $k \leq 2$:
 - ▶ **Banach'23**: Measurable $A, B \Rightarrow \mu(A) = \mu(B)$
 - ▶ **Tarski'25**: Is \mathbb{B}^2 equidecomposable to a square ?
 - ▶ **von Neumann'29**: Yes, with affine transformations
 - ▶ **Dubins-Hirsh-Karush'63**: No, with topological disks

Arbitrary parts, equidecomposing $A, B \subseteq \mathbb{R}^k$

- ▶ $k \geq 3$:
 - ▶ **Banach-Tarski'24**: $A, B \subseteq \mathbb{R}^k$, bounded, non-empty interior $\Rightarrow A \sim B$
 - ▶ $\mathbb{B}^k \sim \mathbb{B}^k \sqcup \mathbb{B}^k$, $\mathbb{B}^k := k$ -dim ball
 - ▶ **Hausdorff 1914**: paradoxes on sphere \mathbb{S}^{k-1}
- ▶ $k \leq 2$:
 - ▶ **Banach'23**: Measurable $A, B \Rightarrow \mu(A) = \mu(B)$
 - ▶ **Tarski'25**: Is \mathbb{B}^2 equidecomposable to a square ?
 - ▶ **von Neumann'29**: Yes, with affine transformations
 - ▶ **Dubins-Hirsh-Karush'63**: No, with topological disks
 - ▶ **Laczkovich'90**: YES!

Measurable pieces?

Measurable pieces?

- ▶ **Necessary** conditions for $A \sim [0, 1]^k$:

Measurable pieces?

- ▶ **Necessary** conditions for $A \sim [0, 1]^k$:
 - ▶ A is measurable

Measurable pieces?

- ▶ **Necessary** conditions for $A \sim [0, 1]^k$:
 - ▶ A is measurable
 - ▶ A is bounded

Measurable pieces?

- ▶ **Necessary** conditions for $A \sim [0, 1]^k$:
 - ▶ A is measurable
 - ▶ A is bounded
 - ▶ $\mu(A) = 1$

Measurable pieces?

- ▶ **Necessary** conditions for $A \sim [0, 1]^k$:
 - ▶ A is measurable
 - ▶ A is bounded
 - ▶ $\mu(A) = 1$
 - ▶ $\bigcup \{\text{finitely many copies of } A\} \supseteq [0, 1]^k$

Measurable pieces?

- ▶ **Necessary** conditions for $A \sim [0, 1]^k$:
 - ▶ A is measurable
 - ▶ A is bounded
 - ▶ $\mu(A) = 1$
 - ▶ $\bigcup \{\text{finitely many copies of } A\} \supseteq [0, 1]^k$
- ▶ **Grabowski-Máthé-P. \geq '17**: sufficient for $k \geq 3$

Measurable pieces?

- ▶ **Necessary** conditions for $A \sim [0, 1]^k$:
 - ▶ A is measurable
 - ▶ A is bounded
 - ▶ $\mu(A) = 1$
 - ▶ $\bigcup \{\text{finitely many copies of } A\} \supseteq [0, 1]^k$
- ▶ **Grabowski-Máthé-P. '17**: sufficient for $k \geq 3$
- ▶ **Grabowski-Máthé-P. '16**: circle squaring with measurable pieces

Measurable pieces?

- ▶ **Necessary** conditions for $A \sim [0, 1]^k$:
 - ▶ A is measurable
 - ▶ A is bounded
 - ▶ $\mu(A) = 1$
 - ▶ $\bigcup \{\text{finitely many copies of } A\} \supseteq [0, 1]^k$
- ▶ **Grabowski-Máthé-P. \geq '17**: sufficient for $k \geq 3$
- ▶ **Grabowski-Máthé-P.'16**: circle squaring with measurable pieces
- ▶ **Marks-Unger \geq '17**: circle squaring with Borel pieces

Equidecompositions via matchings

Equidecompositions via matchings

- ▶ Isometries $\gamma_1, \dots, \gamma_n$

Equidecompositions via matchings

- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Graphing \mathcal{G} :

Equidecompositions via matchings

- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Graphing \mathcal{G} :
 - ▶ $V := A \sqcup B$

Equidecompositions via matchings

- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Graphing \mathcal{G} :
 - ▶ $V := A \sqcup B$
 - ▶ $E = \{xy \in A \times B : \exists i \gamma_i(x) = y\}$

Equidecompositions via matchings

- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Graphing \mathcal{G} :
 - ▶ $V := A \sqcup B$
 - ▶ $E = \{xy \in A \times B : \exists i \gamma_i(x) = y\}$
- ▶ Matching $\mathcal{M} \subseteq E \iff$ disjoint $A_1, \dots, A_n \subseteq A$ st $\gamma_1(A_1), \dots, \gamma_n(A_n) \subseteq B$ are disjoint

Equidecompositions via matchings

- ▶ Isometries $\gamma_1, \dots, \gamma_n$
- ▶ Graphing \mathcal{G} :
 - ▶ $V := A \sqcup B$
 - ▶ $E = \{xy \in A \times B : \exists i \gamma_i(x) = y\}$
- ▶ Matching $\mathcal{M} \subseteq E \iff$ disjoint $A_1, \dots, A_n \subseteq A$ st $\gamma_1(A_1), \dots, \gamma_n(A_n) \subseteq B$ are disjoint
- ▶ \exists perfect matching $\Rightarrow A \sim B$

Measurable maximal matching

Measurable maximal matching

- ▶ Greedy Algorithm:

Measurable maximal matching

- ▶ Greedy Algorithm:

- ▶ $A_1 := A \cap \gamma_1^{-1}(B)$

Measurable maximal matching

- ▶ Greedy Algorithm:
 - ▶ $A_1 := A \cap \gamma_1^{-1}(B)$
 - ▶ $B_1 := \gamma_1(A_1)$

Measurable maximal matching

- ▶ Greedy Algorithm:

- ▶ $A_1 := A \cap \gamma_1^{-1}(B)$
- ▶ $B_1 := \gamma_1(A_1)$
- ▶ $A_2 := (A \setminus A_1) \cap \gamma_2^{-1}(B \setminus B_1)$

Measurable maximal matching

- ▶ Greedy Algorithm:

- ▶ $A_1 := A \cap \gamma_1^{-1}(B)$
- ▶ $B_1 := \gamma_1(A_1)$
- ▶ $A_2 := (A \setminus A_1) \cap \gamma_2^{-1}(B \setminus B_1)$
- ▶ $B_2 := \gamma_2(A_2)$

Measurable maximal matching

- ▶ Greedy Algorithm:

- ▶ $A_1 := A \cap \gamma_1^{-1}(B)$
- ▶ $B_1 := \gamma_1(A_1)$
- ▶ $A_2 := (A \setminus A_1) \cap \gamma_2^{-1}(B \setminus B_1)$
- ▶ $B_2 := \gamma_2(A_2)$
- ▶ $A_3 := (A \setminus (A_1 \cup A_2)) \cap \gamma_3^{-1}(B \setminus (B_1 \cup B_2))$

Measurable maximal matching

- ▶ Greedy Algorithm:

- ▶ $A_1 := A \cap \gamma_1^{-1}(B)$
- ▶ $B_1 := \gamma_1(A_1)$
- ▶ $A_2 := (A \setminus A_1) \cap \gamma_2^{-1}(B \setminus B_1)$
- ▶ $B_2 := \gamma_2(A_2)$
- ▶ $A_3 := (A \setminus (A_1 \cup A_2)) \cap \gamma_3^{-1}(B \setminus (B_1 \cup B_2))$
- ▶ ...

Measurable maximal matching

- ▶ **Greedy Algorithm:**
 - ▶ $A_1 := A \cap \gamma_1^{-1}(B)$
 - ▶ $B_1 := \gamma_1(A_1)$
 - ▶ $A_2 := (A \setminus A_1) \cap \gamma_2^{-1}(B \setminus B_1)$
 - ▶ $B_2 := \gamma_2(A_2)$
 - ▶ $A_3 := (A \setminus (A_1 \cup A_2)) \cap \gamma_3^{-1}(B \setminus (B_1 \cup B_2))$
 - ▶ ...
- ▶ **Elek-Lippner'10:** \exists Borel \mathcal{M}_i without augmenting paths of length $\leq 2i - 1$

Selecting $\gamma_1, \dots, \gamma_n$

Selecting $\gamma_1, \dots, \gamma_n$

- ▶ For $A, B \subseteq \mathbb{S}^k$, $k \geq 2$:

Selecting $\gamma_1, \dots, \gamma_n$

- ▶ For $A, B \subseteq \mathbb{S}^k$, $k \geq 2$:
 - ▶ Margulis'80, Sullivan'81 ($k \geq 4$), Drinfeld'84 ($k = 2, 3$):
 $\exists \rho_1, \dots, \rho_s$ st $(\mathbb{S}^k, E(\rho_1, \dots, \rho_s))$ is expander

Selecting $\gamma_1, \dots, \gamma_n$

- ▶ For $A, B \subseteq \mathbb{S}^k$, $k \geq 2$:
 - ▶ Margulis'80, Sullivan'81 ($k \geq 4$), Drinfeld'84 ($k = 2, 3$):
 $\exists \rho_1, \dots, \rho_s$ st $(\mathbb{S}^k, E(\rho_1, \dots, \rho_s))$ is expander
 - ▶ All ℓ -products of ρ_i 's, large $\ell = \ell(A, B)$

Selecting $\gamma_1, \dots, \gamma_n$

- ▶ For $A, B \subseteq \mathbb{S}^k$, $k \geq 2$:
 - ▶ Margulis'80, Sullivan'81 ($k \geq 4$), Drinfeld'84 ($k = 2, 3$):
 $\exists \rho_1, \dots, \rho_s$ st $(\mathbb{S}^k, E(\rho_1, \dots, \rho_s))$ is expander
 - ▶ All ℓ -products of ρ_i 's, large $\ell = \ell(A, B)$
- ▶ For circle squaring

Selecting $\gamma_1, \dots, \gamma_n$

- ▶ For $A, B \subseteq \mathbb{S}^k$, $k \geq 2$:
 - ▶ Margulis'80, Sullivan'81 ($k \geq 4$), Drinfeld'84 ($k = 2, 3$):
 $\exists \rho_1, \dots, \rho_s$ st $(\mathbb{S}^k, E(\rho_1, \dots, \rho_s))$ is expander
 - ▶ All ℓ -products of ρ_i 's, large $\ell = \ell(A, B)$
- ▶ For circle squaring
 - ▶ Work in $[0, 1)^k \pmod{1}$

Selecting $\gamma_1, \dots, \gamma_n$

- ▶ For $A, B \subseteq \mathbb{S}^k$, $k \geq 2$:
 - ▶ Margulis'80, Sullivan'81 ($k \geq 4$), Drinfeld'84 ($k = 2, 3$):
 $\exists \rho_1, \dots, \rho_s$ st $(\mathbb{S}^k, E(\rho_1, \dots, \rho_s))$ is expander
 - ▶ All ℓ -products of ρ_i 's, large $\ell = \ell(A, B)$
- ▶ For circle squaring
 - ▶ Work in $[0, 1)^k \pmod{1}$
 - ▶ Random $\mathbf{x}_1, \dots, \mathbf{x}_d$, large $d = d(A, B)$

Selecting $\gamma_1, \dots, \gamma_n$

- ▶ For $A, B \subseteq \mathbb{S}^k$, $k \geq 2$:
 - ▶ Margulis'80, Sullivan'81 ($k \geq 4$), Drinfeld'84 ($k = 2, 3$):
 $\exists \rho_1, \dots, \rho_s$ st $(\mathbb{S}^k, E(\rho_1, \dots, \rho_s))$ is expander
 - ▶ All ℓ -products of ρ_i 's, large $\ell = \ell(A, B)$
- ▶ For circle squaring
 - ▶ Work in $[0, 1)^k \pmod{1}$
 - ▶ Random $\mathbf{x}_1, \dots, \mathbf{x}_d$, large $d = d(A, B)$
 - ▶ Translate by $\{n_1 \mathbf{x}_1 + \dots + n_d \mathbf{x}_d : \|\mathbf{n}\|_\infty < M\}$, $M \gg d$

Open Problems

Open Problems

- ▶ **Minimum** number of pieces for circle squaring?

Open Problems

- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92:** *“a rough estimate”* is 10^{40}

Open Problems

- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92**: “*a rough estimate*” is 10^{40}
 - ▶ **Marks-Unger \geq '17**: $\leq 10^{200}$

Open Problems

- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92:** "*a rough estimate*" is 10^{40}
 - ▶ **Marks-Unger \geq '17:** $\leq 10^{200}$
 - ▶ **Lower bound:**

Open Problems

- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92:** “*a rough estimate*” is 10^{40}
 - ▶ **Marks-Unger \geq '17:** $\leq 10^{200}$
 - ▶ **Lower bound:**
 - ▶ 3 (if rotations allowed)

Open Problems

- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92:** “*a rough estimate*” is 10^{40}
 - ▶ **Marks-Unger \geq '17:** $\leq 10^{200}$
 - ▶ **Lower bound:**
 - ▶ 3 (if rotations allowed)
 - ▶ 4 (for translations only)

Open Problems

- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92**: “*a rough estimate*” is 10^{40}
 - ▶ **Marks-Unger \geq '17**: $\leq 10^{200}$
 - ▶ **Lower bound**:
 - ▶ 3 (if rotations allowed)
 - ▶ 4 (for translations only)
- ▶ **Banach-Tarski'24**: 5 parts for $\mathbb{B}^3 \cong \mathbb{B}^3 \sqcup \mathbb{B}^3$

Open Problems

- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92**: “*a rough estimate*” is 10^{40}
 - ▶ **Marks-Unger \geq '17**: $\leq 10^{200}$
 - ▶ **Lower bound**:
 - ▶ 3 (if rotations allowed)
 - ▶ 4 (for translations only)
- ▶ **Banach-Tarski'24**: 5 parts for $\mathbb{B}^3 \cong \mathbb{B}^3 \sqcup \mathbb{B}^3$
- ▶ **Grabowski-Máthé-P. \geq '17**: $\leq 2^{2^{75}}$ for $\mathbb{B}^3 \sim \text{cube}$

Open Problems

- ▶ **Minimum** number of pieces for circle squaring?
 - ▶ **Laczkovich'92**: “a rough estimate” is 10^{40}
 - ▶ **Marks-Unger \geq '17**: $\leq 10^{200}$
 - ▶ **Lower bound**:
 - ▶ 3 (if rotations allowed)
 - ▶ 4 (for translations only)
- ▶ **Banach-Tarski'24**: 5 parts for $\mathbb{B}^3 \cong \mathbb{B}^3 \sqcup \mathbb{B}^3$
- ▶ **Grabowski-Máthé-P. \geq '17**: $\leq 2^{2^{75}}$ for $\mathbb{B}^3 \sim$ cube
- ▶ **Open**: $A \sim B, \mu(A) = \mu(B) \Rightarrow A \sim B$ measurably ?

Thank you!