

# Minimizing the number of triangles in graphs of given order and size

Hong Liu, Oleg Pikhurko, Katherine Staden  
University of Warwick

NSFOCS 2017, Novi Sad

# (Erdős-) Rademacher (-Turán) Problem

# (Erdős-) Rademacher (-Turán) Problem

- ▶ Turán function  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$

# (Erdős-) Rademacher (-Turán) Problem

- ▶ Turán function  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$
- ▶ Turán graph  $T_r(n)$ : maximum  $r$ -partite order- $n$

# (Erdős-) Rademacher (-Turán) Problem

- ▶ Turán function  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$
- ▶ Turán graph  $T_r(n)$ : maximum  $r$ -partite order- $n$ 
  - ▶  $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$

# (Erdős-) Rademacher (-Turán) Problem

- ▶ Turán function  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$
- ▶ Turán graph  $T_r(n)$ : maximum  $r$ -partite order- $n$ 
  - ▶  $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$
- ▶  $t_r(n) = e(T_r(n))$

# (Erdős-) Rademacher (-Turán) Problem

- ▶ Turán function  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$
- ▶ Turán graph  $T_r(n)$ : maximum  $r$ -partite order- $n$ 
  - ▶  $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$
- ▶  $t_r(n) = e(T_r(n))$
- ▶ Turán'41:

# (Erdős-) Rademacher (-Turán) Problem

- ▶ Turán function  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$
- ▶ Turán graph  $T_r(n)$ : maximum  $r$ -partite order- $n$ 
  - ▶  $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$
- ▶  $t_r(n) = e(T_r(n))$
- ▶ Turán'41:
  - ▶  $\text{ex}(n, K_{r+1}) = t_r(n)$



# (Erdős-) Rademacher (-Turán) Problem

- ▶ Turán function  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$
- ▶ Turán graph  $T_r(n)$ : maximum  $r$ -partite order- $n$ 
  - ▶  $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$
- ▶  $t_r(n) = e(T_r(n))$
- ▶ Turán'41:
  - ▶  $\text{ex}(n, K_{r+1}) = t_r(n)$
  - ▶  $T_r(n)$ : unique extremal graph

# (Erdős-) Rademacher (-Turán) Problem

- ▶ Turán function  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$
- ▶ Turán graph  $T_r(n)$ : maximum  $r$ -partite order- $n$ 
  - ▶  $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$
- ▶  $t_r(n) = e(T_r(n))$
- ▶ Turán'41:
  - ▶  $\text{ex}(n, K_{r+1}) = t_r(n)$
  - ▶  $T_r(n)$ : unique extremal graph
  - ▶  $r = 2$  was done by Mantel'06

# (Erdős-) Rademacher (-Turán) Problem

- ▶ **Turán function**  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$
- ▶ **Turán graph**  $T_r(n)$ : maximum  $r$ -partite order- $n$ 
  - ▶  $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$
- ▶  $t_r(n) = e(T_r(n))$
- ▶ **Turán'41:**
  - ▶  $\text{ex}(n, K_{r+1}) = t_r(n)$
  - ▶  $T_r(n)$ : unique extremal graph
  - ▶  $r = 2$  was done by **Mantel'06**
- ▶ **Rademacher'41:**  $e(G_n) \geq t_2(n) + 1$

# (Erdős-) Rademacher (-Turán) Problem

- ▶ **Turán function**  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$
- ▶ **Turán graph**  $T_r(n)$ : maximum  $r$ -partite order- $n$ 
  - ▶  $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$
- ▶  $t_r(n) = e(T_r(n))$
- ▶ **Turán'41:**
  - ▶  $\text{ex}(n, K_{r+1}) = t_r(n)$
  - ▶  $T_r(n)$ : unique extremal graph
  - ▶  $r = 2$  was done by **Mantel'06**
- ▶ **Rademacher'41:**  $e(G_n) \geq t_2(n) + 1 \Rightarrow \geq \lfloor \frac{n}{2} \rfloor K_3$ 's

# (Erdős-) Rademacher (-Turán) Problem

- ▶ Turán function  $\text{ex}(n, F) := \max\{e(G_n) : G_n \not\supseteq F\}$
- ▶ Turán graph  $T_r(n)$ : maximum  $r$ -partite order- $n$ 
  - ▶  $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$
- ▶  $t_r(n) = e(T_r(n))$
- ▶ Turán'41:
  - ▶  $\text{ex}(n, K_{r+1}) = t_r(n)$
  - ▶  $T_r(n)$ : unique extremal graph
  - ▶  $r = 2$  was done by Mantel'06
- ▶ Rademacher'41:  $e(G_n) \geq t_2(n) + 1 \Rightarrow \geq \lfloor \frac{n}{2} \rfloor K_3$ 's
- ▶  $g(n, m) = \min\{\#K_3(G_n) : e(G_n) = m\}$

# (Erdős-) Rademacher (-Turán) Problem

- ▶ Turán function  $\text{ex}(n, F) := \max\{ e(G_n) : G_n \not\supseteq F \}$
- ▶ Turán graph  $T_r(n)$ : maximum  $r$ -partite order- $n$ 
  - ▶  $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$
- ▶  $t_r(n) = e(T_r(n))$
- ▶ Turán'41:
  - ▶  $\text{ex}(n, K_{r+1}) = t_r(n)$
  - ▶  $T_r(n)$ : unique extremal graph
  - ▶  $r = 2$  was done by Mantel'06
- ▶ Rademacher'41:  $e(G_n) \geq t_2(n) + 1 \Rightarrow \geq \lfloor \frac{n}{2} \rfloor K_3$ 's
- ▶  $g(n, m) = \min\{ \#K_3(G_n) : e(G_n) = m \}$
- ▶ Maximising  $\#K_3$  is easy (Kruskal'63, Katona'68)

Just above  $t_2(n)$

Just above  $t_2(n)$

- ▶ When is  $g(n, t_2(n) + q) = q \cdot \lfloor \frac{n}{2} \rfloor$  ?



Just above  $t_2(n)$

- ▶ When is  $g(n, t_2(n) + q) = q \cdot \lfloor \frac{n}{2} \rfloor$  ?
  - ▶ Erdős'55:  $q \leq 3$

## Just above $t_2(n)$

- ▶ When is  $g(n, t_2(n) + q) = q \cdot \lfloor \frac{n}{2} \rfloor$  ?
  - ▶ Erdős'55:  $q \leq 3$
  - ▶ Erdős'62:  $q \leq \varepsilon n$

## Just above $t_2(n)$

- ▶ When is  $g(n, t_2(n) + q) = q \cdot \lfloor \frac{n}{2} \rfloor$  ?
  - ▶ Erdős'55:  $q \leq 3$
  - ▶ Erdős'62:  $q \leq \varepsilon n$
  - ▶ Erdős'55: True for  $q < n/2$  ?

## Just above $t_2(n)$

- ▶ When is  $g(n, t_2(n) + q) = q \cdot \lfloor \frac{n}{2} \rfloor$  ?
  - ▶ Erdős'55:  $q \leq 3$
  - ▶ Erdős'62:  $q \leq \varepsilon n$
  - ▶ Erdős'55: True for  $q < n/2$  ?
    - ▶  $K_{k,k} + q$  edges versus  $K_{k+1,k-1} + (q+1)$  edges

# Just above $t_2(n)$

- ▶ When is  $g(n, t_2(n) + q) = q \cdot \lfloor \frac{n}{2} \rfloor$  ?
  - ▶ Erdős'55:  $q \leq 3$
  - ▶ Erdős'62:  $q \leq \varepsilon n$
  - ▶ Erdős'55: True for  $q < n/2$  ?
    - ▶  $K_{k,k} + q$  edges versus  $K_{k+1,k-1} + (q+1)$  edges
    - ▶  $qk$  versus  $(q+1)(k-1)$  triangles

# Just above $t_2(n)$

- ▶ When is  $g(n, t_2(n) + q) = q \cdot \lfloor \frac{n}{2} \rfloor$  ?
  - ▶ Erdős'55:  $q \leq 3$
  - ▶ Erdős'62:  $q \leq \varepsilon n$
  - ▶ Erdős'55: True for  $q < n/2$  ?
    - ▶  $K_{k,k} + q$  edges versus  $K_{k+1,k-1} + (q+1)$  edges
    - ▶  $qk$  versus  $(q+1)(k-1)$  triangles
- ▶ Lovász-Simonovits'75:  $q < n/2$

## Just above $t_2(n)$

- ▶ When is  $g(n, t_2(n) + q) = q \cdot \lfloor \frac{n}{2} \rfloor$  ?
  - ▶ Erdős'55:  $q \leq 3$
  - ▶ Erdős'62:  $q \leq \varepsilon n$
  - ▶ Erdős'55: True for  $q < n/2$  ?
    - ▶  $K_{k,k} + q$  edges versus  $K_{k+1,k-1} + (q+1)$  edges
    - ▶  $qk$  versus  $(q+1)(k-1)$  triangles
  - ▶ Lovász-Simonovits'75:  $q < n/2$
- ▶ Lovász-Simonovits'83: Solved  $q \leq \varepsilon n^2$

General  $m$



# General $m$

- ▶ **Conjecture (Lovász-Simonovits'75):** For  $n \geq n_0$  and all  $m$ ,  $g(n, m)$  is attained by a graph in  $\mathcal{H}$

# General $m$

- ▶ **Conjecture (Lovász-Simonovits'75):** For  $n \geq n_0$  and all  $m$ ,  $g(n, m)$  is attained by a graph in  $\mathcal{H}$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$

# General $m$

- ▶ **Conjecture (Lovász-Simonovits'75):** For  $n \geq n_0$  and all  $m$ ,  $g(n, m)$  is attained by a graph in  $\mathcal{H}$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $t_2(n) \leq m \leq t_3(n)$ :

# General $m$

- ▶ **Conjecture (Lovász-Simonovits'75):** For  $n \geq n_0$  and all  $m$ ,  $g(n, m)$  is attained by a graph in  $\mathcal{H}$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $t_2(n) \leq m \leq t_3(n)$ :
  - ▶  $g(n, m) = \min\{\#K_3 : \text{complete bipartite} + K_3\text{-free}\}$

# General $m$

- ▶ **Conjecture (Lovász-Simonovits'75):** For  $n \geq n_0$  and all  $m$ ,  $g(n, m)$  is attained by a graph in  $\mathcal{H}$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $t_2(n) \leq m \leq t_3(n)$ :
  - ▶  $g(n, m) = \min\{\#K_3 : \text{complete bipartite} + K_3\text{-free}\}$
  - ▶ **Concave** in  $[t_2(n), t_3(n)]$

# General $m$

- ▶ **Conjecture (Lovász-Simonovits'75):** For  $n \geq n_0$  and all  $m$ ,  $g(n, m)$  is attained by a graph in  $\mathcal{H}$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $t_2(n) \leq m \leq t_3(n)$ :
  - ▶  $g(n, m) = \min\{\#K_3 : \text{complete bipartite} + K_3\text{-free}\}$
  - ▶ **Concave** in  $[t_2(n), t_3(n)]$
- ▶  $t_3(n) \leq m \leq t_4(n)$ :

# General $m$

- ▶ **Conjecture (Lovász-Simonovits'75):** For  $n \geq n_0$  and all  $m$ ,  $g(n, m)$  is attained by a graph in  $\mathcal{H}$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $t_2(n) \leq m \leq t_3(n)$ :
  - ▶  $g(n, m) = \min\{\#K_3 : \text{complete bipartite} + K_3\text{-free}\}$
  - ▶ **Concave** in  $[t_2(n), t_3(n)]$
- ▶  $t_3(n) \leq m \leq t_4(n)$ :
  - ▶ Complete 3-partite +  $K_3$ -free

# General $m$

- ▶ **Conjecture (Lovász-Simonovits'75):** For  $n \geq n_0$  and all  $m$ ,  $g(n, m)$  is attained by a graph in  $\mathcal{H}$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $t_2(n) \leq m \leq t_3(n)$ :
  - ▶  $g(n, m) = \min\{\#K_3 : \text{complete bipartite} + K_3\text{-free}\}$
  - ▶ **Concave** in  $[t_2(n), t_3(n)]$
- ▶  $t_3(n) \leq m \leq t_4(n)$ :
  - ▶ Complete 3-partite +  $K_3$ -free
- ▶ **Singularity:**



# General $m$

- ▶ **Conjecture (Lovász-Simonovits'75):** For  $n \geq n_0$  and all  $m$ ,  $g(n, m)$  is attained by a graph in  $\mathcal{H}$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $t_2(n) \leq m \leq t_3(n)$ :
  - ▶  $g(n, m) = \min\{\#K_3 : \text{complete bipartite} + K_3\text{-free}\}$
  - ▶ **Concave** in  $[t_2(n), t_3(n)]$
- ▶  $t_3(n) \leq m \leq t_4(n)$ :
  - ▶ Complete 3-partite +  $K_3$ -free
- ▶ **Singularity:**
  - ▶ Below  $t_3(n)$ :  $n/3$  triangles per edge

# General $m$

- ▶ **Conjecture (Lovász-Simonovits'75):** For  $n \geq n_0$  and all  $m$ ,  $g(n, m)$  is attained by a graph in  $\mathcal{H}$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $t_2(n) \leq m \leq t_3(n)$ :
  - ▶  $g(n, m) = \min\{\#K_3 : \text{complete bipartite} + K_3\text{-free}\}$
  - ▶ **Concave** in  $[t_2(n), t_3(n)]$
- ▶  $t_3(n) \leq m \leq t_4(n)$ :
  - ▶ Complete 3-partite +  $K_3$ -free
- ▶ **Singularity:**
  - ▶ Below  $t_3(n)$ :  $n/3$  triangles per edge
  - ▶ Above  $t_3(n)$ :  $2n/3$  triangles per edge

# Asymptotic Version

# Asymptotic Version

$$\blacktriangleright g(a) = \lim_{n \rightarrow \infty} \frac{g\left(n, a\binom{n}{2}\right)}{\binom{n}{3}}$$

# Asymptotic Version

- ▶  $g(a) = \lim_{n \rightarrow \infty} \frac{g\left(n, a\binom{n}{2}\right)}{\binom{n}{3}}$
- ▶ Conjecture + calculations:

# Asymptotic Version

- ▶  $g(a) = \lim_{n \rightarrow \infty} \frac{g\left(n, a\binom{n}{2}\right)}{\binom{n}{3}}$
- ▶ Conjecture + calculations:
  - ▶  $g(a): \#K_3$  in  $K_{cn, \dots, cn, (1-tc)n}$

# Asymptotic Version

- ▶  $g(a) = \lim_{n \rightarrow \infty} \frac{g(n, a\binom{n}{2})}{\binom{n}{3}}$
- ▶ Conjecture + calculations:
  - ▶  $g(a): \#K_3$  in  $K_{cn, \dots, cn, (1-tc)n}$
- ▶ Partial results: Goodman'59, Nordhaus-Stewart'63, Bollobás'76, Lovász-Simonovits'83, ...

# Asymptotic Version

- ▶  $g(a) = \lim_{n \rightarrow \infty} \frac{g(n, a \binom{n}{2})}{\binom{n}{3}}$
- ▶ Conjecture + calculations:
  - ▶  $g(a)$ :  $\#K_3$  in  $K_{cn, \dots, cn, (1-tc)n}$
- ▶ Partial results: Goodman'59, Nordhaus-Stewart'63, Bollobás'76, Lovász-Simonovits'83, ...
- ▶ Fisher'89: Determined  $g(a)$  for  $\frac{1}{2} \leq a \leq \frac{2}{3}$




# Asymptotic Version

- ▶  $g(a) = \lim_{n \rightarrow \infty} \frac{g(n, a \binom{n}{2})}{\binom{n}{3}}$
- ▶ Conjecture + calculations:
  - ▶  $g(a): \#K_3$  in  $K_{cn, \dots, cn, (1-tc)n}$
- ▶ Partial results: Goodman'59, Nordhaus-Stewart'63, Bollobás'76, Lovász-Simonovits'83, ...
- ▶ Fisher'89: Determined  $g(a)$  for  $\frac{1}{2} \leq a \leq \frac{2}{3}$
- ▶ Razborov'07,08: Determined  $g(a)$  for all  $a$

# Asymptotic Version

- ▶  $g(a) = \lim_{n \rightarrow \infty} \frac{g(n, a \binom{n}{2})}{\binom{n}{3}}$
- ▶ Conjecture + calculations:
  - ▶  $g(a)$ :  $\#K_3$  in  $K_{cn, \dots, cn, (1-tc)n}$
- ▶ Partial results: Goodman'59, Nordhaus-Stewart'63, Bollobás'76, Lovász-Simonovits'83, ...
- ▶ Fisher'89: Determined  $g(a)$  for  $\frac{1}{2} \leq a \leq \frac{2}{3}$
- ▶ Razborov'07,08: Determined  $g(a)$  for all  $a$
- ▶ Nikiforov'11: different proof

# Possible Edge-Triangle Densities in Limit



# Structure of extremal graphs

# Structure of extremal graphs

- ▶ Recall construction:

# Structure of extremal graphs

► Recall construction:

►  $e(K_{cn, \dots, cn, (1-tc)n}) \approx a\binom{n}{2}$

# Structure of extremal graphs

- ▶ **Recall** construction:
  - ▶  $e(K_{cn, \dots, cn, (1-tc)n}) \approx a \binom{n}{2}$
  - ▶  $H_n^a$ : modify the last two parts, keeping  $K_3$ -free

# Structure of extremal graphs

- ▶ Recall construction:
  - ▶  $e(K_{cn, \dots, cn, (1-tc)n}) \approx a\binom{n}{2}$
  - ▶  $H_n^a$ : modify the last two parts, keeping  $K_3$ -free
  - ▶ Many different graphs



# Structure of extremal graphs

- ▶ **Recall** construction:
  - ▶  $e(K_{cn, \dots, cn, (1-tc)n}) \approx a \binom{n}{2}$
  - ▶  $H_n^a$ : modify the last two parts, keeping  $K_3$ -free
  - ▶ **Many** different graphs
- ▶ **P.-Razborov'17:**  
 $\forall$  almost extremal  $G_n$  is  $o(n^2)$ -close to some  $H_n^a$

Exact Value of  $g(n, m)$

# Exact Value of $g(n, m)$

- ▶ Goodman'59, Nordhaus-Stewart'63:  $m = t_k(n)$

# Exact Value of $g(n, m)$

- ▶ Goodman'59, Nordhaus-Stewart'63:  $m = t_k(n)$
- ▶ Lovász-Simonovits'83:  $t_k(n) \leq m \leq t_k(n) + \varepsilon_k n^2$

# Exact Value of $g(n, m)$

- ▶ Goodman'59, Nordhaus-Stewart'63:  $m = t_k(n)$
- ▶ Lovász-Simonovits'83:  $t_k(n) \leq m \leq t_k(n) + \varepsilon_k n^2$
- ▶ Nikiforov'11:  $n = ta + b, a \geq b, m = e(K_{a, \dots, a, b})$

# Exact Value of $g(n, m)$

- ▶ Goodman'59, Nordhaus-Stewart'63:  $m = t_k(n)$
- ▶ Lovász-Simonovits'83:  $t_k(n) \leq m \leq t_k(n) + \varepsilon_k n^2$
- ▶ Nikiforov'11:  $n = ta + b, a \geq b, m = e(K_{a,\dots,a,b})$
- ▶ Hong-P.-Staden  $\geq$ '17:  $m < \binom{n}{2} - o(n^2)$

# Exact Value of $g(n, m)$

- ▶ Goodman'59, Nordhaus-Stewart'63:  $m = t_k(n)$
- ▶ Lovász-Simonovits'83:  $t_k(n) \leq m \leq t_k(n) + \varepsilon_k n^2$
- ▶ Nikiforov'11:  $n = ta + b$ ,  $a \geq b$ ,  $m = e(K_{a,\dots,a,b})$
- ▶ Hong-P.-Staden  $\geq$ '17:  $m < \binom{n}{2} - o(n^2)$ 
  - ▶  $\forall \varepsilon > 0 \exists n_0 \forall n \geq n_0 \forall m \leq \binom{n}{2} - \varepsilon n^2$ , at least one  $g(n, m)$ -extremal graph is in  $\mathcal{H}$

# Exact Value of $g(n, m)$

- ▶ Goodman'59, Nordhaus-Stewart'63:  $m = t_k(n)$
- ▶ Lovász-Simonovits'83:  $t_k(n) \leq m \leq t_k(n) + \varepsilon_k n^2$
- ▶ Nikiforov'11:  $n = ta + b$ ,  $a \geq b$ ,  $m = e(K_{a,\dots,a,b})$
- ▶ Hong-P.-Staden  $\geq$ '17:  $m < \binom{n}{2} - o(n^2)$ 
  - ▶  $\forall \varepsilon > 0 \exists n_0 \forall n \geq n_0 \forall m \leq \binom{n}{2} - \varepsilon n^2$ , at least one  $g(n, m)$ -extremal graph is in  $\mathcal{H}$
- ▶ Not all extremal graphs are in  $\mathcal{H}$



# Exact Value of $g(n, m)$

- ▶ Goodman'59, Nordhaus-Stewart'63:  $m = t_k(n)$
- ▶ Lovász-Simonovits'83:  $t_k(n) \leq m \leq t_k(n) + \varepsilon_k n^2$
- ▶ Nikiforov'11:  $n = ta + b, a \geq b, m = e(K_{a,\dots,a,b})$
- ▶ Hong-P.-Staden  $\geq$ '17:  $m < \binom{n}{2} - o(n^2)$ 
  - ▶  $\forall \varepsilon > 0 \exists n_0 \forall n \geq n_0 \forall m \leq \binom{n}{2} - \varepsilon n^2$ , at least one  $g(n, m)$ -extremal graph is in  $\mathcal{H}$
- ▶ Not all extremal graphs are in  $\mathcal{H}$ 
  - ▶  $\mathcal{H}_0 := \{\text{complete partite} + K_3\text{-free into some parts}\}$

# Exact Value of $g(n, m)$

- ▶ Goodman'59, Nordhaus-Stewart'63:  $m = t_k(n)$
- ▶ Lovász-Simonovits'83:  $t_k(n) \leq m \leq t_k(n) + \varepsilon_k n^2$
- ▶ Nikiforov'11:  $n = ta + b, a \geq b, m = e(K_{a,\dots,a,b})$
- ▶ Hong-P.-Staden  $\geq$ '17:  $m < \binom{n}{2} - o(n^2)$ 
  - ▶  $\forall \varepsilon > 0 \exists n_0 \forall n \geq n_0 \forall m \leq \binom{n}{2} - \varepsilon n^2$ , at least one  $g(n, m)$ -extremal graph is in  $\mathcal{H}$
- ▶ Not all extremal graphs are in  $\mathcal{H}$ 
  - ▶  $\mathcal{H}_0 := \{\text{complete partite} + K_3\text{-free into some parts}\}$
  - ▶  $\mathcal{H}_1 := \{\text{complete partite} - \text{edges incident to one part}\}$

# Exact Value of $g(n, m)$

- ▶ Goodman'59, Nordhaus-Stewart'63:  $m = t_k(n)$
- ▶ Lovász-Simonovits'83:  $t_k(n) \leq m \leq t_k(n) + \varepsilon_k n^2$
- ▶ Nikiforov'11:  $n = ta + b$ ,  $a \geq b$ ,  $m = e(K_{a,\dots,a,b})$
- ▶ Hong-P.-Staden  $\geq$ '17:  $m < \binom{n}{2} - o(n^2)$ 
  - ▶  $\forall \varepsilon > 0 \exists n_0 \forall n \geq n_0 \forall m \leq \binom{n}{2} - \varepsilon n^2$ , at least one  $g(n, m)$ -extremal graph is in  $\mathcal{H}$
- ▶ Not all extremal graphs are in  $\mathcal{H}$ 
  - ▶  $\mathcal{H}_0 := \{\text{complete partite} + K_3\text{-free into some parts}\}$
  - ▶  $\mathcal{H}_1 := \{\text{complete partite} - \text{edges incident to one part}\}$
- ▶ Hong-P.-Staden  $\geq$ '17: For  $m < \binom{n}{2} - o(n^2)$ , every extremal graph is in  $\mathcal{H}_0 \cup \mathcal{H}_1$

# Exact Value of $g(n, m)$

- ▶ Goodman'59, Nordhaus-Stewart'63:  $m = t_k(n)$
- ▶ Lovász-Simonovits'83:  $t_k(n) \leq m \leq t_k(n) + \varepsilon_k n^2$
- ▶ Nikiforov'11:  $n = ta + b$ ,  $a \geq b$ ,  $m = e(K_{a,\dots,a,b})$
- ▶ Hong-P.-Staden  $\geq$ '17:  $m < \binom{n}{2} - o(n^2)$ 
  - ▶  $\forall \varepsilon > 0 \exists n_0 \forall n \geq n_0 \forall m \leq \binom{n}{2} - \varepsilon n^2$ , at least one  $g(n, m)$ -extremal graph is in  $\mathcal{H}$
- ▶ Not all extremal graphs are in  $\mathcal{H}$ 
  - ▶  $\mathcal{H}_0 := \{\text{complete partite} + K_3\text{-free into some parts}\}$
  - ▶  $\mathcal{H}_1 := \{\text{complete partite} - \text{edges incident to one part}\}$
- ▶ Hong-P.-Staden  $\geq$ '17: For  $m < \binom{n}{2} - o(n^2)$ , every extremal graph is in  $\mathcal{H}_0 \cup \mathcal{H}_1$
- ▶ Open: exact result for all  $m$

# Beginning of Proof

# Beginning of Proof

- ▶  $n \geq n_0$

# Beginning of Proof

- ▶  $n \geq n_0$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$

# Beginning of Proof

- ▶  $n \geq n_0$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $h(n, m) := \min\{ \#K_3(G_n) : G_n \in \mathcal{H}, e(G_n) = m \}$



# Beginning of Proof

- ▶  $n \geq n_0$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $h(n, m) := \min\{ \#K_3(G_n) : G_n \in \mathcal{H}, e(G_n) = m \}$
- ▶ **Optimal** choice:

# Beginning of Proof

- ▶  $n \geq n_0$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $h(n, m) := \min\{\#K_3(G_n) : G_n \in \mathcal{H}, e(G_n) = m\}$
- ▶ **Optimal** choice:
  - ▶ Minimise the number of parts

# Beginning of Proof

- ▶  $n \geq n_0$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $h(n, m) := \min\{\#K_3(G_n) : G_n \in \mathcal{H}, e(G_n) = m\}$
- ▶ **Optimal** choice:
  - ▶ Minimise the number of parts
  - ▶ Minimise the smallest part size

# Beginning of Proof

- ▶  $n \geq n_0$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $h(n, m) := \min\{\#K_3(G_n) : G_n \in \mathcal{H}, e(G_n) = m\}$
- ▶ **Optimal** choice:
  - ▶ Minimise the number of parts
  - ▶ Minimise the smallest part size
- ▶ Consider  $m \in [t_2(n), t_3(n)]$

# Beginning of Proof

- ▶  $n \geq n_0$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $h(n, m) := \min\{\#K_3(G_n) : G_n \in \mathcal{H}, e(G_n) = m\}$
- ▶ **Optimal** choice:
  - ▶ Minimise the number of parts
  - ▶ Minimise the smallest part size
- ▶ Consider  $m \in [t_2(n), t_3(n)]$ 
  - ▶  $m \leq t_2(n)$ :  $g(n, m) = h(n, m) = 0$

# Beginning of Proof

- ▶  $n \geq n_0$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $h(n, m) := \min\{\#K_3(G_n) : G_n \in \mathcal{H}, e(G_n) = m\}$
- ▶ **Optimal** choice:
  - ▶ Minimise the number of parts
  - ▶ Minimise the smallest part size
- ▶ Consider  $m \in [t_2(n), t_3(n)]$ 
  - ▶  $m \leq t_2(n)$ :  $g(n, m) = h(n, m) = 0$
- ▶ **Worst counterexample  $G$** : minimise  $\#K_3(G) - h(n, m)$  with  $n = v(G)$  and  $m = e(G) \leq t_3(n)$

# Beginning of Proof

- ▶  $n \geq n_0$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $h(n, m) := \min\{\#K_3(G_n) : G_n \in \mathcal{H}, e(G_n) = m\}$
- ▶ **Optimal** choice:
  - ▶ Minimise the number of parts
  - ▶ Minimise the smallest part size
- ▶ Consider  $m \in [t_2(n), t_3(n)]$ 
  - ▶  $m \leq t_2(n)$ :  $g(n, m) = h(n, m) = 0$
- ▶ **Worst counterexample  $G$** : minimise  $\#K_3(G) - h(n, m)$  with  $n = v(G)$  and  $m = e(G) \leq t_3(n)$
- ▶ By **Lovász-Simonovits'83**, assume  $m \geq t_2(n) + \varepsilon n^2$ :

# Beginning of Proof

- ▶  $n \geq n_0$
- ▶  $\mathcal{H} := \{\text{complete partite} + K_3\text{-free inside a part}\}$
- ▶  $h(n, m) := \min\{\#K_3(G_n) : G_n \in \mathcal{H}, e(G_n) = m\}$
- ▶ **Optimal** choice:
  - ▶ Minimise the number of parts
  - ▶ Minimise the smallest part size
- ▶ Consider  $m \in [t_2(n), t_3(n)]$ 
  - ▶  $m \leq t_2(n)$ :  $g(n, m) = h(n, m) = 0$
- ▶ **Worst counterexample  $G$** : minimise  $\#K_3(G) - h(n, m)$  with  $n = v(G)$  and  $m = e(G) \leq t_3(n)$
- ▶ By **Lovász-Simonovits'83**, assume  $m \geq t_2(n) + \varepsilon n^2$ :
- ▶ **Case 1**:  $t_2(n) + \varepsilon n^2 \leq m \leq t_3(n) - \Omega(n^2)$



Worst counterexample  $G = (V, E)$

## Worst counterexample $G = (V, E)$

►  $c : e(K_{cn, cn, (1-2c)n}) = m$

## Worst counterexample $G = (V, E)$

- ▶  $c : e(K_{cn, cn, (1-2c)n}) = m$
- ▶  $h(n, -)$ : almost linear with slope  $cn + O(1)$

## Worst counterexample $G = (V, E)$

- ▶  $c : e(K_{cn, cn, (1-2c)n}) = m$
- ▶  $h(n, -)$ : almost linear with slope  $cn + O(1)$
- ▶ Add edge:

# Worst counterexample $G = (V, E)$

- ▶  $c : e(K_{cn, cn, (1-2c)n}) = m$
- ▶  $h(n, -)$ : almost linear with slope  $cn + O(1)$
- ▶ **Add** edge:
  - ▶  $\forall xy \in E \quad \#K_3(G; xy) \leq cn + O(1)$

# Worst counterexample $G = (V, E)$

- ▶  $c : e(K_{cn, cn, (1-2c)n}) = m$
- ▶  $h(n, -)$ : almost linear with slope  $cn + O(1)$
- ▶ **Add** edge:
  - ▶  $\forall xy \in E \quad \#K_3(G; xy) \leq cn + O(1)$
- ▶ **Add** vertex and **remove** another:

# Worst counterexample $G = (V, E)$

- ▶  $c : e(K_{cn, cn, (1-2c)n}) = m$
- ▶  $h(n, -)$ : almost linear with slope  $cn + O(1)$
- ▶ **Add** edge:
  - ▶  $\forall xy \in E \quad \#K_3(G; xy) \leq cn + O(1)$
- ▶ **Add** vertex and **remove** another:
  - ▶  $\forall x \in V \quad f(x) := cn d(x) - \#K_3(G; x) - c^2 n^2 = O(n)$

# Worst counterexample $G = (V, E)$

- ▶  $c : e(K_{cn, cn, (1-2c)n}) = m$
- ▶  $h(n, -)$ : almost linear with slope  $cn + O(1)$
- ▶ **Add** edge:
  - ▶  $\forall xy \in E \quad \#K_3(G; xy) \leq cn + O(1)$
- ▶ **Add** vertex and **remove** another:
  - ▶  $\forall x \in V \quad f(x) := cn d(x) - \#K_3(G; x) - c^2 n^2 = O(n)$
- ▶ **Razborov'07**: limit versions  $\Rightarrow$  asymptotic result



$G$  is close to complete 3-partite

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn,cn,(1-2c)n})| = o(n^2)$

## $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn,cn,(1-2c)n})| = o(n^2)$
- ▶ Suppose **false**

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn,cn,(1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz)$



# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz) \lesssim cn$

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn,cn,(1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz) \lesssim cn$
  - ▶  $d_A \leq |A| \leq cn + o(n)$

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz) \lesssim cn$
  - ▶  $d_A \leq |A| \leq cn + o(n)$
  - ▶  $f(x) \approx (cn - d_A)(d_B - cn) - e(G[N_A]) - e(G[N_B])$

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz) \lesssim cn$
  - ▶  $d_A \leq |A| \leq cn + o(n)$
  - ▶  $f(x) \approx (cn - d_A)(d_B - cn) - e(G[N_A]) - e(G[N_B])$
  - ▶  $\Rightarrow e(G[N_B]) = o(n^2)$

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz) \lesssim cn$
  - ▶  $d_A \leq |A| \leq cn + o(n)$
  - ▶  $f(x) \approx (cn - d_A)(d_B - cn) - e(G[N_A]) - e(G[N_B])$
  - ▶  $\Rightarrow e(G[N_B]) = o(n^2)$
  - ▶  $\Rightarrow d_B \leq cn - \Omega(n)$

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz) \lesssim cn$
  - ▶  $d_A \leq |A| \leq cn + o(n)$
  - ▶  $f(x) \approx (cn - d_A)(d_B - cn) - e(G[N_A]) - e(G[N_B])$
  - ▶  $\Rightarrow e(G[N_B]) = o(n^2)$
  - ▶  $\Rightarrow d_B \leq cn - \Omega(n)$ : o/w Lemma holds

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz) \lesssim cn$
  - ▶  $d_A \leq |A| \leq cn + o(n)$
  - ▶  $f(x) \approx (cn - d_A)(d_B - cn) - e(G[N_A]) - e(G[N_B])$
  - ▶  $\Rightarrow e(G[N_B]) = o(n^2)$
  - ▶  $\Rightarrow d_B \leq cn - \Omega(n)$ : o/w Lemma holds
  - ▶  $\Rightarrow d_A = cn + o(n)$

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz) \lesssim cn$
  - ▶  $d_A \leq |A| \leq cn + o(n)$
  - ▶  $f(x) \approx (cn - d_A)(d_B - cn) - e(G[N_A]) - e(G[N_B])$
  - ▶  $\Rightarrow e(G[N_B]) = o(n^2)$
  - ▶  $\Rightarrow d_B \leq cn - \Omega(n)$ : o/w Lemma holds
  - ▶  $\Rightarrow d_A = cn + o(n)$
  - ▶ **Move**  $x$  to  $B$  and **decrease**  $|E(H) \triangle E(G)| \Rightarrow \Leftarrow$



# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz) \lesssim cn$
  - ▶  $d_A \leq |A| \leq cn + o(n)$
  - ▶  $f(x) \approx (cn - d_A)(d_B - cn) - e(G[N_A]) - e(G[N_B])$
  - ▶  $\Rightarrow e(G[N_B]) = o(n^2)$
  - ▶  $\Rightarrow d_B \leq cn - \Omega(n)$ : o/w Lemma holds
  - ▶  $\Rightarrow d_A = cn + o(n)$
  - ▶ **Move**  $x$  to  $B$  and **decrease**  $|E(H) \triangle E(G)| \Rightarrow \Leftarrow$
- ▶ ...  $\Rightarrow G[A]$  empty,  $G[A, B]$  complete

# $G$ is close to complete 3-partite

- ▶ **Lemma:**  $|E(G) \triangle E(K_{cn, cn, (1-2c)n})| = o(n^2)$
- ▶ Suppose **false**
- ▶  $H := K[A, B] + (K_3\text{-free on } B)$ :  $\min |E(H) \triangle E(G)|$
- ▶ **Claim:**  $\Delta(G[A]) = o(n)$ 
  - ▶ Let  $x \in A$  have  $d_A > \Omega(n)$
  - ▶ typical  $z \in N_A$ :  $d_B \approx \#K_3(xz) \lesssim cn$
  - ▶  $d_A \leq |A| \leq cn + o(n)$
  - ▶  $f(x) \approx (cn - d_A)(d_B - cn) - e(G[N_A]) - e(G[N_B])$
  - ▶  $\Rightarrow e(G[N_B]) = o(n^2)$
  - ▶  $\Rightarrow d_B \leq cn - \Omega(n)$ : o/w Lemma holds
  - ▶  $\Rightarrow d_A = cn + o(n)$
  - ▶ **Move**  $x$  to  $B$  and **decrease**  $|E(H) \triangle E(G)| \Rightarrow \Leftarrow$
- ▶ ...  $\Rightarrow G[A]$  empty,  $G[A, B]$  complete  $\Rightarrow G \in \mathcal{H}$

Sketch of the rest of the proof

# Sketch of the rest of the proof

- ▶ Max-3-cut partition:  $V = V_1 \cup V_2 \cup V_3$

# Sketch of the rest of the proof

- ▶ **Max-3-cut partition:**  $V = V_1 \cup V_2 \cup V_3$
- ▶  $G \approx K[V_1, K_2, V_3]$

# Sketch of the rest of the proof

- ▶ Max-3-cut partition:  $V = V_1 \cup V_2 \cup V_3$
- ▶  $G \approx K[V_1, K_2, V_3]$
- ▶ Local transformations:

# Sketch of the rest of the proof

- ▶ **Max-3-cut partition:**  $V = V_1 \cup V_2 \cup V_3$
- ▶  $G \approx K[V_1, K_2, V_3]$
- ▶ **Local transformations:**
  - ▶  $\sum e(G[V_i]) \ll$  missing edges between parts

# Sketch of the rest of the proof

- ▶ **Max-3-cut partition:**  $V = V_1 \cup V_2 \cup V_3$
- ▶  $G \approx K[V_1, V_2, V_3]$
- ▶ **Local transformations:**
  - ▶  $\sum e(G[V_i]) \ll$  missing edges between parts
  - ▶  $\exists$  small  $Z$  st  $G - Z$  is complete 3-partite



# Sketch of the rest of the proof

- ▶ **Max-3-cut partition:**  $V = V_1 \cup V_2 \cup V_3$
- ▶  $G \approx K[V_1, V_2, V_3]$
- ▶ **Local transformations:**
  - ▶  $\sum e(G[V_i]) \ll$  missing edges between parts
  - ▶  $\exists$  small  $Z$  st  $G - Z$  is complete 3-partite
- ▶ **Global transformation:**

# Sketch of the rest of the proof

- ▶ **Max-3-cut partition:**  $V = V_1 \cup V_2 \cup V_3$
- ▶  $G \approx K[V_1, V_2, V_3]$
- ▶ **Local transformations:**
  - ▶  $\sum e(G[V_i]) \ll$  missing edges between parts
  - ▶  $\exists$  small  $Z$  st  $G - Z$  is complete 3-partite
- ▶ **Global transformation:**
  - ▶  $\exists$  3-partite  $H$  with  $\#K_3(H) - \#K_3(G)$  small

# Sketch of the rest of the proof

- ▶ **Max-3-cut partition:**  $V = V_1 \cup V_2 \cup V_3$
- ▶  $G \approx K[V_1, V_2, V_3]$
- ▶ **Local transformations:**
  - ▶  $\sum e(G[V_i]) \ll$  missing edges between parts
  - ▶  $\exists$  small  $Z$  st  $G - Z$  is complete 3-partite
- ▶ **Global transformation:**
  - ▶  $\exists$  3-partite  $H$  with  $\#K_3(H) - \#K_3(G)$  small
  - ▶ Many missing edges in  $H$

# Sketch of the rest of the proof

- ▶ **Max-3-cut partition:**  $V = V_1 \cup V_2 \cup V_3$
- ▶  $G \approx K[V_1, V_2, V_3]$
- ▶ **Local transformations:**
  - ▶  $\sum e(G[V_i]) \ll$  missing edges between parts
  - ▶  $\exists$  small  $Z$  st  $G - Z$  is complete 3-partite
- ▶ **Global transformation:**
  - ▶  $\exists$  3-partite  $H$  with  $\#K_3(H) - \#K_3(G)$  small
  - ▶ Many missing edges in  $H$
  - ▶ **Compress**  $H$ , losing  $> \#K_3(H) - \#K_3(G)$  triangles

# Sketch of the rest of the proof

- ▶ **Max-3-cut partition:**  $V = V_1 \cup V_2 \cup V_3$
- ▶  $G \approx K[V_1, K_2, V_3]$
- ▶ **Local transformations:**
  - ▶  $\sum e(G[V_i]) \ll$  missing edges between parts
  - ▶  $\exists$  small  $Z$  st  $G - Z$  is complete 3-partite
- ▶ **Global transformation:**
  - ▶  $\exists$  3-partite  $H$  with  $\#K_3(H) - \#K_3(G)$  small
  - ▶ Many missing edges in  $H$
  - ▶ **Compress**  $H$ , losing  $> \#K_3(H) - \#K_3(G)$  triangles
  - ▶  $\Rightarrow \Leftarrow$

# Sketch of the rest of the proof

- ▶ **Max-3-cut partition:**  $V = V_1 \cup V_2 \cup V_3$
- ▶  $G \approx K[V_1, K_2, V_3]$
- ▶ **Local transformations:**
  - ▶  $\sum e(G[V_i]) \ll$  missing edges between parts
  - ▶  $\exists$  small  $Z$  st  $G - Z$  is complete 3-partite
- ▶ **Global transformation:**
  - ▶  $\exists$  3-partite  $H$  with  $\#K_3(H) - \#K_3(G)$  small
  - ▶ Many missing edges in  $H$
  - ▶ **Compress**  $H$ , losing  $> \#K_3(H) - \#K_3(G)$  triangles
  - ▶  $\Rightarrow \Leftarrow$  😊

# Clique minimisation

# Clique minimisation

- ▶ Nikiforov'11: Asymptotic solution for  $K_4$



# Clique minimisation

- ▶ Nikiforov'11: Asymptotic solution for  $K_4$
- ▶ Reiher'17: Asymptotic solution for  $K_r$

# Clique minimisation

- ▶ Nikiforov'11: Asymptotic solution for  $K_4$
- ▶ Reiher'17: Asymptotic solution for  $K_r$
- ▶ Open: structure & exact result

Minimising the number of  $F$ -subgraphs

# Minimising the number of $F$ -subgraphs

- ▶  $F$  bipartite

# Minimising the number of $F$ -subgraphs

- ▶  $F$  bipartite
- ▶ (Erdős-Simonovits)-Sidorenko's Conjecture:

# Minimising the number of $F$ -subgraphs

- ▶  $F$  bipartite
- ▶ (Erdős-Simonovits)-Sidorenko's Conjecture:
  - ▶ Quasi-random graphs are asymptotically optimal

# Minimising the number of $F$ -subgraphs

- ▶  $F$  bipartite
- ▶ (Erdős-Simonovits)-Sidorenko's Conjecture:
  - ▶ Quasi-random graphs are asymptotically optimal
  - ▶ Graphon reformulation

# Minimising the number of $F$ -subgraphs

- ▶  $F$  bipartite
- ▶ (Erdős-Simonovits)-Sidorenko's Conjecture:
  - ▶ Quasi-random graphs are asymptotically optimal
  - ▶ Graphon reformulation
  - ▶  $\geq 2010$ : Conlon, Fox, Hatami, Kim, Lee, Lee, Li, Sudakov, Szegedy, ...



Critical  $F$

# Critical $F$

- ▶  $F$   **$r$ -critical**:  $\chi(F) = r + 1$  &  $\chi(F - e) = r$

# Critical $F$

- ▶  $F$   **$r$ -critical**:  $\chi(F) = r + 1$  &  $\chi(F - e) = r$
- ▶ **Simonovits'68**:  $T_r(n)$  is unique maximum  $F$ -free of order  $n \geq n_0$

# Critical $F$

- ▶  $F$   $r$ -critical:  $\chi(F) = r + 1$  &  $\chi(F - e) = r$
- ▶ Simonovits'68:  $T_r(n)$  is unique maximum  $F$ -free of order  $n \geq n_0$
- ▶  $s_F(n, q) := \min\{\#F(G_n) : e(G_n) = \text{ex}(n, F) + q\}$

# Critical $F$

- ▶  $F$   **$r$ -critical**:  $\chi(F) = r + 1$  &  $\chi(F - e) = r$
- ▶ **Simonovits'68**:  $T_r(n)$  is unique maximum  $F$ -free of order  $n \geq n_0$
- ▶  **$s_F(n, q)$**  :=  $\min\{\#F(G_n) : e(G_n) = \text{ex}(n, F) + q\}$
- ▶ **Erdős'69**:  $g_{C_5}(2m, 1) = m(m-1)(m-2)$

# Critical $F$

- ▶  $F$   **$r$ -critical**:  $\chi(F) = r + 1$  &  $\chi(F - e) = r$
- ▶ **Simonovits'68**:  $T_r(n)$  is unique maximum  $F$ -free of order  $n \geq n_0$
- ▶  **$s_F(n, q)$**  :=  $\min\{\#F(G_n) : e(G_n) = \text{ex}(n, F) + q\}$
- ▶ **Erdős'69**:  $g_{C_5}(2m, 1) = m(m-1)(m-2)$
- ▶  **$s_F^*(n, q)$**  :=  $\min\{\#F : \text{maximum } F\text{-free} + q \text{ edges}\}$

# Optimality of Turán graph + edges

# Optimality of Turán graph + edges

►  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$



# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4$  – edge

# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4$  – edge
- ▶ P.-Yilma'17:

# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4$  – edge
- ▶ P.-Yilma'17:
  - ▶  $\forall$  critical  $F$   $c_1(F) > 0$

# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4$  – edge
- ▶ P.-Yilma'17:
  - ▶  $\forall$  critical  $F$   $c_1(F) > 0$
  - ▶  $c_1(C_{2k+1}) = \frac{1}{2}$

# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4 - \text{edge}$
- ▶ P.-Yilma'17:
  - ▶  $\forall$  critical  $F$   $c_1(F) > 0$
  - ▶  $c_1(C_{2k+1}) = \frac{1}{2}$
  - ▶  $c_1(K_{r+2} - \text{edge}) = \frac{r-1}{r^2}$

# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4 - \text{edge}$
- ▶ P.-Yilma'17:
  - ▶  $\forall$  critical  $F$   $c_1(F) > 0$
  - ▶  $c_1(C_{2k+1}) = \frac{1}{2}$
  - ▶  $c_1(K_{r+2} - \text{edge}) = \frac{r-1}{r^2}$
  - ▶  $c_1(K_{s,t} + e)$

# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4 - \text{edge}$
- ▶ P.-Yilma'17:
  - ▶  $\forall$  critical  $F$   $c_1(F) > 0$
  - ▶  $c_1(C_{2k+1}) = \frac{1}{2}$
  - ▶  $c_1(K_{r+2} - \text{edge}) = \frac{r-1}{r^2}$
  - ▶  $c_1(K_{s,t} + e)$
  - ▶ ...

# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4 - \text{edge}$
- ▶ P.-Yilma'17:
  - ▶  $\forall$  critical  $F$   $c_1(F) > 0$
  - ▶  $c_1(C_{2k+1}) = \frac{1}{2}$
  - ▶  $c_1(K_{r+2} - \text{edge}) = \frac{r-1}{r^2}$
  - ▶  $c_1(K_{s,t} + e)$
  - ▶ ...
- ▶ P.-Yilma'17:  $s_F(n, q) \approx s_F^*(n, q)$  for  $q = o(n^2)$



# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4 - \text{edge}$
- ▶ P.-Yilma'17:
  - ▶  $\forall$  critical  $F$   $c_1(F) > 0$
  - ▶  $c_1(C_{2k+1}) = \frac{1}{2}$
  - ▶  $c_1(K_{r+2} - \text{edge}) = \frac{r-1}{r^2}$
  - ▶  $c_1(K_{s,t} + e)$
  - ▶ ...
- ▶ P.-Yilma'17:  $s_F(n, q) \approx s_F^*(n, q)$  for  $q = o(n^2)$ 
  - ▶ unless  $T_r(n-1) + v$  beats  $T_r(n) + \text{edges}$

# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4 - \text{edge}$
- ▶ P.-Yilma'17:
  - ▶  $\forall$  critical  $F$   $c_1(F) > 0$
  - ▶  $c_1(C_{2k+1}) = \frac{1}{2}$
  - ▶  $c_1(K_{r+2} - \text{edge}) = \frac{r-1}{r^2}$
  - ▶  $c_1(K_{s,t} + e)$
  - ▶ ...
- ▶ P.-Yilma'17:  $s_F(n, q) \approx s_F^*(n, q)$  for  $q = o(n^2)$ 
  - ▶ unless  $T_r(n-1) + v$  beats  $T_r(n) + \text{edges}$
- ▶ Open:  $q = \Omega(n^2)$

# Optimality of Turán graph + edges

- ▶  $c_1(F) := \liminf\{\varepsilon : q \leq \varepsilon n \Rightarrow s_F(n, q) = s_F^*(n, q)\}$
- ▶ Mubayi'10:  $c_1 > 0$  for odd cycles and  $K_4 - \text{edge}$
- ▶ P.-Yilma'17:
  - ▶  $\forall$  critical  $F$   $c_1(F) > 0$
  - ▶  $c_1(C_{2k+1}) = \frac{1}{2}$
  - ▶  $c_1(K_{r+2} - \text{edge}) = \frac{r-1}{r^2}$
  - ▶  $c_1(K_{s,t} + e)$
  - ▶ ...
- ▶ P.-Yilma'17:  $s_F(n, q) \approx s_F^*(n, q)$  for  $q = o(n^2)$ 
  - ▶ unless  $T_r(n-1) + v$  beats  $T_r(n) + \text{edges}$
- ▶ Open:  $q = \Omega(n^2)$
- ▶ Open: conjecture on asymptotics of  $s_F(n, q)$

# Thank you!

Hope to see you at "*Extremal Combinatorics*"  
Warwick, 18-22 September 2017

<http://go.warwick.ac.uk/excomb2017>