


Minimising the Number of Cliques

Oleg Pikhurko and Emil R. Vaughan



C&C 2013

Method: Flag Algebras

Method: Flag Algebras

- ▶ Large graph G

Method: Flag Algebras

- ▶ Large graph G
- ▶ Inequalities between subgraph densities

Method: Flag Algebras

- ▶ Large graph G
- ▶ Inequalities between subgraph densities
- ▶ Razborov'07: Flag algebras

Method: Flag Algebras

- ▶ Large graph G
- ▶ Inequalities between subgraph densities
- ▶ Razborov'07: Flag algebras
 - ▶ Asymptotically true inequalities (as $v(G) \rightarrow \infty$)

Method: Flag Algebras

- ▶ Large graph G
- ▶ Inequalities between subgraph densities
- ▶ Razborov'07: Flag algebras
 - ▶ Asymptotically true inequalities (as $v(G) \rightarrow \infty$)
 - ▶ Formal rules for deriving them

Method: Flag Algebras

- ▶ Large graph G
- ▶ Inequalities between subgraph densities
- ▶ Razborov'07: Flag algebras
 - ▶ Asymptotically true inequalities (as $v(G) \rightarrow \infty$)
 - ▶ Formal rules for deriving them
 - ▶ One aspect: semi-definite programming

Recent Results Obtained with Flag Algebras

Recent Results Obtained with Flag Algebras

- ▶ **Structures:** graphs, digraphs, k -uniform hypergraphs, permutations, subgraphs of the hypercube Q_n , ...

Recent Results Obtained with Flag Algebras

- ▶ **Structures:** graphs, digraphs, k -uniform hypergraphs, permutations, subgraphs of the hypercube Q_n , ...
- ▶ **Contributors:** R.Baber, J.Balogh, J.Cummings, S.Das, V.Falgas-Ravry, R.Glebov, A.Grzesik, H.Hatami, J.Hirst, J.Hladký, P.Hu, H.Huang, T.Klimosova, D.Král', L.Kramer, B.Lidicky, N.Linial, C.-H.Liu, J.Ma, L.Mach, E.Marchant, R.Martin, H.Naves, S.Niess, S.Norine, Y.Peled, F.Pfender, O.Pikhurko, A.Razborov, C.Reiher, J.-S.Sereni, K.Spengler, B.Sudakov, J.Talbot, A.Treglown, E.Vaughan, J.Volec, P.Whalen, Z.Yilma, M.Young, ...

Turán Problem via Flag Algebras

Turán Problem via Flag Algebras

- ▶ Turán function:

$$\text{ex}(n, K_3) = \max\{e(G) : v(G) = n, G \not\supseteq K_3\}$$

Turán Problem via Flag Algebras

- ▶ Turán function:

$$\text{ex}(n, K_3) = \max\{e(G) : v(G) = n, G \not\supseteq K_3\}$$

- ▶ Construction: $\text{ex}(n, K_3) \geq e(T_n^2) = \lfloor n^2/4 \rfloor$

Turán Problem via Flag Algebras

- ▶ Turán function:

$$\text{ex}(n, K_3) = \max\{e(G) : v(G) = n, G \not\supseteq K_3\}$$

- ▶ Construction: $\text{ex}(n, K_3) \geq e(T_n^2) = \lfloor n^2/4 \rfloor$
- ▶ $T_n^2 = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$

Subgraph Densities

Subgraph Densities

- ▶ K_3 -free G of order $n \rightarrow \infty$

Subgraph Densities

- ▶ K_3 -free G of order $n \rightarrow \infty$
- ▶ $\#F = \#(F, G)$

Subgraph Densities

- ▶ K_3 -free G of order $n \rightarrow \infty$
- ▶ $\#F = \#(F, G) = \#$ induced copies of F in G

Subgraph Densities

- ▶ K_3 -free G of order $n \rightarrow \infty$
- ▶ $\#F = \#(F, G) = \#$ induced copies of F in G
 - ▶ $e(G) = \#(K_2, G)$

Subgraph Densities

- ▶ K_3 -free G of order $n \rightarrow \infty$
- ▶ $\#F = \#(F, G) = \#$ induced copies of F in G
 - ▶ $e(G) = \#(K_2, G)$
- ▶ Density of F :

$$\phi(F) = \#F / \binom{n}{v(F)}$$

Subgraph Densities

- ▶ K_3 -free G of order $n \rightarrow \infty$
- ▶ $\#F = \#(F, G) = \#$ induced copies of F in G
 - ▶ $e(G) = \#(K_2, G)$
- ▶ Density of F :

$$\phi(F) = \#F / \binom{n}{v(F)} = \mathbf{Prob}\{G[\text{random } v(F)\text{-set}] \cong F\}$$

Obtaining Inequalities

Obtaining Inequalities

$$\blacktriangleright \mathbf{A} = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq \mathbf{0}$$

Obtaining Inequalities

- ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$

Obtaining Inequalities

- ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ $\mathbf{v}_x = \left(\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1} \right), \quad x \in V(G)$
- ▶ $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$

Obtaining Inequalities

- ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$
- ▶ $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over x (and ignore $o(1)$):

Obtaining Inequalities

- ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$
- ▶ $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over x (and ignore $o(1)$):

$$\frac{1}{n} \sum_x \alpha \frac{d^2(x)}{(n-1)^2}$$

Obtaining Inequalities

- ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$
- ▶ $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over x (and ignore $o(1)$):

$$\frac{1}{n} \sum_x \alpha \frac{d^2(x)}{(n-1)^2} = \frac{\alpha}{n^3} \sum_x \left(\sum_{y \sim x} 1 \right) \left(\sum_{z \sim x} 1 \right)$$

Obtaining Inequalities

- ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ $\mathbf{v}_x = \left(\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1} \right), \quad x \in V(G)$
- ▶ $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over x (and ignore $o(1)$):

$$\begin{aligned} \frac{1}{n} \sum_x \alpha \frac{d^2(x)}{(n-1)^2} &= \frac{\alpha}{n^3} \sum_x \left(\sum_{y \sim x} 1 \right) \left(\sum_{z \sim x} 1 \right) \\ &= \frac{\alpha}{6 \binom{n}{3}} \times 2 \#P_3 \end{aligned}$$

Obtaining Inequalities

- ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ $\mathbf{v}_x = \left(\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1} \right), \quad x \in V(G)$
- ▶ $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over x (and ignore $o(1)$):

$$\begin{aligned} \frac{1}{n} \sum_x \alpha \frac{d^2(x)}{(n-1)^2} &= \frac{\alpha}{n^3} \sum_x \left(\sum_{y \sim x} 1 \right) \left(\sum_{z \sim x} 1 \right) \\ &= \frac{\alpha}{6 \binom{n}{3}} \times 2 \#P_3 = \frac{\alpha}{3} \phi(P_3) \end{aligned}$$

Obtaining Inequalities

- ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$
- ▶ $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over x (and ignore $o(1)$):

$$\begin{aligned} \frac{1}{n} \sum_x \alpha \frac{d^2(x)}{(n-1)^2} &= \frac{\alpha}{n^3} \sum_x \left(\sum_{y \sim x} 1 \right) \left(\sum_{z \sim x} 1 \right) \\ &= \frac{\alpha}{6 \binom{n}{3}} \times 2 \#P_3 = \frac{\alpha}{3} \phi(P_3) \end{aligned}$$

- ▶ $0 \leq (\frac{2\beta}{3} + \frac{\alpha}{3})\phi(P_3) + (\frac{2\beta}{3} + \frac{\gamma}{3})\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$

Bounding Edge Density from Above

Bounding Edge Density from Above

$$\blacktriangleright 0 \leq \left(\frac{2\beta}{3} + \frac{\alpha}{3}\right)\phi(P_3) + \left(\frac{2\beta}{3} + \frac{\gamma}{3}\right)\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$$

Bounding Edge Density from Above

$$\blacktriangleright 0 \leq \left(\frac{2\beta}{3} + \frac{\alpha}{3}\right)\phi(P_3) + \left(\frac{2\beta}{3} + \frac{\gamma}{3}\right)\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$$

$$\phi(K_2)$$

Bounding Edge Density from Above

$$\blacktriangleright 0 \leq \left(\frac{2\beta}{3} + \frac{\alpha}{3}\right)\phi(P_3) + \left(\frac{2\beta}{3} + \frac{\gamma}{3}\right)\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$$

$$\phi(K_2) = \frac{2}{3}\phi(P_3) + \frac{1}{3}\phi(\bar{P}_3)$$

Bounding Edge Density from Above

$$\blacktriangleright 0 \leq \left(\frac{2\beta}{3} + \frac{\alpha}{3}\right)\phi(P_3) + \left(\frac{2\beta}{3} + \frac{\gamma}{3}\right)\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$$

$$\begin{aligned}\phi(K_2) &= \frac{2}{3}\phi(P_3) + \frac{1}{3}\phi(\bar{P}_3) \\ &\leq \left(\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3}\right)\phi(P_3) + \left(\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3}\right)\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)\end{aligned}$$

Bounding Edge Density from Above

$$\blacktriangleright 0 \leq \left(\frac{2\beta}{3} + \frac{\alpha}{3}\right)\phi(P_3) + \left(\frac{2\beta}{3} + \frac{\gamma}{3}\right)\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$$

$$\begin{aligned}\phi(K_2) &= \frac{2}{3}\phi(P_3) + \frac{1}{3}\phi(\bar{P}_3) \\ &\leq \left(\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3}\right)\phi(P_3) + \left(\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3}\right)\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)\end{aligned}$$

$$\blacktriangleright \text{Note: } \phi(P_3) + \phi(\bar{P}_3) + \phi(\bar{K}_3) = 1$$

SDP Formulation

SDP Formulation

$$\blacktriangleright \phi(K_2) \leq \left(\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3}\right)\phi(P_3) + \left(\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3}\right)\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$$

SDP Formulation

- ▶ $\phi(K_2) \leq (\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3})\phi(P_3) + (\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3})\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$
- ▶ **Minimise** δ :

SDP Formulation

- ▶ $\phi(K_2) \leq (\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3})\phi(P_3) + (\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3})\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$
- ▶ **Minimise** δ :
 - ▶ $\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3} \leq \delta$

SDP Formulation

- ▶ $\phi(K_2) \leq (\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3})\phi(P_3) + (\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3})\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$
- ▶ **Minimise** δ :
 - ▶ $\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3} \leq \delta$
 - ▶ $\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3} \leq \delta$

SDP Formulation

- ▶ $\phi(K_2) \leq (\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3})\phi(P_3) + (\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3})\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$
- ▶ **Minimise** δ :
 - ▶ $\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3} \leq \delta$
 - ▶ $\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3} \leq \delta$
 - ▶ $\gamma \leq \delta$

SDP Formulation

- ▶ $\phi(K_2) \leq (\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3})\phi(P_3) + (\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3})\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$
- ▶ **Minimise** δ :
 - ▶ $\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3} \leq \delta$
 - ▶ $\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3} \leq \delta$
 - ▶ $\gamma \leq \delta$
 - ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$

SDP Formulation

- ▶ $\phi(K_2) \leq (\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3})\phi(P_3) + (\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3})\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$
- ▶ **Minimise** δ :
 - ▶ $\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3} \leq \delta$
 - ▶ $\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3} \leq \delta$
 - ▶ $\gamma \leq \delta$
 - ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ **Solution:** $\delta = 1/2$ and $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$

SDP Formulation

- ▶ $\phi(K_2) \leq (\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3})\phi(P_3) + (\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3})\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$
- ▶ **Minimise** δ :
 - ▶ $\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3} \leq \delta$
 - ▶ $\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3} \leq \delta$
 - ▶ $\gamma \leq \delta$
 - ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ **Solution:** $\delta = 1/2$ and $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
- ▶ $\phi(K_2) \leq \frac{1}{2}$

SDP Formulation

- ▶ $\phi(K_2) \leq (\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3})\phi(P_3) + (\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3})\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$
- ▶ **Minimise** δ :
 - ▶ $\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3} \leq \delta$
 - ▶ $\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3} \leq \delta$
 - ▶ $\gamma \leq \delta$
 - ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ **Solution:** $\delta = 1/2$ and $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
- ▶ $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3}\phi(\bar{P}_3)$

SDP Formulation

- ▶ $\phi(K_2) \leq (\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3})\phi(P_3) + (\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3})\phi(\bar{P}_3) + \gamma\phi(\bar{K}_3)$
- ▶ **Minimise** δ :
 - ▶ $\frac{2}{3} + \frac{2\beta}{3} + \frac{\alpha}{3} \leq \delta$
 - ▶ $\frac{1}{3} + \frac{2\beta}{3} + \frac{\gamma}{3} \leq \delta$
 - ▶ $\gamma \leq \delta$
 - ▶ $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ **Solution:** $\delta = 1/2$ and $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
- ▶ $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3} \phi(\bar{P}_3)$
- ▶ **Asymptotic result:** $\text{ex}(n, K_3) \leq (\frac{1}{2} + o(1))\binom{n}{2}$

Interpreting the Solution

Interpreting the Solution

► Recall $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$

Interpreting the Solution

- ▶ Recall $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
- ▶ $\mathbf{v}_x A \mathbf{v}_x^T = \frac{1}{2(n-1)^2} (d(x) - \bar{d}(x))^2$

Interpreting the Solution

- ▶ Recall $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
- ▶ $\mathbf{v}_x A \mathbf{v}_x^T = \frac{1}{2(n-1)^2} (d(x) - \bar{d}(x))^2$
- ▶ **Human proof:**

Interpreting the Solution

- ▶ Recall $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
- ▶ $\mathbf{v}_x A \mathbf{v}_x^T = \frac{1}{2(n-1)^2} (d(x) - \bar{d}(x))^2$
- ▶ **Human proof:**
 - ▶ $0 \leq \sum_x (d(x) - \bar{d}(x))^2$

Interpreting the Solution

- ▶ Recall $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
- ▶ $\mathbf{v}_x A \mathbf{v}_x^T = \frac{1}{2(n-1)^2} (d(x) - \bar{d}(x))^2$
- ▶ **Human proof:**
 - ▶ $0 \leq \sum_x (d(x) - \bar{d}(x))^2$
 - ▶ Expand and re-group terms

Interpreting the Solution

- ▶ Recall $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
- ▶ $\mathbf{v}_x A \mathbf{v}_x^T = \frac{1}{2(n-1)^2} (d(x) - \bar{d}(x))^2$
- ▶ **Human proof:**
 - ▶ $0 \leq \sum_x (d(x) - \bar{d}(x))^2$
 - ▶ Expand and re-group terms
 - ▶ $\#K_2 \leq \frac{n^2}{4} - \frac{1}{n} \# \bar{P}_3 + O(n)$

Structure of Almost Extremal Graphs

Structure of Almost Extremal Graphs

- ▶ $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3} \phi(\bar{P}_3)$

Structure of Almost Extremal Graphs

- ▶ $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3} \phi(\bar{P}_3)$
- ▶ $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\bar{P}_3) = o(1)$

Structure of Almost Extremal Graphs

- ▶ $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3} \phi(\bar{P}_3)$
- ▶ $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\bar{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**
 G is $o(n^2)$ -close in edit distance to \bar{P}_3 -free

Structure of Almost Extremal Graphs

- ▶ $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3} \phi(\bar{P}_3)$
- ▶ $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\bar{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**
 G is $o(n^2)$ -close in edit distance to \bar{P}_3 -free
- ▶ \bar{P}_3 -free \Rightarrow complete multipartite

Structure of Almost Extremal Graphs

- ▶ $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3} \phi(\bar{P}_3)$
- ▶ $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\bar{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**
 G is $o(n^2)$ -close in edit distance to \bar{P}_3 -free
- ▶ \bar{P}_3 -free \Rightarrow complete multipartite
- ▶ K_3 -free \Rightarrow at most 2 parts

Structure of Almost Extremal Graphs

- ▶ $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3} \phi(\bar{P}_3)$
- ▶ $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\bar{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**
 G is $o(n^2)$ -close in edit distance to \bar{P}_3 -free
- ▶ \bar{P}_3 -free \Rightarrow complete multipartite
- ▶ K_3 -free \Rightarrow at most 2 parts
- ▶ $\phi(K_2) \approx \frac{1}{2} \Rightarrow$ parts almost equal

Structure of Almost Extremal Graphs

- ▶ $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3} \phi(\bar{P}_3)$
- ▶ $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\bar{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**
 G is $o(n^2)$ -close in edit distance to \bar{P}_3 -free
- ▶ \bar{P}_3 -free \Rightarrow complete multipartite
- ▶ K_3 -free \Rightarrow at most 2 parts
- ▶ $\phi(K_2) \approx \frac{1}{2} \Rightarrow$ parts almost equal
- ▶ **Stability (Erdős'67, Simonovits'68):**
 \forall almost extremal G_n is $o(n^2)$ -close to T_n^2

Structure of Almost Extremal Graphs

- ▶ $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3} \phi(\bar{P}_3)$
- ▶ $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\bar{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**
 G is $o(n^2)$ -close in edit distance to \bar{P}_3 -free
- ▶ \bar{P}_3 -free \Rightarrow complete multipartite
- ▶ K_3 -free \Rightarrow at most 2 parts
- ▶ $\phi(K_2) \approx \frac{1}{2} \Rightarrow$ parts almost equal
- ▶ **Stability (Erdős'67, Simonovits'68):**
 \forall almost extremal G_n is $o(n^2)$ -close to T_n^2
- ▶ **More work:** exact result for $n \geq n_0$

Further Ways of Obtaining Inequalities

Further Ways of Obtaining Inequalities

- ▶ Recall:

Further Ways of Obtaining Inequalities

- ▶ Recall:

- ▶ $0 \leq \mathbb{E}_x(\mathbf{v}_x A \mathbf{v}_x^T)$

Further Ways of Obtaining Inequalities

- ▶ Recall:

- ▶ $0 \leq \mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T)$

- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1}), \quad x \in V(G)$

Further Ways of Obtaining Inequalities

- ▶ Recall:

- ▶ $0 \leq \mathbb{E}_x(\mathbf{v}_x A \mathbf{v}_x^T)$

- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$

- ▶ Extensions:

Further Ways of Obtaining Inequalities

- ▶ Recall:

- ▶ $0 \leq \mathbb{E}_x(\mathbf{v}_x A \mathbf{v}_x^T)$

- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$

- ▶ Extensions:

- ▶ \mathbf{v}_x : densities of *k*-vertex graphs rooted at x

Further Ways of Obtaining Inequalities

- ▶ Recall:

- ▶ $0 \leq \mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T)$

- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$

- ▶ Extensions:

- ▶ \mathbf{v}_x : densities of **k-vertex** graphs rooted at x

- ▶ \mathbf{v}_{xy} : k -vertex graphs rooted at **edge** xy

Further Ways of Obtaining Inequalities

- ▶ Recall:

- ▶ $0 \leq \mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T)$
- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$

- ▶ Extensions:

- ▶ \mathbf{v}_x : densities of **k-vertex** graphs rooted at x
- ▶ \mathbf{v}_{xy} : k -vertex graphs rooted at **edge** xy
 - ▶ Average over $E(G)$ (assuming $e(G) = \Omega(n^2)$)

Further Ways of Obtaining Inequalities

- ▶ Recall:

- ▶ $0 \leq \mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T)$
- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$

- ▶ Extensions:

- ▶ \mathbf{v}_x : densities of **k-vertex** graphs rooted at x
- ▶ \mathbf{v}_{xy} : k -vertex graphs rooted at **edge** xy
 - ▶ Average over $E(G)$ (assuming $e(G) = \Omega(n^2)$)
- ▶ \mathbf{v}_{xy} : k -vertex graphs rooted at **non-edge** xy

Further Ways of Obtaining Inequalities

- ▶ Recall:

- ▶ $0 \leq \mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T)$
- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$

- ▶ Extensions:

- ▶ \mathbf{v}_x : densities of **k-vertex** graphs rooted at x
- ▶ \mathbf{v}_{xy} : k -vertex graphs rooted at **edge** xy
 - ▶ Average over $E(G)$ (assuming $e(G) = \Omega(n^2)$)
- ▶ \mathbf{v}_{xy} : k -vertex graphs rooted at **non-edge** xy
 - ▶ Average over $E(\bar{G})$ (assuming $e(\bar{G}) = \Omega(n^2)$)

Further Ways of Obtaining Inequalities

- ▶ Recall:

- ▶ $0 \leq \mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T)$
- ▶ $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$, $x \in V(G)$

- ▶ Extensions:

- ▶ \mathbf{v}_x : densities of **k-vertex** graphs rooted at x
- ▶ \mathbf{v}_{xy} : k -vertex graphs rooted at **edge** xy
 - ▶ Average over $E(G)$ (assuming $e(G) = \Omega(n^2)$)
- ▶ \mathbf{v}_{xy} : k -vertex graphs rooted at **non-edge** xy
 - ▶ Average over $E(\bar{G})$ (assuming $e(\bar{G}) = \Omega(n^2)$)
- ▶ ...

Rounding Floating-Point Solution

Rounding Floating-Point Solution

- ▶ $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$

Rounding Floating-Point Solution

- ▶ $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.500007 & -0.499997 \\ -0.499997 & 0.500012 \end{pmatrix} ?$$

Rounding Floating-Point Solution

- ▶ $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.500007 & -0.499997 \\ -0.499997 & 0.500012 \end{pmatrix} ?$$

- ▶ Suppose $A \succeq 0$ **proves** $\phi(K_2) \leq \frac{1}{2}$

Rounding Floating-Point Solution

- ▶ $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.500007 & -0.499997 \\ -0.499997 & 0.500012 \end{pmatrix} ?$$

- ▶ Suppose $A \succeq 0$ **proves** $\phi(K_2) \leq \frac{1}{2}$
- ▶ If $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$:

Rounding Floating-Point Solution

- ▶ $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.500007 & -0.499997 \\ -0.499997 & 0.500012 \end{pmatrix} ?$$

- ▶ Suppose $A \succeq 0$ **proves** $\phi(K_2) \leq \frac{1}{2}$
- ▶ If $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$:
 - ▶ Then $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3}\phi(\bar{P}_3) - \varepsilon\phi(P_3)$

Rounding Floating-Point Solution

- ▶ $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.500007 & -0.499997 \\ -0.499997 & 0.500012 \end{pmatrix} ?$$

- ▶ Suppose $A \succeq 0$ **proves** $\phi(K_2) \leq \frac{1}{2}$
- ▶ If $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$:
 - ▶ Then $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3}\phi(\bar{P}_3) - \varepsilon\phi(P_3)$
 - ▶ **False** for $K_{n/2, n/2}$!

Rounding Floating-Point Solution

- ▶ $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.500007 & -0.499997 \\ -0.499997 & 0.500012 \end{pmatrix} ?$$

- ▶ Suppose $A \succeq 0$ **proves** $\phi(K_2) \leq \frac{1}{2}$
- ▶ If $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$:
 - ▶ Then $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3}\phi(\bar{P}_3) - \varepsilon\phi(P_3)$
 - ▶ **False** for $K_{n/2, n/2}$!
- ▶ **Likewise:** $3\gamma = \frac{1}{2}$

Rounding Floating-Point Solution

- ▶ $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.500007 & -0.499997 \\ -0.499997 & 0.500012 \end{pmatrix} ?$$

- ▶ Suppose $A \succeq 0$ **proves** $\phi(K_2) \leq \frac{1}{2}$
- ▶ If $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$:
 - ▶ Then $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3}\phi(\bar{P}_3) - \varepsilon\phi(P_3)$
 - ▶ **False** for $K_{n/2, n/2}$!
- ▶ **Likewise:** $3\gamma = \frac{1}{2}$
- ▶ $\phi(K_2) + \mathbb{E}_x(\mathbf{v}_x A \mathbf{v}_x^T) = \frac{1}{2} - \frac{1}{3}\phi(\bar{P}_3)$

Rounding Floating-Point Solution

- ▶ $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.500007 & -0.499997 \\ -0.499997 & 0.500012 \end{pmatrix} ?$$

- ▶ Suppose $A \succeq 0$ **proves** $\phi(K_2) \leq \frac{1}{2}$
- ▶ If $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$:
 - ▶ Then $\phi(K_2) \leq \frac{1}{2} - \frac{1}{3}\phi(\bar{P}_3) - \varepsilon\phi(P_3)$
 - ▶ **False** for $K_{n/2, n/2}$!
- ▶ **Likewise:** $3\gamma = \frac{1}{2}$
- ▶ $\phi(K_2) + \mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T) = \frac{1}{2} - \frac{1}{3}\phi(\bar{P}_3) \Rightarrow A \begin{pmatrix} 1/2 \\ 1/2 \end{pmatrix} = 0$

Clique Minimisation Problem

Clique Minimisation Problem

- Independence number:

$$\alpha(G) = \min\{m : \#(\overline{K}_m, G) > 0\}$$

Clique Minimisation Problem

- ▶ Independence number:

$$\alpha(G) = \min\{m : \#(\overline{K}_m, G) > 0\}$$

- ▶ Erdős'62:

$$f(n, k, \ell) = \min\{\#(K_k, G) : \alpha(G) < \ell, \ v(G) = n\}$$

Clique Minimisation Problem

- ▶ Independence number:

$$\alpha(G) = \min\{m : \#(\overline{K}_m, G) > 0\}$$

- ▶ Erdős'62:

$$f(n, k, \ell) = \min\{\#(K_k, G) : \alpha(G) < \ell, \ v(G) = n\}$$

- ▶ Motivation: Ramsey numbers

Clique Minimisation Problem

- ▶ Independence number:

$$\alpha(G) = \min\{m : \#(\overline{K}_m, G) > 0\}$$

- ▶ Erdős'62:

$$f(n, k, \ell) = \min\{\#(K_k, G) : \alpha(G) < \ell, v(G) = n\}$$

- ▶ Motivation: Ramsey numbers

- ▶ $n \geq R(k, \ell) \iff f(n, k, \ell) > 0$

Ramsey Multiplicity Problem

Ramsey Multiplicity Problem

- ▶ Erdős'62: $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$

Ramsey Multiplicity Problem

- ▶ Erdős'62: $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶ $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$

Ramsey Multiplicity Problem

- ▶ Erdős'62: $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶ $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56: $\mu_3 = 1/4$

Ramsey Multiplicity Problem

- ▶ Erdős'62: $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶ $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56: $\mu_3 = 1/4$
- ▶ Erdős'62: Is $\mu_k = 2^{1-\binom{k}{2}}$?

Ramsey Multiplicity Problem

- ▶ Erdős'62: $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶ $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56: $\mu_3 = 1/4$
- ▶ Erdős'62: Is $\mu_k = 2^{1 - \binom{k}{2}}$?
- ▶ Thomason'89: False: $\mu_k \leq 0.936 \cdot 2^{1 - \binom{k}{2}}$

Ramsey Multiplicity Problem

- ▶ Erdős'62: $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶ $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56: $\mu_3 = 1/4$
- ▶ Erdős'62: Is $\mu_k = 2^{1-\binom{k}{2}}$?
- ▶ Thomason'89: False: $\mu_k \leq 0.936 \cdot 2^{1-\binom{k}{2}}$
- ▶ Thomason'89: $\mu_4 \leq \frac{1}{33} < \frac{1}{32}$

Ramsey Multiplicity Problem

- ▶ Erdős'62: $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶ $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56: $\mu_3 = 1/4$
- ▶ Erdős'62: Is $\mu_k = 2^{1-\binom{k}{2}}$?
- ▶ Thomason'89: False: $\mu_k \leq 0.936 \cdot 2^{1-\binom{k}{2}}$
- ▶ Thomason'89: $\mu_4 \leq \frac{1}{33} < \frac{1}{32}$
- ▶ Giraud'79: $\mu_4 \geq \frac{1}{46} = 0.0217\dots$

Ramsey Multiplicity Problem

- ▶ Erdős'62: $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶ $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56: $\mu_3 = 1/4$
- ▶ Erdős'62: Is $\mu_k = 2^{1-\binom{k}{2}}$?
- ▶ Thomason'89: False: $\mu_k \leq 0.936 \cdot 2^{1-\binom{k}{2}}$
- ▶ Thomason'89: $\mu_4 \leq \frac{1}{33} < \frac{1}{32}$
- ▶ Giraud'79: $\mu_4 \geq \frac{1}{46} = 0.0217\dots$
- ▶ Nieß \geq '13, Sperfeld \geq '13: $\mu_4 \geq 0.0287\dots$

Ramsey Multiplicity Problem

- ▶ Erdős'62: $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶ $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56: $\mu_3 = 1/4$
- ▶ Erdős'62: Is $\mu_k = 2^{1-\binom{k}{2}}$?
- ▶ Thomason'89: False: $\mu_k \leq 0.936 \cdot 2^{1-\binom{k}{2}}$
- ▶ Thomason'89: $\mu_4 \leq \frac{1}{33} < \frac{1}{32}$
- ▶ Giraud'79: $\mu_4 \geq \frac{1}{46} = 0.0217\dots$
- ▶ Nieß \geq '13, Sperfeld \geq '13: $\mu_4 \geq 0.0287\dots$
- ▶ Erdős'62: $\mu_k \geq \binom{R(k,k)}{k}^{-1} \geq (4 + o(1))^{-k^2}$

Ramsey Multiplicity Problem

- ▶ Erdős'62: $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶ $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56: $\mu_3 = 1/4$
- ▶ Erdős'62: Is $\mu_k = 2^{1-\binom{k}{2}}$?
- ▶ Thomason'89: False: $\mu_k \leq 0.936 \cdot 2^{1-\binom{k}{2}}$
- ▶ Thomason'89: $\mu_4 \leq \frac{1}{33} < \frac{1}{32}$
- ▶ Giraud'79: $\mu_4 \geq \frac{1}{46} = 0.0217\dots$
- ▶ Nieß \geq '13, Sperfeld \geq '13: $\mu_4 \geq 0.0287\dots$
- ▶ Erdős'62: $\mu_k \geq \binom{R(k,k)}{k}^{-1} \geq (4 + o(1))^{-k^2}$
- ▶ Conlon'12: $\mu_k \geq (2.18 + o(1))^{-k^2}$

Early Results

Early Results

- ▶ $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$

Early Results

- ▶ $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ **Goodman'56:** $f(2m, 3, 3) = \#(K_3, K_m \sqcup K_m)$

Early Results

- ▶ $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ **Goodman'56:** $f(2m, 3, 3) = \#(K_3, K_m \sqcup K_m)$
- ▶ **Lorden'62:** $f(2m+1, 3, 3) = \#(K_3, K_m \sqcup K_{m+1}), m \geq 6$

Early Results

- ▶ $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ **Goodman'56:** $f(2m, 3, 3) = \#(K_3, K_m \sqcup K_m)$
- ▶ **Lorden'62:** $f(2m+1, 3, 3) = \#(K_3, K_m \sqcup K_{m+1})$, $m \geq 6$
- ▶ Complement of T_n^2

Early Results

- ▶ $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ **Goodman'56:** $f(2m, 3, 3) = \#(K_3, K_m \sqcup K_m)$
- ▶ **Lorden'62:** $f(2m+1, 3, 3) = \#(K_3, K_m \sqcup K_{m+1})$, $m \geq 6$
- ▶ Complement of T_n^2
- ▶ **Turán graph T_n^r :** max r -partite order- n graph

Early Results

- ▶ $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ **Goodman'56:** $f(2m, 3, 3) = \#(K_3, K_m \sqcup K_m)$
- ▶ **Lorden'62:** $f(2m+1, 3, 3) = \#(K_3, K_m \sqcup K_{m+1})$, $m \geq 6$
- ▶ Complement of T_n^2
- ▶ **Turán graph T_n^r :** max r -partite order- n graph
- ▶ **Erdős'62:** Is $f(n, k, \ell) = \#(K_3, \overline{T}_n^{\ell-1})$?

Asymptotic densities

Asymptotic densities

- ▶ Nikiforov'01: $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$

Asymptotic densities

- ▶ Nikiforov'01: $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph: $c_{k,\ell} \leq (\ell - 1)^{1-k}$

Asymptotic densities

- ▶ Nikiforov'01: $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph: $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ Nikiforov'01: Strict for all but finitely many (k, ℓ)

Asymptotic densities

- ▶ Nikiforov'01: $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph: $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ Nikiforov'01: Strict for all but finitely many (k, ℓ)
- ▶ Expansion of F :

Asymptotic densities

- ▶ **Nikiforov'01:** $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph: $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many (k, ℓ)
- ▶ **Expansion** of F :
 - ▶ Replace each $ij \in E(F)$ by complete graph $K[V_i, V_j]$

Asymptotic densities

- ▶ **Nikiforov'01:** $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph: $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many (k, ℓ)
- ▶ **Expansion** of F :
 - ▶ Replace each $ij \in E(F)$ by complete graph $K[V_i, V_j]$
 - ▶ Replace each $i \in V(F)$ by clique on $K[V_i]$

Asymptotic densities

- ▶ **Nikiforov'01:** $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph: $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many (k, ℓ)
- ▶ **Expansion** of F :
 - ▶ Replace each $ij \in E(F)$ by complete graph $K[V_i, V_j]$
 - ▶ Replace each $i \in V(F)$ by clique on $K[V_i]$
- ▶ **Uniform expansion** $U_n(F)$: $\forall i, j \quad |V_i - V_j| \leq 1$

Asymptotic densities

- ▶ **Nikiforov'01:** $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph: $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many (k, ℓ)
- ▶ **Expansion** of F :
 - ▶ Replace each $ij \in E(F)$ by complete graph $K[V_i, V_j]$
 - ▶ Replace each $i \in V(F)$ by clique on $K[V_i]$
- ▶ **Uniform expansion** $U_n(F)$: $\forall i, j \quad |V_i - V_j| \leq 1$
- ▶ **Nikiforov'01:** $f(n, 4, 3) \leq \#(K_4, U_n(C_5))$

Asymptotic densities

- ▶ **Nikiforov'01:** $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph: $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many (k, ℓ)
- ▶ **Expansion** of F :
 - ▶ Replace each $ij \in E(F)$ by complete graph $K[V_i, V_j]$
 - ▶ Replace each $i \in V(F)$ by clique on $K[V_i]$
- ▶ **Uniform expansion** $U_n(F)$: $\forall i, j \quad |V_i - V_j| \leq 1$
- ▶ **Nikiforov'01:** $f(n, 4, 3) \leq \#(K_4, U_n(C_5))$
 - ▶ $c_{4,3} \leq 3/25$

Asymptotic densities

- ▶ **Nikiforov'01:** $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph: $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many (k, ℓ)
- ▶ **Expansion** of F :
 - ▶ Replace each $ij \in E(F)$ by complete graph $K[V_i, V_j]$
 - ▶ Replace each $i \in V(F)$ by clique on $K[V_i]$
- ▶ **Uniform expansion** $U_n(F)$: $\forall i, j \quad |V_i - V_j| \leq 1$
- ▶ **Nikiforov'01:** $f(n, 4, 3) \leq \#(K_4, U_n(C_5))$
 - ▶ $c_{4,3} \leq 3/25 < 1/8$

Asymptotic densities

- ▶ **Nikiforov'01:** $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph: $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many (k, ℓ)
- ▶ **Expansion of F :**
 - ▶ Replace each $ij \in E(F)$ by complete graph $K[V_i, V_j]$
 - ▶ Replace each $i \in V(F)$ by clique on $K[V_i]$
- ▶ **Uniform expansion $U_n(F)$:** $\forall i, j \quad |V_i - V_j| \leq 1$
- ▶ **Nikiforov'01:** $f(n, 4, 3) \leq \#(K_4, U_n(C_5))$
 - ▶ $c_{4,3} \leq 3/25 < 1/8$
- ▶ **Nikiforov'05:**
 $\alpha(G_n) < 3$ & **regular** $\Rightarrow \#(K_4, G) \geq (\frac{3}{25} + o(1)) \binom{n}{4}$

Complete Solutions (for $n \geq n_0$)

Complete Solutions (for $n \geq n_0$)

- ▶ Das-Huang-Ma-Naves-Sudakov'13: $(3, 4)$ & $(4, 3)$

Complete Solutions (for $n \geq n_0$)

- ▶ Das-Huang-Ma-Naves-Sudakov'13: $(3, 4)$ & $(4, 3)$
- ▶ P.-Vaughan \geq '13:

$$(k, 3) : \quad 4 \leq k \leq 7$$

$$(3, \ell) : \quad 4 \leq \ell \leq 7$$

New Values of $c_{k,l}$

New Values of $c_{k,\ell}$

- ▶ $c_{3,\ell} = (\ell - 1)^2$ for $4 \leq \ell \leq 7$

New Values of $c_{k,\ell}$

- ▶ $c_{3,\ell} = (\ell - 1)^2$ for $4 \leq \ell \leq 7$
 - ▶ $\overline{T}_n^{\ell-1}$

New Values of $c_{k,\ell}$

- ▶ $c_{3,\ell} = (\ell - 1)^2$ for $4 \leq \ell \leq 7$
 - ▶ $\overline{T}_n^{\ell-1}$
- ▶ $c_{4,3} = 3/25$

New Values of $c_{k,\ell}$

- ▶ $c_{3,\ell} = (\ell - 1)^2$ for $4 \leq \ell \leq 7$
 - ▶ $\overline{T}_n^{\ell-1}$
- ▶ $c_{4,3} = 3/25$
- ▶ $c_{5,3} = 31/625$

New Values of $c_{k,\ell}$

- ▶ $c_{3,\ell} = (\ell - 1)^2$ for $4 \leq \ell \leq 7$
 - ▶ $\overline{T}_n^{\ell-1}$
- ▶ $c_{4,3} = 3/25$
- ▶ $c_{5,3} = 31/625$
 - ▶ Uniform expansion of C_5

New Values of $c_{k,\ell}$

- ▶ $c_{3,\ell} = (\ell - 1)^2$ for $4 \leq \ell \leq 7$
 - ▶ $\overline{T}_n^{\ell-1}$
- ▶ $c_{4,3} = 3/25$
- ▶ $c_{5,3} = 31/625$
 - ▶ Uniform expansion of C_5
- ▶ $c_{6,3} = 19211/2^{20} = 19211/1048576$

New Values of $c_{k,\ell}$

- ▶ $c_{3,\ell} = (\ell - 1)^2$ for $4 \leq \ell \leq 7$
 - ▶ $\overline{T}_n^{\ell-1}$
- ▶ $c_{4,3} = 3/25$
- ▶ $c_{5,3} = 31/625$
 - ▶ Uniform expansion of C_5
- ▶ $c_{6,3} = 19211/2^{20} = 19211/1048576$
- ▶ $c_{7,3} = 98491/2^{24} = 98491/16777216$

Construction for $c_{6,3}$ and $c_{7,3}$

Construction for $c_{6,3}$ and $c_{7,3}$

- ▶ Clebsch graph L :

Construction for $c_{6,3}$ and $c_{7,3}$

- ▶ Clebsch graph L :

- ▶ $V = \{\mathbf{X} \in \{0, 1\}^5 : \text{dist}(\mathbf{X}, \mathbf{0}) \text{ is even}\}$

Construction for $c_{6,3}$ and $c_{7,3}$



- ▶ Clebsch graph L :

- ▶ $V = \{\mathbf{X} \in \{0, 1\}^5 : \text{dist}(\mathbf{X}, \mathbf{0}) \text{ is even}\}$
- ▶ $\mathbf{X} \sim \mathbf{Y}$ iff $\text{dist}(\mathbf{X}, \mathbf{Y}) = 4$

Construction for $c_{6,3}$ and $c_{7,3}$

- ▶ **Clebsch graph L :**

- ▶ $V = \{\mathbf{X} \in \{0, 1\}^5 : \text{dist}(\mathbf{X}, \mathbf{0}) \text{ is even}\}$
- ▶ $\mathbf{X} \sim \mathbf{Y}$ iff $\text{dist}(\mathbf{X}, \mathbf{Y}) = 4$
- ▶ L and \bar{L} :





Construction for $c_{6,3}$ and $c_{7,3}$

- ▶ **Clebsch graph L :**

- ▶ $V = \{\mathbf{X} \in \{0, 1\}^5 : \text{dist}(\mathbf{X}, \mathbf{0}) \text{ is even}\}$
- ▶ $\mathbf{X} \sim \mathbf{Y}$ iff $\text{dist}(\mathbf{X}, \mathbf{Y}) = 4$

- ▶ L and \bar{L} :



- ▶ **Upper bound: $U_n(\bar{L})$**

Computer's Part

Computer's Part

- ▶ $c_{3,7}$: 12,338 constraints

Computer's Part

- ▶ $c_{3,7}$: 12,338 constraints
- ▶ Emil Vaughan's package `Flagmatic`

Computer's Part

- ▶ $c_{3,7}$: 12,338 constraints
- ▶ Emil Vaughan's package `Flagmatic`



Human Part: Stability

Human Part: Stability

- ▶ **Stability:** above ideas + eigenvalues

Human Part: Stability

- ▶ **Stability:** above ideas + eigenvalues
- ▶ Which expansions of \bar{L} minimise K_7 -density?

Human Part: Stability

- ▶ **Stability:** above ideas + eigenvalues
- ▶ Which expansions of \bar{L} minimise K_7 -density?
 - ▶ Part ratios $y_1 + \cdots + y_{16} = 1$

Human Part: Stability

- ▶ **Stability:** above ideas + eigenvalues
- ▶ Which expansions of \bar{L} minimise K_7 -density?
 - ▶ Part ratios $y_1 + \dots + y_{16} = 1$
 - ▶ $\phi(K_7) = p(y_1, \dots, y_{16})$, polynomial of degree 7

Human Part: Stability

- ▶ **Stability:** above ideas + eigenvalues
- ▶ Which expansions of \bar{L} minimise K_7 -density?
 - ▶ Part ratios $y_1 + \dots + y_{16} = 1$
 - ▶ $\phi(K_7) = p(y_1, \dots, y_{16})$, polynomial of degree 7
- ▶ Which expansion of \bar{K}_2 minimise K_3 -density?

Human Part: Stability

- ▶ **Stability:** above ideas + eigenvalues
- ▶ Which expansions of \bar{L} minimise K_7 -density?
 - ▶ Part ratios $y_1 + \dots + y_{16} = 1$
 - ▶ $\phi(K_7) = p(y_1, \dots, y_{16})$, polynomial of degree 7
- ▶ Which expansion of \bar{K}_2 minimise K_3 -density?
 - ▶ $p(y_1, y_2) = y_1^3 + y_2^3$

Human Part: Stability

- ▶ **Stability:** above ideas + eigenvalues
- ▶ Which expansions of \bar{L} minimise K_7 -density?
 - ▶ Part ratios $y_1 + \dots + y_{16} = 1$
 - ▶ $\phi(K_7) = p(y_1, \dots, y_{16})$, polynomial of degree 7
- ▶ Which expansion of \bar{K}_2 minimise K_3 -density?
 - ▶ $p(y_1, y_2) = y_1^3 + y_2^3$
 - ▶ $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$

Human Part: Stability

- ▶ **Stability:** above ideas + eigenvalues
- ▶ Which expansions of \bar{L} minimise K_7 -density?
 - ▶ Part ratios $y_1 + \dots + y_{16} = 1$
 - ▶ $\phi(K_7) = p(y_1, \dots, y_{16})$, polynomial of degree 7
- ▶ Which expansion of \bar{K}_2 minimise K_3 -density?
 - ▶ $p(y_1, y_2) = y_1^3 + y_2^3$
 - ▶ $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
 - ▶ $\mathbf{v}_x = (y_1, y_2)$ or (y_2, y_1)

Human Part: Stability

- ▶ **Stability:** above ideas + eigenvalues
- ▶ Which expansions of \bar{L} minimise K_7 -density?
 - ▶ Part ratios $y_1 + \dots + y_{16} = 1$
 - ▶ $\phi(K_7) = p(y_1, \dots, y_{16})$, polynomial of degree 7
- ▶ Which expansion of \bar{K}_2 minimise K_3 -density?
 - ▶ $p(y_1, y_2) = y_1^3 + y_2^3$
 - ▶ $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
 - ▶ $\mathbf{v}_x = (y_1, y_2)$ or (y_2, y_1)
 - ▶ $\mathbb{E}_x(\mathbf{v}_x A \mathbf{v}_x^T) = 0$

Human Part: Stability

- ▶ **Stability:** above ideas + eigenvalues
- ▶ Which expansions of \bar{L} minimise K_7 -density?
 - ▶ Part ratios $y_1 + \dots + y_{16} = 1$
 - ▶ $\phi(K_7) = p(y_1, \dots, y_{16})$, polynomial of degree 7
- ▶ Which expansion of \bar{K}_2 minimise K_3 -density?
 - ▶ $p(y_1, y_2) = y_1^3 + y_2^3$
 - ▶ $A = \begin{pmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{pmatrix}$
 - ▶ $\mathbf{v}_x = (y_1, y_2)$ or (y_2, y_1)
 - ▶ $\mathbb{E}_x(\mathbf{v}_x A \mathbf{v}_x^T) = 0 \Rightarrow y_1 = y_2$

Human Part: Exact Result

Human Part: Exact Result

- ▶ **Exact result:** One theorem covering all 8 cases

Human Part: Exact Result

- ▶ **Exact result:** One theorem covering all 8 cases
- ▶ **Assumptions:**

Human Part: Exact Result

- ▶ **Exact result:** One theorem covering all 8 cases
- ▶ **Assumptions:**
 - ▶ Stability (\forall almost extremal G is $o(n^2)$ -close to $U_n(F)$)

Human Part: Exact Result

- ▶ **Exact result:** One theorem covering all 8 cases
- ▶ **Assumptions:**
 - ▶ Stability (\forall almost extremal G is $o(n^2)$ -close to $U_n(F)$)
 - ▶ \forall block addition of a vertex x to $U_n(F)$:

Human Part: Exact Result

- ▶ **Exact result:** One theorem covering all 8 cases
- ▶ **Assumptions:**
 - ▶ Stability (\forall almost extremal G is $o(n^2)$ -close to $U_n(F)$)
 - ▶ \forall block addition of a vertex x to $U_n(F)$:
 - ▶ \overline{K}_ℓ is created

Human Part: Exact Result

- ▶ **Exact result:** One theorem covering all 8 cases
- ▶ **Assumptions:**
 - ▶ Stability (\forall almost extremal G is $o(n^2)$ -close to $U_n(F)$)
 - ▶ \forall block addition of a vertex x to $U_n(F)$:
 - ▶ \overline{K}_ℓ is created
 - ▶ **or** x is a clone of an existing vertex

Human Part: Exact Result

- ▶ **Exact result:** One theorem covering all 8 cases
- ▶ **Assumptions:**
 - ▶ Stability (\forall almost extremal G is $o(n^2)$ -close to $U_n(F)$)
 - ▶ \forall block addition of a vertex x to $U_n(F)$:
 - ▶ \overline{K}_ℓ is created
 - ▶ **or** x is a clone of an existing vertex
 - ▶ **or** too many K_k 's are created

Human Part: Exact Result

- ▶ **Exact result:** One theorem covering all 8 cases
- ▶ **Assumptions:**
 - ▶ Stability (\forall almost extremal G is $o(n^2)$ -close to $U_n(F)$)
 - ▶ \forall block addition of a vertex x to $U_n(F)$:
 - ▶ \overline{K}_ℓ is created
 - ▶ **or** x is a clone of an existing vertex
 - ▶ **or** too many K_k 's are created
- ▶ **Conclusion:**

$$f(n, k, \ell) = \#(K_k, \text{expansion of } F), \quad n > n_0$$

Open Problems

Open Problems


- ▶ Conjecture (P.-Vaughan):

$$c_{4,4} = \frac{14 \cdot 2^{1/3} - 11}{192}$$

Open Problems

- Conjecture (P.-Vaughan):


$$c_{4,4} = \frac{14 \cdot 2^{1/3} - 11}{192}$$



Open Problems

- Conjecture (P.-Vaughan):

$$c_{4,4} = \frac{14 \cdot 2^{1/3} - 11}{192}$$




- Das-Huang-Ma-Naves-Sudakov'13:
 $\ell_0 = \max\{\ell : c_{3,\ell} = (\ell - 1)^{-2}\} = ?$

Open Problems

- ▶ Conjecture (P.-Vaughan):

$$c_{4,4} = \frac{14 \cdot 2^{1/3} - 11}{192}$$



- ▶ Das-Huang-Ma-Naves-Sudakov'13:


$$\ell_0 = \max\{\ell : c_{3,\ell} = (\ell - 1)^{-2}\} = ?$$

- ▶ Nikiforov'01: $\ell_0 < \infty$

Open Problems

- ▶ Conjecture (P.-Vaughan):

$$c_{4,4} = \frac{14 \cdot 2^{1/3} - 11}{192}$$



- ▶ Das-Huang-Ma-Naves-Sudakov'13:


$$\ell_0 = \max\{\ell : c_{3,\ell} = (\ell - 1)^{-2}\} = ?$$

- ▶ Nikiforov'01: $\ell_0 < \infty$
- ▶ Das-Huang-Ma-Naves-Sudakov'13: $\ell_0 < 2074$

Open Problems

- ▶ Conjecture (P.-Vaughan):

$$c_{4,4} = \frac{14 \cdot 2^{1/3} - 11}{192}$$



- ▶ Das-Huang-Ma-Naves-Sudakov'13:

$$\ell_0 = \max\{\ell : c_{3,\ell} = (\ell - 1)^{-2}\} = ?$$

- ▶ Nikiforov'01: $\ell_0 < \infty$
- ▶ Das-Huang-Ma-Naves-Sudakov'13: $\ell_0 < 2074$
- ▶ P.-Vaughan \geq '13: $\ell_0 \geq 7$

Thank you!