

# Minimising the Number of Cliques

Oleg Pikhurko and Emil R. Vaughan

TUM, 18 July 2013

# Clique Minimisation Problem

# Clique Minimisation Problem

$$\blacktriangleright \#(F, G) = | \{ X \subseteq V(G) : G[X] \cong F \} |$$

# Clique Minimisation Problem

- ▶  $\#(F, G) = |\{X \subseteq V(G) : G[X] \cong F\}|$
- ▶  $e(G) = \#(K_2, G)$

# Clique Minimisation Problem

- ▶  $\#(F, G) = |\{X \subseteq V(G) : G[X] \cong F\}|$
- ▶  $e(G) = \#(K_2, G)$
- ▶ Independence number  $\alpha(G) = \max\{\ell : \#(\overline{K}_\ell, G) > 0\}$

# Clique Minimisation Problem

- ▶  $\#(F, G) = |\{X \subseteq V(G) : G[X] \cong F\}|$
- ▶  $e(G) = \#(K_2, G)$
- ▶ Independence number  $\alpha(G) = \max\{\ell : \#(\overline{K}_\ell, G) > 0\}$
- ▶ Erdős'62:

$$f(n, k, \ell) = \min\{\#(K_k, G) : \alpha(G) < \ell, \ v(G) = n\}$$

# Motivation

# Motivation

- ▶  $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$



# Motivation

- ▶  $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ Ramsey number

$$R(k, \ell) = \min\{n : \forall G_n \ \alpha(G_n) \geq \ell \text{ or } \#(K_k, G_n) > 0\}$$

# Motivation

- ▶  $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ Ramsey number

$$R(k, \ell) = \min\{n : \forall G_n \ \alpha(G_n) \geq \ell \text{ or } \#(K_k, G_n) > 0\}$$

- ▶  $n \geq R(k, \ell) \iff f(n, k, \ell) > 0$

# Ramsey Multiplicity Problem

# Ramsey Multiplicity Problem

- ▶ Erdős'62:  $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$

# Ramsey Multiplicity Problem

- ▶ Erdős'62:  $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶  $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$

# Ramsey Multiplicity Problem

- ▶ Erdős'62:  $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶  $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56:  $\mu_3 = 1/4$

# Ramsey Multiplicity Problem

- ▶ Erdős'62:  $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶  $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56:  $\mu_3 = 1/4$
- ▶ Erdős'62: Is  $\mu_k = 2^{1-\binom{k}{2}}$  ?

# Ramsey Multiplicity Problem

- ▶ Erdős'62:  $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶  $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56:  $\mu_3 = 1/4$
- ▶ Erdős'62: Is  $\mu_k = 2^{1 - \binom{k}{2}}$  ?
- ▶ Thomason'89: False:  $\mu_k \leq 0.936 \cdot 2^{1 - \binom{k}{2}}$



# Ramsey Multiplicity Problem

- ▶ Erdős'62:  $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶  $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56:  $\mu_3 = 1/4$
- ▶ Erdős'62: Is  $\mu_k = 2^{1-\binom{k}{2}}$  ?
- ▶ Thomason'89: False:  $\mu_k \leq 0.936 \cdot 2^{1-\binom{k}{2}}$
- ▶ Thomason'89:  $\mu_4 \leq \frac{1}{33} < \frac{1}{32}$

# Ramsey Multiplicity Problem

- ▶ Erdős'62:  $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶  $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56:  $\mu_3 = 1/4$
- ▶ Erdős'62: Is  $\mu_k = 2^{1 - \binom{k}{2}}$  ?
- ▶ Thomason'89: False:  $\mu_k \leq 0.936 \cdot 2^{1 - \binom{k}{2}}$
- ▶ Thomason'89:  $\mu_4 \leq \frac{1}{33} < \frac{1}{32}$
- ▶ Giraud'79:  $\mu_4 \geq \frac{1}{46} = 0.0217\dots$

# Ramsey Multiplicity Problem

- ▶ Erdős'62:  $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶  $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56:  $\mu_3 = 1/4$
- ▶ Erdős'62: Is  $\mu_k = 2^{1-\binom{k}{2}}$  ?
- ▶ Thomason'89: False:  $\mu_k \leq 0.936 \cdot 2^{1-\binom{k}{2}}$
- ▶ Thomason'89:  $\mu_4 \leq \frac{1}{33} < \frac{1}{32}$
- ▶ Giraud'79:  $\mu_4 \geq \frac{1}{46} = 0.0217\dots$
- ▶ Nieß  $\geq$ '13, Sperfeld  $\geq$ '13:  $\mu_4 \geq 0.0287\dots$

# Ramsey Multiplicity Problem

- ▶ Erdős'62:  $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶  $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56:  $\mu_3 = 1/4$
- ▶ Erdős'62: Is  $\mu_k = 2^{1-\binom{k}{2}}$  ?
- ▶ Thomason'89: False:  $\mu_k \leq 0.936 \cdot 2^{1-\binom{k}{2}}$
- ▶ Thomason'89:  $\mu_4 \leq \frac{1}{33} < \frac{1}{32}$
- ▶ Giraud'79:  $\mu_4 \geq \frac{1}{46} = 0.0217\dots$
- ▶ Nieß  $\geq$ '13, Sperfeld  $\geq$ '13:  $\mu_4 \geq 0.0287\dots$
- ▶ Erdős'62:  $\mu_k \geq \binom{R(k,k)}{k}^{-1} \geq (4 + o(1))^{-k^2}$

# Ramsey Multiplicity Problem

- ▶ Erdős'62:  $m(k, n) = \min\{\#(K_k, G_n) + \#(\overline{K}_k, G_n)\}$
- ▶  $\mu_k = \lim_{n \rightarrow \infty} m(k, n) / \binom{n}{k}$
- ▶ Goodman'56:  $\mu_3 = 1/4$
- ▶ Erdős'62: Is  $\mu_k = 2^{1-\binom{k}{2}}$  ?
- ▶ Thomason'89: False:  $\mu_k \leq 0.936 \cdot 2^{1-\binom{k}{2}}$
- ▶ Thomason'89:  $\mu_4 \leq \frac{1}{33} < \frac{1}{32}$
- ▶ Giraud'79:  $\mu_4 \geq \frac{1}{46} = 0.0217\dots$
- ▶ Nieß  $\geq$ '13, Sperfeld  $\geq$ '13:  $\mu_4 \geq 0.0287\dots$
- ▶ Erdős'62:  $\mu_k \geq \binom{R(k,k)}{k}^{-1} \geq (4 + o(1))^{-k^2}$
- ▶ Conlon'12:  $\mu_k \geq (2.18 + o(1))^{-k^2}$

# Back to Clique Minimisation

# Back to Clique Minimisation

- ▶  $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$

# Back to Clique Minimisation

- ▶  $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ **Goodman'56:**  $f(2m, 3, 3) = \#(K_3, K_m \sqcup K_m)$



# Back to Clique Minimisation

- ▶  $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ **Goodman'56:**  $f(2m, 3, 3) = \#(K_3, K_m \sqcup K_m)$
- ▶ **Lorden'62:**  $f(2m+1, 3, 3) = \#(K_3, K_m \sqcup K_{m+1})$ ,  $n \geq 6$

# Back to Clique Minimisation

- ▶  $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ **Goodman'56:**  $f(2m, 3, 3) = \#(K_3, K_m \sqcup K_m)$
- ▶ **Lorden'62:**  $f(2m+1, 3, 3) = \#(K_3, K_m \sqcup K_{m+1})$ ,  $n \geq 6$
- ▶ Complement of  $T_n^2$

# Back to Clique Minimisation

- ▶  $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ **Goodman'56:**  $f(2m, 3, 3) = \#(K_3, K_m \sqcup K_m)$
- ▶ **Lorden'62:**  $f(2m+1, 3, 3) = \#(K_3, K_m \sqcup K_{m+1})$ ,  $n \geq 6$
- ▶ Complement of  $T_n^2$
- ▶ **Turán graph  $T_n^r$ :** max  $r$ -partite order- $n$  graph

# Back to Clique Minimisation

- ▶  $f(n, k, \ell) = \min\{\#(K_k, G_n) : \alpha(G_n) < \ell\}$
- ▶ **Goodman'56:**  $f(2m, 3, 3) = \#(K_3, K_m \sqcup K_m)$
- ▶ **Lorden'62:**  $f(2m+1, 3, 3) = \#(K_3, K_m \sqcup K_{m+1})$ ,  $n \geq 6$
- ▶ Complement of  $T_n^2$
- ▶ **Turán graph  $T_n^r$ :** max  $r$ -partite order- $n$  graph
- ▶ **Erdős'62:** Is  $f(n, k, \ell) = \#(K_3, \overline{T}_n^{\ell-1})$  ?

# Asymptotic densities

# Asymptotic densities

- ▶ Nikiforov'01:  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$

# Asymptotic densities

- ▶ Nikiforov'01:  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$

# Asymptotic densities

- ▶ Nikiforov'01:  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ Nikiforov'01: Strict for all but finitely many  $(k, \ell)$



# Asymptotic densities

- ▶ Nikiforov'01:  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ Nikiforov'01: Strict for all but finitely many  $(k, \ell)$
- ▶ Expansion of  $F$ :

# Asymptotic densities

- ▶ Nikiforov'01:  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ Nikiforov'01: Strict for all but finitely many  $(k, \ell)$
- ▶ Expansion of  $F$ :
  - ▶ Replace each  $i \in V(F)$  by clique on  $K[V_i]$

# Asymptotic densities

- ▶ **Nikiforov'01:**  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many  $(k, \ell)$
- ▶ **Expansion** of  $F$ :
  - ▶ Replace each  $i \in V(F)$  by clique on  $K[V_i]$
  - ▶ Replace each  $ij \in E(F)$  by complete graph  $K[V_i, V_j]$

# Asymptotic densities

- ▶ **Nikiforov'01:**  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many  $(k, \ell)$
- ▶ **Expansion** of  $F$ :
  - ▶ Replace each  $i \in V(F)$  by clique on  $K[V_i]$
  - ▶ Replace each  $ij \in E(F)$  by complete graph  $K[V_i, V_j]$
- ▶ **Uniform expansion**  $U_n(F)$ :  $\forall i, j \quad |V_i - V_j| \leq 1$

# Asymptotic densities

- ▶ **Nikiforov'01:**  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many  $(k, \ell)$
- ▶ **Expansion** of  $F$ :
  - ▶ Replace each  $i \in V(F)$  by clique on  $K[V_i]$
  - ▶ Replace each  $ij \in E(F)$  by complete graph  $K[V_i, V_j]$
- ▶ **Uniform expansion**  $U_n(F)$ :  $\forall i, j \mid |V_i - V_j| \leq 1$
- ▶  $\overline{T}_n^m = U_n(\overline{K}_m)$

# Asymptotic densities

- ▶ **Nikiforov'01:**  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many  $(k, \ell)$
- ▶ **Expansion** of  $F$ :
  - ▶ Replace each  $i \in V(F)$  by clique on  $K[V_i]$
  - ▶ Replace each  $ij \in E(F)$  by complete graph  $K[V_i, V_j]$
- ▶ **Uniform expansion**  $U_n(F)$ :  $\forall i, j \mid |V_i - V_j| \leq 1$
- ▶  $\overline{T}_n^m = U_n(\overline{K}_m)$
- ▶ **Nikiforov'01:**  $f(n, 4, 3) \leq \#(K_4, U_n(C_5))$

# Asymptotic densities

- ▶ **Nikiforov'01:**  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many  $(k, \ell)$
- ▶ **Expansion** of  $F$ :
  - ▶ Replace each  $i \in V(F)$  by clique on  $K[V_i]$
  - ▶ Replace each  $ij \in E(F)$  by complete graph  $K[V_i, V_j]$
- ▶ **Uniform expansion**  $U_n(F)$ :  $\forall i, j \mid |V_i - V_j| \leq 1$
- ▶  $\overline{T}_n^m = U_n(\overline{K}_m)$
- ▶ **Nikiforov'01:**  $f(n, 4, 3) \leq \#(K_4, U_n(C_5))$ 
  - ▶  $c_{4,3} \leq 3/25$

# Asymptotic densities

- ▶ **Nikiforov'01:**  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many  $(k, \ell)$
- ▶ **Expansion of  $F$ :**
  - ▶ Replace each  $i \in V(F)$  by clique on  $K[V_i]$
  - ▶ Replace each  $ij \in E(F)$  by complete graph  $K[V_i, V_j]$
- ▶ **Uniform expansion  $U_n(F)$ :**  $\forall i, j \quad |V_i - V_j| \leq 1$
- ▶  $\overline{T}_n^m = U_n(\overline{K}_m)$
- ▶ **Nikiforov'01:**  $f(n, 4, 3) \leq \#(K_4, U_n(C_5))$ 
  - ▶  $c_{4,3} \leq 3/25 < 1/8$



# Asymptotic densities

- ▶ **Nikiforov'01:**  $c_{k,\ell} = \lim_{n \rightarrow \infty} f(n, k, \ell) / \binom{n}{k}$
- ▶ Turán graph:  $c_{k,\ell} \leq (\ell - 1)^{1-k}$
- ▶ **Nikiforov'01:** Strict for all but finitely many  $(k, \ell)$
- ▶ **Expansion of  $F$ :**
  - ▶ Replace each  $i \in V(F)$  by clique on  $K[V_i]$
  - ▶ Replace each  $ij \in E(F)$  by complete graph  $K[V_i, V_j]$
- ▶ **Uniform expansion  $U_n(F)$ :**  $\forall i, j \quad |V_i - V_j| \leq 1$
- ▶  $\overline{T}_n^m = U_n(\overline{K}_m)$
- ▶ **Nikiforov'01:**  $f(n, 4, 3) \leq \#(K_4, U_n(C_5))$ 
  - ▶  $c_{4,3} \leq 3/25 < 1/8$
- ▶ **Nikiforov'05:**  
 $\alpha(G_n) < 3$  & **regular**  $\Rightarrow \#(K_4, G) \geq (\frac{3}{25} + o(1)) \binom{n}{4}$

Complete Solutions (for  $n \geq n_0$ )

# Complete Solutions (for $n \geq n_0$ )

- ▶ Das-Huang-Ma-Naves-Sudakov'13:  $(3, 4)$  &  $(4, 3)$

# Complete Solutions (for $n \geq n_0$ )

- ▶ **Das-Huang-Ma-Naves-Sudakov'13:**  $(3, 4)$  &  $(4, 3)$
- ▶ **P.-Vaughan  $\geq$ '13:**  $(3, 4)$ ,  $(3, 5)$ ,  $(3, 6)$ ,  $(3, 7)$ ,  $(4, 3)$ ,  $(5, 3)$ ,  $(6, 3)$  &  $(7, 3)$

New Values of  $c_{k,l}$

## New Values of $c_{k,\ell}$

- ▶  $c_{3,\ell} = (\ell - 1)^2$  for  $4 \leq \ell \leq 7$

# New Values of $c_{k,\ell}$

- ▶  $c_{3,\ell} = (\ell - 1)^2$  for  $4 \leq \ell \leq 7$ 
  - ▶  $\overline{T}_n^{\ell-1}$

# New Values of $c_{k,\ell}$

- ▶  $c_{3,\ell} = (\ell - 1)^2$  for  $4 \leq \ell \leq 7$ 
  - ▶  $\overline{T}_n^{\ell-1}$
- ▶  $c_{4,3} = 3/25$



# New Values of $c_{k,\ell}$

- ▶  $c_{3,\ell} = (\ell - 1)^2$  for  $4 \leq \ell \leq 7$ 
  - ▶  $\overline{T}_n^{\ell-1}$
- ▶  $c_{4,3} = 3/25$
- ▶  $c_{5,3} = 31/625$

# New Values of $c_{k,\ell}$

- ▶  $c_{3,\ell} = (\ell - 1)^2$  for  $4 \leq \ell \leq 7$ 
  - ▶  $\overline{T}_n^{\ell-1}$
- ▶  $c_{4,3} = 3/25$
- ▶  $c_{5,3} = 31/625$ 
  - ▶ Uniform expansion of  $C_5$

# New Values of $c_{k,\ell}$

- ▶  $c_{3,\ell} = (\ell - 1)^2$  for  $4 \leq \ell \leq 7$ 
  - ▶  $\overline{T}_n^{\ell-1}$
- ▶  $c_{4,3} = 3/25$
- ▶  $c_{5,3} = 31/625$ 
  - ▶ Uniform expansion of  $C_5$
- ▶  $c_{6,3} = 19211/2^{20} = 19211/1048576$

# New Values of $c_{k,\ell}$

- ▶  $c_{3,\ell} = (\ell - 1)^2$  for  $4 \leq \ell \leq 7$ 
  - ▶  $\overline{T}_n^{\ell-1}$
- ▶  $c_{4,3} = 3/25$
- ▶  $c_{5,3} = 31/625$ 
  - ▶ Uniform expansion of  $C_5$
- ▶  $c_{6,3} = 19211/2^{20} = 19211/1048576$
- ▶  $c_{7,3} = 98491/2^{24} = 98491/16777216$

## Construction for $c_{6,3}$ and $c_{7,3}$

# Construction for $c_{6,3}$ and $c_{7,3}$

- ▶ Clebsch graph  $L$ :

# Construction for $c_{6,3}$ and $c_{7,3}$

- ▶ Clebsch graph  $L$ :
  - ▶  $V = \{\mathbf{X} \in \{0, 1\}^5 : \text{dist}(\mathbf{X}, \mathbf{0}) \text{ is even}\}$

# Construction for $c_{6,3}$ and $c_{7,3}$

- ▶ Clebsch graph  $L$ :
  - ▶  $V = \{\mathbf{X} \in \{0, 1\}^5 : \text{dist}(\mathbf{X}, \mathbf{0}) \text{ is even}\}$
  - ▶  $\mathbf{X} \sim \mathbf{Y}$  iff  $\text{dist}(\mathbf{X}, \mathbf{Y}) = 4$





# Construction for $c_{6,3}$ and $c_{7,3}$

- ▶ Clebsch graph  $L$ :
  - ▶  $V = \{\mathbf{X} \in \{0, 1\}^5 : \text{dist}(\mathbf{X}, \mathbf{0}) \text{ is even}\}$
  - ▶  $\mathbf{X} \sim \mathbf{Y}$  iff  $\text{dist}(\mathbf{X}, \mathbf{Y}) = 4$
- ▶ Upper bound:  $U_n(\bar{L})$

# Construction for $c_{6,3}$ and $c_{7,3}$

- ▶ **Clebsch graph  $L$ :**
  - ▶  $V = \{\mathbf{X} \in \{0, 1\}^5 : \text{dist}(\mathbf{X}, \mathbf{0}) \text{ is even}\}$
  - ▶  $\mathbf{X} \sim \mathbf{Y}$  iff  $\text{dist}(\mathbf{X}, \mathbf{Y}) = 4$
- ▶ **Upper bound:**  $U_n(\bar{L})$
- ▶  $L$  and  $\bar{L}$ :



# Flag Algebras

# Flag Algebras

- Subgraph density

$$\phi(F, G) = \frac{\#(F, G)}{\binom{v(G)}{v(F)}} = \mathbf{Prob}\{G[\text{random } v(F)\text{-set}] \cong F\}$$

# Flag Algebras

- ▶ Subgraph density

$$\phi(F, G) = \frac{\#(F, G)}{\binom{v(G)}{v(F)}} = \mathbf{Prob}\{G[\text{random } v(F)\text{-set}] \cong F\}$$

- ▶ Razborov'07: Flag algebras

# Flag Algebras

- ▶ Subgraph density

$$\phi(F, G) = \frac{\#(F, G)}{\binom{v(G)}{v(F)}} = \mathbf{Prob}\{G[\text{random } v(F)\text{-set}] \cong F\}$$

- ▶ Razborov'07: Flag algebras
  - ▶ Asymptotically true inequalities (as  $v(G) \rightarrow \infty$ )

# Flag Algebras

- ▶ Subgraph density

$$\phi(F, G) = \frac{\#(F, G)}{\binom{v(G)}{v(F)}} = \mathbf{Prob}\{G[\text{random } v(F)\text{-set}] \cong F\}$$

- ▶ Razborov'07: Flag algebras

- ▶ Asymptotically true inequalities (as  $v(G) \rightarrow \infty$ )
- ▶ Formal rules for deriving them

# Flag Algebras

- ▶ Subgraph density

$$\phi(F, G) = \frac{\#(F, G)}{\binom{v(G)}{v(F)}} = \mathbf{Prob}\{G[\text{random } v(F)\text{-set}] \cong F\}$$

- ▶ Razborov'07: Flag algebras

- ▶ Asymptotically true inequalities (as  $v(G) \rightarrow \infty$ )
- ▶ Formal rules for deriving them
- ▶ One aspect: semi-definite programming



# Turán Problem via Flag Algebras

# Turán Problem via Flag Algebras

- ▶ Turán function:

$$\text{ex}(n, K_3) = \max\{e(G) : v(G) = n, G \not\supseteq K_3\}$$

# Turán Problem via Flag Algebras

- ▶ Turán function:

$$\text{ex}(n, K_3) = \max\{e(G) : v(G) = n, G \not\supseteq K_3\}$$

- ▶ Construction:  $\text{ex}(n, K_3) \geq e(T_n^2) = \lfloor n^2/4 \rfloor$

# Turán Problem via Flag Algebras

- ▶ Turán function:

$$\text{ex}(n, K_3) = \max\{e(G) : v(G) = n, G \not\supseteq K_3\}$$

- ▶ **Construction:**  $\text{ex}(n, K_3) \geq e(T_n^2) = \lfloor n^2/4 \rfloor$
- ▶  $T_n^2 = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$

# Subgraph Densities

# Subgraph Densities

- ▶  $K_3$ -free  $G$  of "large" order  $n$

# Subgraph Densities

- ▶  $K_3$ -free  $G$  of "large" order  $n$
- ▶  $\#F = \#(F, G) = \#$  induced copies of  $F$  in  $G$

# Subgraph Densities

- ▶  $K_3$ -free  $G$  of "large" order  $n$
- ▶  $\#F = \#(F, G) = \#$  induced copies of  $F$  in  $G$
- ▶  $\phi(F) = \#F / \binom{n}{v(F)}$



# Obtaining Inequalities

# Obtaining Inequalities

$$\blacktriangleright \mathbf{A} = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq \mathbf{0}$$

# Obtaining Inequalities

- ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶  $\mathbf{v}_x = \left( \frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1} \right), \quad x \in V(G)$

# Obtaining Inequalities

- ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶  $\mathbf{v}_x = \left( \frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1} \right), \quad x \in V(G)$
- ▶  $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$

# Obtaining Inequalities

- ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶  $\mathbf{v}_x = \left( \frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1} \right), \quad x \in V(G)$
- ▶  $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over  $x$  (and ignore  $o(1)$ ):

# Obtaining Inequalities

- ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶  $\mathbf{v}_x = \left( \frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1} \right), \quad x \in V(G)$
- ▶  $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over  $x$  (and ignore  $o(1)$ ):

$$\frac{1}{n} \sum_x \alpha \frac{d^2(x)}{(n-1)^2}$$

# Obtaining Inequalities

- ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶  $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$ ,  $x \in V(G)$
- ▶  $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over  $x$  (and ignore  $o(1)$ ):

$$\frac{1}{n} \sum_x \alpha \frac{d^2(x)}{(n-1)^2} = \frac{\alpha}{n^3} \sum_x \left( \sum_{y \sim x} 1 \right) \left( \sum_{z \sim x} 1 \right)$$

# Obtaining Inequalities

- ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶  $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$ ,  $x \in V(G)$
- ▶  $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over  $x$  (and ignore  $o(1)$ ):

$$\begin{aligned} \frac{1}{n} \sum_x \alpha \frac{d^2(x)}{(n-1)^2} &= \frac{\alpha}{n^3} \sum_x \left( \sum_{y \sim x} 1 \right) \left( \sum_{z \sim x} 1 \right) \\ &= \frac{\alpha}{6 \binom{n}{3}} \times 2 \#P_3 \end{aligned}$$



# Obtaining Inequalities

- ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶  $\mathbf{v}_x = (\frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1})$ ,  $x \in V(G)$
- ▶  $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over  $x$  (and ignore  $o(1)$ ):

$$\begin{aligned} \frac{1}{n} \sum_x \alpha \frac{d^2(x)}{(n-1)^2} &= \frac{\alpha}{n^3} \sum_x \left( \sum_{y \sim x} 1 \right) \left( \sum_{z \sim x} 1 \right) \\ &= \frac{\alpha}{6 \binom{n}{3}} \times 2 \#P_3 = \frac{\alpha}{3} \phi(P_3) \end{aligned}$$

# Obtaining Inequalities

- ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶  $\mathbf{v}_x = \left( \frac{d(x)}{n-1}, \frac{\bar{d}(x)}{n-1} \right), \quad x \in V(G)$
- ▶  $0 \leq \mathbf{v}_x A \mathbf{v}_x^T$
- ▶ **Average** over  $x$  (and ignore  $o(1)$ ):

$$\begin{aligned} \frac{1}{n} \sum_x \alpha \frac{d^2(x)}{(n-1)^2} &= \frac{\alpha}{n^3} \sum_x \left( \sum_{y \sim x} 1 \right) \left( \sum_{z \sim x} 1 \right) \\ &= \frac{\alpha}{6 \binom{n}{3}} \times 2 \#P_3 = \frac{\alpha}{3} \phi(P_3) \end{aligned}$$

- ▶  $0 \leq (2\beta + \alpha)\phi(P_3) + (2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$

# Bounding Edge Density from Above

# Bounding Edge Density from Above

- ▶  $0 \leq (2\beta + \alpha)\phi(P_3) + (2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$

# Bounding Edge Density from Above

- ▶  $0 \leq (2\beta + \alpha)\phi(P_3) + (2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶  $\phi(K_2)$

# Bounding Edge Density from Above

- ▶  $0 \leq (2\beta + \alpha)\phi(P_3) + (2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶  $\phi(K_2) = \frac{2}{3}\phi(P_3) + \frac{1}{3}\phi(\overline{P}_3)$

# Bounding Edge Density from Above

- ▶  $0 \leq (2\beta + \alpha)\phi(P_3) + (2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶  $\phi(K_2) = \frac{2}{3}\phi(P_3) + \frac{1}{3}\phi(\overline{P}_3)$   
 $\leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$

# Bounding Edge Density from Above

- ▶  $0 \leq (2\beta + \alpha)\phi(P_3) + (2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶  $\phi(K_2) = \frac{2}{3}\phi(P_3) + \frac{1}{3}\phi(\overline{P}_3)$   
 $\leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶ **Note:**  $\phi(P_3) + \phi(\overline{P}_3) + \phi(\overline{K}_3) = 1$



# SDP Formulation

# SDP Formulation

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$

# SDP Formulation

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶ **Minimise**  $\delta$ :

# SDP Formulation

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶ **Minimise**  $\delta$ :
  - ▶  $\frac{2}{3} + 2\beta + \alpha \leq \delta$

# SDP Formulation

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶ **Minimise**  $\delta$ :
  - ▶  $\frac{2}{3} + 2\beta + \alpha \leq \delta$
  - ▶  $\frac{1}{3} + 2\beta + \gamma \leq \delta$

# SDP Formulation

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶ **Minimise**  $\delta$ :
  - ▶  $\frac{2}{3} + 2\beta + \alpha \leq \delta$
  - ▶  $\frac{1}{3} + 2\beta + \gamma \leq \delta$
  - ▶  $3\gamma \leq \delta$

# SDP Formulation

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶ **Minimise**  $\delta$ :
  - ▶  $\frac{2}{3} + 2\beta + \alpha \leq \delta$
  - ▶  $\frac{1}{3} + 2\beta + \gamma \leq \delta$
  - ▶  $3\gamma \leq \delta$
  - ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$

# SDP Formulation

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶ **Minimise**  $\delta$ :
  - ▶  $\frac{2}{3} + 2\beta + \alpha \leq \delta$
  - ▶  $\frac{1}{3} + 2\beta + \gamma \leq \delta$
  - ▶  $3\gamma \leq \delta$
  - ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ **Solution:**  $\delta = 1/2$  and  $A = \begin{pmatrix} 1/6 & -1/6 \\ -1/6 & 1/6 \end{pmatrix}$



# SDP Formulation

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ **Minimise**  $\delta$ :
  - ▶  $\frac{2}{3} + 2\beta + \alpha \leq \delta$
  - ▶  $\frac{1}{3} + 2\beta + \gamma \leq \delta$
  - ▶  $3\gamma \leq \delta$
  - ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ **Solution:**  $\delta = 1/2$  and  $A = \begin{pmatrix} 1/6 & -1/6 \\ -1/6 & 1/6 \end{pmatrix}$
- ▶  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6} \phi(\bar{P}_3)$

# SDP Formulation

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶ **Minimise**  $\delta$ :
  - ▶  $\frac{2}{3} + 2\beta + \alpha \leq \delta$
  - ▶  $\frac{1}{3} + 2\beta + \gamma \leq \delta$
  - ▶  $3\gamma \leq \delta$
  - ▶  $A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} \succeq 0$
- ▶ **Solution:**  $\delta = 1/2$  and  $A = \begin{pmatrix} 1/6 & -1/6 \\ -1/6 & 1/6 \end{pmatrix}$
- ▶  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6} \phi(\overline{P}_3)$
- ▶ **Asymptotic result:**  $\text{ex}(n, K_3) \leq (\frac{1}{2} + o(1))\binom{n}{2}$

# Interpreting the Solution

# Interpreting the Solution

► Recall  $A = \begin{pmatrix} 1/6 & -1/6 \\ -1/6 & 1/6 \end{pmatrix}$

# Interpreting the Solution

- ▶ Recall  $A = \begin{pmatrix} 1/6 & -1/6 \\ -1/6 & 1/6 \end{pmatrix}$
- ▶  $\mathbf{v}_x A \mathbf{v}_x^T = \frac{1}{6(n-1)^2} (d(x) - \bar{d}(x))^2$

# Interpreting the Solution

- ▶ Recall  $A = \begin{pmatrix} 1/6 & -1/6 \\ -1/6 & 1/6 \end{pmatrix}$
- ▶  $\mathbf{v}_x A \mathbf{v}_x^T = \frac{1}{6(n-1)^2} (d(x) - \bar{d}(x))^2$
- ▶ **Human proof:**

# Interpreting the Solution

- ▶ Recall  $A = \begin{pmatrix} 1/6 & -1/6 \\ -1/6 & 1/6 \end{pmatrix}$
- ▶  $\mathbf{v}_x A \mathbf{v}_x^T = \frac{1}{6(n-1)^2} (d(x) - \bar{d}(x))^2$
- ▶ **Human proof:**
  - ▶  $0 \leq \sum_x (d(x) - \bar{d}(x))^2$

# Interpreting the Solution

- ▶ Recall  $A = \begin{pmatrix} 1/6 & -1/6 \\ -1/6 & 1/6 \end{pmatrix}$
- ▶  $\mathbf{v}_x A \mathbf{v}_x^T = \frac{1}{6(n-1)^2} (d(x) - \bar{d}(x))^2$
- ▶ **Human proof:**
  - ▶  $0 \leq \sum_x (d(x) - \bar{d}(x))^2$
  - ▶ Expand and re-group terms



# Interpreting the Solution

- ▶ Recall  $A = \begin{pmatrix} 1/6 & -1/6 \\ -1/6 & 1/6 \end{pmatrix}$
- ▶  $\mathbf{v}_x A \mathbf{v}_x^T = \frac{1}{6(n-1)^2} (d(x) - \bar{d}(x))^2$
- ▶ **Human proof:**
  - ▶  $0 \leq \sum_x (d(x) - \bar{d}(x))^2$
  - ▶ Expand and re-group terms
  - ▶  $\#K_2 \leq \frac{n^2}{4} - \frac{1}{n} \#P_3 + O(n)$

# Rounding Floating-Point Solution

# Rounding Floating-Point Solution

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$

# Rounding Floating-Point Solution

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.166672 & -0.166665 \\ -0.166665 & 0.166680 \end{pmatrix} ?$$

# Rounding Floating-Point Solution

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.166672 & -0.166665 \\ -0.166665 & 0.166680 \end{pmatrix} ?$$

- ▶ Suppose  $A \succeq 0$  **proves**  $\phi(K_2) \leq \frac{1}{2}$

# Rounding Floating-Point Solution

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.166672 & -0.166665 \\ -0.166665 & 0.166680 \end{pmatrix} ?$$

- ▶ Suppose  $A \succeq 0$  **proves**  $\phi(K_2) \leq \frac{1}{2}$
- ▶ If  $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$ :

# Rounding Floating-Point Solution

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.166672 & -0.166665 \\ -0.166665 & 0.166680 \end{pmatrix} ?$$

- ▶ Suppose  $A \succeq 0$  **proves**  $\phi(K_2) \leq \frac{1}{2}$
- ▶ If  $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$ :
  - ▶ Then  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6}\phi(\bar{P}_3) - \varepsilon\phi(P_3)$

# Rounding Floating-Point Solution

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.166672 & -0.166665 \\ -0.166665 & 0.166680 \end{pmatrix} ?$$

- ▶ Suppose  $A \succeq 0$  **proves**  $\phi(K_2) \leq \frac{1}{2}$
- ▶ If  $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$ :
  - ▶ Then  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6}\phi(\overline{P}_3) - \varepsilon\phi(P_3)$
  - ▶ **False** for  $K_{n/2, n/2}$  !



# Rounding Floating-Point Solution

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\overline{P}_3) + 3\gamma\phi(\overline{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.166672 & -0.166665 \\ -0.166665 & 0.166680 \end{pmatrix} ?$$

- ▶ Suppose  $A \succeq 0$  **proves**  $\phi(K_2) \leq \frac{1}{2}$
- ▶ If  $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$ :
  - ▶ Then  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6}\phi(\overline{P}_3) - \varepsilon\phi(P_3)$
  - ▶ **False** for  $K_{n/2, n/2}$  !
- ▶ **Likewise:**  $3\gamma = \frac{1}{2}$

# Rounding Floating-Point Solution

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.166672 & -0.166665 \\ -0.166665 & 0.166680 \end{pmatrix} ?$$

- ▶ Suppose  $A \succeq 0$  **proves**  $\phi(K_2) \leq \frac{1}{2}$
- ▶ If  $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$ :
  - ▶ Then  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6}\phi(\bar{P}_3) - \varepsilon\phi(P_3)$
  - ▶ **False** for  $K_{n/2, n/2}$  !
- ▶ **Likewise:**  $3\gamma = \frac{1}{2}$
- ▶  $\phi(K_2) + \mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T) \approx \frac{1}{2} - \frac{1}{6}\phi(\bar{P}_3)$

# Rounding Floating-Point Solution

- ▶  $\phi(K_2) \leq (\frac{2}{3} + 2\beta + \alpha)\phi(P_3) + (\frac{1}{3} + 2\beta + \gamma)\phi(\bar{P}_3) + 3\gamma\phi(\bar{K}_3)$
- ▶ How to round

$$A = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix} = \begin{pmatrix} 0.166672 & -0.166665 \\ -0.166665 & 0.166680 \end{pmatrix} ?$$

- ▶ Suppose  $A \succeq 0$  **proves**  $\phi(K_2) \leq \frac{1}{2}$
- ▶ If  $\varepsilon = \frac{1}{2} - (\frac{2}{3} + 2\beta + \alpha) > 0$ :
  - ▶ Then  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6}\phi(\bar{P}_3) - \varepsilon\phi(P_3)$
  - ▶ **False** for  $K_{n/2, n/2}$  !
- ▶ **Likewise:**  $3\gamma = \frac{1}{2}$
- ▶  $\phi(K_2) + \mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T) \approx \frac{1}{2} - \frac{1}{6}\phi(\bar{P}_3) \Rightarrow A \begin{pmatrix} 1/2 \\ 1/2 \end{pmatrix} = 0$

# Asymptotic Structure of Extremal Graphs

# Asymptotic Structure of Extremal Graphs

- ▶ **Aim:** almost extremal graphs looks like  $T_n^2$

# Asymptotic Structure of Extremal Graphs

- ▶ **Aim:** almost extremal graphs looks like  $T_n^2$
- ▶  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6} \phi(\overline{P}_3)$

# Asymptotic Structure of Extremal Graphs

- ▶ **Aim:** almost extremal graphs looks like  $T_n^2$
- ▶  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6} \phi(\overline{P}_3)$
- ▶  $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\overline{P}_3) = o(1)$

# Asymptotic Structure of Extremal Graphs

- ▶ **Aim:** almost extremal graphs looks like  $T_n^2$
- ▶  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6} \phi(\overline{P}_3)$
- ▶  $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\overline{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**  
 $G$  is  $o(n^2)$ -close in edit distance to  $\overline{P}_3$ -free



# Asymptotic Structure of Extremal Graphs

- ▶ **Aim:** almost extremal graphs looks like  $T_n^2$
- ▶  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6} \phi(\overline{P}_3)$
- ▶  $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\overline{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**  
 $G$  is  $o(n^2)$ -close in edit distance to  $\overline{P}_3$ -free
- ▶  $\overline{P}_3$ -free  $\Rightarrow$  complete multipartite

# Asymptotic Structure of Extremal Graphs

- ▶ **Aim:** almost extremal graphs looks like  $T_n^2$
- ▶  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6} \phi(\overline{P}_3)$
- ▶  $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\overline{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**  
 $G$  is  $o(n^2)$ -close in edit distance to  $\overline{P}_3$ -free
- ▶  $\overline{P}_3$ -free  $\Rightarrow$  complete multipartite
- ▶  $K_3$ -free  $\Rightarrow$  at most 2 parts

# Asymptotic Structure of Extremal Graphs

- ▶ **Aim:** almost extremal graphs looks like  $T_n^2$
- ▶  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6} \phi(\overline{P}_3)$
- ▶  $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\overline{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**  
 $G$  is  $o(n^2)$ -close in edit distance to  $\overline{P}_3$ -free
- ▶  $\overline{P}_3$ -free  $\Rightarrow$  complete multipartite
- ▶  $K_3$ -free  $\Rightarrow$  at most 2 parts
- ▶  $\phi(K_2) \approx \frac{1}{4} \Rightarrow$  parts almost equal

# Asymptotic Structure of Extremal Graphs

- ▶ **Aim:** almost extremal graphs looks like  $T_n^2$
- ▶  $\phi(K_2) \leq \frac{1}{2} - \frac{1}{6} \phi(\overline{P}_3)$
- ▶  $\phi(K_2) \approx \frac{1}{2} \Rightarrow \phi(\overline{P}_3) = o(1)$
- ▶ **Induced Removal Lemma:**  
 $G$  is  $o(n^2)$ -close in edit distance to  $\overline{P}_3$ -free
- ▶  $\overline{P}_3$ -free  $\Rightarrow$  complete multipartite
- ▶  $K_3$ -free  $\Rightarrow$  at most 2 parts
- ▶  $\phi(K_2) \approx \frac{1}{4} \Rightarrow$  parts almost equal
- ▶ **Stability (Erdős'67, Simonovits'68):**  
 $\forall$  almost extremal  $G_n$  is  $o(n^2)$ -close to  $T_n^2$

## P.-Vaughan's Proof for $f(n, k, \ell)$

# P.-Vaughan's Proof for $f(n, k, \ell)$

- ▶ Emil Vaughan's package `Flagmatic`

# P.-Vaughan's Proof for $f(n, k, \ell)$

- ▶ Emil Vaughan's package `Flagmatic`
- ▶ **Stability**: above ideas + eigenvalues

# P.-Vaughan's Proof for $f(n, k, \ell)$

- ▶ Emil Vaughan's package `Flagmatic`
- ▶ **Stability**: above ideas + eigenvalues
  - ▶  $\mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T) = o(1) \Rightarrow$  structural information



# P.-Vaughan's Proof for $f(n, k, \ell)$

- ▶ Emil Vaughan's package `Flagmatic`
- ▶ **Stability:** above ideas + eigenvalues
  - ▶  $\mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T) = o(1) \Rightarrow$  structural information
- ▶ **Exact result:** One theorem covering all 8 cases

# P.-Vaughan's Proof for $f(n, k, \ell)$

- ▶ Emil Vaughan's package `Flagmatic`
- ▶ **Stability:** above ideas + eigenvalues
  - ▶  $\mathbb{E}_x(\mathbf{v}_x \mathbf{A} \mathbf{v}_x^T) = o(1) \Rightarrow$  structural information
- ▶ **Exact result:** One theorem covering all 8 cases
  - ▶ Stability &  $\text{grad}(\forall \text{ local move}) > 0 \Rightarrow$  exact result

# Open Problems

# Open Problems


- Conjecture (P.-Vaughan):

$$c_{4,4} = \frac{-11 + 14 \cdot 2^{1/3}}{192}$$

# Open Problems

- Conjecture (P.-Vaughan):


$$c_{4,4} = \frac{-11 + 14 \cdot 2^{1/3}}{192}$$



# Open Problems

- Conjecture (P.-Vaughan):

$$c_{4,4} = \frac{-11 + 14 \cdot 2^{1/3}}{192}$$




- $\ell_0 = \max\{\ell : c_{3,\ell} = (\ell - 1)^{-2}\}$

# Open Problems

- Conjecture (P.-Vaughan):

$$c_{4,4} = \frac{-11 + 14 \cdot 2^{1/3}}{192}$$




- $\ell_0 = \max\{\ell : c_{3,\ell} = (\ell - 1)^{-2}\}$ 
  - Nikiforov'01:  $\ell_0 < \infty$

# Open Problems

- ▶ Conjecture (P.-Vaughan):

$$c_{4,4} = \frac{-11 + 14 \cdot 2^{1/3}}{192}$$




- ▶  $\ell_0 = \max\{\ell : c_{3,\ell} = (\ell - 1)^{-2}\}$ 
  - ▶ Nikiforov'01:  $\ell_0 < \infty$
  - ▶ Das-Huang-Ma-Naves-Sudakov'13:  $\ell_0 < 2074$



# Open Problems

- ▶ Conjecture (P.-Vaughan):

$$c_{4,4} = \frac{-11 + 14 \cdot 2^{1/3}}{192}$$



- ▶  $\ell_0 = \max\{\ell : c_{3,\ell} = (\ell - 1)^{-2}\}$ 
  - ▶ Nikiforov'01:  $\ell_0 < \infty$
  - ▶ Das-Huang-Ma-Naves-Sudakov'13:  $\ell_0 < 2074$
  - ▶ P.-Vaughan  $\geq$ '13:  $\ell_0 \geq 7$

Thank you!