

Math 254A Lecture 20 Notes

Daniel Raban

May 12, 2021

1 Deriving van der Waal's Equation (Part 3)

1.1 Bound on α

Last time, we had a quantity α which depended on various factors. We bounded it by a term $\alpha(m, r, \varepsilon)$ which only depended on these 3 quantities. We will compare this to integral of $\int \varphi = \int \varphi^r$.

$$\left| \alpha(m, r, \varepsilon) - \int \varphi^r \right| = \left| (\varepsilon m)^3 \sum_{v \in \varepsilon m \mathbb{Z}^3} \varphi^r(v) - \int \varphi^r \right|$$

Let Q_v be the cube with side length εm and center v .

$$\begin{aligned} &= \left| (\varepsilon m)^3 \sum_{v \in \varepsilon m \mathbb{Z}^3} \left(\varphi^n(v) - \int_{Q_v} \varphi^n \right) \right| \\ &= \left| \sum_v \left(|Q_v| \varphi^r(v) - \int_{Q_v} \varphi^r \right) \right| \\ &\leq \sum_v \int_{Q_v} |\varphi^r(v) - \varphi^r(x)| dx \\ &\leq \sum_{v: Q_v \cap \overline{B_r} \neq \emptyset} (\varepsilon m)^3 \frac{\sqrt{3}}{2} \varepsilon m \cdot \frac{\|\nabla \varphi\|}{r^4} \xrightarrow{O(1)} \\ &= O\left(\frac{r^3}{(\varepsilon m)^3} \cdot \frac{(\varepsilon m)^4}{r^4}\right) \\ &= O\left(\frac{\varepsilon m}{r}\right). \end{aligned}$$

So

$$\alpha(m, r, \varepsilon) = \alpha + o\left(\frac{\varepsilon m}{r}\right)$$

for some constant α .

1.2 Maximizing the entropy term

Now consider maximizing


$$\frac{1}{(n/m^3)} \sum_{k \in \mathcal{C}_n} [H(\rho_k, 1 - \rho_k) + \gamma \rho_k^2],$$

where $\gamma = \beta \alpha(m, \varepsilon, r) / \varepsilon^3 = (\beta / \varepsilon^3)(\alpha + o(\varepsilon m / r))$. Now, we want to try to maximize this over $\rho \in \tilde{\Omega}_n$ with $|\rho| = N_n / m^3$ (recall $N_n / n^3 \rightarrow \varepsilon^3 / v$ as $n \rightarrow \infty$). We can also write this expression as

$$\frac{1}{|\mathcal{C}_n|} \sum_{k \in \mathcal{C}_n} f_\gamma(\rho_k),$$

where $f_\gamma(x) = H(x, 1 - x) + \gamma x^2$ for $0 \leq x \leq 1$.

What does f_γ look like? Here is a picture:



There is a critical value of γ which is the largest γ for which this is still concave. Check using calculus that the critical γ equals 2.


If $\gamma \leq 2$, f_γ is concave, so Jensen's inequality gives

$$\begin{aligned} \frac{1}{|\mathcal{C}_n|} \sum_{k \in \mathcal{C}_n} f_\gamma(\rho_k) &\leq f_\gamma \left(\frac{1}{|\mathcal{C}_n|} \sum_k \rho_k \right) \\ &= f_\gamma \left(\frac{N_n}{m^3} \right) \end{aligned}$$

$$\begin{aligned}
&= f_\gamma \left(\frac{N_n}{m^3} \right) \\
&\xrightarrow{n \rightarrow \infty} f_\gamma \left(\frac{\varepsilon^3}{v} \right).
\end{aligned}$$

We can make this close to tight by taking $\rho_k \approx \varepsilon^3/v$ for all k .

What if $\gamma > 2$? Use the *concave envelope* F_γ of f_γ :



Then

$$\begin{aligned}
\frac{1}{|\mathcal{C}_n|} \sum_{k \in \mathcal{C}_n} f_\gamma(\rho_k) &\leq \frac{1}{|\mathcal{C}_n|} \sum_{k \in \mathcal{C}_n} F_\gamma(\rho_k) \\
&\leq F_\gamma \left(\frac{1}{|\mathcal{C}_n|} \sum_{k \in \mathcal{C}_n} \rho_k \right) \\
&\approx F_\gamma \left(\frac{\varepsilon^3}{v} \right)
\end{aligned}$$

as $n \rightarrow \infty$. This also can be brought as close as we like once $n \rightarrow \infty$ and m is large.

- If $\varepsilon^3/v \notin (a, b)$, then $F_\gamma(\varepsilon^3/v) = f_\gamma(\varepsilon^3/v)$. Then just take $\rho_k \approx \varepsilon^3/v$ for all k .
- If $a < \varepsilon^3/v < b$, then $\rho_k \approx \varepsilon^3/v$ for all k will give you

$$\frac{1}{|\mathcal{C}_n|} \sum_{k \in \mathcal{C}_n} f_\gamma(\rho_k) = f_\gamma\left(\frac{\varepsilon^3}{v}\right) < F_\gamma\left(\frac{\varepsilon^3}{v}\right).$$

Instead, express $\varepsilon^3/v = ta + (1-t)b$. Now choose the values ρ_k so that $\rho_k \approx a$ for $\approx t|\mathcal{C}_n|$ many k s and $\rho_k \approx b$ for $\approx (1-t)|\mathcal{C}_n|$ many k s. Then

$$\frac{1}{|\mathcal{C}_n|} \sum_{k \in \mathcal{C}_n} f_\gamma(\rho_k) \approx tf_\gamma(a) + (1-t)f_\gamma(b) = F_\gamma\left(\frac{\varepsilon^3}{v}\right)$$

The conclusion is that

$$\frac{1}{|\mathcal{C}_n|} \sum_{k \in \mathcal{C}_n} f_\gamma(\rho_k) = F_\gamma\left(\frac{\varepsilon^3}{v}\right) + O\left(\frac{1}{m}\right)$$

as $n \rightarrow \infty$.

1.3 Maximizing the effective partition function

Is the maximization problem for $\log \widehat{Z}_n^r$ close to the same value? Yes!

Go back to

$$\max_{|\rho|=N_n/m^3} \left\{ W(\rho) - \frac{\beta}{n^3} \widetilde{\Phi}_n^r(\rho) \right\}.$$

Can we get this close to the same value? Yes. Since f_γ is strictly convex near a and b , we must have roughly a t fraction of ρ_k s close to a and roughly a $(1-t)$ fraction of ρ_k s close to b


When is the above maximum close to the average of f_γ ? We had the bound via AM-GM:

$$\sum_k H(\rho_k, 1-\rho_k) + \beta m^3 \sum_{k,\ell} \varphi^r(\varepsilon(k-\ell)) \rho_k \rho_\ell \leq \sum_k H(\rho_k, 1-\rho_k) + \beta m^3 \sum_{k,\ell} \varphi^r(\varepsilon(k-\ell)) \frac{\rho_k^2 + \rho_\ell^2}{2}.$$

The difference equals

$$\beta m^3 \sum_{k,\ell} \varphi^r(\varepsilon(k-\ell)) \frac{1}{2} (\rho_k - \rho_\ell)^2.$$

This is small if $\rho_k \approx \rho_\ell$ for most pairs (k, ℓ) where $\varphi^r(\varepsilon(k-\ell))$ is not negligible. What kinds of ρ achieve all these requirements? Choose it according to this picture:



Then we do get

$$W(\rho) - \frac{\beta}{n^3} \tilde{\Phi}_n^r(\rho) \approx \frac{1}{|C_n|} \sum_k f_\gamma(\rho_k) \approx F_\gamma\left(\frac{\varepsilon^3}{v}\right).$$

We will finish off this story next time.