

# Excitation and de-excitation rates from a relativistic reactor antineutrino source

Working derivation based on Appendix B of `main_v4.tex`  
(Dated: July 23, 2026)

## I. SCOPE AND ASSUMPTIONS

This note repeats the excitation and de-excitation calculation of Appendix B for a reactor source. We make the following assumptions.

1. The source contains electron antineutrinos with  $E \sim \text{MeV} \gg m_i$ , so all neutrino masses are neglected in external-state kinematics and spinor traces.
2. The produced antineutrinos have the active, positive-helicity state. Contributions from the wrong helicity are of relative order  $m_i^2/E^2$ .
3. Only neutral-current scattering from a nuclear spin is included. The neutral-current axial coupling  $g_A$  is flavour universal, diagonal in the neutrino mass basis, and the nucleus is treated as pointlike on the scale of the momentum transfer relevant here.
4. The nuclear spin is fixed in the sample. Target recoil and energy transfer to phonons are neglected, as requested. Consequently the nuclear spin can absorb momentum without a recoil-energy term in the energy-conservation delta function.
5. The reactor occupation is extremely dilute, so final-state Pauli blocking is set to one.
6. The incident field is first treated locally as a beam of plane waves. A finite core is included by integrating the result over its angular flux distribution. The additional spatial-coherence assumptions needed to apply a plane-wave  $N^2$  form factor to a macroscopic sample are stated explicitly in Sec. VI.

The external magnetic field defines the spin quantisation axis  $\hat{z}$ . As in the main text, the spin-dependent interaction is

$$H_{\text{int}} = \Sigma_z J_z + \Sigma_- J_- + \Sigma_+ J_+, \quad (1)$$

where

$$\begin{aligned} \Sigma_z &= \sqrt{2} G_F g_A \sum_i \bar{\nu}_i \gamma^3 (1 - \gamma^5) \nu_i, \\ \Sigma_{\pm} &= \frac{G_F g_A}{\sqrt{2}} \sum_i \bar{\nu}_i (\gamma^1 \mp i \gamma^2) (1 - \gamma^5) \nu_i. \end{aligned} \quad (2)$$

The upper sign denotes spin excitation and the lower sign spin de-excitation. The spin splitting is  $\omega_0 > 0$ .

## II. REACTOR FLUX

Let

$$\phi(E, \hat{p}) \equiv \frac{d^3 \Phi_{\bar{\nu}_e}}{dE d\Omega_{\hat{p}}} \quad (3)$$

be the differential antineutrino flux at the sample. It is a physical one-helicity flux; there is no average over two initial helicities. For a pointlike, isotropic  $^{235}\text{U}$  source at distance  $L$ ,

$$\phi(E, \hat{p}) = \frac{\dot{N}_f}{4\pi L^2} S_{235}(E) \delta^{(2)}(\hat{p} - \hat{n}_R), \quad \dot{N}_f = \frac{P_{\text{th}}}{Q_f}, \quad (4)$$

where  $S_{235}(E)$  is the yield per unit energy per fission,  $P_{\text{th}}$  is the thermal power, and  $Q_f$  is the energy released per fission. For an extended core, Eq. (4) is replaced by the sum or volume integral of the contributions from each core element. All subsequent formulae remain valid with the general angular flux Eq. (3).

A produced electron antineutrino is a superposition of mass eigenstates. The neutral-current interaction preserves the mass index. After summing over the unobserved final mass eigenstate, the leading rate is proportional to  $\sum_i |U_{ei}|^2 = 1$ . Reactor oscillations therefore do not alter the neutral-current rate in the limit  $m_i^2/E^2 \rightarrow 0$ . This statement would not apply to flavour-dependent or charged-current couplings.

### III. SINGLE-SPIN RATE

#### A. Kinematics

Write the incoming and outgoing massless four-momenta as

$$p^\mu = E(1, \hat{p}), \quad p'^\mu = E'(1, \hat{p}'). \quad (5)$$

With recoil neglected, energy conservation gives

$$\begin{aligned} E'_+ &= E - \omega_0, & E &\geq \omega_0 & \text{(excitation),} \\ E'_- &= E + \omega_0, & E &\geq 0 & \text{(de-excitation).} \end{aligned} \quad (6)$$

The momentum transferred to the fixed target is  $\vec{q}_\pm = \vec{p} - \vec{p}'_\pm$ .

#### B. Definite-helicity spinor traces

For an incoming active reactor antineutrino, summing over the allowed outgoing antineutrino state gives

$$\begin{aligned} \mathcal{S}_+ &\equiv \sum_c |\bar{v}_+(p)(\gamma^1 - i\gamma^2)(1 - \gamma^5)v_c(p')|^2, \\ \mathcal{S}_- &\equiv \sum_c |\bar{v}_+(p)(\gamma^1 + i\gamma^2)(1 - \gamma^5)v_c(p')|^2. \end{aligned} \quad (7)$$

The subscript on  $v_+$  denotes the positive-helicity antineutrino, not a spin excitation. Using  $\sum_c v_c(p')\bar{v}_c(p') = \not{p}'$  in the massless limit, the two traces evaluate to

$$\begin{aligned} \mathcal{S}_+ &= 16EE'(1 + \hat{p} \cdot \hat{z})(1 - \hat{p}' \cdot \hat{z}), \\ \mathcal{S}_- &= 16EE'(1 - \hat{p} \cdot \hat{z})(1 + \hat{p}' \cdot \hat{z}). \end{aligned} \quad (8)$$

Equation (8) is the main difference from an isotropic, helicity-independent bath. The parity-violating weak interaction retains the direction of the reactor beam relative to the quantisation axis.

#### C. Golden-rule result

Starting with the invariant phase-space measure  $d\tilde{p} = d^3p/[(2\pi)^3 2E]$ , the antineutrino contribution is

$$\gamma_\pm = \pi G_F^2 g_A^2 \int d\tilde{p} d\tilde{p}' (2\pi)^3 \frac{d^3 n_{\bar{\nu}}}{d^3 p} \delta(E'_\pm - E') \mathcal{S}_\pm. \quad (9)$$

For a relativistic beam,  $d^3 n_{\bar{\nu}}/(dE d\Omega) = \phi(E, \hat{p})$ . Substitution of Eq. (8) then yields

$$\begin{aligned} \gamma_+ &= \frac{G_F^2 g_A^2}{2\pi^2} \int_{\omega_0}^{\infty} dE \int d\Omega_{\hat{p}} \phi(E, \hat{p}) (E - \omega_0)^2 \\ &\quad \times \int d\Omega_{\hat{p}'} (1 + \hat{p} \cdot \hat{z})(1 - \hat{p}' \cdot \hat{z}), \\ \gamma_- &= \frac{G_F^2 g_A^2}{2\pi^2} \int_0^{\infty} dE \int d\Omega_{\hat{p}} \phi(E, \hat{p}) (E + \omega_0)^2 \\ &\quad \times \int d\Omega_{\hat{p}'} (1 - \hat{p} \cdot \hat{z})(1 + \hat{p}' \cdot \hat{z}). \end{aligned} \quad (10)$$

The normalisation in Eq. (10) is for the physical active-helicity flux. In particular, it contains no factor 1/2 from averaging over an unoccupied initial helicity.

For a pencil beam along  $\hat{n}_R$ , define

$$c \equiv \cos \alpha = \hat{n}_R \cdot \hat{z}, \quad \phi_R(E) \equiv \int d\Omega_{\hat{p}} \phi(E, \hat{p}). \quad (11)$$

The unrestricted outgoing angular integral gives the compact result

$$\boxed{\begin{aligned} \gamma_+ &= \frac{2G_F^2 g_A^2}{\pi} (1+c) \int_{\omega_0}^{\infty} dE \phi_R(E) (E - \omega_0)^2, \\ \gamma_- &= \frac{2G_F^2 g_A^2}{\pi} (1-c) \int_0^{\infty} dE \phi_R(E) (E + \omega_0)^2. \end{aligned}} \quad (12)$$

Thus  $\gamma_+ = 0$  for a beam antiparallel to  $+\hat{z}$ , while  $\gamma_- = 0$  for a beam parallel to  $+\hat{z}$ , within the massless and recoil-free approximations. Reversing either the beam or magnetic-field direction interchanges the two rates.

#### D. Small- $\omega_0/E$ expansion

For a transverse beam,  $c = 0$ , define the flux moments

$$I_n \equiv \int_0^{\infty} dE E^n \phi_R(E). \quad (13)$$

The negligible interval  $0 < E < \omega_0$  aside, Eq. (12) becomes

$$\begin{aligned} \gamma_+ &= \mathcal{C}(I_2 - 2\omega_0 I_1 + \omega_0^2 I_0), \\ \gamma_- &= \mathcal{C}(I_2 + 2\omega_0 I_1 + \omega_0^2 I_0), \quad \mathcal{C} \equiv \frac{2G_F^2 g_A^2}{\pi}. \end{aligned} \quad (14)$$

If  $\gamma_0 \equiv \mathcal{C}I_2$ , the leading kinematic imbalance is

$$\boxed{\frac{\gamma_- - \gamma_+}{\gamma_0} = 4\omega_0 \frac{I_1}{I_2} + \mathcal{O}\left(\frac{\omega_0^2}{E^2}\right)}. \quad (15)$$

This replaces the nonrelativistic suppression  $\sim 2m_\nu \omega_0 / p_\nu^2$ . For a non-transverse beam, the  $\mathcal{O}(1)$  factors  $1 \pm c$  dominate over the minute kinematic effect in Eq. (15).

#### IV. $^{235}\text{U}$ SPECTRUM BENCHMARK

The file `../235U_nspec.csv` contains a piecewise-constant spectrum  $S_{235}(E) = dN_{\bar{\nu}}/(dE \text{ dfission})$ . Defining

$$M_n = \int dE E^n S_{235}(E), \quad (16)$$

direct integration gives

$$\begin{aligned} M_0 &= 6.077743311 \text{ fission}^{-1}, \\ M_1 &= 8.815892881 \text{ MeV fission}^{-1}, \\ M_2 &= 21.54413023 \text{ MeV}^2 \text{ fission}^{-1}. \end{aligned} \quad (17)$$

The mean and root-mean-square energies are 1.45052 MeV and 1.88275 MeV, respectively, while the energy controlling Eq. (15) is

$$E_{\text{eff}} \equiv \frac{M_2}{M_1} = 2.44378 \text{ MeV}. \quad (18)$$

For  $\omega_0 = 10^{-8}$  eV,

$$4\omega_0 \frac{M_1}{M_2} = 1.6368 \times 10^{-14}. \quad (19)$$

As an illustrative normalisation, take  $g_A = 1/2$ ,  $P_{\text{th}} = 1$  GW,  $L = 10$  m, and  $Q_f = 202.36$  MeV. The point-source flux and the spectrum-averaged transverse single-spin cross section are

$$\begin{aligned} \Phi_R &= 1.49175 \times 10^{13} \text{ cm}^{-2}\text{s}^{-1}, \\ \langle \sigma_{\pm} \rangle &= 2.98854 \times 10^{-44} \text{ cm}^2, \end{aligned} \quad (20)$$

which give

$$\gamma_+ \simeq \gamma_- = 4.45817 \times 10^{-31} \text{ s}^{-1} \quad (21)$$

at the displayed precision. These numbers scale as  $g_A^2 P_{\text{th}}/L^2$ . They are not a claim about a particular reactor; the power, distance, fission energy, coupling, and angle are exposed as inputs in `reactor_rates.py`.

## V. N-SPIN ENSEMBLE

For spins at positions  $\vec{x}_\alpha$ , define

$$F_{\pm}(\vec{q}) = \sum_{\mathbf{f}_{\text{spin}}} \left| \langle \mathbf{f}_{\text{spin}} | \sum_{\alpha} J_{\pm}^{\alpha} e^{-i\vec{q} \cdot \vec{x}_{\alpha}} | \mathbf{i}_{\text{spin}} \rangle \right|^2. \quad (22)$$

Repeating the single-spin calculation without performing the outgoing angular integral gives

$$\begin{aligned} \Gamma_+ &= \frac{G_F^2 g_A^2}{2\pi^2} \int_{\omega_0}^{\infty} dE \int d\Omega_{\hat{p}} \phi(E, \hat{p}) (E - \omega_0)^2 \\ &\quad \times \int d\Omega_{\hat{p}'} (1 + \hat{p} \cdot \hat{z})(1 - \hat{p}' \cdot \hat{z}) F_+(\vec{q}_+), \\ \Gamma_- &= \frac{G_F^2 g_A^2}{2\pi^2} \int_0^{\infty} dE \int d\Omega_{\hat{p}} \phi(E, \hat{p}) (E + \omega_0)^2 \\ &\quad \times \int d\Omega_{\hat{p}'} (1 - \hat{p} \cdot \hat{z})(1 + \hat{p}' \cdot \hat{z}) F_-(\vec{q}_-). \end{aligned} \quad (23)$$

This is the reactor-source replacement for Eq. (B.18) of the main manuscript. Unlike the isotropic expression, the weak angular kernel cannot generally be pulled outside the form-factor integral.

For the initial transverse product state  $|\rightarrow\rangle^{\otimes N}$ , the continuum form factor of a uniform sphere of radius  $R$  is

$$F_{\pm}^P(q) = \frac{N}{2} + \frac{N^2}{4} \left[ \frac{3j_1(qR)}{qR} \right]^2. \quad (24)$$

The  $N/2$  term is the incoherent contribution; the  $N^2$  term contains the collective contribution. For a pencil beam, let  $\mu = \hat{p} \cdot \hat{p}'$ . Averaging over the outgoing azimuth about  $\hat{p}$  gives  $\langle \hat{p}' \cdot \hat{z} \rangle_{\text{az}} = c\mu$ , and hence

$$\Gamma_{\pm} = \frac{G_F^2 g_A^2}{\pi} \int dE \phi_R(E) E_{\pm}'^2 (1 \pm c) \int_{-1}^1 d\mu (1 \mp c\mu) F_{\pm}(q_{\pm}), \quad (25)$$

with the excitation threshold understood and

$$\begin{aligned} q_{\pm}^2 &= E^2 + E_{\pm}'^2 - 2EE'_{\pm}\mu \\ &= \omega_0^2 + 2EE'_{\pm}(1 - \mu). \end{aligned} \quad (26)$$

For  $F_{\pm} = 1$ , Eq. (25) reduces to Eq. (12).

## VI. RELATIVISTIC COHERENCE LIMITS

In the exactly forward direction,

$$q_{\pm, \min} = \omega_0, \quad (27)$$

instead of the nonrelativistic result  $q_{\min} \sim m_\nu \omega_0 / p_\nu$ . At small scattering angle  $\theta$ ,

$$q_{\pm}^2 \simeq \omega_0^2 + EE'_{\pm} \theta^2. \quad (28)$$

The forward collective region therefore requires

$$\omega_0 R \ll 1, \quad \theta \lesssim \frac{1}{ER}. \quad (29)$$

For  $R = 1$  cm,  $\omega_0 = 10^{-8}$  eV gives  $\omega_0 R = 5.07 \times 10^{-4}$ , so longitudinal coherence is possible. Using the root-mean-square energy of the supplied spectrum,

$$ER = 9.54 \times 10^{10}, \quad \theta_{\text{coh}} \simeq 1.05 \times 10^{-11}. \quad (30)$$

Thus the whole solid angle is never coherent for a centimetre-scale reactor target; only a very narrow forward cone contributes to the  $N^2$  term.

For  $ER \gg 1$ ,  $\omega_0 R \ll 1$ , and a spherical sample, the angular integral can be done asymptotically:

$$\begin{aligned} \int d\Omega_{\hat{p}'} \left[ \frac{3j_1(qR)}{qR} \right]^2 &\simeq \frac{18\pi}{E^2 R^2} \int_0^\infty \frac{dx}{x} j_1^2(x) \\ &= \frac{9\pi}{2E^2 R^2}, \end{aligned} \quad (31)$$

where  $\int_0^\infty dx j_1^2(x)/x = 1/4$ . Inside this cone  $\hat{p}' \simeq \hat{p}$ , so both circular-current kernels approach  $1 - c^2$ . The leading collective cross sections are consequently

$$\boxed{\sigma_{\text{coh},+} \simeq \sigma_{\text{coh},-} \simeq \frac{9G_F^2 g_A^2 N^2}{16\pi R^2} (1 - c^2).} \quad (32)$$

Here  $R$  is expressed in natural units. The leading result is independent of  $E$ : the elementary  $E^2$  scaling is cancelled by the  $(ER)^{-2}$  forward-cone fraction.

The incoherent terms for the transverse product state are

$$\sigma_{\text{inc},\pm} \simeq \frac{NG_F^2 g_A^2}{\pi} E^2 (1 \pm c). \quad (33)$$

For  $c = 0$ , their spectrum-integrated ratio is

$$\frac{\Gamma_{\text{coh}}}{\Gamma_{\text{inc}}} \simeq \frac{9N}{16R^2} \frac{M_0}{M_2}. \quad (34)$$

As a plane-wave benchmark, a sphere with  $R = 1$  cm and spin density  $n_s = 3 \times 10^{22} \text{ cm}^{-3}$  contains  $N = 1.25664 \times 10^{23}$  spins. Equations (32) and (34) give

$$\begin{aligned} \sigma_{\text{coh},\pm} &= 1.4580 \times 10^{-20} \text{ cm}^2, \\ \Gamma_{\text{coh},\pm} &= 2.1750 \times 10^{-7} \text{ s}^{-1}, \\ \frac{\Gamma_{\text{coh}}}{\Gamma_{\text{inc}}} &= 7.7646 \end{aligned} \quad (35)$$

for the illustrative 1-GW, 10-m, transverse-beam normalisation of Sec. IV. These are ensemble transition rates, not single-spin rates.

Two qualifications are important.

1. When  $c = \pm 1$ , the leading forward kernel  $1 - c^2$  vanishes. One of the two rates vanishes identically in the massless limit, while the other receives only a subleading contribution from nonzero scattering angles. Equation (32) must not be used to form a 0/0 ratio in this aligned geometry.

2. The  $N^2$  benchmark assumes that the incident neutrino state is sufficiently coherent across the sample for the plane-wave phase in Eq. (22) to apply. A real reactor has a finite core, finite source distance, and neutrino wave packets. If those effects limit the mutual coherence length below  $R$ , Eq. (24) must be replaced by the corresponding two-point source correlation function. The single-spin result and the incoherent  $N/2$  term are unaffected. The numerical  $N^2$  result above should therefore be read as the plane-wave upper benchmark, not yet as a realistic reactor prediction.

## VII. REPRODUCIBILITY

The numerical sources are stored alongside this document:

- `check_spinor_traces.py` constructs explicit Dirac gamma matrices and verifies Eq. (8) for  $10^4$  random direction pairs.
- `reactor_rates.py` integrates `../235U_nspec.csv`, evaluates the flux moments, single-spin rates, coherence angle, and the plane-wave ensemble benchmark. Its power, distance,  $Q_f$ ,  $g_A$ ,  $\omega_0$ , beam angle, radius, and spin density are command-line options.