

submitted to *Geophys. J. Int.*

Magnetic radial inversion for 3-D source geometry estimation

L.B. Vital^{1*}, V.C. Oliveira Jr.¹ and V.C.F. Barbosa¹

¹ *Observatório Nacional, Gal. José Cristino, 77, São Cristóvão, Rio de Janeiro, 20921-400, Brazil*

SUMMARY

We present a method for inverting total-field anomaly data to estimate the geometry of a uniformly magnetized 3-D geological source in subsurface. The method assumes the total-magnetization direction is known. We approximate the source by an ensemble of vertically juxtaposed right prisms, all of them with the same total-magnetization vector and depth extent. The horizontal cross-section of each prism is defined by a polygon having the same number of vertices equally spaced from 0° to 360° . The position of these vertices, the horizontal location of each prism and the depth extent of all prisms are the parameters to be estimated by solving a constrained nonlinear inverse problem for minimizing a goal function. We run successive inversions for a range of tentative total-magnetization intensities and depths to the top of the shallowest prism. The estimated models producing the lowest values of the goal function form the set of candidate solutions. To obtain stable solutions, we impose the zeroth- and first-order Tikhonov regularizations on the shape of the prisms. The method allows estimating the geometry of both vertical and inclined sources by the suitable use of first-order Tikhonov regularization. Tests with synthetic data produced by simple and complex simulated bodies show the efficiency of our method in retrieving the shape of the magnetic source. Results obtained by inverting airborne total-field anomaly data over the Anitápolis alkaline-carbonatic complex, in the Santa Catarina state, Brazil, suggest that the emplacement of the magnetic sources was controlled by NW-SE-trending faults at depth, in accordance with known structural features at the study area.

Key words: Magnetic anomalies: modelling and interpretation; Inverse theory; Numerical solutions.

1 INTRODUCTION

The interpretation of total-field anomalies on the surface of the Earth is an important challenge in exploration geophysics due to the nonuniqueness of 3-D magnetic inversion. It is well-known that several magnetization distribution in subsurface can reproduce the same magnetic data with the same accuracy. To overcome this inherent ambiguity, a priori information need to be introduced for reducing the number of possible solutions coherent with the local geology. There are basically two groups of 3-D magnetic inversion methods. The available a priori information determines the suitable approach for each case.

The first group is formed methods approximating the source by a geometrically simple causative body having its geometry defined by a small number of parameters (e.g., Ballantyne 1980; Bhattacharyya 1980; Silva & Hohmann 1983). These methods estimate both the geometry and the physical property of the source by solving a nonlinear inverse problem. Due to the very restrictive parametrization, such methods usually do not have severe problems with ambiguity.

The second group is formed by the vast majority of methods. They approximate the subsurface by a grid of juxtaposed rectangular prisms having a constant total-magnetization direction. Some methods presume that a purely induced magnetization (e.g., Cribb 1976; Li & Oldenburg 1996; Pilkington 1997) and the isotropic magnetic susceptibility of the prisms is the quantity estimated by solving a linear inverse problem. Some of these methods allow a magnetization direction different from that of the local main field (e.g., Pignatelli et al. 2006). In this case, the parameters to be estimated are the total-magnetization intensities of the prisms. In all these methods, the geometry of the magnetic sources is estimated indirectly, by interpreting the estimated susceptibility or total-magnetization intensity distribution. Theoretically, these methods are capable of recovering the geometry of complex sources. However, they require a lot of a priori information to overcome their nonuniqueness and instability due to the large number of parameters to be estimated by inversion. Additionally, they are characterized by a high computational cost associated with the solution of large linear systems.

The third group of 3-D magnetic inversion methods presume some knowledge about the physical property distribution and estimate the geometry of the sources. They are usually formulated as nonlinear inverse problems. Wang & Hansen (1990) approximate the source by a polyhedron and estimate the position of its vertices in the Fourier domain. Li et al. (2017) have developed a multiple level-set method to estimate geometry of a set of causative bodies with uniform magnetic susceptibility. These methods have a small number of parameters to be estimated by inversion and much less ambiguity problems if compared to those methods the second group.

We present a magnetic inversion method to estimate the geometry of an isolated and uniformly 3-D magnetic source having known total-magnetization direction. Our method is an extension for total-

field anomaly data of those methods presented by Oliveira Jr. et al. (2011) and Oliveira Jr. & Barbosa (2013) for inverting gravity and gravity-gradient data, respectively. We approximate the source by an interpretation model formed vertically juxtaposed right prisms having horizontal cross-sections defined by polygons, all of them with the same number of vertices. For convenience, all prisms have the same depth extension and total-magnetization intensity. Differently from Oliveira Jr. et al. (2011) and Oliveira Jr. & Barbosa (2013), our method estimates not only the position of the vertices defining the horizontal cross-sections and the horizontal position, but also the depth extension of all prisms forming the interpretation model. Additionally, we run the inversion for different values of total-magnetization intensities and depths to the top of the shallowest prism. Among the estimated models, those producing the lowest values of goal function form the set of candidate models. To obtain a stable solutions, use the same set of regularizing functions proposed by Oliveira Jr. et al. (2011) and also propose a new one for constraining the depth extent of the prisms.

A test with synthetic data produced by a simple symmetric model shows the performance of our method in an ideal case. We also applied our method to interpret the synthetic data produced by a complex sources having variable shape and dip along depth. The results show that our method can be a very useful tool for interpreting magnetic data in real situations. Based on these results with synthetic data, we applied our method to interpret airborne data located over the alkaline-carbonatitic complex of Anitápolis, in the Santa Catarina state, Brazil. Our results bring some light on the debate about the structural control of the Anitápolis complex. Our results suggest that the complex follows a NW-SE direction at depth, in agreement with known geological structures not only located at, but also northward and southward the study area. These results attest that our method is a promising tool for interpreting total-field anomaly data.

2 METHODOLOGY

2.1 Forward problem

Let \mathbf{d}^o be the observed data vector, whose i th element d_i^o , $i = 1, \dots, N$, is the total-field anomaly produced by a 3-D source (Fig. 1a) at the point (x_i, y_i, z_i) of a Cartesian coordinate system with x , y and z axes pointing to north, east and down, respectively. We assume that the direction of the total magnetization vector of the source is constant and known. We approximate the volume of the source by a set of L vertically juxtaposed 3-D prisms (Fig. 1b) by following the same approach of Oliveira Jr. et al. (2011) and Oliveira Jr. & Barbosa (2013). The depth to the top of the shallowest prism is defined by z_0 and m_0 is the constant total-magnetization intensity of all prisms. The horizontal cross-section of each prism is described by a polygon with a fixed number V of vertices equally spaced from 0° to 360° , which are described in polar coordinates referred to an internal origin O^k . The radii of the vertices $(r_j^k, j = 1, \dots, V, k = 1, \dots, L)$, the horizontal coordinates $(x_0^k$ and $y_0^k, k = 1, \dots, L)$ of the origins $O^k, k = 1, \dots, L$, and the depth extent dz of the L vertically stacked prisms (Fig. 1b) are arranged in a $M \times 1$ parameter vector \mathbf{p} , $M = L(V + 2) + 1$, given by

$$\mathbf{p} = \begin{bmatrix} \mathbf{r}^{1\top} & x_0^1 & y_0^1 & \dots & \mathbf{r}^{L\top} & x_0^L & y_0^L & dz \end{bmatrix}^\top, \quad (1)$$

where “ \top ” denotes transposition and \mathbf{r}^k is a $V \times 1$ vector containing the radii r_j^k of the k th prism. Let $\mathbf{d}(\mathbf{p})$ be the predicted data vector, whose i th element

$$d_i(\mathbf{p}) \equiv \sum_{k=1}^L f_i^k(\mathbf{r}^k, x_0^k, y_0^k, dz, z_1^k, m_0), \quad i = 1, \dots, N, \quad (2)$$

is the total-field anomaly produced by the ensemble of L prisms at the i th observation point (x_i, y_i, z_i) . In eq. 2, $f_i^k(\mathbf{r}^k, x_0^k, y_0^k, dz, z_1^k, m_0)$ is the total-field anomaly produced, at the observation point (x_i, y_i, z_i) , by the k th prism, with depth to the top $z_1^k = z_0 + (k - 1)dz$. We calculate $d_i(\mathbf{p})$ (eq. 2) by using the Python package Fatiando a Terra (Uieda et al. 2013), which implements the formulas proposed by Plouff (1976).

2.2 Inverse problem formulation

The total-magnetization of the source m_0 and depth to the top of the shallowest prism z_0 are hyperparameters of the inversion, i. e., they are not estimated during the inversion, but their value influences the final solution. Given a set of tentative values for m_0 and z_0 , we solve a constrained non-linear problem to estimate the parameter vector \mathbf{p} (eq. 1) by minimizing the goal function

$$\Gamma(\mathbf{p}) = \phi(\mathbf{p}) + \sum_{\ell=1}^7 \alpha_\ell \varphi_\ell(\mathbf{p}), \quad (3)$$

subject to

$$p_l^{min} < p_l < p_l^{max}, \quad l = 1, \dots, M, \quad (4)$$

where $\varphi(\mathbf{p})$ is the data-misfit function given by

$$\phi(\mathbf{p}) = \frac{1}{N} \|\mathbf{d}^o - \mathbf{d}(\mathbf{p})\|_2^2, \quad (5)$$

which represents the normalized squared Euclidean norm of the difference between the observed data vector \mathbf{d}^o and the predicted data vector $\mathbf{d}(\mathbf{p})$, α_ℓ is a positive number representing the weight of the ℓ th constraint function $\varphi_\ell(\mathbf{p})$ and p_l^{min} and p_l^{max} are, respectively, the lower and upper limits for the l th element p_l of the parameter vector \mathbf{p} (eq. 1). These limits are defined by the interpreter based on both the horizontal extent of the magnetic anomaly and the knowledge about the source.

To solve our nonlinear inverse problem, we use a gradient-based method and, consequently, we need to define the gradient vector $\nabla\Gamma(\mathbf{p})$ and Hessian matrix $\mathbf{H}(\mathbf{p})$ of the goal function $\Gamma(\mathbf{p})$ (eq. 3):

$$\nabla\Gamma(\mathbf{p}) = \nabla\phi(\mathbf{p}) + \sum_{\ell=1}^7 \alpha_\ell \nabla\varphi_\ell(\mathbf{p}) \quad (6)$$

and

$$\mathbf{H}(\mathbf{p}) = \mathbf{H}_\phi(\mathbf{p}) + \sum_{\ell=1}^7 \alpha_\ell \mathbf{H}_\ell, \quad (7)$$

where

$$\nabla\phi(\mathbf{p}) = -\frac{2}{N} \mathbf{G}(\mathbf{p})^\top [\mathbf{d}^o - \mathbf{d}(\mathbf{p})] \quad (8)$$

and

$$\mathbf{H}_\phi(\mathbf{p}) = \frac{2}{N} \mathbf{G}(\mathbf{p})^\top \mathbf{G}(\mathbf{p}) \quad (9)$$

are the gradient vector of the Hessian matrix of the misfit function $\phi(\mathbf{p})$ (eq. 5), respectively, the terms $\nabla\varphi_\ell(\mathbf{p})$ and \mathbf{H}_ℓ , $\ell = 1, \dots, 7$, are the gradient vectors and Hessian matrices of the constraint functions, respectively, and $\mathbf{G}(\mathbf{p})$ is an $N \times M$ matrix whose element ij is the derivative of the predicted data $d_i(\mathbf{p})$ (eq. 2) with respect to the j element p_j of the parameter vector \mathbf{p} (eq. 1). Details about the constraint functions $\varphi_\ell(\mathbf{p})$, $\ell = 1, \dots, 7$, as well as the numerical procedure to solve this nonlinear inverse problem are given in the following sections.

2.3 Constraint functions

We have divided the constraint functions $\varphi_\ell(\mathbf{p})$ (eq. 3), $\ell = 1, \dots, 7$, used here to obtain stable solutions and introduce a priori information about the magnetic source into three groups.

2.3.1 Smoothness constraints

This group is formed by variations of the first-order Tikhonov regularization (Aster et al. 2019, p. 103) and impose smoothness on the radii r_j^k and the Cartesian coordinates x_0^k and y_0^k of the origin O^k , $j = 1, \dots, V$, $k = 1, \dots, L$, defining the horizontal section of each prism (Fig.1b). They were proposed by Oliveira Jr. et al. (2011) and Oliveira Jr. & Barbosa (2013) and play a very role in introducing a prior information about the shape of the source.

The first constraint of this group is the *Smoothness constraint on the adjacent radii defining the horizontal section of each vertical prism*. This constraint imposes that adjacent radii r_j^k and r_{j+1}^k within each prism must be close to each other. It forces the estimated prism to be approximately cylindrical. Mathematically, the constraint is given by

$$\begin{aligned}\varphi_1(\mathbf{p}) &= \sum_{k=1}^L \left[(r_V^k - r_1^k)^2 + \sum_{j=1}^{V-1} (r_j^k - r_{j+1}^k)^2 \right] \\ &= \mathbf{p}^\top \mathbf{R}_1^\top \mathbf{R}_1 \mathbf{p} ,\end{aligned}\quad (10)$$

where

$$\mathbf{R}_1 = \mathbf{I}_L \otimes \begin{bmatrix} (\mathbf{I}_V - \mathbf{D}_V^\top) & \mathbf{0}_{V \times 2} \end{bmatrix} , \quad (11)$$

$\mathbf{0}_{LV \times 1}$ is an $LV \times 1$ vector with null elements, \mathbf{I}_L is the identity matrix of order L , “ \otimes ” denotes the Kronecker product (Horn & Johnson 1991, p. 243), $\mathbf{0}_{V \times 2}$ is a $V \times 2$ matrix with null elements, \mathbf{I}_V is the identity matrix of order V and \mathbf{D}_V^\top is the upshift permutation matrix of order V (Golub & Loan 2013, p. 20). The gradient and Hessian of function $\varphi_1(\mathbf{p})$ (eq. 10) are given by:

$$\nabla \varphi_1(\mathbf{p}) = 2\mathbf{R}_1^\top \mathbf{R}_1 \mathbf{p} , \quad (12)$$

and

$$\mathbf{H}_1(\mathbf{p}) = 2\mathbf{R}_1^\top \mathbf{R}_1 . \quad (13)$$

The second constraint of this group is the *Smoothness constraint on the adjacent radii of the vertically adjacent prisms*, which imposes that adjacent radii r_j^k and r_j^{k+1} within vertically adjacent prisms must be close to each other. This constraint forces the shape of all prisms to be similar to each other and is given by

$$\begin{aligned}\varphi_2(\mathbf{p}) &= \sum_{k=1}^{L-1} \left[\sum_{j=1}^V (r_j^{k+1} - r_j^k)^2 \right] \\ &= \mathbf{p}^\top \mathbf{R}_2^\top \mathbf{R}_2 \mathbf{p}\end{aligned}\quad (14)$$

where

$$\mathbf{R}_2 = \begin{bmatrix} \mathbf{S}_2 & \mathbf{0}_{(L-1)V \times 1} \end{bmatrix}_{(L-1)V \times M}, \quad (15)$$

$$\mathbf{S}_2 = \left(\begin{bmatrix} \mathbf{I}_{L-1} & \mathbf{0}_{(L-1) \times 1} \end{bmatrix} - \begin{bmatrix} \mathbf{0}_{(L-1) \times 1} & \mathbf{I}_{L-1} \end{bmatrix} \right) \otimes \begin{bmatrix} \mathbf{I}_V & \mathbf{0}_{V \times 2} \end{bmatrix}, \quad (16)$$

$\mathbf{0}_{(L-1)V \times 1}$ is an $(L-1)V \times 1$ vector with null elements, $\mathbf{0}_{(L-1) \times 1}$ is an $(L-1) \times 1$ vector with null elements and \mathbf{I}_{L-1} is the identity matrix of order $L-1$. The gradient and Hessian of function $\varphi_2(\mathbf{p})$ (eq. 14) are given by:

$$\nabla \varphi_2(\mathbf{p}) = 2\mathbf{R}_2^\top \mathbf{R}_2 \mathbf{p}, \quad (17)$$

and

$$\mathbf{H}_2(\mathbf{p}) = 2\mathbf{R}_2^\top \mathbf{R}_2. \quad (18)$$

The last constraint of this group is the *Smoothness constraint on the horizontal position of the arbitrary origins of the vertically adjacent prisms*. This constraint imposes that the estimated horizontal Cartesian coordinates (x_0^k, y_0^k) and (x_0^{k+1}, y_0^{k+1}) of the origins O^k and O^{k+1} of adjacent prisms must be close to each other. It forces the prisms to be vertically aligned. This constraint is given by

$$\begin{aligned} \varphi_3(\mathbf{p}) &= \sum_{k=1}^{L-1} \left[(x_0^{k+1} - x_0^k)^2 + (y_0^{k+1} - y_0^k)^2 \right], \\ &= \mathbf{p}^\top \mathbf{R}_3^\top \mathbf{R}_3 \mathbf{p} \end{aligned}, \quad (19)$$

where

$$\mathbf{R}_3 = \begin{bmatrix} \mathbf{S}_3 & \mathbf{0}_{(L-1)2 \times 1} \end{bmatrix}_{(L-1)2 \times M}, \quad (20)$$

$$\mathbf{S}_3 = \left(\begin{bmatrix} \mathbf{I}_{L-1} & \mathbf{0}_{(L-1) \times 1} \end{bmatrix} - \begin{bmatrix} \mathbf{0}_{(L-1) \times 1} & \mathbf{I}_{L-1} \end{bmatrix} \right) \otimes \begin{bmatrix} \mathbf{0}_{2 \times V} & \mathbf{I}_2 \end{bmatrix}, \quad (21)$$

$\mathbf{0}_{(L-1)2 \times 1}$ is an $(L-1)2 \times 1$ vector with null elements, $\mathbf{0}_{2 \times V}$ is a $2 \times V$ matrix with null elements and \mathbf{I}_2 is the identity matrix of order 2. The gradient and Hessian of function $\varphi_3(\mathbf{p})$ (eq. 19) are given by:

$$\nabla \varphi_3(\mathbf{p}) = 2\mathbf{R}_3^\top \mathbf{R}_3 \mathbf{p}, \quad (22)$$

and

$$\mathbf{H}_3(\mathbf{p}) = 2\mathbf{R}_3^\top \mathbf{R}_3. \quad (23)$$

2.3.2 Equality constraints

This group is formed by two constraints that were proposed by Oliveira Jr. et al. (2011) and Oliveira Jr. & Barbosa (2013) by following the same approach proposed Barbosa et al. (1997) and Barbosa

et al. (1999a). They introduce a priori information about the shallowest prism and are suitable for outcropping sources.

The *Source's outcrop constraint* imposes that the horizontal cross-section of the shallowest prism must be close to the intersection of the geologic source with the known outcropping boundary. The matrix form of the this constraint is given by

$$\begin{aligned}\varphi_4(\mathbf{p}) &= \left[(x_0^1 - x_0^0)^2 + (y_0^1 - y_0^0)^2 + \sum_{j=1}^V (r_j^1 - r_j^0)^2 \right] , \\ &= (\mathbf{R}_4 \mathbf{p} - \mathbf{a})^\top (\mathbf{R}_4 \mathbf{p} - \mathbf{a})\end{aligned}\quad (24)$$

where \mathbf{a} is a vector containing the radii and the horizontal Cartesian coordinates of the polygon defining the outcropping boundary

$$\mathbf{a} = \begin{bmatrix} \tilde{r}_1^0 & \dots & \tilde{r}_V^0 & \tilde{x}_0^0 & \tilde{y}_0^0 \end{bmatrix}^\top , \quad (25)$$

and

$$\mathbf{R}_4 = \begin{bmatrix} \mathbf{I}_{V+2} & \mathbf{0}_{(V+2) \times (M-V-2)} \end{bmatrix}_{(V+2) \times M} , \quad (26)$$

where \mathbf{I}_{V+2} is the identity matrix of order $V+2$ and $\mathbf{0}_{(V+2) \times (M-V-2)}$ is a matrix with null elements. The gradient and Hessian of function $\varphi_4(\mathbf{p})$ (eq. 24) are given by:

$$\nabla \varphi_4(\mathbf{p}) = 2\mathbf{R}_4^\top (\mathbf{R}_4 \mathbf{p} - \mathbf{a}) , \quad (27)$$

and

$$\mathbf{H}_4(\mathbf{p}) = 2\mathbf{R}_4^\top \mathbf{R}_4 . \quad (28)$$

The *Source's horizontal location constraint* imposes that the horizontal Cartesian coordinates of the origin within the shallowest prism must be as close as possible to a known outcropping point. The matrix form of the this constraint is given by

$$\begin{aligned}\varphi_5(\mathbf{p}) &= \left[(x_0^1 - x_0^0)^2 + (y_0^1 - y_0^0)^2 \right] , \\ &= (\mathbf{R}_5 \mathbf{p} - \mathbf{b})^\top (\mathbf{R}_5 \mathbf{p} - \mathbf{b})\end{aligned}\quad (29)$$

where \mathbf{b} is a vector containing the horizontal Cartesian coordinates of the outcropping point

$$\mathbf{b} = \begin{bmatrix} \tilde{x}_0^0 & \tilde{y}_0^0 \end{bmatrix}^\top , \quad (30)$$

and

$$\mathbf{R}_5 = \begin{bmatrix} \mathbf{0}_{2 \times V} & \mathbf{I}_2 & \mathbf{0}_{2 \times (M-V-2)} \end{bmatrix}_{2 \times M} , \quad (31)$$

where \mathbf{I}_2 is the identity matrix of order 2 and $\mathbf{0}_{2 \times (M-V-2)}$ and $\mathbf{0}_{2 \times V}$ are matrices with null elements.

The gradient and Hessian of function $\varphi_5(\mathbf{p})$ (eq. 29) are given by:

$$\nabla \varphi_5(\mathbf{p}) = 2\mathbf{R}_5^T (\mathbf{R}_5 \mathbf{p} - \mathbf{b}) , \quad (32)$$

and

$$\mathbf{H}_5(\mathbf{p}) = 2\mathbf{R}_5^T \mathbf{R}_5 . \quad (33)$$

2.3.3 Minimum Euclidean norm constraints

Two constraints use the zeroth-order Tikhonov regularization with the purpose of obtaining stable solutions without necessarily introducing significant a priori information about the source.

The *Minimum Euclidean norm of the radii* imposes that all estimated radii within each prism must be close to null values. This constraint was proposed by Oliveira Jr. et al. (2011) and Oliveira Jr. & Barbosa (2013) and can be rewritten in matrix form as follows

$$\begin{aligned} \varphi_6(\mathbf{p}) &= \sum_{k=1}^L \sum_{j=1}^V \left(r_j^k \right)^2 , \\ &= \mathbf{p}^T \mathbf{R}_6^T \mathbf{R}_6 \mathbf{p} \end{aligned} \quad (34)$$

where

$$\mathbf{R}_6 = \begin{bmatrix} \mathbf{S}_6 & \mathbf{0}_{(M-1) \times 1} \\ \mathbf{0}_{1 \times (M-1)} & 0 \end{bmatrix}_{M \times M} , \quad (35)$$

and

$$\mathbf{S}_6 = \begin{bmatrix} \mathbf{I}_V & \mathbf{0}_{V \times 2} \\ \mathbf{0}_{2 \times V} & \mathbf{I}_2 \end{bmatrix}_{(V+2) \times (V+2)} . \quad (36)$$

The gradient and Hessian of function $\varphi_6(\mathbf{p})$ (eq. 34) are given by:

$$\nabla \varphi_6(\mathbf{p}) = 2\mathbf{R}_6^T \mathbf{R}_6 \mathbf{p} , \quad (37)$$

and

$$\mathbf{H}_6(\mathbf{p}) = 2\mathbf{R}_6^T \mathbf{R}_6 . \quad (38)$$

The other constraint, the *Minimum Euclidean norm of the depth extent*, imposes that the depth extent of all prisms must be close to zero. We present this constraint to introduce a priori information about the maximum depth of the source. It is given by

$$\begin{aligned} \varphi_7(\mathbf{p}) &= dz^2 , \\ &= \mathbf{p}^T \mathbf{R}_7^T \mathbf{R}_7 \mathbf{p} \end{aligned} \quad (39)$$

where

$$\mathbf{R}_7 = \begin{bmatrix} \mathbf{0}_{(M-1) \times (M-1)} & \mathbf{0}_{(M-1) \times 1} \\ \mathbf{0}_{1 \times (M-1)} & 1 \end{bmatrix}_{M \times M}. \quad (40)$$

The gradient and Hessian of function $\varphi_7(\mathbf{p})$ (eq. 39) are given by:

$$\nabla \varphi_7(\mathbf{p}) = 2\mathbf{R}_7^T \mathbf{R}_7 \mathbf{p}, \quad (41)$$

and

$$\mathbf{H}_7(\mathbf{p}) = 2\mathbf{R}_7^T \mathbf{R}_7. \quad (42)$$

2.4 Computational procedures

To estimate the parameter vector \mathbf{p} (eq. 1) that minimizes the goal function $\Gamma(\mathbf{p})$ (eq. 3), subjected to the inequality constraint (eq. 4), we use the Levenberg-Marquardt method (e.g., Aster et al. 2019, p. 240). This is an iterative gradient-based method that, at each iteration k , updates the estimated parameter vector $\hat{\mathbf{p}}_{(k)}$ (where the superscript hat “ $\hat{\cdot}$ ” denotes estimated) to obtain new a estimated parameter vector $\hat{\mathbf{p}}_{(k+1)}$. We compute this update by following the same strategy of Barbosa et al. (1999b), Oliveira Jr. et al. (2011) and Oliveira Jr. & Barbosa (2013) to incorporate the inequality constraint (eq. 4). This strategy consists in transforming each element p_l of the estimated parameter vector $\hat{\mathbf{p}}_{(k)}$ into the element p_l^\dagger of a new vector $\hat{\mathbf{p}}_{(k)}^\dagger$ as follows:

$$p_l^\dagger = -\ln \left(\frac{p_l^{max} - p_l}{p_l - p_l^{min}} \right), \quad (43)$$

where p_l^{min} and p_l^{max} are defined in the inequality constraint (eq. 4). Then, we compute a correction $\Delta \hat{\mathbf{p}}_{(k)}^\dagger$ and a new vector $\hat{\mathbf{p}}_{(k+1)}^\dagger = \hat{\mathbf{p}}_{(k)}^\dagger + \Delta \hat{\mathbf{p}}_{(k)}^\dagger$. Finally, we transform each element p_l^\dagger of $\hat{\mathbf{p}}_{(k+1)}^\dagger$ into the element p_l of the new estimated parameter vector $\hat{\mathbf{p}}_{(k+1)}$ as follows:

$$p_l = p_l^{min} + \left(\frac{p_l^{max} - p_l^{min}}{1 + e^{-p_l^\dagger}} \right). \quad (44)$$

2.4.1 Considerations about the weights $\alpha_1 - \alpha_7$

Attributing values to the weights α_ℓ (eq. 3) is an important feature of our method. However, there is no analytical rule to define them and their values can be dependent on the particular characteristics of the interpretation model. To overcome this problem, we normalize the α_ℓ values as follows:

$$\alpha_\ell = \tilde{\alpha}_\ell \frac{E_\phi}{E_\ell}, \quad \ell = 1, \dots, 7, \quad (45)$$

where $\tilde{\alpha}_\ell$ is a positive scalar and E_ϕ/E_ℓ is a normalizing factor. In this equation, E_ℓ represents the trace of the Hessian matrix \mathbf{H}_ℓ (eqs 13, 18, 23, 28, 33, 38, and 42) of the ℓ th constraining function $\varphi_\ell(\mathbf{p})$

(eqs 10, 14, 19, 24, 29, 34, and 39). The constant E_ϕ is the trace of the Hessian matrix $\mathbf{H}_\phi(\mathbf{p}_0)$ (eq. 9) of the misfit function $\phi(\mathbf{p})$ (eq. 5) computed with the initial approximation $\hat{\mathbf{p}}_{(0)}$ for the parameter vector \mathbf{p} (eq. 1) at the beginning of the inversion algorithm. According to this empirical strategy, the weights α_ℓ are defined using the positive scalars $\tilde{\alpha}_\ell$ (eq. 45), which are less dependent on the particular characteristics of the interpretation model.

2.4.2 Inversion algorithm

At each iteration k of our algorithm, the correction $\Delta\hat{\mathbf{p}}_{(k)}^\dagger$ is computed by solving the following linear system:

$$\mathbf{D}_{(k)} \left[\mathbf{D}_{(k)} \mathbf{H}^\dagger(\hat{\mathbf{p}}_{(k)}) \mathbf{D}_{(k)} + \lambda_{(k)} \mathbf{I} \right] \mathbf{D}_{(k)} \Delta\hat{\mathbf{p}}_{(k)}^\dagger = -\nabla\Gamma(\hat{\mathbf{p}}_{(k)}), \quad (46)$$

where $\lambda_{(k)}$ is a positive scalar which is adjusted at each iteration and is associated with the Levenberg-Marquardt method (e.g., Aster et al. 2019, p. 240), \mathbf{I} is the identity matrix with order M , $\nabla\Gamma(\hat{\mathbf{p}})$ is the gradient of the goal function (eq. 6) and $\mathbf{H}^\dagger(\hat{\mathbf{p}}_{(k)})$ is a matrix given by

$$\mathbf{H}^\dagger(\hat{\mathbf{p}}_{(k)}) = \mathbf{H}(\hat{\mathbf{p}}_{(k)}) \mathbf{T}(\hat{\mathbf{p}}_{(k)}), \quad (47)$$

where $\mathbf{H}(\hat{\mathbf{p}}_{(k)})$ is the Hessian matrix of the goal function (eq. 7) and $\mathbf{T}(\hat{\mathbf{p}}_{(k)})$ is a diagonal matrix whose element ll is given by

$$t(p_l) = \frac{(p_l^{max} - p_l)(p_l - p_l^{min})}{p_l^{max} - p_l^{min}}, \quad l = 1, \dots, M, \quad (48)$$

with p_l being the l th element of the estimated parameter vector $\hat{\mathbf{p}}_{(k)}$. In eq. 46, $\mathbf{D}_{(k)}$ is a diagonal matrix proposed by Marquardt (1963) for scaling the parameter $\lambda_{(k)}$ at each iteration and improving the convergence of the algorithm. The element ll of this diagonal matrix is given by

$$d_{ll} = \frac{1}{\sqrt{h_{ll}^\dagger}}, \quad (49)$$

where h_{ll}^\dagger is the element ll of the matrix $\mathbf{H}^\dagger(\hat{\mathbf{p}}_{(k)})$ (eq. 47).

2.4.3 Practical considerations

Our algorithm depends on several parameters that significantly impact the estimated models and cannot be automatically set without the interpreter's judgment. They are the parameters $\tilde{\alpha}_1 - \tilde{\alpha}_7$ (eq. 45). Based on our practical experience, we suggest some empirical procedures for setting these parameters.

The parameters $\tilde{\alpha}_1$ and $\tilde{\alpha}_2$ impose priori information on the shape of the horizontal cross-section of the prisms. The first one forces all prisms to have a circular horizontal cross-section, while the second forces all prisms to have a similar horizontal cross-section. Generally, their values vary from

10^{-5} to 10^{-3} and differs from each other by one order of magnitude, at most. The parameter $\tilde{\alpha}_3$ also varies from 10^{-5} to 10^{-3} and controls the relative position of adjacent prisms forming the model. A high value privileges a vertical estimated body, whereas a small value tends to generate an inclined estimated body.

In comparison to $\tilde{\alpha}_1$, $\tilde{\alpha}_2$ and $\tilde{\alpha}_3$, the other parameters usually have smaller values varying from 10^{-7} to 10^{-5} . The parameters $\tilde{\alpha}_4$ and $\tilde{\alpha}_5$ are used when a priori information about the source is available at the study area. The parameter $\tilde{\alpha}_6$ has a purely mathematical meaning and it is used only to obtain stable solutions for the inverse problem. Its value is set to be as small as possible. The parameter $\tilde{\alpha}_7$ controls the total-vertical extension of the the estimated body. The greater its value, the smaller the estimated total-vertical extension and vice versa. A general rule is starting with values $\tilde{\alpha}_1 = 10^{-4}$, $\tilde{\alpha}_2 = 10^{-4}$, $\tilde{\alpha}_3 = 10^{-4}$, $\tilde{\alpha}_4 = 0$, $\tilde{\alpha}_5 = 0$, $\tilde{\alpha}_6 = 10^{-7}$, $\tilde{\alpha}_7 = 10^{-5}$ and change them to refine the results.

Another important aspect of our method is the initial approximation $\hat{p}_{(0)}$. We set the same constant radii for all prisms, so that they approximate a cylinder and enclose both positive and negative parts of the observed anomaly. Also, we set the same constant depth extent dz for all prisms with the purpose of generating a bottom depth greater than that we expect for the true source. Finally, we adjust the parameters of the initial approximation $\hat{p}_{(0)}$ in order to obtain a preliminary fit of the observed data.

3 APPLICATION TO SYNTHETIC DATA

3.1 Simple model test

We have simulated a funnel-shaped source with simple geometry (blue prisms in Figs 3b and 5), which extends from $z_0 = 0$ m to 1600 m along depth and satisfies most of the constraints described in subsection 2.3. It is formed by $L = 8$ prisms, all of them with the same number of vertices $V = 20$, depth extent $dz = 200$ m and horizontal coordinates $(x_0^k, y_0^k) = (0, 0)$ m of the origins O^k , $k = 1, \dots, L$. The radii of all vertices are equal to each other within the same prism and decrease linearly with depth, varying from $r_j^0 = 1920$ m, at the shallowest prism, $r_j^L = 800$ m, at the deepest prism, $j = 1, \dots, V$. All prisms have the same total-magnetization direction with inclination -21.5° , declination -18.7° and intensity $m_0 = 9$ A/m. We calculated the total-field anomaly produced by this simple model on an 100 km^2 area, simulating an airborne survey composed of 21 flight lines that are equally spaced 500 m apart along the y axis, at a constant vertical coordinate $z = -150$ m. At each line, there are 100 observation points spaced 101 m apart along x axis. The total-field anomaly is corrupted with a pseudorandom Gaussian noise having mean $\mu_0 = 0$ nT and standard deviation $\sigma_0 = 5$ nT (Fig. 3a).

We have inverted the synthetic total-field anomaly (Fig. 3a) produced by the simple model and obtained 36 different models. Each model was obtained by using a different pair of depth to the top z_0 and total-magnetization intensity m_0 (Fig. 4). All models were generated by using the true values of total-magnetization inclination and declination, the same interpretation model formed by $L = 5$ prisms, each one with $V = 20$ vertices, and the same weights for the constraining functions: $\tilde{\alpha}_1 = 10^{-5}$, $\tilde{\alpha}_2 = 10^{-4}$, $\tilde{\alpha}_3 = 10^{-4}$, $\tilde{\alpha}_4 = 0$, $\tilde{\alpha}_5 = 0$, $\tilde{\alpha}_6 = 10^{-7}$, and $\tilde{\alpha}_7 = 10^{-6}$. The initial approximation for all models have the same constant radii $r_j^k = 2000$ m, $k = 1, \dots, L$, $j = 1, \dots, V$, the same depth extent $dz = 350$ m and the same origin $(x_0^k, y_0^k) = (0, 0)$ m for all prisms.

Fig. 4 shows that the estimated model obtained by using the true values for depth to the top z_0 and total-magnetization intensity m_0 (represented by the red triangle in Fig. 4) produces the lowest value of goal function $\Gamma(p)$ (eq. 3). Fig. 5a shows that this estimated model (red prisms in Figs 5c and d) not only fits the noise-corrupted data, but also retrieves the geometry of the true model (blue prisms). The inset in Fig. 5a shows that the residuals follow a normal distribution with mean μ and standard deviation σ compatible with those values used to generate the synthetic noise. The estimated depth extent of each prism is $dz = 297.65$ m and results in a total depth extent (1485 m) very close to the true one (1600 m). These results illustrate the good performance of our method in an ideal case.

3.2 Complex model test

We have simulated a complex inclined body (blue prisms in Figs 6 and 8), which extends from $z_0 = -300$ m to 5700 m along depth and violates most of the constraints described in subsection 2.3. It is formed by $L = 10$ prisms, all of them with the same number of vertices $V = 30$ and depth extent $dz = 600$ m. The horizontal coordinates of the origins O^k vary linearly from $(x_0^0, y_0^0) = (-250, 750)$ m, at the shallowest prism, to $(x_L^0, y_L^0) = (250, -750)$ m resulting a dip in the direction NW-SE, at the deepest prism. The radii $r_j^k, k = 1, \dots, L, j = 1, \dots, V$, defining the vertices vary from 240 m to 1540 m and also differ from each other within the same prism. All prisms have a constant total magnetization with inclination -50° , declination 9° and intensity $m_0 = 12$ A/m. This total magnetization is based on Zhang et al. (2018) which have estimated the total-magnetization vector for magnetic sources on the Goiás alkaline province (GAP). So, we are simulating an alkaline vertical dipping intrusion. We have calculated the total-field anomaly produced by this complex model on an 100 km^2 area, simulating an airborne survey composed of 18 north-south flight lines distributed from -5000 m to 5000 m along the y axis and a east-west tie line approximately located at $x = 0$ m. The data points are located on the undulated surface shown in Fig. 6a. Notice that both flight and tie lines are not perfectly straight. We added a pseudorandom Gaussian noise having mean μ_0 nT and standard deviation σ_0 nT to the produced total-field anomaly (Fig. 3a).

Actually, they simulate the real survey presented in the following section. To compute the synthetic total-field anomaly, we consider a constant main field with inclination -21.5° and declination -18.7° , which is significantly different from the total-magnetization direction of the complex model. Finally, we have contaminated the synthetic total-field anomaly with a pseudorandom Gaussian noise having mean and standard deviation equal to 0 nT and 5 nT, respectively (Fig. 6a). We have inverted the synthetic total-field anomaly (Fig. 6a) produced by the complex model and to obtain 36 different models. Each model was obtained by using a specific pair of depth to the top z_0 and total-magnetization intensity m_0 (Fig. 7). Differently from the previous simulation with a simple model, the present grid of m_0 and z_0 does not contain the true values (represented by the red triangle in Fig. 7). All models were generated by using the true values of total-magnetization inclination and declination, the same interpretation model formed by $L = 8$ prisms, each one with $V = 15$ vertices, and the same weights for the constraining functions: $\tilde{\alpha}_1 = 10^{-5}$, $\tilde{\alpha}_2 = 10^{-4}$, $\tilde{\alpha}_3 = 10^{-4}$, $\tilde{\alpha}_4 = 0$, $\tilde{\alpha}_5 = 0$, $\tilde{\alpha}_6 = 10^{-7}$, and $\tilde{\alpha}_7 = 10^{-6}$. The initial approximation for all models have the same constant radii $r_j^k = 800$ m, $k = 1, \dots, L, j = 1, \dots, V$, the same depth extent $dz = 650$ m and the same origin $(x_0^k, y_0^k) = (-300, 300)$ m for all prisms.

Fig. 7 shows that, compared to that shown in Fig. 4, the color map obtained for the complex model presents an elongated region containing candidate solutions, all of them producing small values for

the goal function $\Gamma(\mathbf{p})$ (eq. 3), with different total-magnetization intensity m_0 and depth-to-the-top z_0 . This result illustrates the inherent ambiguity of potential-field methods in retrieving both the physical-property distribution and volume of the sources. In this case, some a priori information must be used to constraint the range of reliable solutions.

The estimated model obtained by using the total-magnetization intensity $m_0 = 11.4 \text{ A/m}$ and depth-to-the-top $z_0 = -320 \text{ m}$ (represented by the cyan diamond in Fig. 7), close to the true values represented by the red triangle in Fig. 7, produces the lowest value for the goal function $\Gamma(\mathbf{p})$ (eq. 3). Fig. 8 shows that this estimated model (red prisms in Figs 8c and d) fits the noise-corrupted data and also retrieves the geometry of the true source (blue prisms). Note that the red prisms edges accurately matches the blue prisms ones. The inset in Fig. 5a shows that the residuals follow a normal distribution with mean μ and standard deviation σ compatible with those values used to generate the noise-corrupted data. The estimated total depth extent (5597.7 m) and volume (11.0 km³) are underestimated, but still close to the true values (6000 m and 12.60 km³). These results show that our method can also be very useful to interpret complex sources, even if they do not perfectly satisfy the constraints imposed to solve the nonlinear inverse problem.

4 APPLICATION TO FIELD DATA

We have applied our method to interpret airborne magnetic data provided by Geological Survey of Brazil (CPRM) over the Anitápolis complex, in the Santa Catarina state, Brazil. The survey was at an elevation of 100 m above the terrain, with N-S and E-W lines spaced by 500 m and 10,000 m from each other, respectively. The total-field anomaly data were corrected from daytime variation and subtracted from the main magnetic field using the IGRF. The inclination, declination and intensity of the main field at the study area, for the period of the survey, are -37.05° , -18.17° and $\approx 22\,768$ nT, respectively. To isolate the target total-field anomaly, we have applied a regional separation using a second-order polynomial fit. We have also continued the anomaly upward to a constant height $z = -2000$ m (Fig. 9a) by using the equivalent-layer technique (Dampney 1969; Emilia 1973; Oliveira Jr. & Barbosa 2013). The upward-continued data were calculated on a regular grid of 50×50 points equally spaced from 6916 to 6926 km along the x axis and from 683 to 693 km along the y axis. Fig. 9b shows the geometric height (referred to the WGS84 ellipsoid) and UTM horizontal coordinates of the data.

The Anitápolis alkaline-carbonatitic complex forms a circular concentric body (≈ 6 km 2 in area) containing magnetite as part of its mineralogical composition. It intruded the Late Proterozoic leucogranites of the Dom Feliciano mobile belt in the Early Cretaceous (132 Ma), apparently concomitant with the voluminous flood tholeiitic basalts of the Serra Geral Formation (133-130 Ma) at the southern side of the Paraná Basin (Gibson et al. 1999; Scheibe et al. 2005). As pointed out by Gomes et al. (2018), there is still some debate about the emplacement of the Anitápolis alkaline-carbonatitic complex. Melcher & Coutinho (1966) pointed out the influence of N-S-trending faults. Scheibe et al. (2005) considered that it is roughly emplaced along the E-W Rio Uruguay Lineament. According to Riccomini et al. (2005), the Anitápolis complex does not show a clear structural control.

To define the total-magnetization direction of the interpretation model, we applied the method proposed by Oliveira Jr. et al. (2015) and verified the result using the reduction to the pole technique (RTP). The results (not shown) indicate that the total magnetization of the source has the same direction of the main field in the study area, suggesting a purely induced magnetization. Measurements of the magnetic susceptibility χ made at the Jacupiranga complex, another alkaline complex located northward of the study area, with the same age as the Anitápolis complex, show values varying from $\chi = 1.06 \times 10^{-3}$ SI to $\chi = 161.05 \times 10^{-3}$ SI (Alva-Valdivia et al. 2009, tb. 1). Hence, we used these values and the main magnetic field intensity in the study area ($\approx 22\,768$ nT) as a priori information to constraint the total-magnetization intensity m_0 values used to interpret the magnetic data at the Anitápolis complex. The magnetic susceptibility (χ) and the corresponding total-magnetization intensity m_0 ranges used in our application are shown in Fig. 10.

We used an interpretation model formed by $L = 8$ prisms, each one with $V = 30$ ($k = 1, \dots, 10$) vertices defining their horizontal cross-sections. We inverted the observed total-field anomaly (Fig. 9a) for each pair of m_0 and z_0 shown in Fig. 10, resulting in 100 estimated models. For all models, we set an initial approximation $\hat{\mathbf{p}}_{(0)}$ with origin at $(x_0^k, y_0^k) = (6921, 688)$ km, constant radii $r_j^k = 1500$ m for all vertices forming all prisms and the same constant depth extent $dz = 700$ m. We also used the same weights $\tilde{\alpha}_1 = 10^{-3}$, $\tilde{\alpha}_2 = 10^{-4}$, $\tilde{\alpha}_3 = 10^{-3}$, $\tilde{\alpha}_4 = 0$, $\tilde{\alpha}_5 = 0$, $\tilde{\alpha}_6 = 10^{-6}$, and $\tilde{\alpha}_7 = 10^{-5}$ (eq. 45).

Fig. 10 shows that, similarly to the results obtained with the complex model (Fig. 7), there is an elongated region containing candidate solutions producing small values for the goal function $\Gamma(\mathbf{p})$ (eq. 3), with different values of m_0 and z_0 . The magenta diamond in Fig. 10 represents the estimated model shown in Fig. 11. This model produces the lowest value for $\Gamma(\mathbf{p})$ (3), it has a volume 48.37 km^3 , total depth extent of 5709.20 m ($dz = 713.65$), depth-to-the-top $z_0 = -1150$ m and total-magnetization intensity $m_0 = 3.0 \text{ A/m}$. This value corresponds to a magnetic susceptibility $\chi = 0.16 \text{ SI}$. Fig. 10a shows that this model produces a very good data fit. This estimated depth-to-the-top z_0 indicates a non-outcropping source, which is compatible with a priori information about the study area. We do not have evidences of an outcropping source for this anomaly, although there are outcropping intrusions in the area of the Anitápolis complex (Gibson et al. 1999). The estimated magnetic susceptibility χ is also compatible with the available a priori information. It is close to the upper limit found by Alva-Valdivia et al. (2009) in the Jacupiranga complex.

The cyan diamond in Fig. 10 represents the alternative model shown in Fig. 12. This model is similar to that shown in Fig. 11. It has a volume 52.89 km^3 , total depth extent 5434.35 m ($dz = 679.29$), depth-to-the-top $z_0 = -1200$ m and total-magnetization intensity $m_0 = 2.5 \text{ A/m}$, which corresponds to a magnetic susceptibility $\chi = 0.13 \text{ SI}$. In comparison to the model shown in Fig. 11, the alternative model shown in Fig. 12 has a shallower (but still non-outcropping) top and a smaller estimated magnetic susceptibility. As a consequence, the alternative model has a greater volume in order to produce an equally good data fit (Fig. 12a).

Both estimated models show an NW-SE elongated body with variable dip along depth (Figs 11 and 12). This is the same direction associated with the Serra Geral Lineament, crossing the study area, the Ponta Grossa Arch and the Torres syncline, which are prominent structural features located, respectively, northward and southward of the study area (e.g., Scheibe et al. 2005, p. 535). The estimated magnetic susceptibilities and depths-to-the-top are compatible with the available a priori information and produce very good data fits. Hence, both models represent the possible geometry of the Anitápolis complex.

5 CONCLUSIONS

We have developed a total-field anomaly nonlinear inversion to estimate the shape of an isolated 3-D geological body assuming the knowledge about its total-magnetization direction. We approximate the body by a set of vertically stacked right prisms. The horizontal cross-section of each prism is a polygon defined by a given number of equally spaced vertices from 0° to 360° . We run our inversion for a set of given depths to the top and total-magnetization intensities. For each depth to the top and total-magnetization intensity, our method estimates the geometry of the cross-sections, the depth extent and the horizontal position of the prisms approximating the 3-D geological body by solving a constrained nonlinear inversion. The estimated bodies producing the lowest values of the goal function form the set of candidate solutions. To stabilize the inversions, we introduce a set of seven constraints on the source shape. Our method is an extension of previous works developed for retrieving the geometry of 3-D bodies by inverting gravity and gravity-gradient data. We not only adapted the previous methods for interpreting total-field anomaly data, but also generalize them to include the depth to the top and depth extension of the prisms among the estimated parameters.

Applications to synthetic data produced by a simple symmetric source illustrate the efficiency of our method in an ideal case. Moreover, the results obtained with synthetic data produced by a complex source, with variable dip and shape along depth, show that our method can also be used to interpret magnetic data produced by a realistic geological source. Both tests with synthetic data show that our method is able to retrieve the source's shape and fit the data.

We applied our method to interpret a total-field anomaly data over the alkaline-carbonatitic complex of Anitápolis, Santa Catarina state, Brazil. We obtained two candidate models having similar shapes, depths to the top and total-magnetic intensities, all of them in close agreement with the available a priori information. The histogram of the residuals produced by both models show that they equally fit the observed total-field anomaly. Both models suggest that the emplacement of the Anitápolis complex seems to be controlled by NW-SE-trending faults at depth. This is the same direction associated with the Serra Geral Lineament, crossing the study area, and the prominent structural features Ponta Grossa Arch and Torres syncline, which are located northward and southward of the study area, respectively. Possible extensions of this work is the inversion of elongated and/or multiple sources. In addition, an the combination of gradient-based and heuristic optimization methods could be applied to estimate optimal regularization weights, overcoming problems with local minima.

ACKNOWLEDGMENTS

REFERENCES

- Alva-Valdivia, L. M., Perrin, M., Rivas-Sánchez, M. L., Goguitchaichvili, A., López-Loera, H., Lopes, O. F., & Bonás, T. B., 2009. Rock magnetism and microscopy of the jacupiranga alkaline-carbonatitic complex, southern brazil, *Earth, Planets and Space*, **61**(1), 161–171.
- Aster, R. C., Borchers, B., & Thurber, C. H., 2019. *Parameter Estimation and Inverse Problems*, Elsevier, 3rd edn.
- Ballantyne, E. J., 1980. Magnetic curve fit for a thin dike; calculator program (TI 59), *Geophysics*, **45**(3), 447–455.
- Barbosa, V. C. F., Silva, J. B. C., & Medeiros, W. E., 1997. Gravity inversion of basement relief using approximate equality constraints on depths, *Geophysics*, **62**, 1745–1757.
- Barbosa, V. C. F., Silva, J. B. C., & Medeiros, W. E., 1999a. Gravity inversion of a discontinuous relief stabilized by weighted smoothness constraints on depth, *Geophysics*, **64**(5), 1429–1437.
- Barbosa, V. C. F., Silva, J. B. C., & Medeiros, W. E., 1999b. Stable inversion of gravity anomalies of sedimentary basins with nonsmooth basement reliefs and arbitrary density contrast variations, *Geophysics*, **64**(3), 754–764.
- Bhattacharyya, B. K., 1980. A generalized multibody model for inversion of magnetic anomalies, *Geophysics*, **45**(2), 255–270.
- Cribb, J., 1976. Application Of The Generalized Linear Inverse To The Inversion Of Static Potential Data, *Geophysics*, **41**(6), 1365–1369.
- Dampney, C. N. G., 1969. The equivalent source technique, *Geophysics*, **34**(1), 39–53.
- Emilia, D. A., 1973. Equivalent sources used as an analytic base for processing total magnetic field profiles, *Geophysics*, **38**(2), 339–348.
- Gibson, S. A., Thompson, R. N., Leonardos, O. H., Dickin, A. P., & Mitchell, J. G., 1999. The limited extent of plume-lithosphere interactions during continental flood-basalt genesis: geochemical evidence from cretaceous magmatism in southern Brazil, *Contributions to Mineralogy and Petrology*, **137**(1), 147–169.
- Golub, G. H. & Loan, C. F. V., 2013. *Matrix Computations (Johns Hopkins Studies in the Mathematical Sciences)*, Johns Hopkins University Press, 4th edn.
- Gomes, C. d. B., Comin-Chiaromonti, P., Azzone, R. G., Ruberti, E., & Rojas, G. E. E., 2018. Cretaceous carbonatites of the southeastern Brazilian Platform: a review, *Brazilian Journal of Geology*, **48**, 317 – 345.
- Horn, R. A. & Johnson, C. R., 1991. *Topics in Matrix Analysis*, Cambridge University Press, 1st edn.
- Li, W., Lu, W., Qian, J., & Li, Y., 2017. A multiple level-set method for 3d inversion of magnetic data, *GEOPHYSICS*, **82**(5), J61–J81.
- Li, Y. & Oldenburg, D. W., 1996. 3-D inversion of magnetic data, *Geophysics*, **61**(2), 394–408.
- Marquardt, D. W., 1963. An algorithm for least-squares estimation of nonlinear parameters, *Journal of the society for Industrial and Applied Mathematics*, **11**(2), 431–441.
- Melcher, G. C. & Coutinho, J. M. V., 1966. Rochas alcalinas e carbonatito de Anitápolis, estado de Santa

- Catarina, *Boletim da Sociedade Brasileira de Geologia*, **15**(1), 59–93.
- Oliveira Jr., V. C. & Barbosa, V. C. F., 2013. 3-D radial gravity gradient inversion, *Geophysical Journal International*, **195**(2), 883–902.
- Oliveira Jr., V. C., Barbosa, V. C. F., & Silva, J. B. C., 2011. Source geometry estimation using the mass excess criterion to constrain 3-D radial inversion of gravity data, *Geophysical Journal International*, **187**(2), 754–772.
- Oliveira Jr., V. C., Sales, D., Barbosa, V. C. F., & Uieda, L., 2015. Estimation of the total magnetization direction of approximately spherical bodies, *Nonlinear Processes in Geophysics*, **22**(2).
- Pignatelli, A., Nicolosi, I., & Chiappini, M., 2006. An alternative 3d inversion method for magnetic anomalies with depth resolution, *Annals of geophysics = Annali di geofisica*.
- Pilkington, M., 1997. 3-D magnetic imaging using conjugate gradients, *Geophysics*, **62**(4), 1132–1142.
- Plouff, D., 1976. Gravity and magnetic fields of polygonal prisms and application to magnetic terrain corrections, *Geophysics*, **41**(4), 727–741.
- Riccomini, C., Velázquez, V., & Gomes, C., 2005. Tectonic controls of the Mesozoic and Cenozoic alkaline magmatism in the central-southeastern brazilian platform, in *Mesozoic to Cenozoic Alkaline Magmatism in the Brazilian Platform*, chap. 2, pp. 31–55, eds Comin-Chiaromonti, P. & Gomes, C., Edusp/Fapesp, São Paulo.
- Scheibe, L., Furtado, S., Comin-Chiaromonti, P., & Gomes, C., 2005. Cretaceous alkaline magmatism from Santa Catarina state, southern brazil, in *Mesozoic to Cenozoic Alkaline Magmatism in the Brazilian Platform*, chap. 16, pp. 523–571, eds Comin-Chiaromonti, P. & Gomes, C., Edusp/Fapesp, São Paulo.
- Silva, J. B. C. & Hohmann, G. W., 1983. Nonlinear magnetic inversion using a random search method, *Geophysics*, **48**(12), 1645–1658.
- Uieda, L., Oliveira Jr., V. C., & Barbosa, V. C. F., 2013. Modeling the earth with fatiando a terra, in *Proceedings of the 12th Python in Science Conference*, pp. 96 – 103.
- Wang, X. & Hansen, R. O., 1990. Inversion for magnetic anomalies of arbitrary three-dimensional bodies, *Geophysics*, **55**(10), 1321–1326.
- Zhang, H., Ravat, D., Marangoni, Y. R., Chen, G., & Hu, X., 2018. Improved total magnetization direction determination by correlation of the normalized source strength derivative and the reduced-to-pole fields, *Geophysics*, **83**(6), J75–J85.




Figure 1. Schematic representation of (a) total-field anomaly (gray surface) produced by (b) a 3-D anomalous source (dark gray volume). The interpretation model in (b) consists of a set of L vertical, juxtaposed 3-D prisms P^k , $k = 1, \dots, L$, (light gray prisms) in the vertical direction of a right-handed coordinate system.




Figure 2. Polygonal cross-section of the k th vertical prism P^k described by V vertices (white dots) with polar coordinates (r_j^k, θ_j^k) , $j = 1, \dots, V$, $k = 1, \dots, L$, referred to an arbitrary origin O^k (grey dot) with horizontal Cartesian coordinates (x_0^k, y_0^k) , $k = 1, \dots, L$.




Figure 3. Simple model simulation. (a) Noise-corrupted total-field anomaly produced by the simple model (blue prisms) in (b) with a pseudorandom Gaussian distribution having mean $\mu_0 = 0$ nT and standard deviation $\sigma_0 = 5$ nT, the black dots represent the observation points. The connected red dots are the vertices of the initial approximate horizontally projected at the data map. (b) Perspective view of the simple model represented by the blue prisms.




Figure 4. Application to the simple model data. Goal function $\Gamma(\mathbf{p})$ (eq. 3), in nT, produced by estimated models with different depths-to-the-top (z_0) and total-magnetization intensities (m_0). The red triangle represents the m_0 and z_0 of the true source.




Figure 5. Application to the simple model data. (a) Residuals between the noise-corrupted data (Fig. 3a) and the predicted data (not shown) produced by the estimated model. The inset in (a) shows the histogram of the residuals and the Gaussian curve (dashed line) whose mean and standard deviation are, respectively, $\mu = 0.04$ nT and $\sigma = 7.20$ nT. (b) Perspective view of the initial approximate (red prisms) and the true model (blue prisms). (c) and (d) Comparison between the estimated source (red prisms) and the true model (blue prisms) in perspective views.




Figure 6. Complex model simulation. (a) Noise-corrupted total-field anomaly with a pseudorandom Gaussian distribution having mean $\mu_0 = 0$ nT and standard deviation $\sigma_0 = 5$ nT produced by the complex model, the black dots represent the observation points. The connected red dots are the vertices of the initial approximate horizontally projected at the data map. (b) Vertical coordinates of the observations simulating an airborne survey. (c) and (d) Perspective views of the complex model represented by the blue prisms.




Figure 7. Application to the complex model data. Goal function $\Gamma(\mathbf{p})$ (eq. 3), in nT, produced by estimated models with different depths-to-the-top (z_0) and total-magnetization intensities (m_0). The red triangle represents the m_0 and z_0 of the true source. The cyan diamond represents the estimated model that produces the lowest value for $\Gamma(\mathbf{p})$.




Figure 8. Application to complex model data. (a) Residuals between the noise-corrupted data (Fig. 6a) and the predicted data (not shown) produced by the estimated model. The inset in (a) shows the histogram of the residuals and the Gaussian curve (dashed line) whose mean and standard deviation are, respectively, $\mu = 0.09$ nT and $\sigma = 6.66$ nT. (b) Perspective view of the initial approximate (red prisms) and the true model (blue prisms). (c) and (d) Comparison between the estimated source (red prisms) and the true model (blue prisms) in perspective views.




Figure 9. (a) Residual total-field anomaly (in nT) over the Anitápolis complex, upward-continued to $z = -2000$ m. The horizontal UTM coordinates are referred to the central meridian 51° W. (b) Geometric height (referred to the WGS84 ellipsoid) of the observation points in m. The black dots are the observation points. The connected red dots in both maps are the projected vertices of the initial approximation $\hat{\mathbf{p}}_{(0)}$ used in the inversion.




Figure 10. Application to the field data over the Anitápolis complex, Brazil. Goal function $\Gamma(\mathbf{p})$ (eq. 3), in nT, produced by estimated models with different depths-to-the-top (z_0) and total-magnetization intensities (m_0). The corresponding range of magnetic susceptibility χ is represented in the upper axis by considering a purely induced magnetization with inducing field $\approx 22\,768$ nT. The magenta diamond represents the estimated model that produces the lowest value of $\Gamma(\mathbf{p})$. The cyan diamond represents an alternative model whose magnetic susceptibility is also compatible with values found at the Jacupiranga complex in a previous work (Alva-Valdivia et al. 2009, tb. 1).




Figure 11. Application to the field data over the Anitápolis complex, Brazil. Estimated model producing the lowest goal function value, represented by the magenta diamond in Fig. 10. (a) Residuals between the observed data (Fig. 9a) and the predicted data (not shown) produced by the estimated model. The inset shows the histogram of the residuals and a normal Gaussian curve (dashed line) with mean and standard deviation $\mu = -0.59$ nT and $\sigma = 5.80$ nT, respectively. The light-gray polygons represent the projection of the estimated model on the horizontal plane. (b) Perspective view of the initial approximate (red prisms). (c) and (d) Perspective views of the estimated model (red prisms).




Figure 12. Application to the field data over the Anitápolis complex, Brazil. Estimated model with magnetic susceptibility value compatible with a priori information close to the study area, represented by the cyan diamond in Fig. 10. (a) Residuals between the observed data (Fig. 9a) and the predicted data (not shown) produced by the estimated model. The inset shows the histogram of the residuals and a normal Gaussian curve (dashed line) with mean and standard deviation $\mu = 0.91$ nT and $\sigma = 6.95$ nT, respectively. The light-gray polygons represent the projection of the estimated model on the horizontal plane. (b) Perspective view of the initial approximate (red prisms). (c) and (d) Perspective views of the estimated model (red prisms).