

# 2D gravity inversion with isostatic constraint applied to passive rifted margins

B. Marcela S. Bastos\* and Vanderlei C. Oliveira Jr\*

*\*Observatório Nacional,*

*Department of Geophysics,*

*Rio de Janeiro, Brazil*

(August 6, 2018)

**GEO-XXXX**

Running head: **2D gravity inversion for passive rifted margins**

## ABSTRACT

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

## INTRODUCTION

Several methods have been proposed for using gravity data to estimate the relief of an interface separating two layers.

These methods usually presume that the two layers have constant and known density values. Additionally, they also presume that the interface oscillates around a known reference depth.

For different combinations of density values and reference depths, it is possible to find different interfaces producing the same gravity data.

To deal with this inherent ambiguity (Roy, 1962; Skeels, 1947) and obtain meaningful solutions, the interpreter must use a priori information obtained from seismic data and/or boreholes, for example, in order to constrain the range of possible models.

One of the first automated methods for estimating the interface separating two homogeneous layers was presented by Bott (1960), in space domain.

Since then, several other approaches have been developed in space domain (Tanner, 1967; Dyrelius and Vogel, 1972; Pedersen, 1977; Richardson and MacInnes, 1989; Barbosa et al., 1997, 1999b,a; Chakravarthi and Sundararajan, 2007; Silva et al., 2010; Camacho et al., 2011; Lima et al., 2011; Martins et al., 2011; Barnes and Barraud, 2012; Silva et al., 2014; Santos et al., 2015; Silva and Santos, 2017) and also in the Fourier domain (Oldenburg, 1974; Granser, 1987; Reamer and Ferguson, 1989; Guspí, 1993; Braitenberg et al., 1997; Braitenberg and Zadro, 1999).

Methods in Fourier domain use Parker's formula (Parker, 1973).

Faltam os papers do pilkington, o paper do salem e os papers de satelite regional (Leo

e Sampietro) e os globais (pegar no trabalho do sampietro e do reguzzoni)

Dentre estes trabalhos, tem que dividir quais impem equilibrio isosttico e como eles fazem isso.

## METHODOLOGY

### Forward problem

Let  $\mathbf{d}^o$  be the observed data vector, whose  $i$ -th element  $d_i^o$ ,  $i = 1, \dots, N$ , represent the observed gravity disturbance at the point  $(x_i, y_i, z_i)$ , on a profile located over a rifted passive margin. The coordinates are referred to a topocentric Cartesian system, with  $z$  axis pointing down,  $y$ -axis along the profile and  $x$ -axis perpendicular to the profile. We assume that the observed gravity disturbance is produced by an anomalous mass distribution defined as the difference between the actual mass distribution in the subsurface, which is schematically represented in Figure ??, and a reference mass distribution (Figure ??). In doing it, we implicitly assume that Figure ?? represents the outer layers of a global mass distribution producing the normal gravity field.

The anomalous mass distribution producing the observed data is approximated by an interpretation model (Figure ??) formed by  $N$  adjacent columns. For convenience, we presume that the observed data are regularly spaced, so that there is one observation at the centre of the top of each column forming the interpretation model. The  $i$ -th column is formed by four vertically adjacent layers, , which in turn are composed of vertically adjacent prisms having infinite length along the  $x$ -axis. The first and shallowest layer represents the water layer, is formed by a single prism, has thickness  $t_i^w$  and a constant density contrast  $\Delta\rho^w = \rho^w - \rho^r$ , where  $\rho^w$  and  $\rho^r$  represents, respectively, the densities of water and the

reference mass distribution (Figure ??) at the same point. The third layer represents the crust, it is also formed by a single prism, has thickness  $t_i^c$  and density contrast  $\Delta\rho_i^c = \rho^c - \rho^r$ , with  $\rho^c$  being the crust density. For simplicity, we presume that the crust density  $\rho_i^c$  may be equal to  $\rho^{cc}$ , for  $y_i \leq y_{COT}$ , which represents continental crust, or equal to  $\rho^{oc}$ , for  $y_i > y_{COT}$ , which represents oceanic crust. The crust density depends on the position of the  $i$ -th column with respect to  $y_{COT}$ , which defines an abrupt Crust-Ocean Transition (COT). Consequently, the crust may have two possible density contrasts:  $\Delta\rho_i^c = \rho^{cc} - \rho^r$  or  $\Delta\rho_i^c = \rho^{oc} - \rho^r$ . The top of this layer defines the basement relief and its bottom the relief of the Moho. The fourth layer represents the mantle, it is divided into two parts, each one formed by a single prism having the same density  $\rho^m$  and, consequently, the same density contrast  $\Delta\rho^m = \rho^m - \rho^r$ . The shallowest portion of this layer has thickness  $t_i^m$ . Its top and bottom define, respectively, the depths of Moho and the planar isostatic compensation layer  $S_0$ . The deepest portion of the fourth layer has thickness  $\Delta S_0$ , top at the surface  $S_0$  and bottom at the planar surface  $S_0 + \Delta S_0$ , which defines the Moho in the reference mass distribution model (Figure ??). Finally, the second layer forming the  $t$ -th column of the interpretation model is defined by the interpreter, according to the geological environment to be studied and the a priori information availability. As a general rule, this layer can be defined by a set of  $Q$  vertically adjacent prisms, each one with thickness  $t_i^q$ , density  $\rho^q$  and density contrast  $\Delta\rho^q = \rho^q - \rho^r$ ,  $q = 1, \dots, Q$ .

Given the density contrasts, the COT position  $y_{COT}$ , the isostatic compensation surface  $S_0$ , the thickness of the water layer and of the  $Q - 1$  prisms forming the shallowest portion of the second layer, it is possible to describe the interpretation model in terms of an  $M \times 1$

parameter vector  $\mathbf{p}$ ,  $M = 2N + 1$ , defined as follows:

$$\mathbf{p} = \begin{bmatrix} \mathbf{t}^Q \\ \mathbf{t}^m \\ \Delta S_0 \end{bmatrix}, \quad (1)$$

where  $\mathbf{t}^Q$  and  $\mathbf{t}^m$  are  $N \times 1$  vectors whose  $i$ -th elements  $t_i^Q$  and  $t_i^m$  represent, respectively, the thickness of the prism forming the deepest portion of the second layer and the thickness of the prism forming the shallowest portion of the fourth layer of the interpretation model. In this case, the gravity disturbance produced by the interpretation model (the predicted gravity disturbance) at the position  $(x_i, y_i, z_i)$  can be written as the sum of the vertical component of the gravitational attraction exerted by the  $L$  prisms forming the interpretation model as follows:

$$d_i(\mathbf{p}) = k_g G \sum_{j=1}^L f_{ij}(\mathbf{p}), \quad (2)$$

where  $f_{ij}(\mathbf{p})$  represents an integral over the volume of the  $j$ -th prism. Here, these volume integrals are computed with the expressions proposed by Nagy et al. (2000), by using the open-source Python package *Fatiando a Terra* (Uieda et al., 2013).

## Inverse problem

Let  $\mathbf{d}(\mathbf{p})$  be the predicted data vector, whose  $i$ -th element  $d_i(\mathbf{p})$  is defined by Equation 2. Estimating the particular parameter vector  $\mathbf{p} = \hat{\mathbf{p}}$  producing a predicted data vector  $\mathbf{d}(\mathbf{p})$  as close as possible to the observed data vector  $\mathbf{d}^o$  can be formulated as the problem of minimizing the goal function

$$\Gamma(\mathbf{p}) = \Phi(\mathbf{p}) + \mu \sum_{k=0}^3 \alpha_k \Psi_k(\mathbf{p}), \quad (3)$$

subject to all elements of  $\hat{\mathbf{p}}$  be positive. In Equation 3,  $\mu$  represents the regularizing parameter,  $\Phi(\mathbf{p})$  represents the misfit function given by

$$\Phi(\mathbf{p}) = \frac{1}{N} \|\mathbf{d}^o - \mathbf{d}(\mathbf{p})\|_2^2, \quad (4)$$

where  $\|\cdot\|_2^2$  represents the squared Euclidean norm,  $\alpha_k$  represent the weights assigned to the regularizing functions  $\Psi_k(\mathbf{p})$ , with define the constraints on the parameters to be estimated,  $k = 0, 1, 2, 3$ .

### Airy constraint

Consider that the interpretation model is in isostatic equilibrium according to the Airy model (Turcotte and Schubert, 2002; Hofmann-Wellenhof and Moritz, 2005; Lowrie, 2007). In this case, the pressure (or lithostatic stress) exerted by the model is constant on the isostatic compensation surface  $S_0$ . The pressure per unit area exerted by the  $i$ -th column of the model on  $S_0$ , divided by gravity, is given by:

$$t_i^w \rho^w + t_i^1 \rho_i^1 + \cdots + t_i^Q \rho_i^Q + t_i^c \rho_i^c + t_i^m \rho^m = \sigma_0, \quad (5)$$

where  $\sigma_0$  is an arbitrary positive constant. Rearranging terms in Equation 5 and using the relation

$$S_0 = t_i^w + t_i^1 + \cdots + t_i^Q + t_i^c + t_i^m, \quad (6)$$

it is possible to show that:

$$(\rho_i^Q - \rho_i^c) t_i^Q + (\rho^m - \rho_i^c) t_i^m + (\rho^w - \rho_i^c) t_i^w + (\rho_i^1 - \rho_i^c) t_i^1 + \cdots + (\rho_i^{Q-1} - \rho_i^c) t_i^{Q-1} + \rho_i^c S_0 = \sigma_0. \quad (7)$$

In order to describe the pressure exerted by all columns forming the interpretation model on the surface  $S_0$ , Equation 7 can be written, in matrix notation, as follows:

$$\mathbf{M}^Q \mathbf{t}^Q + \mathbf{M}^m \mathbf{t}^m + \mathbf{M}^w \mathbf{t}^w + \mathbf{M}^1 \mathbf{t}^1 + \cdots + \mathbf{M}^{Q-1} \mathbf{t}^{Q-1} + \boldsymbol{\rho}^c S_0 = \sigma_0 \mathbf{1}, \quad (8)$$

where  $\mathbf{1}$  is an  $N \times 1$  vector with all elements equal to one,  $\mathbf{t}^\alpha$  are  $N \times 1$  vectors with  $i$ -th element defined by the thickness  $t_i^\alpha$  of a prism forming the  $i$ -th column,  $\alpha = w, 1, \dots, Q - 1, Q, m$ , and  $\mathbf{M}^Q, \mathbf{M}^m, \mathbf{M}^w, \mathbf{M}^1, \dots, \mathbf{M}^{Q-1}$  are  $N \times N$  diagonal matrices with elements  $ii$  of main diagonal are given by density contrasts  $(\rho_i^Q - \rho_i^c), (\rho^m - \rho_i^c), (\rho^w - \rho_i^c), (\rho_i^1 - \rho_i^c)$  and  $\dots, (\rho_i^{Q-1} - \rho_i^c)$ , respectively, and  $\boldsymbol{\rho}^c$  is an  $N \times 1$  vector containing the densities of the prisms representing the crust. By applying the first-order Tikhonov regularization (Aster et al., 2005) to the constant vector  $\sigma_0 \mathbf{1}$ , we obtain the following expression:

$$\mathbf{R}(\mathbf{C}\mathbf{p} + \mathbf{D}\mathbf{t}) = \mathbf{0}, \quad (9)$$

where  $\mathbf{0}$  is a vector with null elements and the remaining terms are given by:

$$\mathbf{C} = \begin{bmatrix} \mathbf{M}^Q & \mathbf{M}^m & \mathbf{0} \end{bmatrix}_{N \times M}, \quad (10)$$

$$\mathbf{D} = \begin{bmatrix} \mathbf{M}^w & \mathbf{M}^1 & \dots & \mathbf{M}^{Q-1} & \boldsymbol{\rho}^c \end{bmatrix}_{N \times (QN+1)}, \quad (11)$$

$$\mathbf{t} = \begin{bmatrix} \mathbf{t}^w \\ \mathbf{t}^1 \\ \vdots \\ \mathbf{t}^{Q-1} \\ S_0 \end{bmatrix}_{(QN+1) \times 1}, \quad (12)$$

$\mathbf{p}$  is the parameter vector (Equation 1) and  $\mathbf{R}$  is an  $(N - 1) \times N$  matrix, whose element  $ij$

is defined as follows:

$$[\mathbf{R}]_{ij} = \begin{cases} 1 & , \quad j = i \\ -1 & , \quad j = i + 1 \\ 0 & , \quad \text{otherwise} \end{cases} \quad (13)$$

Finally, from Equation 9, it is possible to define the regularizing function  $\Psi_0(\mathbf{p})$  (Equation 3):

$$\Psi_0(\mathbf{p}) = \|\mathbf{R}(\mathbf{C}\mathbf{p} + \mathbf{D}\mathbf{t})\|_2^2. \quad (14)$$

We call this function as *Airy constraint*. Notice that minimizing this function imposes smoothness on the pressure exerted by the interpretation model on the isostatic compensation surface  $S_0$ .

### Smoothness constraint

This constraint imposes smoothness on the adjacent thickness of the prisms forming the deepest portion of the second layer and the shallowest part of the fourth layer of the interpretation model by applying the first-order Tikhonov regularization (Aster et al., 2005) to the vectors  $\mathbf{t}^Q$  and  $\mathbf{t}^m$  (Equation 1). Mathematically, this constraint is represented by the regularizing function  $\Psi_1(\mathbf{p})$  (Equation 3):

$$\Psi_1(\mathbf{p}) = \|\mathbf{S}\mathbf{p}\|_2^2, \quad (15)$$

where  $\mathbf{S}$  is an  $(N - 1) \times M$  matrix given by:

$$\mathbf{S} = \begin{bmatrix} \mathbf{R} & \mathbf{R} & \mathbf{0} \end{bmatrix}, \quad (16)$$

where  $\mathbf{R}$  is defined by Equation 13 and  $\mathbf{0}$  is a vector with all elements equal to zero.

### Equality constraint

#### *Equality constraint on basement depths*

Let  $\mathbf{a}$  be a vector whose  $k$ -th element  $a_k$ ,  $k = 1, \dots, A$ , is the known basement depth at the horizontal coordinate  $y_k^A$  of the profile. These known basement depth values are used to



define the regularizing function  $\Psi_2(\mathbf{p})$  (Equation 3):

$$\Psi_2(\mathbf{p}) = \|\mathbf{A}\mathbf{p} - \mathbf{a}\|_2^2, \quad (17)$$

where  $\mathbf{A}$  is an  $A \times M$  matrix whose  $k$ -th line has one element equal to one and all the remaining elements equal to zero. The location of the single non-null element in the  $k$ -th line of  $\mathbf{A}$  depends on the coordinate  $y_k^A$  of the known basement depth  $a_k$ . Let us consider, for example, an interpretation model formed by  $N = 10$  columns. Consider also that the basement depth at the coordinates  $y_1^A = y_4$  and  $y_2^A = y_9$  of the profile are equal to 25 and 35.7 km, respectively. In this case,  $A = 2$ ,  $\mathbf{a}$  is a  $2 \times 1$  vector with elements  $a_1 = 25$  and  $a_2 = 35.7$  and  $\mathbf{A}$  is a  $2 \times M$  matrix ( $M = 2N + 1 = 21$ ). The element 4 of the first line and the element 9 of the second line of  $\mathbf{A}$  are equal to 1 and all its remaining elements are equal to zero.

#### *Equality constraint on Moho depths*

Let  $\mathbf{b}$  be a vector whose  $k$ -th element  $b_k$ ,  $k = 1, \dots, B$ , is the difference between the isostatic compensation depth  $S_0$  and the known Moho depth at the horizontal coordinate  $y_k^B$  of the profile. These differences, which must be positive, are used to define the regularizing function  $\Psi_3(\mathbf{p})$  (Equation 3):

$$\Psi_3(\mathbf{p}) = \|\mathbf{B}\mathbf{p} - \mathbf{b}\|_2^2, \quad (18)$$

where  $\mathbf{B}$  is a  $B \times M$  matrix whose  $k$ -th line has one element equal to one and all the remaining elements equal to zero. This matrix is defined in the same way as matrix  $\mathbf{A}$  (Equation 17).

## CONCLUSIONS

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend conse-

quat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

## ACKNOWLEDGMENTS

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

## REFERENCES

- Aster, R. C., B. Borchers, and C. H. Thurber, 2005, Parameter estimation and inverse problems (international geophysics): Academic Press.
- Barbosa, V. C. F., J. ao B. C. Silva, and W. E. Medeiros, 1997, Gravity inversion of basement relief using approximate equality constraints on depths: *Geophysics*, **62**, 1745–1757.
- , 1999a, Gravity inversion of a discontinuous relief stabilized by weighted smoothness constraints on depth: *GEOPHYSICS*, **64**, 1429–1437.
- , 1999b, Stable inversion of gravity anomalies of sedimentary basins with nonsmooth basement reliefs and arbitrary density contrast variations: *GEOPHYSICS*, **64**, 754–764.
- Barnes, G., and J. Barraud, 2012, Imaging geologic surfaces by inverting gravity gradient data with depth horizons: *Geophysics*, **77**, G1–G11.
- Bott, M. H. P., 1960, The use of rapid digital computing methods for direct gravity interpretation of sedimentary basins: *Geophysical Journal International*, **3**, 63–67.
- Braitenberg, C., F. Pettenati, and M. Zadro, 1997, Spectral and classical methods in the evaluation of moho undulations from gravity data: The ne italian alps and isostasy: *Journal of Geodynamics*, **23**, 5 – 22.
- Braitenberg, C., and M. Zadro, 1999, Iterative 3d gravity inversion with integration of seismologic data: *Bollettino di Geofisica Teorica ed Applicata*, **40**, 469–475.
- Camacho, A. G., J. Fernndez, and J. Gottsmann, 2011, A new gravity inversion method for multiple subhorizontal discontinuity interfaces and shallow basins: *Journal of Geophysical Research: Solid Earth*, **116**.
- Chakravarthi, V., and N. Sundararajan, 2007, 3d gravity inversion of basement relief a depth-dependent density approach: *GEOPHYSICS*, **72**, I23–I32.
- Dyrelius, D., and A. Vogel, 1972, Improvement of convergency in iterative gravity interpre-

- tation: *Geophysical Journal of the Royal Astronomical Society*, **27**, 195–205.
- Granser, H., 1987, Three-dimensional interpretation of gravity data from sedimentary basins using an exponential density-depth function: *Geophysical Prospecting*, **35**, 1030–1041.
- Guspi, F., 1993, Noniterative nonlinear gravity inversion: *Geophysics*, **58**, 935–940.
- Hofmann-Wellenhof, B., and H. Moritz, 2005, *Physical geodesy*: Springer.
- Lima, W. A., C. M. Martins, J. B. Silva, and V. C. Barbosa, 2011, Total variation regularization for depth-to-basement estimate: Part 2 physicogeologic meaning and comparisons with previous inversion methods: *Geophysics*, **76**, I13–I20.
- Lowrie, W., 2007, *Fundamentals of geophysics*: Cambridge University Press. (A second edition of this classic textbook on fundamental principles of geophysics for geoscience undergraduates.).
- Martins, C. M., W. A. Lima, V. C. Barbosa, and J. B. Silva, 2011, Total variation regularization for depth-to-basement estimate: Part 1 - mathematical details and applications: *Geophysics*, **76**, I1–I12.
- Nagy, D., G. Papp, and J. Benedek, 2000, The gravitational potential and its derivatives for the prism: *Journal of Geodesy*, **74**, 311–326.
- Oldenburg, D. W., 1974, The inversion and interpretation of gravity anomalies: *Geophysics*, **39**, 526–536.
- Parker, R. L., 1973, The rapid calculation of potential anomalies: *Geophysical Journal of the Royal Astronomical Society*, **31**, 447–455.
- Pedersen, L. B., 1977, Interpretation of potential field data a generalized inverse approach: *Geophysical Prospecting*, **25**, 199–230.
- Reamer, S. K., and J. F. Ferguson, 1989, Regularized twodimensional fourier gravity inversion method with application to the silent canyon caldera, nevada: *Geophysics*, **54**,

486–496.

Richardson, R. M., and S. C. MacInnes, 1989, The inversion of gravity data into three-dimensional polyhedral models: *Journal of Geophysical Research: Solid Earth*, **94**, 7555–7562.

Roy, A., 1962, Ambiguity in geophysical interpretation: *Geophysics*, **27**, 90–99.

Santos, D. F., J. B. C. Silva, C. M. Martins, R. D. C. S. dos Santos, L. C. Ramos, and A. C. M. de Araújo, 2015, Efficient gravity inversion of discontinuous basement relief: *GEOPHYSICS*, **80**, G95–G106.

Silva, J. B., A. S. Oliveira, and V. C. Barbosa, 2010, Gravity inversion of 2d basement relief using entropic regularization: *Geophysics*, **75**, I29–I35.

Silva, J. B. C., and D. F. Santos, 2017, Efficient gravity inversion of basement relief using a versatile modeling algorithm: *GEOPHYSICS*, **82**, G23–G34.

Silva, J. B. C., D. F. Santos, and K. P. Gomes, 2014, Fast gravity inversion of basement relief: *Geophysics*, **79**, G79–G91.

Skeels, D. C., 1947, Ambiguity in gravity interpretation: *Geophysics*, **12**, 43–56.

Tanner, J. G., 1967, An automated method of gravity interpretation: *Geophysical Journal of the Royal Astronomical Society*, **13**, 339–347.

Turcotte, D. L., and G. Schubert, 2002, *Geodynamics*, 2. ed. ed.: Cambridge Univ. Press.

Uieda, L., V. C. Oliveira Jr., and V. C. F. Barbosa, 2013, Modeling the earth with fatiando a terra: *Proceedings of the 12th Python in Science Conference*, 96 – 103.