

Introduction to Machine Learning

Introduction to Probabilistic Methods

Varun Chandola

Computer Science & Engineering
State University of New York at Buffalo
Buffalo, NY, USA
chandola@buffalo.edu



University at Buffalo
Department of Computer Science
and Engineering
School of Engineering and Applied Sciences



Introduction to Probability

Random Variables

Bayes Rule

More About Conditional Independence

Continuous Random Variables

Different Types of Distributions

Handling Multivariate Distributions

Transformations of Random Variables

Information Theory - Introduction

What is Probability? [3, 1]

- ▶ Probability that a coin will land heads is 50%¹
- ▶ **What does this mean?**

¹Dr. Persi Diaconis showed that a coin is 51% likely to land facing the same way up as it is started.

FREQUENTISTS



BAYESIANS

Frequentist Interpretation

- ▶ Number of times an event will be observed in n trials

Frequentist Interpretation

- ▶ Number of times an event will be observed in n trials
- ▶ What if the event can only occur once?
 - ▶ *My winning the next month's powerball.*
 - ▶ *Polar ice caps melting by year 2020.*



Bayesian Interpretation

- ▶ **Uncertainty** of the event
- ▶ Use for making decisions
 - ▶ Should I put in an offer for a sports car?

What is a Random Variable (X)?

- ▶ Can take any value from \mathcal{X}
- ▶ **Discrete Random Variable** - \mathcal{X} is finite/countably finite
 - ▶ Categorical??
- ▶ **Continuous Random Variable** - \mathcal{X} is infinite
- ▶ $P(X = x)$ or $P(x)$ is the probability of X taking value x
 - ▶ an **event**
- ▶ What is a distribution?

Examples

1. Coin toss ($\mathcal{X} = \{heads, tails\}$)
2. Six sided dice ($\mathcal{X} = \{1, 2, 3, 4, 5, 6\}$)

Notation, Notation, Notation

- ▶ X - random variable (**\mathbf{X}** if multivariate)
- ▶ x - a specific value taken by the random variable ((**\mathbf{x}** if multivariate))
- ▶ $P(X = x)$ or $P(x)$ is the probability of the event $X = x$
- ▶ $p(x)$ is either the **probability mass function** (discrete) or **probability density function** (continuous) for the random variable X at x
 - ▶ Probability mass (or density) at x

Basic Rules - Quick Review

- ▶ For two events A and B:
 - ▶ $P(A \vee B) = P(A) + P(B) - P(A \wedge B)$
 - ▶ **Joint Probability**
 - ▶ $P(A, B) = P(A \wedge B) = P(A|B)P(B)$
 - ▶ Also known as the *product rule*
 - ▶ **Conditional Probability**
 - ▶ $P(A|B) = \frac{P(A, B)}{P(B)}$

Chain Rule of Probability

- ▶ Given D random variables, $\{X_1, X_2, \dots, X_D\}$

$$P(X_1, X_2, \dots, X_D) = P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots P(X_D|X_1, X_2, \dots, X_D)$$

Marginal Distribution

- ▶ Given $P(A, B)$ what is $P(A)$?
 - ▶ Sum $P(A, B)$ over all values for B

$$P(A) = \sum_b P(A, B) = \sum_b P(A|B = b)P(B = b)$$

- ▶ **Sum rule**

Bayes Rule or Bayes Theorem

- ▶ Computing $P(X = x|Y = y)$:

Bayes Theorem

$$\begin{aligned}P(X = x|Y = y) &= \frac{P(X = x, Y = y)}{P(Y = y)} \\&= \frac{P(X = x)P(Y = y|X = x)}{\sum_{x'} P(X = x')P(Y = y|X = x')}\end{aligned}$$

Example

- ▶ Medical Diagnosis
- ▶ Random event 1: A *test* is positive or negative (X)
- ▶ Random event 2: A person has cancer (Y) – yes or no
- ▶ What we know:
 1. Test has accuracy of 80%
 2. Number of times the test is positive when the person has cancer

$$P(X = 1|Y = 1) = 0.8$$

3. Prior probability of having cancer is 0.4%

$$P(Y = 1) = 0.004$$

Question?

If I test positive, does it mean that I have 80% rate of cancer?

Base Rate Fallacy

- ▶ Ignored the prior information
- ▶ What we need is:

$$P(Y = 1|X = 1) = ?$$

- ▶ More information:
 - ▶ False positive (alarm) rate for the test
 - ▶ $P(X = 1|Y = 0) = 0.1$

$$P(Y = 1|X = 1) = \frac{P(X = 1|Y = 1)P(Y = 1)}{P(X = 1|Y = 1)P(Y = 1) + P(X = 1|Y = 0)P(Y = 0)}$$

Classification Using Bayes Rules

- ▶ Given input example \mathbf{X} , find the true class

$$P(Y = c|\mathbf{X})$$

- ▶ Y is the random variable denoting the true class
- ▶ Assuming the **class-conditional** probability is known

$$P(\mathbf{X}|Y = c)$$

- ▶ Applying Bayes Rule

$$P(Y = c|\mathbf{X}) = \frac{P(Y = c)P(\mathbf{X}|Y = c)}{\sum_c P(Y = c')P(\mathbf{X}|Y = c')}$$

- ▶ One random variable does not depend on another
- ▶ $A \perp B \iff P(A, B) = P(A)P(B)$
- ▶ Joint written as a product of marginals
- ▶ **Conditional Independence**

$$A \perp B|C \iff P(A, B|C) = P(A|C)P(B|C)$$

More About Conditional Independence

- ▶ Alice and Bob live in the same town but far away from each other
- ▶ Alice drives to work and Bob takes the bus
- ▶ Event A - Alice comes late to work
- ▶ Event B - Bob comes late to work
- ▶ Event C - A snow storm has hit the town
- ▶ $P(A|C)$ - Probability that Alice comes late to work given there is a snowstorm

More About Conditional Independence

- ▶ Alice and Bob live in the same town but far away from each other
- ▶ Alice drives to work and Bob takes the bus
- ▶ Event A - Alice comes late to work
- ▶ Event B - Bob comes late to work
- ▶ Event C - A snow storm has hit the town
- ▶ $P(A|C)$ - Probability that Alice comes late to work given there is a snowstorm
- ▶ Now if I know that Bob has also come late to work, will it change the probability that Alice comes late to work?

More About Conditional Independence

- ▶ Alice and Bob live in the same town but far away from each other
- ▶ Alice drives to work and Bob takes the bus
- ▶ Event A - Alice comes late to work
- ▶ Event B - Bob comes late to work
- ▶ Event C - A snow storm has hit the town
- ▶ $P(A|C)$ - Probability that Alice comes late to work given there is a snowstorm
- ▶ Now if I know that Bob has also come late to work, will it change the probability that Alice comes late to work?
- ▶ What if I do not observe C ? Will B have any impact on probability of A happening?

Continuous Random Variables

- ▶ X is continuous
- ▶ Can take any value
- ▶ How does one define probability?
 - ▶ $\sum_x P(X = x) = 1$

Continuous Random Variables

- ▶ X is continuous
- ▶ Can take any value
- ▶ How does one define probability?
 - ▶ $\sum_x P(X = x) = 1$
- ▶ Probability that X lies in an interval $[a, b]$?
 - ▶ $P(a < X \leq b) = P(x \leq b) - P(x \leq a)$
 - ▶ $F(q) = P(x \leq q)$ is the **cumulative distribution function**
 - ▶ $P(a < X \leq b) = F(b) - F(a)$

Probability Density Function

$$p(x) = \frac{\partial}{\partial x} F(x)$$

$$P(a < X \leq b) = \int_a^b p(x) dx$$

- Can $p(x)$ be greater than 1?

Expectation

- ▶ Expected value of a random variable

$$\mathbb{E}[X]$$

- ▶ What is most likely to happen in terms of X ?
- ▶ For discrete variables

$$\mathbb{E}[X] \triangleq \sum_{x \in \mathcal{X}} xP(X = x)$$

- ▶ For continuous variables

$$\mathbb{E}[X] \triangleq \int_{\mathcal{X}} xp(x)dx$$

- ▶ **Mean** of X (μ)

Expectation of Functions of Random Variable

- ▶ Let $g(X)$ be a function of X
- ▶ If X is discrete:

$$\mathbb{E}[g(X)] \triangleq \sum_{x \in \mathcal{X}} g(x)P(X = x)$$

- ▶ If X is continuous:

$$\mathbb{E}[g(X)] \triangleq \int_{\mathcal{X}} g(x)p(x)dx$$

Properties

- ▶ $\mathbb{E}[c] = c$, c - constant
- ▶ If $X \leq Y$, then $\mathbb{E}[X] \leq \mathbb{E}[Y]$
- ▶ $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- ▶ $\mathbb{E}[aX] = a\mathbb{E}[X]$
- ▶ $\text{var}[X] = \mathbb{E}[(X - \mu)^2] = \mathbb{E}[X^2] - \mu^2$
- ▶ $\text{Cov}[X, Y] = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$
- ▶ Jensen's inequality: If $\varphi(X)$ is convex,

$$\varphi(\mathbb{E}[X]) \leq \mathbb{E}[\varphi(X)]$$

What is a Probability Distribution?

Discrete

- ▶ Binomial, *Bernoulli*
- ▶ Multinomial, *Multinomial*
- ▶ Poisson
- ▶ Empirical

Continuous

- ▶ Gaussian (Normal)
- ▶ Degenerate pdf
- ▶ Laplace
- ▶ Gamma
- ▶ Beta
- ▶ Pareto

Binomial Distribution

- ▶ X = Number of heads observed in n coin tosses
- ▶ Parameters: n, θ
- ▶ $X \sim \text{Bin}(n, \theta)$
- ▶ Probability mass function (*pmf*)

$$\text{Bin}(k|n, \theta) \triangleq \binom{n}{k} \theta^k (1 - \theta)^{n-k}$$

Bernoulli Distribution

- ▶ Binomial distribution with $n = 1$
- ▶ Only one parameter (θ)

Multinomial Distribution

- ▶ Simulates a K sided die
- ▶ Random variable $\mathbf{x} = (x_1, x_2, \dots, x_K)$
- ▶ Parameters: n, θ
- ▶ $\theta \leftarrow \Re^K$
- ▶ θ_j - probability that j^{th} side shows up

$$Mu(\mathbf{x}|n, \theta) \triangleq \binom{n}{x_1, x_2, \dots, x_K} \prod_{j=1}^K \theta_j^{x_j}$$

Multinoulli Distribution

- ▶ Multinomial distribution with $n = 1$
- ▶ \mathbf{x} is a vector of 0s and 1s with only one bit set to 1
- ▶ Only one parameter (θ)

Gaussian (Normal) Distribution



$$\mathcal{N}(x|\mu, \sigma^2) \triangleq \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2\sigma^2}(x-\mu)^2}$$

► Parameters:

1. $\mu = \mathbb{E}[X]$
2. $\sigma^2 = \text{var}[X] = \mathbb{E}[(X - \mu)^2]$

- $X \sim \mathcal{N}(\mu, \sigma^2) \Leftrightarrow p(X = x) = \mathcal{N}(\mu, \sigma^2)$
- $X \sim \mathcal{N}(0, 1) \Leftrightarrow X$ is a **standard normal random variable**
- **Cumulative distribution function:**

$$\Phi(x; \mu, \sigma^2) \triangleq \int_{-\infty}^x \mathcal{N}(z|\mu, \sigma^2) dz$$



Joint Probability Distributions

- ▶ Multiple *related* random variables
- ▶ $p(x_1, x_2, \dots, x_D)$ for $D > 1$ variables (X_1, X_2, \dots, X_D)
- ▶ Discrete random variables?
- ▶ Continuous random variables?
- ▶ What do we measure?

Covariance

- ▶ How does X vary with respect to Y
- ▶ For linear relationship:

$$\text{cov}[X, Y] \triangleq \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])] = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$$

Covariance and Correlation

- ▶ \mathbf{x} is a d -dimensional random vector

$$\begin{aligned} \text{cov}[\mathbf{X}] &\triangleq \mathbb{E}[(\mathbf{X} - \mathbb{E}[\mathbf{X}])(\mathbf{X} - \mathbb{E}[\mathbf{X}])^\top] \\ &= \begin{pmatrix} \text{var}[X_1] & \text{cov}[X_1, X_2] & \cdots & \text{cov}[X_1, X_d] \\ \text{cov}[X_2, X_1] & \text{var}[X_2] & \cdots & \text{cov}[X_2, X_d] \\ \vdots & \vdots & \ddots & \vdots \\ \text{cov}[X_d, X_1] & \text{cov}[X_d, X_2] & \cdots & \text{var}[X_d] \end{pmatrix} \end{aligned}$$

- ▶ Covariances can be between 0 and ∞
- ▶ Normalized covariance \Rightarrow **Correlation**

- ▶ *Pearson Correlation Coefficient*

$$\text{corr}[X, Y] \triangleq \frac{\text{cov}[X, Y]}{\sqrt{\text{var}[X]\text{var}[Y]}}$$

- ▶ What is $\text{corr}[X, X]$?
- ▶ $-1 \leq \text{corr}[X, Y] \leq 1$
- ▶ When is $\text{corr}[X, Y] = 1$?

- ▶ *Pearson Correlation Coefficient*

$$\text{corr}[X, Y] \triangleq \frac{\text{cov}[X, Y]}{\sqrt{\text{var}[X]\text{var}[Y]}}$$

- ▶ What is $\text{corr}[X, X]$?
- ▶ $-1 \leq \text{corr}[X, Y] \leq 1$
- ▶ When is $\text{corr}[X, Y] = 1$?
 - ▶ $Y = aX + b$

Multivariate Gaussian Distribution

- ▶ Most widely used joint probability distribution

$$\mathcal{N}(\mathbf{X}|\mu, \Sigma) \triangleq \frac{1}{(2\pi)^{D/2}|\Sigma|^{D/2}} \exp \left[-\frac{1}{2}(\mathbf{x} - \mu)^\top \Sigma^{-1}(\mathbf{x} - \mu) \right]$$

Linear Transformations of Random Variables

- ▶ What is the distribution of $f(\mathbf{X})$ ($\mathbf{X} \sim p()$)?
 - ▶ Linear transformation:

$$Y = \mathbf{a}^\top \mathbf{X} + b$$

- ▶ $\mathbb{E}[Y]$?
- ▶ $\text{var}[Y]$?

$$\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$$

- ▶ $\mathbb{E}[\mathbf{Y}]$?
- ▶ $\text{cov}[\mathbf{Y}]$?

- ▶ The Matrix Cookbook [2]
- ▶ <http://orion.uwaterloo.ca/~hwolkowi/matrixcookbook.pdf>
- ▶ Available on Piazza

General Transformations

- ▶ $f()$ is **not linear**
- ▶ Example: X - discrete

$$Y = f(X) = \begin{cases} 1 & \text{if } X \text{ is even} \\ 0 & \text{if } X \text{ is odd} \end{cases}$$

General Transformations for Continuous Variables

- ▶ For continuous variables, work with cdf

$$F_Y(y) \triangleq P(Y \leq y) = P(f(X) \leq y) = P(X \leq f^{-1}(y)) = F_X(f^{-1}(y))$$

- ▶ For pdf

$$p_Y(y) \triangleq \frac{d}{dy} F_Y(y) = \frac{d}{dy} F_X(f^{-1}(y)) = \frac{dx}{dy} \frac{d}{dx} F_X(x) = \frac{dx}{dy} p_X(x)$$

- ▶ $x = f^{-1}(y)$

Example

- ▶ Let X be $Uniform(-1, 1)$
- ▶ Let $Y = X^2$
- ▶ $p_Y(y) = \frac{1}{2}y^{-\frac{1}{2}}$

Monte Carlo Approximation

- ▶ Generate N samples from distribution for X
- ▶ For each sample, $x_i, i \in [1, N]$, compute $f(x_i)$
- ▶ Use empirical distribution as *approximate* true distribution

Approximate Expectation

$$\mathbb{E}[f(X)] = \int f(x)p(x)dx \approx \frac{1}{N} \sum_{i=1}^N f(x_i)$$

Introduction to Information Theory


- ▶ Quantifying uncertainty of a random variable

Entropy

- ▶ $\mathbb{H}(X)$ or $\mathbb{H}(p)$

$$\mathbb{H}(X) \triangleq - \sum_{k=1}^K p(X = k) \log_2 p(X = k)$$

- ▶ Variable with maximum entropy?
- ▶ Lowest entropy?



Comparing Two Distributions

- ▶ **Kullback-Leibler Divergence** (or *KL Divergence* or *relative entropy*)

$$\begin{aligned}\mathbb{KL}(p||q) &\triangleq \sum_{k=1}^K p(k) \log \frac{p_k}{q_k} \\ &= \sum_k p(k) \log p(k) - \sum_k p(k) \log q(k) \\ &= -\mathbb{H}(p) + \mathbb{H}(p, q)\end{aligned}$$

- ▶ $\mathbb{H}(p, q)$ is the *cross-entropy*
- ▶ Is KL-divergence symmetric?
- ▶ *Important fact:* $\mathbb{H}(p, q) \geq \mathbb{H}(p)$

Mutual Information

- ▶ What does learning about one variable X tell us about another, Y ?
 - ▶ Correlation?

Mutual Information

$$\mathbb{I}(X; Y) = \sum_x \sum_y p(x, y) \log \frac{p(x, y)}{p(x)p(y)}$$

- ▶ $\mathbb{I}(X; Y) = \mathbb{I}(Y; X)$
- ▶ $\mathbb{I}(X; Y) \geq 0$, equality iff $X \perp Y$

References



E. Jaynes and G. Bretthorst.

Probability Theory: The Logic of Science.

Cambridge University Press Cambridge:, 2003.



K. B. Petersen and M. S. Pedersen.

The matrix cookbook, nov 2012.

Version 20121115.



L. Wasserman.

All of Statistics: A Concise Course in Statistical Inference (Springer Texts in Statistics).

Springer, Oct. 2004.